

This report is based on research conducted by the Vanderbilt University Evidence-based Practice Center (EPC) under contract to the Agency for Healthcare Research and Quality (AHRQ), Rockville, MD (Contract No. HHS 290-2007-10065-I). The findings and conclusions in this document are those of the authors, who are responsible for its content, and do not necessarily represent the views of AHRQ. No statement in this report should be construed as an official position of AHRQ or of the U.S. Department of Health and Human Services.

The information in this report is intended to help clinicians, employers, policymakers, and others make informed decisions about the provision of health care services. This report is intended as a reference and not as a substitute for clinical judgment.

This report may be used, in whole or in part, as the basis for the development of clinical practice guidelines and other quality enhancement tools, or as a basis for reimbursement and coverage policies. AHRQ or U.S. Department of Health and Human Services endorsement of such derivative products may not be stated or implied.

Treatment of Overactive Bladder in Women

Prepared for:

Agency for Healthcare Research and Quality
U.S. Department of Health and Human Services
540 Gaither Road
Rockville, MD 20850
www.ahrq.gov

Contract No. 290-2007-10065-I

Prepared by:

Vanderbilt Evidence-based Practice Center
Institute for Medicine and Public Health
Vanderbilt University Medical Center
2525 West End Avenue
Nashville, Tennessee 37203-1738

Investigators

Katherine E. Hartmann, M.D., Ph.D.
Melissa L. McPheeters, Ph.D., M.P.H.
Daniel H. Biller, M.D.
Renée M. Ward, M.D.
J. Nikki McKoy, B.A.
Rebecca N. Jerome, M.L.I.S., M.P.H.
Sandra R. Micucci, M.Sc.
Laura Meints, M.D., M.B.A.
Jill A. Fisher, Ph.D.
Theresa A. Scott, M.S.
James C. Slaughter, Dr.P.H.
Jeffrey D. Blume, Ph.D.

AHRQ Publication No. 09-E017
August 2009

This document is in the public domain and may be used and reprinted without permission except those copyrighted materials noted for which further reproduction is prohibited without the specific permission of copyright holders.

Suggested Citation:

Hartmann KE, McPheeters ML, Biller DH, Ward RM, McKoy JN, Jerome RN, Micucci SR, Meints L, Fisher JA, Scott T, Slaughter JC, Blume JD. Treatment of Overactive Bladder in Women. Evidence Report/Technology Assessment No. 187 (Prepared by the Vanderbilt Evidence-based Practice Center under Contract No. 290-2007-10065-I.) AHRQ Publication No. 09-E017. Rockville, MD: Agency for Healthcare Research and Quality. August. 2009

No investigators have any affiliations or financial involvement (e.g., employment, consultancies, honoraria, stock options, expert testimony, grants or patents received or pending, or royalties) that conflict with material presented in this report.

Preface

The Agency for Healthcare Research and Quality (AHRQ), through its Evidence-Based Practice Centers (EPCs), sponsors the development of evidence reports and technology assessments to assist public- and private-sector organizations in their efforts to improve the quality of health care in the United States. This report was requested by the American Urological Association in preparation for clinical guideline development. These reports and assessments provide organizations with comprehensive, science-based information on common, costly medical conditions and new health care technologies. The EPCs systematically review the relevant scientific literature on topics assigned to them by AHRQ and conduct additional analyses when appropriate prior to developing their reports and assessments.

To bring the broadest range of experts into the development of evidence reports and health technology assessments, AHRQ encourages the EPCs to form partnerships and enter into collaborations with other medical and research organizations. The EPCs work with these partner organizations to ensure that the evidence reports and technology assessments they produce will become building blocks for health care quality improvement projects throughout the Nation. The reports undergo peer review prior to their release.

AHRQ expects that the EPC evidence reports and technology assessments will inform individual health plans, providers, and purchasers as well as the health care system as a whole by providing important information to help improve health care quality.

We welcome comments on this evidence report. They may be sent by mail to the Task Order Officer named below at: Agency for Healthcare Research and Quality, 540 Gaither Road, Rockville, MD 20850, or by e-mail to epc@ahrq.gov.

Carolyn M. Clancy, M.D.
Director
Agency for Healthcare Research and Quality

Jean Slutsky, P.A., M.S.P.H.
Director, Center for Outcomes and Evidence
Agency for Healthcare Research and Quality

Beth A. Collins Sharp, Ph.D.
Acting Director, EPC Program
Agency for Healthcare Research and Quality

Shilpa H. Amin, M.D, M.Bsc, FAAFP
EPC Program Task Order Officer
Agency for Healthcare Research and Quality

Acknowledgments

We are indebted to a tireless group of colleagues who made this report possible. Each step of systematic reviews draws on the skills and attention of an entire team.

Dr. Shilpa Amin has been an insightful task order officer who brought special considerations and decisions into focus. She was patient with our many questions, created a strong link with our project partner, and provided practical advice as we tackled challenges.

Ms. Rachel Bazan was our energetic student worker. She spent hours helping to track and file documents, always positive, and always ready to ensure that the project investigators had what they needed to do their work.

Dr. Liana Castel served as a quality grader and brought to the project her expertise in study design and analytic methods. We thank her for her thoughtful attention to the process of quality grading and willingness to help. Her comments on approaches to study reporting, particularly as they related to the implications of loss to followup and drop out, were especially valued.

Ms. Allison Glasser provided research assistance on this report, including combing papers for specific data points requested by the investigators and formatting summary and evidence tables. We appreciate her eagerness to learn and employ systematic review methods and her commitment to ensuring that tables were formatted exactly to specification. Ms. Glasser demonstrated great flexibility under pressure filling many needs.

Dr. Mark Hartmann brought his extraordinary attention to detail – and his commitment to perfection – to completion of the evidence tables. He spent many, many hours checking and rechecking tables both for formatting and for content. His ability to point out inconsistencies and enhance uniformity was key to ensuring smooth development of the evidence tables.

Ms. Sarah McLellan was a source of energy for all as she enthusiastically went about formatting evidence tables and completing and organizing appendices. We are grateful to lean on such detail oriented and focused research staff with a gift for encouraging others.

Ms. Toyce Spencer oversaw the completion and formatting of the summary tables, and other critical elements of the report as a whole. She transferred data from investigator-developed spreadsheets into more tables than we imagined – and made sure that formatting was consistent, keeping track of the innumerable sets of changes that came her way.

Ms. Rachel Walden is a key member of the library science staff – her support of Ms. Jerome, including her creative insights and detailed approach to literature searching was invaluable. Her thoughtful attention to detail in reviewing abstracts and articles for inclusion in the review was a generous gift to the project and to the EPC as a whole.

Structured Abstract

Objectives: The Vanderbilt Evidence-based Practice Center systematically reviewed evidence on treatment of overactive bladder (OAB), urge urinary incontinence, and related symptoms. We focused on prevalence and incidence, treatment outcomes, comparisons of treatments, modifiers of outcomes, and costs.

Data: We searched PubMed, MEDLINE®, EMBASE, and CINAHL.

Review Methods: We included studies published in English from January 1966 to October 2008. We excluded studies with fewer than 50 participants, fewer than 75 percent women, or lack of relevance to OAB. Of 232 included publications, 20 were good quality, 145 were fair, and 67 poor. We calculated weighted averages of outcome effects and conducted a mixed-effects meta-analysis to investigate outcomes of pharmacologic treatments across studies.

Results: OAB affects more than 10 to 15 percent of adult women, with 5 to 10 percent experiencing urge urinary incontinence (UUI) monthly or more often. Six available medications are effective in short term studies: estimates from meta-analysis models suggest extended release forms (taken once a day) reduce UUI by 1.78 (95 percent confidence interval (CI): 1.61, 1.94) episodes per day, and voids by 2.24 (95 percent CI: 2.03, 2.46) per day. Immediate release forms (taken twice or more a day) reduce UUI by 1.46 (95 percent CI: 1.28, 1.64), and voids by 2.17 (95 percent CI: 1.81, 2.54). As context, placebo reduces UUI episodes by 1.08 (95 percent CI: 0.86, 1.30), and voids by 1.48 (95 percent CI: 1.19, 1.71) per day. No one drug was definitively superior to others, including comparison of newer more selective agents to older antimuscarinics.

Current evidence is insufficient to guide choice of other therapies including sacral neuromodulation, instillation of oxybutynin, and injections of botulinum toxin. Acupuncture was the sole complementary and alternative medicine treatment, among reflexology and hypnosis, with early evidence of benefit. The strength of the evidence is insufficient to fully inform choice of these treatments. Select behavioral interventions were associated with symptom improvements comparable to medications. Limited evidence suggests no clear benefit from adding behavioral interventions at the time of initiation of pharmacologic treatment.

Conclusions: OAB and associated symptoms are common. Treatment effects are modest. Quality of life and treatment satisfaction measures suggest such improvements can be important to women. The amount of high quality literature available is meager for helping guide women's choices. Gaps include weak or absent data about long-term followup, poorly characterized and potentially concerning harms, information about best choices to minimize side effects, and study of how combinations of approaches may best be used. This is problematic since the condition is chronic and a single treatment modality is unlikely to fully resolve symptoms for most women.

Contents

- Executive Summary 1
- Evidence Report 13**
- Chapter 1. Introduction 15
 - Importance of Overactive Bladder Treatment..... 15
 - Defining OAB..... 15
 - OAB and Awareness of the General Public..... 16
 - Treatment Options..... 17
 - This Evidence Report 17
 - Scope of the Report..... 17
 - Key Questions..... 17
 - Analytic Framework 18
 - Organization of this Evidence Report..... 19
 - Technical Expert Panel (TEP) 19
 - Uses of this Report 20
- Chapter 2. Methods.....21
 - Literature Review Methods 21
 - Inclusion and Exclusion Criteria..... 21
 - Literature Search and Retrieval Process 22
 - Literature Synthesis..... 25
 - Development of Evidence Tables and Data Abstraction Process 25
 - Synthesis of the Evidence 25
 - Quality Rating of Individual Studies 27
 - Strength of Available Evidence..... 30
 - External Peer Review 31
- Chapter 3. Results33
 - KQ1. Prevalence and Incidence of Overactive Bladder..... 33
 - Prevalence of Overactive Bladder 33
 - Prevalence of Urge Urinary Incontinence..... 37
 - Other Measures 38
 - Incidence of Overactive Bladder and Urge Urinary Incontinence..... 38
 - KQ2. Outcomes of Treatment of Overactive Bladder..... 39
 - Pharmacologic Treatments..... 39
 - Harms of Pharmacologic Treatments..... 61
 - Procedural and Surgical Treatments 64
 - Behavioral Treatments 72
 - Complementary and Alternative Medicine Treatments 78

KQ3: Comparisons of Treatments.....	79
Comparisons between pharmacologic treatments.....	79
Comparisons between procedural and pharmaceutical treatments.....	82
Comparisons between behavioral and pharmacologic treatments.....	82
Comparisons of combined behavioral and pharmacologic treatment to pharmacologic treatment alone.....	85
Comparisons of combined behavioral and pharmacologic treatment to behavioral treatment alone.....	87
KQ4: Modifiers of Treatment Outcomes.....	88
Identified modifiers.....	88
KQ5: Costs of Overactive Bladder Treatment.....	92
Chapter 4. Discussion.....	97
Strength of Evidence.....	97
Principal Findings and Considerations.....	97
KQ1: Incidence and Prevalence of Overactive Bladder.....	97
KQ2: Outcomes of Treatments for Overactive Bladder.....	99
KQ3: Comparisons of Treatments.....	102
KQ4: Modifiers of Treatment Outcomes.....	103
KQ5: Costs of Overactive Bladder Treatment.....	104
Future Research.....	105
State of the Literature.....	105
Future Research Directions.....	107
Conclusions.....	108
References and Included Studies.....	109
List of Acronyms/Abbreviations.....	118

Tables

Table 1. Executive Summary. Estimates of mean reductions in incontinent episodes and voids per day.....	5
Table 1. Inclusion and Exclusion Criteria.....	21
Table 2. PubMed search strategies (last updated October 1, 2008).....	22
Table 3. EMBASE search (OVID) (last updated October 1, 2008).....	23
Table 4. CINAHL search (EBSCO) (last updated October 1, 2008).....	24
Table 5. Scoring algorithm for quality rating of individual studies.....	30
Table 6. Study characteristics of the prevalence and incidence literature.....	34
Table 7. Prevalence of OAB.....	36
Table 8. RCT arms for oxybutynin chloride effect on urge incontinence.....	41
Table 9. RCT arms for oxybutynin chloride effect on voids per day.....	42
Table 10. RCT arms for tolterodine tartrate effect on urge incontinence.....	44
Table 11. RCT arms for tolterodine tartrate effect on voids per day.....	46
Table 12. RCT arms for fesoterodine fumarate effect on urge incontinence and voids.....	50
Table 13. RCT arms for solifenacin succinate effect on urge incontinence and voids.....	51

Table 14. RCT arms for darifenacin effect on urge incontinence.....	54
Table 15. RCT arms for darifenacin effect on voids per day.....	54
Table 16. RCT arms for trospium chloride effect on urge incontinence	56
Table 17. RCT arms for trospium chloride effect on voids per day	56
Table 18. Effect on quality of life and satisfaction for pharmacologic treatment	58
Table 19. Side effects and harms of pharmacologic treatment.....	63
Table 20. Outcomes of clinical trials of procedures	66
Table 21. Effect on quality of life and satisfaction of procedural treatments.....	69
Table 22. Outcomes of behavioral treatment trials.....	74
Table 23. Effects on quality of life and satisfaction of behavioral treatments	77
Table 24. Direct comparisons of pharmaceutical treatments on urge incontinence	81
Table 25. Direct comparisons of pharmaceutical treatments on voids per day	81
Table 26. Direct comparisons between pharmacologic and behavioral interventions.....	82
Table 27. Comparisons of pharmaceuticals with and without behavioral interventions	85
Table 28. Effect on quality of life and satisfaction of combination treatment	87
Table 29. Study definitions used for cost determination	93
Table 30. Total cost differences in annual medical care among persons filling prescriptions for OAB drug treatment	94
Table 31. Estimates of mean reductions in incontinent episodes and voids per day	100
Table 32. Funding sources and conflict of interest by decade of publication	107

Figures

Figure 1. Analytic framework for the treatment of OAB in women	17
Figure 2. Disposition of articles for the treatment of OAB	24

Appendixes

Appendix A: Exact Search Strings
Appendix B: Sample Data Abstraction Forms
Appendix C: Evidence Tables
Appendix D: List of Excluded Studies
Appendix E: Peer Reviewers

Appendixes and Evidence Tables for this report are provided electronically at <http://www.ahrq.gov/downloads/pub/evidence/pdf/bladder/bladder.pdf>.

Executive Summary

Introduction

Importance of Overactive Bladder Treatment

At minimum, 11 to 16 million women in the United States cope on a daily basis with symptoms that include sudden strong urges to urinate, difficulty delaying voids, frequent trips to the bathroom, and in many cases involuntary loss of urine when urgency strikes.¹⁻⁴ They may wear pads for accidents, plan ahead for access to bathrooms, and modify their social and work lives to accommodate their symptoms. Some are very distressed by the symptoms whether mild or severe, and others find mechanisms to adapt, reporting little trouble with symptoms or interference with normal routines. Others report their symptoms negatively influence quality of life factors as varied as self-esteem, self-assessment of attractiveness, and sexual function. Many women believe that some amount of urinary incontinence is inevitable with aging. The majority of women with these symptoms do not talk with their health care providers concerning their bladder dysfunction, and providers may not systematically inquire. As a result, a small minority receive treatment.

Popular wisdom encourages self-management of symptoms of OAB through reduction of fluid intake, cutting back on caffeine, modifying voiding habits, and taking note of what individual factors influence severity of symptoms.

For this review, we operationally defined OAB as “idiopathic urinary urgency and frequency with or without associated urge urinary incontinence in adult females, not related to neurogenic conditions or as a result of (stress incontinence) surgery.” The report is focused on treatments that are prescribed or provided by a healthcare practitioner and have been formally investigated including:

- Pharmacologic treatments, including prescription medications, both pills and patches
- Surgeries and procedures, such as sacral neuromodulation and botulinum injections
- Behavioral interventions, such as behavior modification programs and bladder training
- Complementary and alternative medicine, such as acupuncture and reflexology

OAB management is usually individualized to address the component symptom(s) that the patient finds most bothersome. Where possible, we have tried to address treatments with respect to the primary component symptoms of OAB: urge urinary incontinence, urgency, and frequency, so that the women, their health care providers, payors, policy-makers, and others have a detailed picture of the expected outcomes of available treatments.

Key Questions

In preparing this report, we have answered the following key questions:

KQ1. What is the prevalence and incidence of overactive bladder as estimated in representative populations?

KQ2. Among women with overactive bladder, what are the short and long-term outcomes of the following treatment approaches, or combinations of treatment approaches?

- a. Pharmacologic treatments
 - b. Procedural and surgical treatments
 - c. Behavioral and physical therapy treatments
 - d. Complementary and alternative medicine treatments
- KQ3.** Where direct comparisons have been made between or among treatment modalities of interest, which modalities achieve superior outcomes with respect to benefits, short and long-term risks, and quality of life?
- KQ4.** Are the short and long term outcomes of these treatment approaches modified by clinical presentation, physical exam findings, urodynamic findings, menopausal status, age or other factors?
- KQ5.** What are the costs associated with these treatment approaches?

Methods

Literature search. We employed multi-term search strategies to retrieve research about treatment of overactive bladder in women, including exploration of three databases: PubMed, MEDLINE®, EMBASE, and the Cumulative Index to Nursing and Allied Health Literature (CINAHL). We also hand-searched the reference lists of relevant articles to identify additional studies. Controlled vocabulary terms served as the foundation of our searches in each database, complemented by additional keyword phrases to represent the myriad ways in which overactive bladder is referred to in the clinical literature. We also employed indexing terms within each of the databases to exclude undesired publication types (e.g., reviews, case reports, CME handouts) and items published in languages other than English. We excluded studies that (1) were published in languages other than English; (2) did not report information pertinent to the key questions; (3) had fewer than 50 participants [at enrollment]; (4) were not original studies; and (5) provided data only for stress or mixed urinary incontinence.

Study selection. Two reviewers separately evaluated abstracts for inclusion or exclusion. If one reviewer concluded the abstract should be included for full review of the article, it was retained. For the full article review, two reviewers read each article and decided whether it met our inclusion criteria. Discordance was resolved by third-party adjudication.

Quality assessment. The research team used a quality assessment approach that ensured capture of key points most relevant to this literature. Quality was assessed by two reviewers independently, who resolved differences through discussion, review of the publications and coming to consensus with the team.

Data extraction. All team members shared the task of entering information into the evidence tables. After initial data extraction, another member of the team reviewed the article and checked all table entries for accuracy, completeness, and consistency. The two abstractors reconciled disagreements concerning the information reported in the evidence tables.

Evidence synthesis. The information included in tables reflects those outcomes most consistently reported in the literature: urge incontinence episodes and number of voids per day. (Studies with weekly or other metrics that could be converted to daily metric are included.) When common measures were available across studies using roughly comparable assessments (i.e., index questions, time intervals, etc.), we compiled tables to summarize outcomes of treatments. Measures of quality of life, interference with daily activities, degree of distress from symptoms, and satisfaction with the outcomes of treatment were also common and helpful metrics in this literature.

For behavioral, surgical, and complementary and alternative treatments, we produced evidence tables, summary tables of common outcomes where possible, and provided analysis in text form. For pharmacologic treatment, we produced similar summary tables and conducted a limited meta-analysis.

Conduct of meta-analysis. Descriptive statistics were computed and examined for homogeneity among studies. Studies that reported weekly rates for UUI (urinary urge incontinence) episodes and voids were standardized to daily rates. When only ranges of continuous variables were reported (instead of standard deviations), we estimated the standard deviations by dividing the range by four.⁵ Study results were combined and summarized using two meta-analysis techniques, weighted averages and fixed effects regression models.⁶ Minimum variance weighted averages of the mean daily decrease in UUI and voids per arm were computed using weights that were inversely proportional to their standard errors. To borrow strength across arms, we used fixed effect regression models with robust standard errors (to account for the clustering by study), and weighted the study arms inversely proportional to their standard errors of the mean. Each arm was treated as a fixed effect, and study was not included in the model except in the sense that the clustering was addressed by the robust standard errors. Fixed effects models were also adjusted for mean age and proportion of women in each arm.

Literature search yield. As a result of the search, 2,559 non-duplicate articles were identified. Two hundred thirty-two articles were included in the review, representing one hundred seventy-nine distinct studies, with 75 articles pertaining to KQ1, 150 to KQ2, 34 to KQ3, 32 to KQ4, and 5 to KQ5. Reasons and process for exclusions are described in the full report.

Conflict of interest. We used a two-step process to describe the contribution of industry-sponsored research to the evidence base. The first step was to calculate the proportion of publications that explicitly state the source of the funding for research, and the second step was to calculate the proportion supported by industry *among those that report funding source*. These counts are important because several studies have shown that industry-sponsored research tends to produce results that are favorable to the drug manufacturer even when the research is conducted in academic medical centers.⁷⁻⁹

Results

KQ1. Prevalence and incidence of overactive bladder

Content of the literature. We identified 75 publications,^{2-4, 10-81} from 60 distinct study populations. Fifteen studies were conducted in United States populations; 24 in European populations; 13 Asian; and 8 in other countries. Thirty-eight of these studies (63 percent) appeared in print after the 2002 International Continence Society (ICS) revised definitions and 37 percent of these reported specifically about incidence or prevalence of OAB. One study prior to the standardized definitions used fully comparable definitions and the term “overactive bladder”. The strength of the evidence for understanding prevalence of OAB is moderate and for understanding incidence is weak.

Prevalence estimates. A total of 15 studies provided information specifically about OAB prevalence in adult women. These studies included a total of 64,528 women in 16 distinct populations. Estimates of prevalence ranged from 7.7 to 31.3 percent. Across all studies the

average prevalence of OAB was 16.1 percent. Excluding the highest and lowest estimates, an estimated 15.1 percent of women meet criteria for OAB, with 8.2 percent of those surveyed, having OAB with a component of urge incontinence. Combined estimates from the two populations from the United States are similar 15.1 percent with OAB and 11.0 percent with a component of urge incontinence.

A larger number of studies (n=48) examined urge urinary incontinence as the primary prevalence estimate of interest. Twenty-eight appeared after the ICS 2002 standardization of definitions. Across populations, prevalence of urge incontinence ranged from 1.5 to 22.0 percent. Average urge incontinence prevalence was 9.7 percent in the United States, 10.6 percent in Europe, and 9.6 percent in Asia.

Incidence and resolution. Ten studies provided incidence data and two reported on resolution of symptoms. Estimates for annual incidence of OAB ranged from 2.6 to 143 cases per thousand, with higher estimates in the oldest population groups. A proportion of cases, 23 percent to 35 percent, resolve over a year's time; however the majority of women have symptoms for years. No studies evaluated lifetime natural history.

KQ2: Outcomes of Treatment for Overactive Bladder

Pharmacologic treatment.

Content of the literature. We identified 13 randomized controlled trials (RCTs) of oxybutynin for treatment of OAB. A total of 22 study arms compared oxybutynin at varied doses, formulations, and intervals. These studies included five placebo arms. Most participants were recruited from specialty populations with seven studies performed in the United States,⁸²⁻⁸⁸ three in Europe,⁸⁹⁻⁹¹ and one each in Japan,⁹² Taiwan,⁹³ and South Korea.⁹⁴ These studies included a total of 2,575 women in treatment arms, and 383 women in the placebo arms.

For tolterodine there were 19 RCTs, including 29 drug arms and 13 placebo arms. Most were multinational studies conducted at centers in Europe, the United States, Australia, and Asia. A total of 6,564 women were in the treatment arms, with 3,109 women in the placebo arms.

Two RCTs compared fesoterodine at 4 and 8 mg to placebo for reducing symptoms of OAB and met criteria for inclusion in the systematic review.^{95,96} These studies included a total of 1,017 women in the treatment arms, and a total of 518 women in the placebo arms.

Three RCTs investigated solifenacin compared to placebo for reducing symptoms of OAB.⁹⁷⁻⁹⁹ These studies included a total of 1,541 women in the solifenacin treatment arms, and a total of 638 women in the placebo arms.

Four RCTs provided data on the effectiveness of darifenacin.^{82,100-102} These four studies included a total of 690 women in the darifenacin treatment arms and a total of 304 women in the placebo arms.

Five RCTs evaluated trospium for reduction of symptoms of OAB. Four trials compared trospium to placebo, and one compared trospium to oxybutynin. Four were conducted in the United States,¹⁰³⁻¹⁰⁶ and the fifth at multiple centers in Europe and Asia.⁹⁰ These studies included a total of 1,309 women in the trospium treatment arms, and a total of 1,130 women in the placebo arms.¹⁰³⁻¹⁰⁶

Three studies assessed the role of oral estrogen therapy in different doses and formulations in the alleviation of OAB symptoms.¹⁰⁷⁻¹⁰⁹ All studies were performed in Europe and Scandinavia with a total of 514 women. Two were RCTs and one was a prospective cohort. After review and analysis of all 110 studies, of which four were good quality, 75 fair and 31 poor, including 68

RCTs, the strength of the evidence for managing OAB with pharmacologic treatment is weak to moderate for short term outcomes and weak for long term outcomes and harms.

Outcomes of treatment. All pharmacologic treatments were effective at improving one or more OAB symptoms when compared to placebo. Reductions ranged from 0.9 to 4.6 in incontinence episodes per day across all drug treatments and from 0.7 to 4.2 in voids per day. Study by study, extended release formulations achieved modestly better effects than immediate release, statistical significance varied. No one drug was definitively superior to others by preponderance of evidence, including more recently approved drugs. As estimated by meta-analysis, extended release forms (taken once a day) reduce UUI by 1.78 (95 percent CI: 1.61, 1.94) episodes per day, and voids by 2.24 (95 percent CI: 2.03, 2.46) per day. Immediate release forms (taken twice or more a day) reduce UUI episodes by 1.46 (95 percent CI: 1.28, 1.64) per day, and voids by 2.17 (95 percent CI: 1.81, 2.54) per day. Of note, placebo reduces UUI episodes by 1.08 (95 percent CI: 0.86, 1.30), and voids by 1.48 (95 percent CI: 1.19, 1.71) per day. Even in the context of small to moderate affect on symptoms, pharmacologic treatments were generally associated with increased quality of life and reductions in measures of impact or distress, compared to baseline and to placebo.

Table 1 below provides estimates of treatment effects for pharmacologic treatments represented by more than one trial arm. Some drugs and doses of drugs are not reported because the publications with trial arms for that treatment did not provide sufficient information to estimate variance in meta-analysis models.

Table 1. Estimates of mean reductions in incontinent episodes and voids per day

Drug	Decrease in Incontinent Episodes per Day			Decrease in Voids per Day		
	Estimate	95% CI		Estimate	95% CI	
Single drug estimates						
Placebo	1.08	0.86	1.30	1.48	1.19	1.71
Oxybutynin IR	1.49	1.18	1.80	2.18	1.75	2.61
Oxybutynin ER	*	*	*	*	*	*
Tolterodine IR	1.45	1.24	1.66	2.19	1.76	2.61
Tolterodine ER	1.75	1.65	1.85	2.48	1.94	3.02
Fesoterodine	2.03	1.74	2.31	1.84	1.64	2.03
Darifenacin	*	*	*	*	*	*
Solifenacin	1.46	1.32	1.59	2.19	1.94	3.02
Trospium IR	*	*	*	*	*	*
Trospium ER	2.45	2.19	2.70	2.68	2.38	2.98
Combined comparison of extended versus immediate release formulations						
Placebo	1.08	0.86	1.30	1.48	1.19	1.71
Extended Release	1.78	1.61	1.94	2.24	2.03	2.46
Immediate Release	1.46	1.28	1.64	2.17	1.81	2.54

*Estimates could not be calculated for these formulations because publications did not include adequate data on variance for weighting of the raw data.

Since baseline episodes of UUI per day ranged from 1.6 to 5.3, and voids per day from 7.2 to 13.7, these reductions (Table 1) reflect modest margins of benefit from baseline above placebo. Data was not consistently provided across studies to estimate the proportion of women who became symptom free.

Procedural and surgical treatment.

Content of the literature. We identified 18 studies about surgical treatments and procedures for OAB. Eleven were of sacral neuromodulation, one of peripheral neuromodulation, and one of electromagnetic therapy. Three studied bladder instillation or injection of drugs; one was on bladder distention; and one about bladder transection. This literature included 13 case series studies: nine prospective¹¹⁰⁻¹¹⁸ and three retrospective.¹¹⁹⁻¹²¹ One study had both retrospective and prospective components.¹²² Given consideration of 18 studies, of which 11 were fair quality and seven poor, including five RCTs, the strength of the evidence for managing OAB with procedural and surgical treatment is weak for all aspects of understanding outcomes of care.

One study was a prospective cohort that compared outcomes among participants receiving sacral neuromodulation to participants who had lead placement without activation of electrical stimulation.¹²³ Four studies were RCTs: one looked at sacral neuromodulation versus medical therapy,¹²⁴ one evaluated transcutaneous electromagnetic stimulation versus sham,¹²⁵ and two evaluated instillation of a drug into the bladder versus placebo – one using oxybutynin¹²⁶ and one using resiniferatoxin.¹²⁷

Outcomes of treatment. Among the trials of procedures and surgery, one study demonstrated a statistically significant benefit of sacral neuromodulation over usual care for the reduction of episodes of incontinence per day (average reduction of 7.1 compared to 2.1 increase among subjects who had failed medical management).¹²⁴ One trial demonstrated benefit of instillation of oxybutynin compared to sterile water in the reduction of voids per day (average reduction of 6.8 compared to 2.4)¹²⁶ and another reported benefits from botulinum toxin treatment which is compatible with the findings of a recent Cochrane Collaboration literature review and meta-analysis.¹²⁸

Behavioral interventions.

Content of the literature. We identified nine studies that included only behavioral approaches; no two studies compared the same set of approaches. They included assessment of bladder training, multicomponent behavioral training, with or without biofeedback, pelvic muscle exercises or training, vaginal electrical stimulation, and caffeine reduction. The literature base included three retrospective case series.¹²⁹⁻¹³¹ All three were conducted in community-based clinical settings.

One prospective cohort study compared three approaches to providing bladder training: self-administered, coaching, and cognitive strategies.¹³² Five studies were RCTs. One compared bladder training to a “control” condition.¹³³ One compared bladder training to pelvic muscle exercises.¹³⁴ One included three arms: pelvic floor muscle training, pelvic floor muscle training assisted with biofeedback, and vaginal electrical stimulation.⁹³ Another compared three different approaches to multicomponent behavioral training: biofeedback, verbal feedback and self-administered.¹³⁵ A last study compared bladder training to bladder training with caffeine reduction.¹³⁶ After review and analysis of 29 studies, of which 14 were fair quality and 15 poor, including 17 RCTs, the strength of the evidence for managing OAB with behavioral approaches treatment is moderate to weak for short term outcomes and weak for long term outcomes and harms.

Outcomes of treatment. No two studies could be combined to produce summary data. Overall, behavioral approaches can be effective in reducing episodes of incontinence and daily voids. Multicomponent approaches are most effective, and they perform relatively equivalently to pharmacologic treatment. Generally speaking, improvements were modest, with decreases in

incontinence episodes of up to 1.9 per day, and reductions in voids per day of up to about four. The addition of caffeine reduction reduced frequency, but made no difference in reduction of incontinence episodes. There is no evidence that behavioral approaches enhance the effectiveness of pharmacologic therapy for reducing episodes of incontinence or voiding, although they may improve patient satisfaction and quality of life measures.

Complementary and alternative therapies.

Content of the literature. We identified three studies that used complementary and alternative medicine therapies to treat OAB: a fair quality trial of acupuncture,¹³⁷ a fair quality trial of foot reflexology,¹³⁸ and a poor quality prospective case series of hypnotherapy.¹³⁹ There is weak to no evidence for complementary and alternative approaches to managing OAB.

Outcomes of treatment. The small trial of acupuncture has intriguing results related to decreased frequency of voiding and reduced symptoms of urgency which are associated with changes in cystometrics related to improved bladder capacity that are logical intermediates of the improvement in symptoms. Women felt they were improved as measured by scales that capture bother and quality of life. Evidence is insufficient to support definitive choice of acupuncture but offers preliminary information that promises modest improvements similar to those reported in many pharmacologic trials.

Reflexology is represented by a small trial with unmasking of participants that could have biased the results. No evidence supports choice of this modality. Likewise, hypnotherapy is not supported by the scant information provided by one case series with little detail, patient reported outcomes, or statistical assessment. Given the scope of placebo effects demonstrated in other well-conducted studies of OAB treatment, it is difficult to know whether to attribute any effect to hypnotherapy.

Conflict of interest. Changing trends in editorial standards have resulted in more complete reporting of funding source and author conflict of interest over the period in which the OAB literature was developing. Sources of funding were not reported for the majority of publications that appeared in the 1980s; and no publications in that decade reported on author conflict of interest. In the 1990s through the end of 2008, nine (56 percent) studies of procedural treatments (including sacral neuromodulation and bladder instillation or injection) reported source of funding, and six of the nine studies (67 percent) were industry sponsored. The other three were institutionally funded. Among studies of medications, 89 (77 percent) reported source of funding and among those, 82 (92 percent) were industry sponsored. Among studies that had a behavioral component, 13 (68 percent) reported on funding source and four had industry support. Author conflict of interest was poorly reported until the current decade, with fewer than half of all publications providing information. Within papers that did report conflict of interest, more than half of the authors (272 of 407) indicated that they had a financial relationship with one or more companies relevant to the research.

KQ3: Comparisons of Treatments

Comparisons between pharmacologic treatments.

Content of the literature. Twelve RCTs included direct comparisons of pharmacologic approaches. Specific comparisons have been made in the literature for the following pairs of drugs to identify differences in reduction in urge urinary incontinence or voids per day:

- Oxybutynin ER to Tolterodine ER⁸⁴
- Oxybutynin ER to Tolterodine IR⁸³

- Oxybutynin IR to Tolterodine IR^{89, 91, 94}
- Oxybutynin IR to Darifenacin⁸²
- Oxybutynin IR to Trospium IR⁹⁰
- Oxybutynin TDS to Tolterodine ER⁸⁸
- Tolterodine ER to Tolterodine IR^{140, 141}
- Tolterodine ER to Solifenacin⁹⁹
- Tolterodine ER to Fesoterodine⁹⁶
- Tolterodine IR to Solifenacin⁹⁷

Outcomes of treatment. In the majority of comparisons, neither drug was reported more effective at reducing either incontinence episodes or voids per day with a few exceptions. Both oxybutynin and tolterodine in their extended release forms demonstrated superiority in reducing urge incontinence episodes over tolterodine immediate release.^{83, 140, 142} Oxybutynin extended release was more effective at reducing voids per day than tolterodine in immediate⁸³ or extended release formulations.⁸⁴

Comparisons between procedural and pharmaceutical treatments.

Content of the literature. One RCT compared sacral neuromodulation to medical therapy. In this study, 98 participants were randomized to immediate sacral nerve stimulation or delayed sacral nerve stimulation. The delay group continued unspecified medical management for a six month period before having the procedure.

Outcomes of treatment. This study, which randomized after successful test stimulation, found a reduction in daily urge incontinence episodes from 9.7 to 2.6 in the sacral neuromodulation group, compared to an increase of 9.3 to 11.3 in the medical management group at 6 months ($p < 0.01$) for patients with refractory OAB.¹²⁴ At 18 months, 76 percent of participants receiving sacral neuromodulation reported that they were completely dry or had experienced a reduction in symptoms of 50 percent or greater. Note that the comparison is not ideal, as subjects continuing to receive medical therapy had already failed medical management.

Comparisons between behavioral and pharmacologic treatments.

Content of the literature. Seven studies (nine papers) included behavioral and pharmacologic arms in direct comparisons.^{93, 143-150} This literature included one prospective cohort study,¹⁴⁷ and six RCTs.^{93, 143-146, 148-150}

The behavioral approaches examined included bladder training,^{151 146-150} multicomponent behavioral approaches,¹⁴³⁻¹⁴⁵ and electrical stimulation.⁹³

One study reported significantly greater reductions in incontinence episodes with multicomponent behavioral modification compared to oxybutynin,¹⁴³ but no studies found a difference in reductions in voids per day. In the same study that found greater reductions in incontinence with behavioral treatment, participants reported higher satisfaction with behavioral compared to pharmacologic interventions.

Comparison of combined behavioral and pharmacologic treatment to pharmacologic alone.

Content of the literature. Six studies examined the effect of adding a behavioral intervention to drug compared to drug alone.¹⁵⁰⁻¹⁵⁵ In all but one study, the drug was tolterodine. The literature included five RCTs^{150-152, 154, 155} and one randomized open-label trial.¹⁵³

Outcomes of treatment. No added benefit for reducing incontinence episodes or voids per day was found by adding behavioral treatments to pharmacologic approaches for reduction in

incontinence. Two studies associated the addition of behavioral modification with reductions in voids per day, but two reported no difference.

KQ4: Modifiers of Treatments

Age. Eight publications examined the relationship of age to response to pharmacologic treatment.^{83, 102, 156-161} Tolterodine was the focus of four of these studies;^{156-158, 160} and one compared oxybutynin to tolterodine.⁸³ The largest study of tolterodine using clinically adjusted doses, was from an open-label clinical cohort of 2,250 patients from 462 urology practices in Germany. Participants on average were taking 2 mg per day and age did not predict response.¹⁵⁷ No studies reported lack of efficacy for reducing UUI, voids per day, or symptom distress among older participants. While effect was in some cases reduced, it was not the case that age predicted worse treatment response. Older individuals do benefit from treatment.

Prior treatment. Eight publications investigated whether prior treatment with antimuscarinics predicted treatment response.^{85, 86, 102, 162-165} In two placebo-controlled trials of oxybutynin transdermal system⁸⁵ and tolterodine IR,¹⁶² participants who had previously been on medication for OAB had comparable outcomes to those who were treatment naïve. The tolterodine study specifically commented on prior treatment failures, noting improvements among those who had failed prior treatments that were above placebo but not statistically significant; few participants were in this category making conclusions difficult to reach.¹⁶²

Baseline severity. Two studies contrasted those with UUI at baseline to those without. In an open-label study of 3,824 participants with nine months of treatment, urgency, frequency, nocturia, and OAB scales were similarly improved regardless of UUI baseline status.¹⁶⁶ In the other, open-label, study of solifenacin, with the exception of nocturia, the point estimates for improvement in individual OAB indicators were better in the group with UUI at baseline though not in each instance statistically significantly better than among those without UUI.¹⁶⁷

Severity of UUI, grouped as severe for those with 21 or more episodes per week, did not prevent tolterodine from having an effect above placebo, and both severe and non-severe groups had a high percentage reduction in incontinence episodes per day: a decrease of 67.6 percent among those with severe UUI and 71.4 percent among those with mild to moderate UUI.¹⁶⁸ In another study, those with severe baseline symptoms were less likely than others to be symptom free at 12 weeks.¹⁵⁷ Cure rates were reported to be comparable when comparing participants with UUI to those with MUI.¹⁶⁹

Urodynamic findings. Five publications related baseline urodynamic findings to outcomes of treatment.^{89, 170-173} Three did not identify urodynamic findings that predicted poor response or non-response to treatment. Both women with and without detrusor instability had comparable benefits from treatment.¹⁷¹ Classification of participants as having detrusor overactivity or sensory urgency based on urodynamics was not statistically significant as an effect modifier or predictor of treatment outcomes.¹⁷⁰ The third study to group participants by urodynamic findings had group sizes (n = 6, 25, 36, 40) too small to make definitive assessments but suggested in the two larger groups that those with low volume and high pressure profiles had comparable results to those with low volume and low pressure profiles. Among women with proven detrusor instability, those with detrusor activity in response to provocations like washing hands or running water, were less likely to improve,¹⁷² and women who reported coital incontinence were more likely to be non-responders to treatment.¹⁷³

Urodynamic findings. Five publications related baseline urodynamic findings to outcomes of treatment.^{17, 98-101} Three did not identify urodynamic findings that predicted poor response or non-response to treatment. Both women with and without detrusor instability had comparable benefits from treatment.⁹⁹ Classification of participants as having detrusor overactivity or sensory urgency based on urodynamics was not statistically significant as an effect modifier or predictor of treatment outcomes.⁹⁸ The third study to group participants by urodynamic findings had group sizes (n = 6, 25, 36, 40) too small to make definitive assessments but suggested in the two larger groups that those with low volume and high pressure profiles had comparable results to those with low volume and low pressure profiles. Among women with proven detrusor instability, those with detrusor activity in response to provocations like washing hands or running water, were less likely to improve,¹⁰⁰ and women who reported coital incontinence were more likely to be non-responders to treatment.¹⁰¹

BMI. A single trial of tolterodine ER, tolterodine IR, and placebo (n=1,235) reported that women who were above the mean for BMI (> 27kg/m²) were more likely to have UUI at baseline but achieved comparable results after 12 weeks of treatment.

Gender. Authors frequently reported that men, especially older men, fared less well in resolution of symptoms of OAB.^{85, 106, 156, 157, 174} This evidence review was focused on outcomes of treatment among women. However, in order to retain landmark studies we included a number of studies that enrolled men as long as the proportion of women in the study was 75 percent or more. This means that treatment effects may be attenuated when men are included.

KQ5: Costs of Treatments

Content of the literature. We identified five studies on financial costs related to OAB that met criteria for inclusion.¹⁷⁵⁻¹⁷⁹ All studies included assessment of direct medical costs, and two included costs due to lost productivity. One study additionally assessed financial implications for pain and suffering.

Costs of treatment. Total direct health care costs for women with OAB in 2000 were estimated at \$6.9 billion, of which \$1.1 billion were for pharmacologic treatment and \$550 million for surgical treatment. The rest was estimated for “consequences” costs, which would include things like falls, longer hospital length of stay and skin conditions. Medication costs for OAB with the two most commonly used drugs (oxybutynin and tolterodine) ranged from \$56 to \$360 over a twelve month period for newly diagnosed patients. However, total health care costs were highest for patients who take oxybutynin, relative to tolterodine in any formulation, with costs lowest for patients on tolterodine ER. Explanations for this were not apparent.

Discussion

The study of OAB as a syndrome is entering its second decade. As is typical of advancing areas of research, publications based on case series are giving way to observational cohorts. Trials, beyond those required for FDA approval of indication for OAB, are appearing in the literature and health services researchers are investigating population-level factors such as cost of care and risk of rare but serious side effects of treatments.

The 2002 ICS standardization of terminology¹⁸⁰ was associated with a productive trend toward greater attention to and clarity of operational definitions in research. Documentation of inclusion and exclusion criteria, baseline characteristics, and change in symptom profiles have

become more detailed and nuanced in the last five to seven years. Improved clarity about research definitions for conducting the study and analyzing data was the case even when authors departed from ICS definitions. Simultaneously important research gains have been made in crafting, refining, and validating questionnaire and interview instruments for classifying symptoms, assessing severity of symptoms, describing impact of OAB on quality of life, and measuring satisfaction with outcomes. Despite this momentum the overall content of the current literature is fair to poor with a preponderance of study designs that do not provide strong evidence.

We find a concerning lack of high-quality evidence to inform clinical decision-making for millions of women in the United States. Medications can provide symptom relief which is often not complete, but valued by women who struggle with OAB. Well-conducted trials of greater duration and sophistication, separate from drug development and marketing efforts, are crucial. Because benefits of current treatments are modest, opportunities exist to study how to gain synergy from combinations of types of treatments. We must note that lack of evidence of benefits is not equivalent to evidence of no benefit. A number of treatments that are potentially promising warrant continued investigation. Cross-cutting concerns about the quality of research must be addressed to achieve literature that can be meaningfully synthesized. Current literature does not permit definitive conclusions about relative benefit, harm, or costs to achieve similar results. Given how common and concerning OAB is, a priority on promoting high-quality research in the United States is imperative. Women and their care providers deserve better information to guide their choices.

Evidence Report

Chapter 1. Introduction

Importance of Overactive Bladder Treatment

At minimum, 11 to 16 million women in the United States cope on a daily basis with symptoms that include sudden strong urges to urinate, difficulty delaying voids, frequent trips to the bathroom, and in many cases involuntary loss of urine when urgency strikes.¹ They may wear pads for accidents, plan ahead for access to bathrooms, and modify their social and work lives to accommodate their symptoms. Some women are very distressed by the symptoms whether mild or severe, and some find mechanisms to adapt, reporting little trouble with symptoms or interference with normal routines. Others report their symptoms negatively influence factors as varied as self-esteem, self-assessment of attractiveness, and sexual function. Many women believe that some amount of incontinence is inevitable with aging and the majority of women with these symptoms do not talk with their health care providers about their concerns with bladder function. As a result, a small minority receive treatment.

Defining OAB

Overactive bladder syndrome, referred to as OAB in this report, is formally defined as:

- urgency, which is the complaint of sudden need to void;
- with or without urge incontinence, involuntary loss of urine with urgency symptoms;
- usually with frequency, which is the individual's perception that she voids too often during the day, and is often defined as more than eight voids during waking hours;
- usually with nocturia, which is awakening from sleep to empty the bladder.

This operational definition was formally standardized as part of a consensus process of experts, in 2002 by the International Continence Society (ICS) as part of an effort to promote healthcare professionals' and researchers' use of common terminology in the care and study of women with OAB. Components of the syndrome have had varied, and at times conflicting, nomenclature that include detrusor (bladder muscle) instability; detrusor dysfunction; detrusor dysynergia; detrusor overactivity, and irritable bladder. In each case, these terms shared a causal model that hypothesizes that mistimed or poorly regulated bladder contractions create the sensation of sudden need to void with or without leakage. However clinical study of bladder muscle function using urodynamic testing to measure characteristics like bladder capacity, pattern and timing of bladder contractions, and bladder volume at which women first experienced the urge to void, did not reveal uniform test results among women who had identical complaints. Lacking a reliable biologic marker to define and describe the severity of the condition with objective tests of the bladder itself, clinicians, researchers, pharmaceutical companies, and others came to conceptualize the symptoms of OAB, which often appeared in combination, as a syndrome.

Syndromes are medical conditions defined by the symptoms, which are the sensations (urgency), changes (frequency), or events (incontinence episodes) experienced by the individual. A syndrome is not defined by a known biologic cause. The pathophysiology of OAB is incompletely characterized and the syndrome is a diagnosis of exclusion reached when other causes of the symptoms, like urinary tract infection, urethral inflammation, or neurologic causes of incontinence are ruled out as the cause. OAB symptoms may be life long, relapsing and

remitting; or may completely resolve; which manifestations of the syndrome a woman has may also vary over time. However followup studies that track women who have OAB find that on average it is a chronic condition that women experience for a year or more at a time, and may have into the indefinite future.

Little is known about causes and most physiology and clinical research aimed at understanding etiology is now focused at the descriptive and hypothesis development and testing phase of investigation. The most promising theories postulate abnormalities in control of bladder function resulting from aberrations in neurologic signals from the bladder (sensation) and in central and peripheral nervous system regulation.

OAB and Awareness of the General Public

Prior to the mid-1990s research and clinical care focused on describing and managing frank incontinence. The term “overactive bladder” was introduced into the lexicon in the mid-1990s by Pharmacia (acquired by Pfizer in 2002) to describe the frequent urge to urinate as part of its advanced marketing campaign of Detrol (tolterodine). The company framed this as an opportunity to “destigmatize” a range of symptoms encompassing urgency, frequent voiding, and urge incontinence, so that patients would not be afraid to speak with their doctors about the problem.¹⁸¹ The construct – and subsequent marketing success – of OAB medications revolved around encouraging women to reflect on how their symptoms influenced their quality of life and the degree to which the symptoms caused inconvenience, emotional distress, withdrawal from activities, or sexual problems.

As a result, a broader spectrum of women, extending beyond those with incontinence, became candidates for treatment. This included those who were inconvenienced by or worried about frequent urination or who engaged in what has been referred to as “defensive voiding”, emptying the bladder in an attempt to extend the interval between symptoms or to reduce the amount of urine that leaks with incontinence, and “toilet mapping,” being aware of where bathrooms are and canvassing new locations to be sure the options are known. The group of those encouraged to consider treatment also came to include women who perceive they urinate more than “normal” and women whose jobs or lifestyles do not accommodate frequent, strong urges to urinate. Early advertisements featured school crossing guards and jurors who could not readily take a break. Momentum toward a very broad definition in marketing was a factor in the updated consensus definition of OAB by the ICS in 2002, so that the biomedical community would have an opportunity to formally standardize the definition with specificity.

Marketing of drug and the drug indication to physicians occurred through the usual channels, such as paid educational trips, speaking engagements, outsourcing drug studies, etc.^{182, 183} Nearly simultaneously, as a result of less restrictive rules about direct to consumer marketing, women were reached directly through the power of television and print media in new ways.¹⁸⁴ Use of “buzz drivers,” or people paid to promote the drug during news broadcasts or celebrity interviews came into play as new marketing techniques.¹⁸⁵ By 2006, the first drug to specifically target the broader definition of OAB hit the blockbuster mark of \$1 billion in sales for the year.¹⁸⁶ Two drugs, tolterodine tartrate and oxybutynin, were the only drugs approved in the United States specifically for OAB until 2004 when trospium, darifenacin, and solifenacin were introduced.¹⁸⁷ Fesoterodine, a metabolite of tolterodine was approved in October 2008. Oxybutynin is now available in a transdermal formulation. Thus over roughly a decade – a very short time window in clinical medicine – both the condition of OAB and pharmaceutical treatments for OAB became part of the consciousness of the public and the general medical community alike.

Treatment Options

Popular wisdom encourages self-management of symptoms of OAB through reduction of fluid intake, cutting back on caffeine, modifying voiding habits, and taking note of what factors like phase of the menstrual cycle, food choices, or contraceptives may influence severity of symptoms in order to adapt or reduce the impact of OAB. Over-the-counter remedies like cranberry capsules and herbal preparations are reported to promote bladder health, reduce bladder irritation, or reduce the urgency associated with bladder infections while also taking antibiotics, have crossed-over into use by women who have the symptoms without an infection. While perhaps quite common, these strategies are not well-reflected in the scientific literature.

This report is focused on those treatments that have been formally investigated including:

- Pharmacologic treatments, including prescription medications, both pills and patches
- Surgeries and procedures, such as sacral neuromodulation and botulinum injections
- Behavioral interventions, such as behavior modification programs and bladder training
- Complementary and alternative medicine, such as acupuncture and reflexology

Note that when initiated, treatment should be prompted by distress over symptoms and their influence on quality of life. The symptoms are not *de facto* harmful, though consequences such as sleep interruption or risks of falls and fractures from rushing to the toilet may be harmful.¹⁸⁸

¹⁸⁹ As a result OAB management is usually individualized to address the component symptom(s) that the individual finds most bothersome. Where possible we have tried to address treatments with respect to the primary component symptoms of OAB: urgency, frequency (daytime and nighttime), and urge urinary incontinence, so that the women, their health care providers, payors, policy-makers and others have a detailed picture of the expected outcomes of available treatments.

This Evidence Report

Scope of the Report

Evidence reviews of therapeutics seek to identify and systematically summarize objective information about the evidence related to:

- Effectiveness of specific, well-defined treatments
- Relative benefit of one treatment over another
- Common side effects and serious risks of a treatment
- Whether individual characteristics help predict who will benefit or be harmed
- Degree to which individuals find the treatment acceptable or satisfactory
- Costs of care or risk-benefit assessments

Key Questions

For this review, we operationally defined OAB as “idiopathic urinary urgency and frequency with or without associated incontinence in adult females, not related to neurogenic conditions or as a result of (incontinence) surgery.” This review is restricted to OAB, rather than exclusively mixed incontinence, stress incontinence, painful bladder syndrome, and other lower urinary tract symptoms (LUTS).

We have synthesized evidence in the published literature to address these key questions:

KQ1. What is the prevalence and incidence of overactive bladder as estimated in representative populations?

KQ2. Among women with overactive bladder, what are the short and long-term outcomes of the following treatments, or combinations of treatment approaches:

- a. Pharmacologic treatments
- b. Surgical and procedural treatments
- c. Behavioral and physical therapy treatments
- d. Complementary and alternative medicine treatments

KQ3. Where direct comparisons have been made between or among treatment modalities of interest, which modalities achieve superior outcomes with respect to benefits, short and long-term risks, and quality of life?


KQ4. Are the short and long term outcomes of these treatment approaches modified by clinical presentation, physical exam findings, urodynamic findings, menopausal status, age, or other factors?

KQ5. What are the costs associated with these treatment approaches?

Analytic Framework for the Treatment of OAB Women

The analytic framework in Figure 1 summarizes the conceptual model used to guide this systematic review by focusing the key questions on critical health care-related pathways and decision points. We recognize a number of other factors like provider prescribing preferences and types of testing performed for a patient presenting with symptoms are part of this pathway. However little literature was available to inform other nodes in the process of care.

Figure 1. Analytic framework for the treatment of OAB in women



Organization of this Evidence Report

Chapter 2 describes our methods including our search strategy, inclusion and exclusion criteria, approach to review of abstracts, to review of full publications, and for extraction of data into evidence tables, compiling evidence, and when possible conducting meta-analysis. We also describe the approach to grading of the quality of the literature and to describing the strength of the literature.

Chapter 3 presents the results of the evidence report by key question, synthesizing the findings across treatment type. We report the number and type of studies identified and we differentiate between total numbers of publications and unique studies to bring into focus the number of duplicate publications in this literature in which multiple publications are derived from the same study population. We emphasize the effect of treatment on the core symptom complex of OAB. Chapter 4 discusses the results in Chapter 3 and enlarges on methodologic considerations relevant to each key question. We also outline the current state of the literature and challenges for future research on OAB.

We have prioritized reporting on clinically relevant commonalities for United States care settings, being aware of the fact that primary care generalists and specialists alike are called upon to evaluate and treat OAB patients. We placed greatest value on the studies, and the content within studies, that is most likely to be applicable to help guide patient care, such as treatment selection, as well as inform anticipatory guidance about likely magnitude of treatment effects and risk of both nuisance side effects and serious harms.

Technical Expert Panel (TEP)

We identified technical experts on the topic of OAB in the fields of urology, urogynecology, gynecology, primary care, nursing, and patient advocacy to provide assistance during the project. The TEP (see Appendix E) was expected to contribute to AHRQ's broader goals of (1) creating and maintaining science partnerships as well as public-private partnerships and (2) meeting the needs of an array of potential customers and users of its products. Thus, the TEP was both an additional resource and a sounding board during the project. The TEP included twelve members serving as technical or clinical experts, including an AUA representative. To ensure robust, scientifically relevant work, we called on the TEP to provide reactions to work in progress and advice on substantive issues or possibly overlooked areas of research. TEP members participated in conference calls and discussions through e-mail to:

- Refine the analytic framework and key questions at the beginning of the project;
- Discuss the preliminary assessment of the literature, including inclusion/exclusion criteria;
- Provide input on the information and categories included in evidence tables;
- Develop a hierarchy of participant characteristics and outcomes to systematically assess;
- Advise about the clinical availability, use, and most common doses for therapeutics.

Because of their extensive knowledge of the literature, including numerous articles authored by TEP members themselves, and their active involvement in professional societies and as practitioners in the field, we also asked TEP members to participate in the external peer review of the draft report.

Uses of This Report

This evidence report addresses the key questions outlined above using methods described in Chapter 2 to conduct a systematic review of published literature including a meta-analysis of effects of pharmacologic treatment. We anticipate that the report will be of value to all urologic and women's health care providers, including AUA (our partner), the American College of Obstetrician Gynecologists, the American Urogynecologic Society, the American Academy of Family Physicians, American Academy of Nurse Practitioners, and other clinical groups who care for women from menarche through the remainder of their lives, such as the American Geriatrics Society. In addition, this review will be of use to the National Institutes of Health, Centers for Disease Control and Prevention, Centers for Medicare & Medicaid Services, and the Health Resources and Services Administration – all of which have offices or bureaus devoted to women's health issues. This report can bring practitioners up to date about the current state of evidence, and it provides an assessment of the quality of studies that aim to determine the outcomes of therapeutic options for the management of OAB. It will be of interest to individual women and the general public because of the high prevalence of OAB and the recurring need for women and their health care providers to make the best possible decisions among numerous options. We also anticipate it will be of use to private sector organizations concerned with women's health, such as Our Bodies Ourselves, the National Women's Health Network, the National Association for Continence, the Society of Urodynamics and Female Urology (SUFU), and the Simon Foundation for Continence.

Researchers can obtain a concise analysis of the current state of knowledge in this field. They will be poised to pursue further investigations that are needed to understand the prevalence and natural history of OAB, to clarify risk factors, develop prevention strategies, develop new treatment options, and optimize the effectiveness and safety of clinical care for those with OAB.

Chapter 2. Methods

In this chapter, we document the procedures that the Vanderbilt Evidence-based Practice Center used to develop this comprehensive evidence report on the treatment of OAB in women. We first describe our strategy for identifying articles relevant to our five key questions, our inclusion/exclusion criteria, and the process we used to abstract relevant information from the eligible articles and generate our evidence tables. We also discuss our criteria for grading the quality of individual articles and for rating the strength of the evidence as a whole. Finally, we describe the peer review process.

Literature Review Methods

Inclusion and Exclusion Criteria

Our inclusion/exclusion criteria were developed in consultation with the TEP, to capture the literature most tightly related to the key questions. Criteria are summarized below.

Table 1. Inclusion and Exclusion Criteria

Category	Criteria
Study population	Adult, community-dwelling females
Publication languages	English only
Admissible evidence (study design and other criteria)	<u>Admissible designs</u> Randomized controlled trials, cohorts with comparison, case-control, and case series <u>Other criteria</u> Original research studies that provide sufficient detail regarding methods and results to enable use and aggregation of the data and results Patient populations must include women with overactive bladder Studies must have relevant population ≥ 50 participants Studies must address one or more of the following for overactive bladder: Treatment modality Symptom management approach Short- and long-term outcomes and quality of life Prevalence and/or incidence Relevant outcomes must be able to be abstracted from data in the papers

We excluded studies that (1) were not published in English; (2) did not report information pertinent to the key questions; (3) had fewer than 50 female participants [at enrollment]; (4) had less than 75 percent female participants or failed to report results by gender; and (5) were not original studies.

For this review, the relevant population for all key questions was women with overactive bladder, defined as “idiopathic urinary urgency and frequency with or without associated urge urinary incontinence, not related to neurogenic conditions or as a result of (incontinence) surgery.” The same inclusion/exclusion criteria were applied to identify papers for treatment-related key questions. We applied additional restrictions for inclusion and exclusion of incidence and prevalence publications for KQ1. To inform this question about the epidemiology of OAB and/or its component symptoms, we required that the study methods specify a population base *a*

Appendixes and Evidence Tables for this report are provided electronically at <http://www.ahrq.gov/downloads/pub/evidence/pdf/bladder/bladder.pdf>

priori from which a sample of individuals was drawn as a representative selection to estimate the true proportion of prevalent and incident cases of OAB in the larger population. Additional information is provided in the results for KQ1. For KQ5, we required that publications provide data on direct costs in United States dollars for treatments reviewed in this report.

Treatment studies that included at least 75 percent women were included. This level was selected with expert input to avoid restricting to studies with only female participants as a large proportion of this literature includes both men and women, and to establish a threshold below which the difference in underlying processes (e.g., BPH in men) might substantially influence treatment effects. Studies with lower proportions of women were included if they presented results separately for women; or indicated that an interaction with gender was tested and found not to exist; or gender was controlled in the analysis. Publications about incontinence that did not distinguish between, or present results by urge, stress, and/or mixed incontinence were excluded.

Literature Search and Retrieval Process

Databases. We employed multi-term search strategies to retrieve research about treatment of overactive bladder in women, including exploration of three databases: PubMed, MEDLINE®, EMBASE, and the Cumulative Index to Nursing and Allied Health Literature (CINAHL). We also hand-searched the reference lists of relevant articles to identify additional studies for review.

Search terms. Controlled vocabulary terms served as the foundation of our search in each database, complemented by additional keyword phrases to represent the myriad ways in which overactive bladder is referred to in the clinical literature. We also employed indexing terms within each of the databases to exclude undesired publication types (e.g., reviews, case reports, CME handouts) and items published in languages other than English.

Tables 2 to 4 outline our search terms and the yield from each database. Our searches were executed between April and October, 2008, and were not limited by date. From PubMed, we identified 2,400 items for further review; EMBASE yielded 318 items, including 310 already identified in PubMed and 8 unique items; CINAHL yielded 264 citations, including 240 duplicates with PubMed and 24 new articles for review.

Table 2. PubMed search strategies (last updated October 1, 2008)

	Search terms	Search results
#1	("Urinary Bladder, Overactive"[Mh] OR "overactive bladder" OR "urge incontinence" OR urinary incontinence, urge[mh] OR "detrusor instability" OR "overactive detrusor" OR "urinary urgency" OR "urinary frequency" OR "irritable bladder" OR "detrusor overactivity") AND "female"[MeSH Terms] AND "humans"[MeSH Terms] AND English[lang]	2,886
#2	#1 AND editorial[pt]	10
#3	#1 AND letter[pt]	30
#4	#1 AND case reports[pt]	164
#5	#1 AND review[pt]	299
#6	#1 AND practice guideline[pt]	2
#7	#1 NOT (#2 OR #3 OR #4 OR #5 OR #6)	2,400*†

* Approximately 250 of these citations represent pediatric literature (due to variability in indexing for this topic, we were unable to exclude pediatric literature at the search strategy level).

† Numbers do not total due to exclusions in more than one category; 5 items were indexed as both letters and case reports and 14 items were indexed as both reviews and case reports

Table 3. EMBASE search (OVID) (last updated October 1, 2008)

Search Terms	Search Results
#1 *overactive bladder/ or *urinary urgency/ or *urge incontinence/ or *urinary frequency/ or *detrusor dyssynergia/ or *bladder irritation/	1624
#2 limit 1 to (human and female and english language and (adult <18 to 64 years> or aged <65+ years>))	363
#3 #2 and review.pt.	12
#4 #2 and conference paper.pt.	4
#5 #2 and editorial.pt.	1
#6 #2 and letter.pt.	0
#7 #2 and note.pt.	3
#8 #2 and short survey.pt.	4
#9 #2 and case report/	18
#10 #2 and practice guideline/	4
#11 #2 and "systematic review"/	1
#12 #2 and meta analysis/	1
#13 #2 not (3 or 4 or 5 or 6 or 7 or 8 or 9 or 10 or 11 or 12)	318*†

* Overlap with PubMed: 310 citations; 8 new citations retrieved for inclusion.

† Numbers do not total due to exclusions in more than one category: 1 item was indexed as both a case report and review; 1 item was indexed as both a case report and a note; and 1 item was indexed as both a review and a systematic review.


Table 4. CINAHL search (EBSCO) (last updated October 1, 2008)

Search Terms	Search Results
#1 (MH "Urge Incontinence") or (MH "Overactive Bladder") or "overactive bladder" or "urge incontinence" or "urge urinary incontinence" or "detrusor instability" or "overactive detrusor" or "urinary urgency" or "urinary frequency" or "detrusor overactivity" and (MH "Adult+") and (ZL "ENGLISH") and (PT "Journal Article")	305
#2 #1 and case reports	18
#3 #1 and review	6
#4 #1 and CE material	5
#5 #1 and abstract/commentary	7
#6 #1 and consumer literature	5
#7 1 not (2 or 3 or 4 or 5 or 6)	264*

* Overlap with PubMed: 240 citations; 24 new citations retrieved for inclusion.

Yield of literature searches. Figure 2 presents the yield and results from our searches. Beginning with a yield of 2,559 articles, we retained 232 articles covering 221 studies that we determined were relevant to answer our key questions and met our inclusion/exclusion criteria.

Figure 2. Disposition of articles for the treatment of overactive bladder (OAB)



KQ = key question

*The number of articles addressing each key question and those excluded exceed the total number of articles in each category because some of articles fit into multiple exclusion categories or addressed more than one key question. The excluded papers focused exclusively on populations of individuals with movement disorders, spinal cord injuries, or multiple sclerosis, for example. As a result, they did not meet the criteria for an “idiopathic” syndrome which was a requirement of the report.

Article selection process. Once we identified articles through the electronic database searches, review articles, and bibliographies, we examined abstracts of articles to determine whether studies met our criteria. Two reviewers separately evaluated the abstracts for inclusion or exclusion, using an Abstract Review Form (Appendix B). If one reviewer concluded that the article could be eligible for the review based on the abstract, we retained it. The group included three physicians (KH, DB, RW), and two senior health services researchers (MM, SM).

Of the entire group of 2,559 articles, 586 required full text review. For the full article review, two reviewers read each article and decided whether it met our inclusion criteria, using a Full Text Inclusion/Exclusion form. Reasons for article exclusion are listed in Appendix D.

Literature Synthesis

Development of Evidence Tables and Data Abstraction Process

The staff members and clinical experts who conducted this review jointly developed the evidence tables. We designed the tables to provide sufficient information to enable readers to understand the studies and to determine their quality; we gave particular emphasis to essential information related to our KQs. We based the format of our evidence tables on successful designs used for prior systematic reviews.

The team was trained to abstract by abstracting several articles into evidence tables and then reconvening as a group to discuss the utility of the table design. We repeated this process through several iterations until we decided that the tables included the appropriate categories for gathering the information contained in the articles. A priori, with the technical expert panel, a hierarchy of baseline characteristics and outcome measures was developed: UUI episodes, urgency, incontinence, voids, nocturia, QoL, urodynamic measures, and adverse events. All team members shared the task of initially entering information into the evidence tables. Another member of the team also reviewed the articles and edited all initial table entries for accuracy, completeness, and consistency. The two abstractors reconciled disagreements concerning the information reported in the evidence tables. The full research team met regularly during the article abstraction period and discussed global issues related to the data abstraction process. In addition to outcomes related to treatment effectiveness, we abstracted all data available on harms. Harms encompasses the full range of specific negative effects, including the narrower definition of adverse events.

The final evidence tables are presented in their entirety in Appendix C. Studies are presented in the evidence tables alphabetically by the last name of the first author. When possible, studies resulting from the same study population were grouped into a single evidence table. A list of abbreviations and acronyms used in the tables appears at the beginning of that appendix.

Synthesis of the Evidence

A series of spreadsheets was created to support systematic tabulation and assessment of study characteristics including key study population characteristics, number of participants by group, treatment received, length of followup, age of participants by group, outcomes measured and outcomes. This allowed us to identify common threads in reporting across publications.

Within the pharmaceutical treatment studies all unique trial arms were entered into a spreadsheet with exact values to two-decimal points as available for baseline measures, followup measures, difference from baseline, and for the related statistical indicators of precision (such as

standard deviations, standard errors, or confidence bounds). This was done to facilitate calculation of weighted averages and to support meta-analysis.

Conduct of meta-analysis. Descriptive statistics were computed and examined for homogeneity among studies. Studies that reported weekly rates for UIIs and voids were standardized to daily rates. When only ranges of continuous variables were reported (instead of standard deviations), we estimated the standard deviations by dividing the range by four.⁵ Study results were combined and summarized using two meta-analysis techniques, weighted averages and fixed effects regression models.⁶ In particular, minimum variance weighted averages of the mean daily decrease in UII and voids per arm were computed using weights that were inversely proportional to their standard errors. To borrow strength across arms, we used fixed effect regression models with robust standard errors (to account for the clustering by study) and weighted the study arms inversely proportional to their standard errors of the mean. Each arm was treated as a fixed effect and study was not included in the model except in the sense that the clustering was addressed by the robust standard errors. Fixed effects models were also adjusted for mean age and proportion of women in each arm. We used STATA 10.0 and R statistical packages for computations.

Summary tables within this report. Each of the pharmacologic agents in this report is available in a least one dose form for clinical use. As part of the process of Food and Drug Administration (FDA) approval each has been determined superior to placebo for at least one facet of the syndrome, e.g., urge urinary incontinence, frequency of urination, symptoms of urgency, or nocturia. The experience of having overactive bladder is a constellation of these self-reported events, symptoms, and the impact that they have on an individual's life. Thus measures of quality of life, interference with daily activities, degree of distress from symptoms, and satisfaction with the outcomes of treatment are also common and helpful metrics in this literature. Where common measures are available across studies using roughly comparable assessments (i.e., similar index questions, time intervals, etc), we have compiled tables to summarize outcomes of treatments.

Given the content of the literature, this means that the majority of the information included in tables is for the outcomes of number of urge incontinence episodes per day and number of voids per day. (Studies with weekly or other metrics that could be converted to daily metric are included.) Because momentum in drug development within related classes of drugs has been toward daily dosing, the pharmacologic treatments are arranged from highest dose at lowest frequency of administration (daily) to lower doses and greater administration frequency (twice or more daily). Placebo arms from the same trials in which the drugs were evaluated are included within the table, or for areas in which the literature is large, in companion placebo outcome tables. The number of weeks of treatment and timing of followup outcome assessment were the same in these trials. The weeks of treatment column is therefore comparable to weeks at evaluation of outcomes.

Summary tables include data from distinct clinical trial arms in which the drug and dose or other type of treatment were evaluated for the related outcome. As a result, a single study may contribute more than one treatment arm as well as a placebo arm to the summary tables. For pharmacologic treatment we included only study arms in which no dose adjustment was allowed. Because many studies are dose finding with multiple drug arms or are direct comparisons of pharmacologic agents, there are more drug arms than placebo arms for virtually all drugs and treatment types.

Quality Rating of the Individual Studies

Rating the quality of individual articles. We developed our approach to assessing the quality of individual articles based on our prior experience with conducting systematic reviews.

Internal validity. The criteria for assessing internal validity were as follows:

Randomized allocation to treatment. This assessment combines randomization and method of randomization into a single criterion with a three-point scale.

Rationale: By randomly assigning groups to the intervention of interest, other factors that may confound the results are equally distributed between groups (assuming a large enough sample size). This equal distribution minimizes the chances of over- or underestimation of treatment effect based on unequal distribution of confounding factors.

If randomized, we also evaluated the study for randomization methods, using the rationale described in Matchar and colleagues, 2001.¹⁹⁰

Rationale: “Pseudo-randomization” methods may be susceptible to bias, as demonstrated by evidence of unequal distribution of subject characteristics¹⁹¹ and larger effect sizes compared with studies using more rigorous methods.¹⁹² In addition, methods of allocation concealment are also important in preventing bias (e.g., use of prepared sealed envelopes).

We combined these elements into a single operational definition, as described below:

Operational definition: Criterion met if randomization methods were not susceptible to bias, such as computer-generated numbers in sealed sequentially numbered envelopes (+). Criterion not met by studies that either used methods more prone to bias, such as alternate medical record numbers, or did not describe randomization methods or methods of allocation concealment (-). Criterion not applicable if treatment was not randomly allocated (NA).

Masking.

Rationale: Masking, also known as blinding, refers to the concealment of treatment allocation from the care provider, the assessor, and the patient.¹⁹³ In certain trials, particularly surgical trials, masking the patient or the surgeon from the treatment allocation can be challenging or impossible. Similarly, masking the assessor assigned to record immediate post-procedural outcomes such as wound healing can also be difficult. Nevertheless, when possible, masking prevents expectations from influencing findings.

Operational definition: Criterion was met if assessors and participants were masked to treatment or group (+). Criterion was not met if either care provider, assessor, or patient were not masked (-). Criterion not applicable if treatment was not randomly allocated.

Adequate description of patients and control selection criteria.

Rationale: Patient characteristics that might affect outcomes (such as severity of symptoms, duration of symptoms, failure of prior treatment, or medical comorbidities) are likely to differ between two interventions. If these differences are not characterized, then erroneous conclusions may be drawn.

Operational definition: Criterion met if (a) inclusion and exclusion criteria for participation in the study were well described.

We expected that the study population should be adequately described to make clear the potential for confounding in the analysis. We expected the study authors to adequately describe the study population such that it could theoretically be reproducible by another investigator. We expected comparable methods to be used to identify and screen participants across exposure or treatment groups.

Description of loss to followup.

Rationale: Failing to account for patients lost to followup may lead to erroneous conclusions, especially if the loss to followup is related to either the underlying disease or the intervention (e.g., patients seeking care elsewhere because of continuing symptoms or unacceptable side effects of treatment).

Operational definition: Criterion met for adequate followup (+) if (a) loss to followup was explicitly reported and (b) no more than 20 percent of any study arm was lost to followup. Those studies with less than 10 percent lost to followup were given an extra (+). Studies with greater than 20 percent lost to followup were considered inadequate for this measure (-).

Description of dropout rates.

Rationale: Dropout rates may reflect differences in clinically important variables, such as side effects or treatment response. Failure to account for dropouts may result in erroneous conclusions similar to those seen with failure to account for loss to followup.

Operational definition: Criterion met if (a) patients dropping out of the study prior to completion were reported and (b) no more than 10 percent in any study arm left the study for reasons related to the study intervention or withdrawal of consent.

Power calculation provided.

Rationale: Many studies, especially case series, lack sufficient power to detect clinically important differences in outcomes or patient characteristics.

Operational Definition: Criterion met if a power calculation (pre or post) was provided.

Recognition and description of statistical issues.

Rationale: Use of inappropriate tests may lead to misleading conclusions. For example, variables such as number of voids per day or costs are often not normally distributed; use of means instead of medians when data may be affected by outlying observations can be misleading.

Operational definition: Criterion met if (a) appropriate statistical tests were used (e.g., nonparametric methods for variables with nonnormal distributions, or survival analysis techniques to account for loss to followup and dropouts) and (b) potential study limitations regarding design and analysis were discussed. Criterion not met if (a) inappropriate statistical tests were used or (b) study limitations were not discussed. An intention-to-treat (ITT) analysis was required of clinical trials.

External validity. The criteria for assessing external validity were as follows:

Baseline characteristics: We created a composite score for adequacy of the description of baseline characteristics. At minimum, we expected age and baseline OAB status to be presented. If either of these were omitted, criteria were not met. If the authors provided additional information above and beyond age and OAB status at baseline on any of the following, they were awarded an additional +: race/ethnicity, BMI, parity, menopausal status, prior treatment/surgery, duration of symptoms.

Required elements:

Description of age of study population.

Rationale: The outcomes of many interventions are affected by patient age. Age is especially important in studies related to reproductive health in women and associated with rates of overactive bladder.

Operational definition: Criterion met if summary statistics of subject age were given by comparison group. Criterion not met if summary statistics were not given.

Baseline OAB status.

Rationale: The baseline level of severity of OAB could affect the likelihood of successful treatment. Furthermore, definitions of OAB are not consistent across studies and may include different combinations of urgency, frequency, and incontinence that could affect interpretation of the outcomes. Therefore, we sought to determine whether studies defined OAB status by ICS or other criteria, by UUI alone or by combinations of UUI, urgency and frequency.

Operational definition: Criterion met if symptoms of OAB were presented by study group.

Length of followup.

Rationale: Outcome measures may vary depending on when they are obtained. Description of when outcomes were measured facilitates comparison between studies. We considered three months to be a minimally acceptable period of followup for observing effectiveness of treatment for OAB.

Operational definition: Criterion met if the study followed participants for at least three months, with an extra point provided for greater than or equal to six months.

Adequate description of methods used for outcome measurement.

Rationale: Comparison between studies requires common methods of measurement, which in turn requires adequate description of the methods used to assess comparability.

Operational definition: Criterion met if (a) methods used to measure outcomes were adequately described or referenced (e.g., 2002 ICS; QoL scales), (b) definitions were given (e.g., description of outcomes classified as “adverse events”), or (c) outcomes were unambiguous (e.g., reduction in number of voids per day). Criterion not met if (a), (b), or (c) was not present.

Adequate description of validity and reliability of outcome measurement.

Rationale: Measurements of outcomes are only useful if changes in the outcome being measured are reflected in changes in the measurement (validity) and if these changes are reasonably consistent between the same observer measuring at different times or between different observers (reliability). For example, changes in a scale to assess menstrual blood flow should correlate with some other physiological measure of menstrual blood loss, and this correlation should be consistent when different women apply the same scale.

Operational definition: Criterion met if (a) a description of the methods used to assess validity and reliability of at least one outcome measure was provided, (b) a reference to another article documenting validity and reliability was provided, or (c) only unambiguous outcomes were included as primary outcomes. Criterion not met if (a), (b), or (c) was not present.

Adequate description of the intervention provided to subjects.

Rationale: The ability to replicate study results is dependent on adequate description of methods. Additionally, readers should be aware of aspects of clinical care that might influence outcomes.

Operational definition: Criterion met if (a) a detailed description of the therapy (dose, dosing schedule, protocols for behavioral interventions, and route of administration for medications and/or techniques for invasive therapies) was provided; (b) a reference to another publication describing the procedure was provided; or (c) statistical adjustment was made for likely sources of variation in clinical care (e.g., site where care was given, type of specialist providing care, individual provider, dose and timing).

Criterion not met if (a), (b), or (c) was not provided.

Table 5. Scoring algorithm for quality rating of individual studies

Definition and Scoring Algorithm	Rating
Score Algorithm for Internal Validity Quality Rating	
<ul style="list-style-type: none"> No negative scores, lowest loss-to-followup score, and lowest dropout rate One negative score or intermediate loss-to-followup High loss-to-followup score, or high dropout rate OR Two negative scores OR One negative score and intermediate loss-to-followup score 	<p>Good internal validity</p> <p>Fair internal validity</p> <p>Poor internal validity</p>
Score Algorithm for External Validity Quality Rating	
<ul style="list-style-type: none"> No negative scores One or two negative scores Three or more negative scores 	<p>Good external validity</p> <p>Fair external validity</p> <p>Poor external validity</p>
Score Algorithm for Overall Quality Rating	
<ul style="list-style-type: none"> Good internal validity and good external validity Fair internal validity and fair external validity OR Good internal validity and fair external validity OR Good internal validity and poor external validity OR Fair internal validity and good external validity OR Poor internal validity and good external validity Poor internal validity and poor external validity OR Fair internal validity and poor external validity OR Poor internal validity and fair external validity 	<p>Good overall</p> <p>Fair overall</p> <p>Poor overall</p>

Strength of Available Evidence

Our scheme follows the criteria applied in earlier systematic reviews of systems for rating the strength of a body of evidence.^{194, 195} That system includes three domains: quality of the research, quantity of studies (including number of studies and adequacy of the sample size), and consistency of findings. Two senior investigators assigned grades by consensus.

We graded the body of literature for each key question and present those ratings as part of the discussion in Chapter 4. The possible grades were:

I. Strong: The evidence is from studies of strong design; results are both clinically important and consistent with minor exceptions at most; results are free from serious doubts about generalizability, bias, or flaws in research design. Studies with negative results have sufficiently large samples to have adequate statistical power.

II. Moderate: The evidence is from studies of strong design, but some uncertainty remains because of inconsistencies or concern about generalizability, bias, research design flaws, or adequate sample size. Alternatively, the evidence is consistent but derives from studies of weaker design.

III. Weak: The evidence is from a limited number of studies of weaker design. Studies with strong design either have not been done or are inconclusive.

IV. No evidence: No published literature.

External Peer Review

As is customary for all systematic evidence reviews done for AHRQ, this report was reviewed by a wide array of individual outside experts in the field, including our TEP, and from relevant professional societies and public organizations. AHRQ also requested review from its own staff. The Scientific Resource Center sent 11 invitations for peer review. Reviewers included clinicians (e.g., urologists, urogynecologists, gynecologists, geriatricians, family medicine physicians, and nurse practitioners), representatives of federal agencies, advocacy groups, and potential users of the report.

The Scientific Resource Center charged peer reviewers with commenting on the content, structure, and format of the evidence report, providing additional relevant citations, and pointing out issues related to how we had conceptualized and defined the topic and KQs. We also asked reviewers to complete a peer review checklist. The Scientific Resource Center received eight responses in addition to comments from AHRQ staff. The individuals listed in Appendix E gave us permission to acknowledge their review of the draft. We compiled all comments and addressed each one individually, revising the text as appropriate.

Chapter 3. Results

KQ1. Prevalence and Incidence of Overactive Bladder

To understand the epidemiology of OAB and associated symptoms (frequency, urgency, and urge incontinence), we sought publications that provided estimates of prevalence, which is the proportion of the population with the condition, or those that examined the proportion with onset of new symptoms over time, which is termed incidence. The strongest estimates come from a well-specified population-base that allows sampling to generate a representative group.

We applied operational criteria to select those studies best suited to estimate prevalence and incidence. The key criterion for inclusion was that the study authors specified a population base *a priori* from which a sample of individuals was drawn as a representative selection of individuals to estimate the true proportion in the larger population. The population was required to be clearly defined and capable of being enumerated such as the population of a region, the participants in a health plan or individuals on voter registration roles. This “roster” defines the sampling frame from which stratified samples are drawn. Often the research team has an explicit statistical goal of randomly selecting participants with aggregate characteristics (range of age, race/ethnicity, income level, etc.) that approximate the entire population of a city, region, or country. Common approaches include random digit dialing within a specific geographical region, or mailings of questionnaires to a sample of individuals who are in a pool of registered voters, registered residents, or on the patient panels of a national health systems or large healthcare organizations. Outside the United States, household canvassing and administered interviews were also common.

The characteristic of being able to define a sampling frame from which participants were subsequently drawn was a required characteristic for inclusion in the prevalence and incidence summary. Other approaches to the study of large groups, such as enrolling individuals who are presenting for clinical care, or who have just had a health event like childbirth or surgery, can be informative, however, such studies do not by virtue of their design, generate participant pools who are representative of a larger population. They represent those with access to care, who have sought care for a problem visit, chronic condition, or preventive care, or who have a medical condition or experience in common. Such designs may also reflect characteristics of the site of care at which the cohort was recruited, such as a high proportion of indigent patients or specialty referral patients. For this reason, cohort studies and clinical samples, without a population-based sampling frame were excluded in order to emphasize the broadest picture of the epidemiology and natural history of OAB is as possible.

Prevalence of Overactive Bladder

Using this approach, we identified 75 publications,^{2-4, 10-81} from 60 distinct study populations. Detailed summaries for all studies are included in Appendix C and a summary is provided in Table 6.

Table 6. Study characteristics of the prevalence and incidence literature

Study Characteristic	Population Studied				
	US (n=15)	European (n=24)	Asian (n=13)	Other (n=8)	Total (n=60)
Publications after 2002 ICS Definitions					
Distinct populations	(n=13)	(n=11)	(n=7)	(n=7)	(n=38)
Prevalence Only	10	7	7	6	30
Incidence Only	2	--	--	--	2
Both	1	4	--	1	6
Assessment Method					
Mailed	3	7	2	--	12
Telephone	2	2	1	3	8
Administered	6	1	4	3	14
Other	2	1	--	1	4
Conditions Assessed*					
Urge Incontinence	11	7	5	5	28
Overactive Bladder	2	4	5	3	14
Urgency	4	3	4	3	14
Frequency	3	3	5	3	14
Funding Source					
National (NIH, MRC, etc.)	8	4	1	--	13
Foundation/Other	--	2	--	1	3
Industry	4	3	1	2	10
Not Reported	1	2	5	4	12
Publications before 2002 ICS Definitions					
Distinct populations	(n=2)	(n=13)	(n=6)	(n=1)	(n=22)
Prevalence Only	--	13	6	1	20
Incidence Only	--	--	--	--	--
Both	2	--	--	--	2
Assessment Method					
Mailed	--	8	2	1	11
Telephone	1	2	2	--	5
Administered	1	2	2	--	5
Other	--	1	--	--	1
Conditions Assessed*					
Urge Incontinence	2	11	6	1	20
Overactive Bladder	--	1	--	--	1
Urgency	--	3	2	--	5
Frequency	--	3	2	--	5
Funding Source					
National (NIH, MRC, etc.)	2	3	--	--	5
Foundation/Other	--	1	1	1	3
Industry	--	4	1	--	5
Not Reported	--	5	4	--	9

* Inclusive: Total is greater than number of studies because some publications report multiple conditions.

Fifteen studies were conducted in United States populations; 24 in European populations; 13, Asian; and 8 other countries. Thirty-eight of these studies appeared in print after the 2002 International Continence Society revised definitions and 37 percent of these reported specifically about incidence or prevalence of OAB. One study prior to the consensus definitions used fully comparable definitions and the term “overactive bladder”.

A total of 15 studies provided information about OAB prevalence (Table 7). These studies included a total of 64,528 women in 16 distinct populations. The majority of respondents completed questionnaires returned by mail. The highest estimates of OAB prevalence was 31.3 percent in a telephone questionnaire study conducted in Korea in which the average age of participants was 59 years; the lowest was 7.7 percent in a mailed questionnaire conducted in the United Kingdom among women of nearly identical average age. Across all studies the weighted average prevalence of OAB was 13.7 percent. Excluding the highest and lowest estimates, an estimated 14.8 percent of women meet criteria for OAB, with 8.0 percent of those surveyed having OAB with a component of urge incontinence. Combined estimates from the two United States populations are similar: 14.7 percent with OAB and 11.3 percent with a component of urge incontinence.

No clear pattern of higher or lower estimates for prevalence of OAB was associated with survey response rates. The direction of bias is therefore difficult to estimate: researchers have noted both that those with symptoms may be more likely to be interested in the topic and to respond and that the social stigma or embarrassment associated with bladder control symptoms may prompt under-response. In each case in which the authors addressed non-response, they report the demographic characteristics of those who did not respond were similar to those who did with the exception of several authors who noted modest under-representation of the very oldest residents.

Age and prevalence of OAB. The relationship between OAB and age was fairly uniform across studies with a trend to increase with age.^{2, 3, 12-14, 38, 42} As discussed below, this increase appears to be more notable for OAB than UUI. Because OAB criteria can be met with urgency combined with incontinence, frequency, or nocturia, the “amplification” of age effects for OAB risk may in part be related to well-documented increases in stress urinary incontinence and therefore also mixed stress and urge incontinence with age. Prevalence of OAB for women in their 20s was in the range of 4.6 to 5.9 percent^{2, 3, 11, 13} while uniformly double digits, 11.7 to 19 percent^{2, 3, 11, 13} for women older than 60. Several researchers noted a threshold effect such that prevalence was not statistically different until an inflection in the 60s or 70s.^{12, 14, 38, 42} Few data were provided, however discussion materials often noted that all component symptoms: urgency, frequency, nocturia, and urge and mixed incontinence, increased with age therefore contributing to the rise in OAB.

Table 7. Prevalence of OAB

Author, Year Country	Respondents Response*	Measurement	Age of Respondents† Mean, Range	OAB Prevalence (%)
Lawrence et al. ^{2, 3, 13} 2008 US	4,103 34.0%	Mailed EPIQ	56.5 ± 15.8 (25, 84)	Any: 13.3 Wet: 12.7
Wagner et al. ^{4, 42} 2002 US	2,735 83.9%	Telephone	54.2 (18, ≥ 75)	Any: 16.9 Wet: 9.3
Herschorn et al. ¹² 2007 Canada	518 NR	Telephone	44.5 ± 17.2 (18, 90)	Any: 14.7 Wet: 7.1
Irwin et al. ¹⁸ 2006 Europe & Canada	9,000^ 33.0%	Telephone	(18, ≥ 70)	Any: 12.8 Wet: 6.3
Corcos et al. ³⁸ 2004 Canada	1,683 53.7%	Telephone	50.9 (35, ≥ 75)	Any: 21.3 Wet: 6.5
McGrother et al. ²⁶ 2006 UK	12,570 65.3%	Mailed	59.5 ± 13.0 (40, 98)	Any: 7.7
Dallosso et al. ³⁹ 2004 UK	12,568 30.2%	Mailed	Median: 61	Any: 15.9
Van Der Vaart et al. ⁴⁸ 2002 Netherlands	933 67.0%	Mailed UDI	34.2 ± 3.2 (20, 45)	Any: 11.9
Choo et al. ¹⁴ 2007 Korea	1,005 NR	Telephone	59.4 ± 11.6 (40, 89)	Any: 31.3 Wet: 15.0
Yu et al. ³⁰ 2006 Taiwan	925 33.5%	Administered	49.9 (30, 79)	Any: 18.3
Homma et al. ³¹ 2005 Japan	2,380 45.0%	Mailed	61 (41, 100)	Any: 11.0 Wet: 7.0
Song et al. ^{29, 33, 34} 2005 China	4,684 77.2%	Mailed BFLUTS	40.4 ± 11.1	Any: 8.0 Wet: 5.6
Chen et al. ^{41, 47} 2003 Taiwan	1,247 78.7%	Administered Bristol	43.2 ± 15.1 (20, ≥ 65)	Any: 18.6 Wet: 9.1
Teloken et al. ²³ 2006 Brazil	449 NR	Written	(15, 55)	Any: 23.2
Milsom et al. ⁵¹ 2001 Europe	9,728 NR	Telephone	(40, ≥ 75)	Any: 17.4
	Total: 64,528 Weighted average: 49.5%			Weighted averages: Any: 13.7 Any (without extremes): 14.8 Wet: 8.0

* Proportion of eligible sample who responded to survey and had responses included in analysis.

^Exact number of women not reported.

† Data provided is that included by the authors. Hierarchy was to provide mean if no indication of skew; median if author provided only median or data to show that median better captured a skewed distribution, and then range. Range was only provided when other options were not available. Complete information for each study is included in evidence tables; OAB-wet indicates OAB with incontinence.

Prevalence of Urge Urinary Incontinence

A larger number of studies (n=48) examined urge urinary incontinence as the primary prevalence estimate of interest. Twenty-six distinct study populations, with 36 publications, appeared after the ICS 2002 consensus definitions.^{2, 3, 10-14, 17-19, 21, 25, 27, 29, 31-38, 70, 72, 75, 80} These are presented here in greater detail because the definitions used are more similar to those that define “OAB-wet” and therefore are more likely to have comparability in the measures. Across all these populations, prevalence of urge incontinence ranged from a low of 1.5 percent in a large European study of all adult women from ages 18 to 70,¹⁸ up to a high estimate of 22.0 percent in a United States mall-based consumer survey³⁶ and 26.4 percent among a household sample of Jordanian women age 50 to 65.³⁵ Each of the latter had somewhat ambiguous survey items.

Average urge incontinence prevalence by region was:

United States: 9.7 percent among 37,596 respondents from nine populations^{2-4, 10, 11, 13, 17, 19, 25, 27, 32, 36, 40, 42, 72}

Europe: 10.6 percent among 68,051 respondents from seven studies including nine countries^{18, 37, 43, 45, 48, 75, 81}

Asia: 9.6 percent among 14,537 respondents from five populations in three countries^{14, 21, 29, 31, 33, 34, 41, 47, 80}

Other: 12.5 percent among 6,219 respondents in five populations from five countries^{12, 35, 38, 49, 70}

Though more than half of studies did not report the frequency of urge incontinence episodes required to meet the case definition (e.g., weekly, monthly), or used nonspecific terms such as “mostly” and “sometimes” in definition, these estimates are concordant with the range of estimates for the prevalence of OAB with urge incontinence features, suggesting that the measurement instruments captured similar features. These estimates are consistent with the AHRQ report on the Prevention of Fecal and Urinary Incontinence in Adults, which reported prevalence of UUI in community dwelling adults as increasing “from 5 percent in younger women to 10 percent in women 45-64 (32 studies), and to 12 percent in women older than 65 years (28 studies)”¹⁹⁶.

In the nine studies in which the interval of occurrence required to meet the definition of urge incontinence was specified, there was not a clear pattern of relationship between length of the interval and proportion of women classified as having urge incontinence.^{10, 17, 27, 31, 34, 37, 38, 72, 80} Overall the average prevalence of UUI across these nine studies was 8.9 percent. The two studies that required weekly symptoms reported a prevalence of 7.0 and 6.5 percent (in populations which had an average age of 61, and an age range from 35 to 75 without a mean reported, respectively).^{21, 31, 38} The average across six studies that reflect a wide age range and required at least monthly episodes was 9.4 percent, and the sole study reporting any urge incontinence within the year was 7.9 percent with an average respondent age of 51.⁷² This lack of a pattern associated with the operational definition of frequency of urge incontinence episodes may result from variation in the methods of measurement or characteristics of the respondents that obscures any trend, or, more likely given large population-based samples, similarity in estimates may reflect that most women who are affected have fairly frequent symptoms and are detected and properly classified regardless of interval definition required.

Other Measures

Urgency symptoms. Individually, the symptoms of urgency and frequency are common. The range of prevalence for urgency symptoms (in post ICS studies) was from 8.0 percent in a young population of pre-menopausal Indian women,⁵⁰ to the highest estimate of 45.4 percent among an even younger population of European women (mean age 34).⁴⁸ Despite this wide range, which may reflect varied operational definitions used by researchers, nine of ten estimates for the prevalence of urgency are 10 percent or above.^{12, 14, 18, 34, 38, 47, 48, 50, 73}

Frequency. Thirteen studies with fourteen groups reported on frequency. Most defined frequency as more than eight voids per day (or more). The range of estimates for women with daily frequency was 5.2 percent among Thai women ages 20 to 59,²⁰ to 34 percent among Danish women ages 20 to 45,⁴⁸ and 49 percent among Japanese women with an average age of 61.³¹ The average proportion of women with frequent voiding across study groups was 16.0 percent, with five estimates between 5 and 15 percent,^{12, 18, 20, 45, 50} and four estimates between 15 and 25 percent.^{14, 34, 38, 47}

Age as a predictor of UUI. The relationship between urge urinary incontinence and increasing age was less consistent across studies than that for OAB and age, with some authors reporting nearly identical prevalence across all age brackets and no statistical trend^{15, 53} and others noting increases parallel to OAB,^{11, 12, 14, 42} more modest increases⁶⁴ or threshold ages as in OAB which are inflection points at which prevalence was meaningfully increased, typically among the oldest age strata: >60,¹¹ >65,⁴² >70,⁵⁵ and >75.³⁸ In aggregate, the trend was less pronounced or not apparent for a strong and continuous influence of age on prevalence of UUI. Thresholds in older age seem more likely with risk being similar across wide ranges of younger ages. We must also note that such effects may not be results of age per se but of comorbidities and medication use that change with age.

Other predictors. Other factors noted across studies were the influence of race and ethnicity in United States populations. Three studies (one of high quality and two of fair quality), several with more than one related publication, documented statistical association showing black women were more likely than white to have urge urinary incontinence (while noting higher risk of mixed and stress among white women compared to black).^{11, 17, 25, 197} Another high quality United States study found no difference with adjusted estimates of prevalence of 3.5 percent among black women and 3.6 percent among white women.¹⁰

Incidence of Overactive Bladder and Urge Urinary Incontinence

Ten studies provide incidence data. Three investigated the occurrence of new OAB symptoms over varied periods of time.^{22, 26, 39} Seven investigated the onset of urge urinary incontinence.^{15, 17, 49, 64, 74, 77, 81} Two provide information about resolution of symptoms among those with symptoms at baseline.^{64, 77}

Estimates for annual incidence of OAB ranged from 2.6 to 143 cases per thousand. The lowest estimates came from a population-based estimate using the national health services database of the United Kingdom and including all adults; the highest is from the top age bracket (≥ 80 year) in another UK study.³⁹ In that study, by decade beginning at 40, the annual incidence was 78, 65, 100, 117 and 143 per 1,000; the remaining report had participants with a mean of 59 years and an estimate of 54 new cases per 1,000 women per year.³⁹

The largest study to address urge urinary incontinence was conducted in the United States within the Nurses Health Study. They report followup from 64,650 respondents with two-year data. In each five year age bracket from 35 to 55 years of age, the annual incidence among those without symptoms at baseline was 4 per 1,000 with the exception of the 46 to 50 category at 5

per 1,000.¹⁵ A separate analysis within the Nurses Health Study, using supplemental data collection to gather additional details, reported two-year incidence of urge incontinence of 14 per 1,000, which annualized to approximately 7 new cases per 1,000 women per year.^{69, 74} A six state urban sample that recruited 16,065 women and followed a subset of 3,032 for five years, reported 5-year incidence of urge incontinence of 15.9 percent or 3 cases per thousand per year.¹⁷ Work in Finland in a much smaller population sample of adults who were in their 50s and 60s at baseline and followed into 60s and 70s report the equivalent of 1.1 case per 1,000 per year, and if mortality is taken into account, 1.7 cases per year. A single study in Southern Australia, reports estimates that are meaningfully higher than these. Defining urge incontinence as that which occurs at least occasionally, and without providing a specific definition of how urge was queried, they report annual incidence of 226 cases per thousand. Of note the entire study population was 70 or older and no information about adjustment for competing morbidity is provided.⁴⁹

A study of 2,025 women older than 65 who lived in rural Iowa provides additional information. This interview-based study was conducted prior to ICS consensus definitions, the research team used the index item “How often do you have difficulty holding your urine until you can get to a toilet?”, classifying those who answered “never” and “hardly ever” as free of urge incontinence. Estimated 3-year incidence of urge incontinence, over two rounds of followup was 20.4 and 24 percent, which annualizes to 6.8 to 8.0 per 1,000 among older women.⁶⁴

The authors also report on remission: among those with UUI, 31.7 to 34.9 percent had resolution of their symptoms over the 3-year followup windows.⁶⁴ Other research in a single United States county, among women 60 and older, found annual incidence of 17 per 1000, with 22.7 percent of women with urge incontinence symptoms reporting resolution within a year.⁷⁷

KQ2: Outcomes of Treatment of OAB

This section presents results of our literature search and findings about outcomes of pharmacologic treatments for OAB. We review the basic mechanism of action of the family of medications classified as antimuscarinic agents. Six specific agents are available to United States practitioners: oxybutynin, tolterodine, fesoterodine, solifenacin, darifenacin, and trospium. Because a number of the studies are dose ranging and safety studies, we have included doses and preparations that may not be available. We also summarize studies of estrogen treatment for OAB. Each pharmacologic agent is presented individually with a thorough description of the content of the literature followed by the findings from trials and information from cohorts and case series, when those studies provided additional information beyond that provided by trials. Side effect and harms of treatment are reviewed together at the end of this section.

Pharmacologic Treatments

Pharmacologic treatments for OAB include antimuscarinic agents which have differing affinities for multiple subtypes of muscarinic receptors found both in the bladder as well as throughout the body. These agents generally bind to muscarinic receptors on the bladder muscle blocking the input required for contraction of the muscle. In short, such drugs prevent or decrease the intensity of involuntary bladder contractions.

Because muscarinic receptors are present throughout the body, nonselective agents affect other processes explaining the occurrence of side effects such as dry mouth (reduced action on salivation), constipation (slowing gut contractions), dry eyes (affecting tear ducts), and altered cognition (affecting central nervous system). In an attempt to decrease the effect on other organs

and improve tolerability, the development of agents purported to be more specific in targeting subtypes of muscarinic receptors found in the bladder has been undertaken.

Medication is often initiated with the lowest dose of an agent and adjusted to the desired clinical effect while minimizing adverse effects. The two most frequently prescribed pharmacologic treatments for OAB in the United States are oxybutynin and tolterodine. Oxybutynin is available in an oral immediate-release (IR) and an extended-release (ER) form, a transdermal patch, and a topical gel, approved by the FDA in January 2009. Tolterodine is also available in both immediate and extended-release pills. Fesoterodine was approved by the FDA in October 2008. Fesoterodine is a first order metabolite of tolterodine with similar selectivity. Newer agents available in the United States include solifenacin, darifenacin, and trospium, which is available in both immediate and extended-release formulations.

We reviewed 110 studies,^{82-96, 98-109, 140-172, 174, 198-255} of which four were good quality, 75 fair and 31 poor. This section of the report presents summary data for each pharmacologic treatment, including varied doses, intervals, and delivery mechanism. Direct comparisons across or among types of treatment (including pharmacologic, behavioral and procedural, and others) are presented in KQ3. Additional information is also provided in that section about comparisons across extended release versus immediate release. Detailed information on all studies relating to pharmacologic treatment of OAB can be found in evidence tables in Appendix C.

The summary tables included here compile unique arms of randomized trials that did not allow dose adjustment within the arm. Each drug and dose combination, as well as the related placebo arms, is presented here if data for the outcome in the summary table was provided in the publication. When necessary, weekly or monthly measures were converted to daily in order to allow ready assessment of the baseline similarity, outcomes, and effect size across studies. Each of these pharmacologic agents has been shown to be statistically superior to placebo for some facet of OAB symptoms. Complete details of measurement approach and statistical comparisons are available in Appendix C; comparisons among doses of the same drug are discussed in this section, and direct comparisons between or among drugs are presented with KQ3.

Treatment with oxybutynin.

Content of the literature. We identified 13 randomized controlled studies of oxybutynin for treatment of OAB. A total of 22 study arms compared oxybutynin at various doses and intervals, and included five placebo arms (Tables 8 and 9). Most participants were recruited from non-primary care populations with seven studies performed in the United States,⁸²⁻⁸⁸ three in Europe,⁸⁹⁻⁹¹ and one each in Japan,⁹² Taiwan,⁹³ and South Korea.⁹⁴ These studies included a total of 2,575 women in treatment arms, and 383 women in the placebo arms. Participants had an average age of 59.3 and 60.9, in the treatment and placebo groups, respectively.

Outcomes of treatment. Baseline numbers of incontinence episodes per day in the oxybutynin treatment groups ranged from 1.0 to 7.2, and in the placebo arms from 0.0 to 3.0. Because many trials were drug to drug comparisons, this difference in range does not reflect marked differences between placebo and comparison groups within studies, rather the higher severity of symptoms in the drug to drug comparison studies which were more likely to include more individuals with multiple urge incontinence episodes per day. Study participants using oral oxybutynin achieved a reduction in mean episodes of urge urinary incontinence of between 0.0 and 4.6 over the course of placebo-controlled studies that ranged in followup from 2 to 52 weeks. Of the formulations studied, 10 mg per day appeared to achieve the greatest reduction with a range of 2.9 to 7.2

(Tables 8 and 9). However, all estimates in this section of the report should be assessed with reference to the placebo arms, in which participants also reported reductions in incontinence

episodes. In oxybutynin trials, in the placebo arms, reduction in UUI episodes ranged from 0 to 1.4.

Oxybutynin was also associated with a decrease in voids per day in both oral and transdermal formulations. Baseline voids per day in drug and placebo arms were similar (7.2 to 11.7 overall). Voids per day were lowered by 1.8 to 2.3 in the transdermal arms, 0.7 to 4.2 in oral oxybutynin IR, 3.4 to 4.1 in oral oxybutynin ER and 0.8 to 1.7 in the placebo arms.

Meta-analysis was possible for immediate release but not extended release. Immediate release reduced UUI by 1.49 episodes per day (95 percent CI: 1.18, 1.80); and voids by 2.18 episodes per day (95 percent CI: 1.75, 2.61).

Table 8. RCT arms for oxybutynin chloride effect on urge incontinence

Author Year	N (% Women)	Mean age	Episodes per day baseline	Episodes per day on treatment	Decrease in episodes per day	Weeks treated
Oxybutynin 10 mg once a day						
Sand et al. ⁸³ 2004	147 (100.0)	58.4	3.6	0.9	2.7	12
Diokno et al. ⁸⁴ 2003	339 (100.0)	60.0	5.3	1.5	3.8	12
Oxybutynin 5 mg twice a day						
Halaska et al. ⁹⁰ 2003	66 (87.0)	52.2	2.1	1.1	1.0	52
Lee et al. ⁹⁴ 2002	90 (79.0)	52.0	2.4	1.0	1.4	8
Oxybutynin 5 mg three times a day						
Zinner et al. ⁸² 2005	13 (93.4)	59.9	2.9	1.4	1.6	2
Abrams et al. ⁹¹ 1998	117 (74.5)	58.0	2.6	0.9	1.7	12
Oxybutynin 2.5 mg two or three times a day						
Wang et al. ⁹³ 2006	23 (NR)	NR	1.0	1.0	0.0	12
Davila et al. ⁸⁷ 2001	36 (97.0)	63.0	7.2	2.6	4.6	6
Transdermal Oxybutynin 3.9 mg (or cm ²)						
Homma and Koyama ⁹² 2006	164 (81)	62.7	2.9	0.9	1.0	8
Transdermal Oxybutynin 3.9 mg (or cm ²)						
Dmochowski et al. ⁸⁸ 2003	121 (90.1)	63.1	4.7	1.9	2.8	12

Table 8. RCT arms for oxybutynin chloride effect on urge incontinence (continued)

Author Year	N (% Women)	Mean age	Episodes per day baseline	Episodes per day on treatment	Decrease in episodes per day	Weeks treated
Oral Placebo						
Wang et al. ⁹³ 2006	23 (NR)	NR	0.0	0.0	0.0	12
Abrams et al. ⁹¹ 1998	56 (75.4)	58.0	3.3	2.4	0.9	12
Transdermal Placebo						
Homma et al. ⁹² 2006	161 (72.7)	62.9	3.0	1.6	1.4	8

Table 9. RCT arms for oxybutynin chloride effect on voids per day

Author Year	N (% Women)	Mean age	Voids per day baseline	Voids per day on treatment	Decrease in voids per day	Weeks treated
Oxybutynin 10 mg once a day						
Sand et al. ⁸³ 2004	147 (100.0)	58.4	13.1	9.7	3.4	12
Diokno et al. ⁸⁴ 2003	339 (100.0)	60.0	13.6	9.5	4.1	12
Oxybutynin 5 mg twice a day						
Halaska et al. ⁹⁰ 2003	66 (87.0)	52.2	12.5	8.3	4.2	52
Lee et al. ⁹⁴ 2002	90 (79.0)	52.2	12.4	10.6	1.8	8
Oxybutynin 5 mg three times a day						
Abrams et al. ⁹¹ 1998	117 (74.5)	58.0	10.7	8.4	2.3	12
Giannitsas et al. ⁸⁹ 2004	6 (100.0)	53.0	7.2	6.1	1.1	6
Giannitsas et al. ⁸⁹ 2004	25 (100.0)	57.0	8.0	7.3	0.7	6
Giannitsas et al. ⁸⁹ 2004	36 (100.0)	57.0	8.3	7.5	0.8	6
Giannitsas et al. ⁸⁹ 2004	40 (100.0)	54.0	9.3	8.3	1.0	6
Zinner et al. ⁸² 2005	13 (93.4)	59.9	10.4	9.2	1.2	2
Transdermal Oxybutynin 3.9 mg patch						
Dmochowski et al. ⁸⁸ 2003	121 (90.1)	63.1	12.4	10.4	2.0	12
Transdermal Oxybutynin 3.9 mg patch						
Dmochowski et al. ⁸⁵ 2002	123 (94.4)	59.4	12.3	10.0	2.3	12

Table 9. RCT arms for oxybutynin chloride effect on voids per day (continued)

Author Year	N (% Women)	Mean age	Voids per day baseline	Voids per day on treatment	Decrease in voids per day	Weeks treated
Oral Placebo						
Abrams et al. ⁹¹ 1998	56 (75.4)	58.0	11.7	10.1	1.6	12
Zinner et al. ⁸² 2005	15 (93.4)	59.9	10.4	9.6	0.8	2
Transdermal Placebo						
Dmochowski et al. ⁸⁵ 2002	130 (89.4)	62.7	NR	NR	1.7	12

Quality of life outcomes in these studies were measured with several validated tools, including the Kings Health Questionnaire, Incontinence Inventory Questionnaire (IIQ), and Urinary Distress Inventory (UDI) (Table 18). In all studies, statistically significant improvements were observed relative to baseline, and in drug groups compared to placebo.^{85, 86, 88, 92, 93} The MATRIX study assessed HRQoL in patients treated with transdermal oxybutynin.^{245, 246} This patient population included 2,508 women with an average age of 62.5 years, and followed patients for a minimum of six months. Sand and colleagues used the KHQ and the PPBC to assess general quality of life improvement, and reported improvement in nine of ten domains versus baseline ($p < 0.001$).²⁴⁵ A second paper by Sand and colleagues employed the Beck-Depression Inventory (BDI-II) and the KHQ, and demonstrated improvement in embarrassment scores, effect on sex life, and relationships with partners, all $p < 0.001$. The BDI-II also showed improvement in interest in sex from baseline to study end ($p < 0.001$).²⁴⁶

Treatment with tolterodine.

Content of the literature. We included 20 RCTs^{83, 84, 86, 88, 89, 91, 94, 96, 97, 99, 140, 141, 154, 162, 168, 209, 223, 233, 240, 241} and 11 prospective cohort studies^{157, 160, 166, 169, 172, 212, 224, 227, 232, 239, 243} of tolterodine. Among the RCTs were 30 tolterodine arms and 14 placebo arms (Tables 10 and 11). Most were multinational studies conducted at centers in Europe, the United States, Australia, and Asia. A total of 6,746 women were in the treatment arms, with 3,298 women in the placebo arms. The average ages were 58.1 and 59.9, respectively.

Outcomes. At baseline, women reported between 1.6 and 5.0 episodes of urge urinary incontinence per day. Treatment with 4 mg of drug once a day reduced episodes of urge incontinence an average of 0.9 to 3.7 episodes each day, compared to a reduction of 1.3 to 2.4 for 2 mg taken twice a day. These doses and intervals are common in clinical practice; other doses and intervals also are included in Tables 10 and 11. Overall, reductions in urge urinary incontinence episodes on active therapy ranged from 0.9 to 3.7, compared to reductions in the placebo arms of 0.6 to 2.1.

With respect to frequency of voiding, tolterodine was associated with 0.9 to 3.6 fewer voids per day in study populations that had baseline voiding frequencies of 7.2 to 13.7. The range of response for placebo was 0.4 to 2.2 fewer voids per day, in study populations that had baseline voiding frequencies of 10.3 to 12.3 per day. Reductions differed by treatment dosage and timing, with a reduction of 1.7 to 3.6 with 4 mg once daily, 0.9 to 1.1 for 4 mg twice daily, and 1.4 to 3.3 for 2 mg twice daily.

Estimates were developed in the meta-analysis for decreases in UII episodes and voids per day in tolterodine immediate release and tolterodine extended release formulations. Tolterodine immediate release reduced UII by 1.45 episodes per day (95 percent CI: 1.24, 1.66) and voids

per day by 2.19 (95 percent CI: 1.76, 2.61). Tolterodine extended release reduced UUI by 1.75 episodes per day (95 percent CI: 1.65, 1.85) and voids per day by 2.48 (95 percent CI: 1.94, 3.02). The aggregate effect of placebo across all available study arms was a decrease in UUI episodes per day of 1.08 (95 percent CI: 0.86, 1.30) and voids by 1.48 (95 percent CI: 1.19, 1.71) per day.

Two RCTs included urodynamic parameters to assess intermediate effects of tolterodine treatment.^{221, 238} In a dose-finding study, tolterodine 2 mg twice a day demonstrated statistically significant increase in the volume at first contraction (p=0.03), and an increase in maximal cystometric capacity following four weeks of treatment compared to baseline urodynamic parameters.²²¹ In contrast, mean volume at first contraction was significantly increased after two weeks of treatment compared to baseline (p=0.046), but no improved outcome effect was seen for maximum cystometric capacity, bladder compliance, or number of detrusor contractions.²³⁸

Table 10. RCT arms for tolterodine tartrate effect on urge incontinence

Author Year	N (% Women)	Mean age	Episodes per day baseline	Episodes per day on treatment	Decrease in episodes per day	Weeks treated
Tolterodine 4 mg once a day						
Rogers et al. ²⁴¹ 2008	182 (100)	49.0	2.5	NR	1.8	12
Chapple et al. ⁹⁹ 2007	297 (87.1)	56.9	NR	NR	0.9	4
Chapple et al. ⁹⁶ 2007	253 (78.0)	57.7	3.8	2.1	1.7	12
Robinson et al. ²⁴⁰ 2007	53 (100)	NR	2.3	0.7	1.7	8
Landis et al. ¹⁶⁸ 2004	321 (81.6)	60.9	1.6	0.5	1.1	12
Landis et al. ¹⁶⁸ 2004	284 (83.6)	60.0	4.7	1.6	3.3	12
Millard et al. ¹⁵⁴ 2004	205 (75.4)	53.6	3.2	1.1	2.2	12
Millard et al. ¹⁵⁴ 2004	191 (75.4)	53.6	3.2	1.0	2.3	24
Diokno et al. ⁸⁴ 2003	357 (100)	60.0	5.3	1.6	3.7	12
Tolterodine 4 mg once a day						
Dmochowski et al. ⁸⁸ 2003	123 (95.1)	62.9	5.0	1.9	3.1	12
Swift et al. ¹⁴⁰ 2003	417 (100)	59.0	3.2	1.5	1.7	12
Van Kerrebroeck et al. ¹⁴¹ 2001	507 (82.0)	60.0	3.2	1.5	1.2	12

Table 10. RCT arms for tolterodine tartrate effect on urge incontinence (continued)

Tolterodine 2 mg twice a day						
Sand et al. ⁸³ 2004	161 (100)	58.8	3.6	1.2	2.4	12
Chapple et al. ⁹⁷ 2003	279 (72.9)	58.1	2.3	0.9	1.4	12
Swift et al. ¹⁴⁰ 2003	408 (100)	59.0	3.3	1.8	1.4	12
Lee et al. ⁹⁴ 2002	97 (74.0)	52.0	2.6	0.4	2.2	8
Jacquetin ¹⁶² 2001	103 (81.6)	58.0	3.2	1.9	1.3	4
Van Kerrebroeck et al. ¹⁴¹ 2001	514 (79)	60.0	3.3	1.8	1.5	12
Millard et al. ²³³ 1999	116 (77.0)	60.2	3.6	1.9	1.7	12
Abrams et al. ⁹¹ 1998	118 (77.1)	55.0	2.9	1.6	1.3	12
Tolterodine 2 mg three times a day						
Chancellor et al. ²⁰⁹ 2000	514 (80.0)	60.0	3.3	1.8	1.5	12
Tolterodine 1 mg twice a day						
Jacquetin et al. ¹⁶² 2001	97 (76.3)	53.0	3.7	2.6	1.1	4
Millard et al. ²³³ 1999	114 (78.0)	60.1	3.9	2.2	1.7	12
Placebo						
Chapple et al. ⁹⁶ 2007	252 (81.0)	56.0	3.7	2.6	1.1	12
Landis et al. ¹⁶⁸ 2004	171 (78.5)	60.6	1.6	1.0	0.6	12
Landis et al. ¹⁶⁸ 2004	210 (84.8)	61.8	4.5	3.2	1.3	12

Table 10. RCT arms for tolterodine tartrate effect on urge incontinence (continued)

Author Year	N (% Women)	Mean age	Episodes per day baseline	Episodes per day on treatment	Decrease in episodes per day	Weeks treated
Placebo						
Rogers et al. ²⁴¹ 2008	189 (100)	47.0	2.2	NR	1.4	12
Robinson et al. ²⁴⁰ 2007	59 (100)	NR	2.9	2.2	0.7	8
Chapple et al. ⁹⁷ 2003	267 (76.3)	57.7	2.0	1.4	0.6	12
Dmochowski et al. ⁸⁸ 2003	117 (93.2)	64.5	5.0	2.9	2.1	12
Swift et al. ¹⁴⁰ 2003	410 (100)	60.0	3.4	2.4	1.0	12
Van Kerrebroeck et al. ¹⁴¹ 2001	507 (81.0)	61.0	3.3	2.3	1.0	12

Table 11. RCT arms for tolterodine tartrate effect on voids per day

Author Year	N (% Women)	Mean age	Voids per day baseline	Voids per day on treatment	Decrease in voids per day	Weeks treated
Tolterodine 4 mg once a day						
Rogers et al. ²⁴¹ 2008	182 (100)	49.0	13.0	NR	3.3	12
Chapple et al. ⁹⁶ 2007	253 (78.0)	57.7	11.5	9.8	1.7	12
Chapple et al. ⁹⁹ 2007	267 (87.1)	56.9	NR	NR	1.5	4
Robinson et al. ²⁴⁰ 2007	53 (100)	NR	13.2	10.6	2.6	8
Landis et al. ¹⁶⁸ 2004	321 (81.6)	60.9	9.6	NR	1.2	12
Landis et al. ¹⁶⁸ 2004	171 (78.5)	60.6	10.3	NR	0.9	12
Millard et al. ¹⁵⁴ 2004	191 (75.4)	53.6	12.8	9.2	3.6	24
Millard et al. ¹⁵⁴ 2004	205 (75.4)	53.6	12.8	9.4	3.4	12
Diokno et al. ⁸⁴ 2003	357 (100)	60.0	13.7	10.2	3.6	12
Dmochowski et al. ⁸⁸ 2003	123 (95.1)	62.9	12.1	9.9	2.2	12

Table 11. RCT arms for tolterodine tartrate effect on voids per day (continued)

Author Year	N (% Women)	Mean age	Voids per day baseline	Voids per day on treatment	Decrease in voids per day	Weeks treated
Tolterodine 4 mg once a day						
Swift et al. ¹⁴⁰ 2003	417 (100)	59.0	10.8	9.0	1.8	12
Van Kerrebroeck et al. ¹⁴¹ 2001	507 (82.2)	60.0	10.9	7.4	3.5	12
Tolterodine 4 mg twice a day						
Giannitsas et al. ⁸⁹ 2004	6 (100)	53.0	7.2	6.3	0.9	6
Giannitsas et al. ⁸⁹ 2004	25 (100)	57.0	8.0	7.0	1.0	6
Giannitsas et al. ⁸⁹ 2004	36 (100)	57.0	8.3	7.2	1.1	6
Giannitsas et al. ⁸⁹ 2004	40 (100)	54.0	8.5	7.6	0.9	6
Tolterodine 2 mg twice a day						
Sand et al. ⁸³ 2004	161 (100)	58.8	13.1	10.2	2.9	12
Chapple et al. ⁹⁷ 2003	279 (72.9)	58.1	12.1	9.9	2.2	12
Swift et al. ¹⁴⁰ 2003	408 (100)	59.0	11.1	9.3	1.8	12
Lee et al. ⁹⁴ 2002	97 (74.0)	52.0	12.2	9.6	2.6	8
Tolterodine 2 mg twice a day (continued)						
Jacquetin et al. ¹⁶² 2001	103 (81.6)	58.0	10.8	9.4	1.4	4
Van Kerrebroeck et al. ¹⁴¹ 2001	514 (79)	60.0	11.1	7.8	3.3	12
Millard et al. ²³³ 1999	116 (77.0)	60.2	11.2	8.9	2.3	12
Abrams et al. ⁹¹ 1998	118 (77.1)	55.0	11.5	8.8	2.7	12
Tolterodine 2 mg three times a day						
Chancellor et al. ²⁰⁹ 2000	514 (80.0)	60.0	11.1	9.4	1.7	12
Tolterodine 1 mg twice a day						
Jacquetin et al. ¹⁶² 2001	97 (76.3)	53.0	10.7	9.3	1.4	4
Millard et al. ²³³ 1999	114 (78.0)	60.1	11.5	9.2	2.3	12

Table 11. RCT arms for tolterodine tartrate effect on voids per day (continued)

Author Year	N (% Women)	Mean age	Voids per day baseline	Voids per day on treatment	Decrease in voids per day	Weeks treated
Placebo						
Rogers et al. ²⁴¹ 2008	189 (100)	47.0	12.5	NR	2.3	12
Chapple et al. ⁹⁶ 2007	252 (81.0)	56.0	11.5	11.1	1.0	12
Robinson et al. ²⁴⁰ 2007	59 (100)	NR	11.9	10.1	1.8	8
Landis et al. ¹⁶⁸ 2004	210 (84.8)	61.8	11.1	NR	0.4	12
Landis et al. ¹⁶⁸ 2004	284 (83.6)	60.0	10.7	NR	1.9	12
Chapple et al. ⁹⁷ 2003	267 (76.3)	57.7	12.2	11.0	1.2	12
Dmochowski et al. ⁸⁸ 2003	117 (93.2)	64.5	12.3	10.9	1.4	12
Swift et al. ¹⁴⁰ 2003	410 (100)	60.0	11.2	9.9	1.3	12
Van Kerrebroeck et al. ¹⁴¹ 2001	507 (81.0)	61.0	11.3	9.1	2.2	12

Findings from cohort studies and case series were compatible with these results. Twelve such studies demonstrated improvement in urinary urgency, frequency, and urge incontinence with tolterodine treatment.^{142, 157, 160, 166, 169, 172, 212, 224, 227, 232, 239, 243, 247}

Twelve studies including RCTs, prospective cohorts and extension studies evaluated the effect of tolterodine on patient reported outcomes, including quality of life in at least one arm.^{86, 88, 91, 99, 142, 158, 160, 174, 223, 240, 241, 247} Participants in tolterodine arms consistently reported greater changes in quality of life, including on various domains of the Kings Health Questionnaire, IIQ, and UDI-6 when compared to placebo that were similar to those experienced by participants of oxybutynin arms in the studies.

One study evaluated the impact of treatment with tolterodine ER 4 mg once daily on emotional and sexual health. Sexual Quality of Life Questionnaire- Female (SQoL-F), Pelvic Organ Prolapse/Urinary Incontinence Sexual Questionnaire (PISQ), and the Hospital Anxiety and Depression scale (HAD) were used to assess patient reported outcomes. Treatment with tolterodine resulted in improved scores versus placebo in SQoL-F; PISQ- total score, and HAD anxiety (p=0.004, p=0.009, p=0.03 respectively).²⁴¹

One study stratified outcomes by treatment in patients with prolapse and found no significant difference in changes in quality of life between those with anterior vaginal wall prolapse and those without.²⁴³ Another study stratified patients by those with urge incontinence and those with OAB syndrome without incontinence, and found similar bother and severity scores in each group in an open label study of tolterodine ER.¹⁵⁹

Most outcomes were equivalent in studies comparing tolterodine IR versus ER; however, in two RCTs, extended release tolterodine resulted in a greater reduction in incontinence episodes per day, p=0.036 and p=0.05, respectively.^{140, 141} Other outcome parameters including voids per

day and pads per day showed a similar reduction compared to placebo with no statistical difference seen between the two formulations.

Treatment with fesoterodine.

Content of the literature. Two randomized controlled trials compared fesoterodine at 4 and 8 mg to placebo for reducing symptoms of OAB in this literature that met criteria for inclusion in the systematic review (Table 12).^{95,96} These studies included a total of 1,017 women with a mean age of 57.7 in the treatment arms, and a total of 518 women with a mean age of 57.5 in the placebo arms. Both studies reported outcome data at 12 weeks of treatment. One study included a tolterodine 4 mg arm, discussed in KQ3.⁹⁶ There is one paper included which is a post hoc analysis of this study population⁹⁶ assessing treatment effect on health-related quality of life.¹⁴²

Outcomes. At baseline, women in the treatment arms reported between 2.2 and 3.2 episodes of urge urinary incontinence per day. Treatment with drug resulted in reductions of 1.95 and 3.2 episodes per day. In the placebo arms at baseline, participants had an average of 3.7 episodes of urge incontinence per day. UUI in the placebo group was reduced by 1.14 to 1.48 episodes per day after 12 weeks of treatment.^{95,96}

At baseline, women in the treatment arms reported between 11.6 and 12.9 voids per day. Treatment with drug resulted in reductions of 1.4 to 1.9 voids per day. Women in the placebo arms reported between 12.0 to 12.2 voids per day at baseline, and reported a reduction of voids per day ranging from 0.7 to 0.95.^{95,96}

In the meta-analysis, fesoterodine decreased UUI per day by 2.03 (95 percent CI: 1.74, 2.31) episodes and voids per day by 1.84 (95 percent CI: 1.64, 2.03) episodes. placebo reduces UUI episodes by 1.08 (95 percent CI: 0.86, 1.30), and voids by 1.48 (95 percent CI: 1.19, 1.71) per day. The aggregate effect of placebo across all available study arms was a decrease in UUI episodes per day of 1.08 (95 percent CI: 0.86, 1.30) and voids by 1.48 (95 percent CI: 1.19, 1.71) per day.

The post hoc analysis of fesoterodine arms showed significant improvement in HRQoL as assessed by KHQ and ICIQ-SF in both fesoterodine 8 mg arm and the tolterodine ER 4 mg arm versus placebo, but there was no significant difference between arms. The fesoterodine arm showed significant improvement in eight of nine domains (non significant in General Health domain). A subset of patients who reported incontinence at baseline showed similar improvements versus baseline; however, no significant difference was seen between arms as well as between subsets of those reporting incontinence at baseline versus those without.¹⁴²

Table 12. RCT arms for fesoterodine fumarate effect on urge incontinence and voids

Author Year	N (% Women)	Mean age	Episodes per day baseline	Episodes per day on treatment	Decrease in episodes per day	Weeks treated
Episodes of urge incontinence						
Fesoterodine 8 mg once a day						
Chapple et al. ⁹⁶ 2007	252 (81.0)	55.6	3.7	1.5	2.2	12
Nitti et al. ⁹⁵ 2007	267 (78.0)	59.0	3.9	0.7	3.2	12
Fesoterodine 4 mg once a day						
Chapple et al. ⁹⁶ 2007	231 (82.0)	57.1	3.8	1.9	2.0	12
Nitti et al. ⁹⁵ 2007	267 (76.0)	59.0	3.9	1.3	2.6	12
Voids per day						
Fesoterodine 8 mg once a day						
Chapple et al. ⁹⁶ 2007	252 (81.0)	55.6	11.9	10.0	1.9	12
Nitti et al. ⁹⁵ 2007	267 (78.0)	59.0	12.0	10.1	1.9	12
Fesoterodine 4 mg once a day						
Chapple, et al. ⁹⁶ 2007	231 (82.0)	57.1	11.6	9.8	1.8	12
Nitti et al. ⁹⁵ 2007	267 (76.0)	59.0	12.9	11.5	1.4	12

Treatment with solifenacin.

Content of the literature. Three RCTs investigated solifenacin compared to placebo for reducing symptoms of OAB (Table 13).⁹⁷⁻⁹⁹ These studies included a total of 1,541 women, with a mean age of 58.2 years in the solifenacin treatment arms, and a total of 638 women with a mean age of 59.3 in the placebo arms. Two were conducted at multiple centers across Europe, and the other in Japan at academic health centers. One of these trials compared solifenacin to tolterodine, without a placebo arm, and is described in KQ3.^{99, 211} A newly published RCT performed at multiple centers in the United States comparing solifenacin at various doses to placebo over twelve weeks focused on outcomes of urgency and “warning time” showing significant decrease in urgency episodes per day ($p < 0.001$), and an increase of 31.5 seconds in median warning time ($p = 0.008$).²²² Voids per day and UUI episodes per day were significantly reduced compared to placebo (both $p < 0.001$).²²²

Outcomes. The first study compared solifenacin 5 mg and 10 mg daily to placebo and propiverine 10 mg.⁹⁸ Women in the two solifenacin arms (713 women with a mean age of 60.2) had an average decrease in the number of UUI episodes per of 1.45 and 1.52 at 12 weeks compared to a reduction in the placebo arm of 0.89 ($p < 0.001$ for comparisons of both dosages to placebo).⁹⁸ Participants had a significant decrease in the number of voids per day of 1.9 to 2.9 ($p < 0.001$ for both doses)

The second study also studied 5 mg and 10 mg doses, compared to placebo. Women in the two solifenacin arms, 5 mg and 10 mg, (531 women with a mean age of 57.1) experienced reductions in number of UUI episodes per day of 1.4 and 0.9, respectively at 12 weeks. The placebo arm (267 women with a mean age of 57.7) demonstrated a reduction in urge urinary incontinence episodes per day of 0.6.⁹⁷

Solifenacin was associated with a decrease of 1.46 episodes of UUI per day (95 percent CI: 1.32, 1.59) and of 2.19 voids per day (95 percent CI: 1.94, 3.02) in the meta-analysis.

Quality of life was assessed in four studies of solifenacin.^{98, 99, 218, 225} Quality of life measures improved significantly in the solifenacin arms²¹⁸ including when compared to placebo.^{98, 225} Significant improvement in quality of life was maintained upon sub-analysis for African Americans and Hispanics.^{205, 229} Perception of bladder condition was better in one solifenacin arm that was compared to tolterodine⁹⁹ (p=0.006).

Two prospective case series with multiple publications evaluated solifenacin in the treatment of OAB.^{205, 218, 220, 229} One study included 1,280 women with a mean age of 56.4 years.²²⁰ Participants had a 66 percent reduction in incontinence episodes per day (from 2.66 to 0.93), a 23 percent reduction in voids per day (12.16 to 9.18), a 63 percent reduction in urgency episodes per day (5.76 to 2.28), and a 32 percent reduction in nocturia episodes (1.95 to 1.25) as assessed by urinary diary at one-year followup.

A dose ranging study evaluating treatment with solifenacin 5 and 10 mg daily after washout from tolterodine 4 mg ER, darifenacin or trospium found significant improvement in UUI episodes per day and voids per day over a 12 week treatment period after 14 day washout period (p<0.001). Patient reported outcomes assessed by the PPBC and OAB-q questionnaires indicated significant subjective improvement (p<0.001 in all domains).²⁰⁸

Another prospective randomized trial compared Solifenacin 5 and 10 mg daily to placebo. This study used the PPIUS (Patient Perception of Intensity of Urgency Scale) to assess treatment effect on urgency, and demonstrated that Solifenacin at the two doses demonstrated significant improvement in severe urgency episodes over placebo (p<0.001). In addition, patient reported outcomes of urgency bother, patient PBC score, and treatment satisfaction were significantly reduced compared to placebo (p<0.001).²⁰⁶

Table 13. RCT arms for solifenacin succinate effect on urge incontinence and voids

Author Year	N (% Women)	Mean age	Episodes per day baseline	Episodes per day on treatment	Decrease in episodes per day	Weeks treated
Episodes of urge incontinence						
Solifenacin 10 mg once a day						
Yamaguchi et al. ⁹⁸ 2007	349 (85.7)	59.9	1.9	0.4	1.5	12

Table 13. RCT arms for solifenacin succinate effect on urge incontinence and voids (continued)

Author Year	N (% Women)	Mean age	Episodes per day baseline	Episodes per day on treatment	Decrease in episodes per day	Weeks treated
Episodes of urge incontinence						
Solifenacin 10 mg once a day						
Chapple et al. ⁹⁷ 2003	263 (80.0)	56.9	1.9	1.0	0.9	12
Solifenacin 5 mg once a day						
Yamaguchi et al. ⁹⁸ 2007	364 (83.0)	60.4	2.0	0.4	1.5	12
Chapple et al. ⁹⁹ 2007	297 (87.5)	56.6	NR	NR	NR	4
Chapple et al. ⁹⁷ 2003	268 (71.2)	57.2	2.1	0.8	1.4	12
Placebo						
Yamaguchi et al. ⁹⁸ 2007	371 (84.3)	60.8	1.7	0.8	0.9	12
Chapple et al. ⁹⁷ 2003	267 (76.3)	57.7	2.0	1.4	0.6	12
Voids per day						
Solifenacin 10 mg once a day						
Chapple et al. ⁹⁷ 2003	263 (80.0)	56.9	12.1	10.2	1.9	12
Solifenacin 5 mg once a day						
Chapple et al. ⁹⁹ 2007	297 (87.5)	56.6	NR	NR	1.7	4
Chapple et al. ⁹⁷ 2003	268 (71.2)	57.2	12.3	9.7	2.6	12
Placebo						
Chapple et al. ⁹⁷ 2003	267 (76.3)	57.7	12.2	11.0	1.2	12

Treatment with darifenacin.

Content of the literature. Five RCTs provided data on the effectiveness of darifenacin.^{82, 100-102, 210} Three compared 15 and 30 mg daily relative to placebo (Tables 14 and 15).^{82, 100, 101} These three studies included a total of 690 women, with a mean age of 57.6 years in the darifenacin treatment arms and a total of 304 women with a mean age of 57.6 in the placebo arms. Two of the three were conducted in the United States in a non-primary care population.^{82, 101} The other was performed at multiple centers in Europe including community-based ambulatory populations, and also included a 7.5 mg arm.¹⁰⁰ The fourth study compared only darifenacin 7.5 mg to placebo and included 269 women with mean age of 57.5 in the treatment arm, and 129 women with an average of 58.5 in the placebo arm. This was a dose change study without report of outcomes by dosage.¹⁰² A fifth RCT also compared darifenacin 7.5 mg to placebo in a multinational, multicenter trial. This was a dose changing study without reporting of outcome by dosage. The treatment arm included 206 women with an average age of 72 years and a placebo

arm including 100 women with an average age of 73 years.²¹⁰ One of these studies⁸² also compared darifenacin to oxybutynin; the darifenacin arm is presented in Tables 14 and 15, but the comparison is described in the chapter on KQ3.

Outcomes. Baseline urge urinary incontinence episodes per day in the darifenacin arms ranged from 2.0 to 2.9. Urge urinary incontinence episodes per day on treatment ranged from 0.8 to 1.6, for a decrease in episodes of UUI per day ranging from 1.2 to 1.8. This is compared to an average reduction in the placebo groups of 0.8 to 1.0, with similar baseline measures of 2.3 to 3.0.

At baseline, participants reported an average of 10.3 to 11.0 voids per day. The treatment groups reported reductions ranging from 1.1 to 2.2 per day, compared to 0.8 to 1.8 in the placebo group. Study followup ranged from 2 to 12 weeks. Daily episodes of the symptom of urinary urgency also were reported to be reduced after treatment with darifenacin (range of reduction: 1.4 to 3.0 compared to 0.6 to 1.2 in placebo). The one study that reported reduction in urgency on a weekly basis, however, found no significant difference between treatment and placebo.¹⁰¹

Because of a lack of uniform information provided in the studies on variance measures, we could not use meta-analysis techniques to estimate effects of darifenacin on UUI or voids per day.

In the first dose-adjustment trial, urge urinary incontinence episodes were not reported, but voiding frequency was significantly reduced in the treatment arm versus placebo, $p=0.001$.¹⁰² In the second dose changing trial changes in urge urinary incontinence episodes per week were not statistically different between the two arms ($p=0.328$), but reductions in voids per day were statistically greater in the treatment arm versus placebo ($p=0.006$). This study also assessed urgency episodes per day, but found no statistically significant difference in effect ($p=0.174$).²¹⁰

Three studies assessed quality of life (QoL) with validated questionnaires using Visual Analog Scale (VAS), OAB-q, King's Health Questionnaire (KHQ), and ICIQ. The arms using 15 and 30 mg showed statistically significant improvement versus placebo, $p=0.045$ and $p=0.011$, respectively.¹⁰⁰ Zinner evaluated 15 mg versus placebo and showed significant improvement in quality of life questionnaires, OAB-q ($p<0.001$), ICIQ ($p<0.001$) and KHQ ($p<0.05$).¹⁰¹ Chapple demonstrated improvement in all domains of the OAB-q versus placebo ($p<0.001$).²¹⁰ A single prospective cohort study evaluated the efficacy and tolerability of darifenacin.²¹⁹ This prospective, non-comparative, open-label extension study was conducted in multiple countries in Europe, Australia, and North America.¹⁰² The authors stratified results by age. All outcome parameters were similar between groups showing similar reductions in voids per day, urgency episodes per day, and severity of urgency in those over and under 65 years old.²¹⁹ A second paper describing the open labeled extension study of those patients²¹⁹ demonstrated significant improvement from baseline in eight of nine domains ($p<0.001$) in health related quality of life outcomes as measured by the KHQ at 24 months. Significant reductions were confirmed in the subgroup of subjects 65 years old and greater. The only domain that did not show improvement was the General Health Perceptions.²¹⁷

Table 14. RCT arms for darifenacin effect on urge incontinence

Author Year	N (% Women)	Mean age	Episodes per day baseline	Episodes per day on treatment	Decrease in episodes per day	Weeks treated
Darifenacin 30 mg once a day						
Hill et al. ¹⁰⁰ 2006	96 (86.0)	54.0	2.7	1.1	1.6	12
Zinner et al. ⁸² 2005	15 (93.4)	59.9	2.9	1.3	1.7	2
Darifenacin 15 mg once a day						
Hill et al. ¹⁰⁰ 2006	93 (86.0)	55.1	2.5	1.0	1.5	12
Zinner et al. ¹⁰¹ 2006	185 (86.4)	59.1	2.7	0.9	1.8	12
Zinner et al. ¹⁰¹ 2006	185 (86.4)	59.1	2.7	1.4	1.3	2
Zinner et al. ⁸² 2005	17 (93.4)	59.9	2.9	1.6	1.4	2
Darifenacin 7.5 mg once a day						
Hill et al. ¹⁰⁰ 2006	99 (87.0)	56.1	2.0	0.8	1.2	12
Placebo						
Hill et al. ¹⁰⁰ 2006	101 (83.0)	53.7	2.3	1.5	0.8	12
Zinner et al. ¹⁰¹ 2006	188 (88.0)	59.1	3.0	2.0	1.0	2
Zinner et al. ⁸² 2005	15 (93.4)	59.9	2.9	2.1	0.8	2

Table 15. RCT arms for darifenacin effect on voids per day

Author Year	N (% Women)	Mean age	Voids per day baseline	Voids per day on treatment	Decrease in voids per day	Weeks treated
Darifenacin 7.5 mg once a day						
Hill et al. ¹⁰⁰ 2006	99 (87.0)	56.1	10.3	8.6	1.7	12
Darifenacin 15 mg once a day						
Hill et al. ¹⁰⁰ 2006	93 (86.0)	55.1	11.0	9.1	1.9	12
Zinner et al. ¹⁰¹ 2006	185 (86.4)	59.1	11.0	8.8	2.2	12
Zinner et al. ⁸² 2005	17 (93.4)	59.9	10.4	9.3	1.1	2

Table 15. RCT arms for darifenacin effect on voids per day (continued)

Author Year	N (% Women)	Mean age	Voids per day baseline	Voids per day on treatment	Decrease in voids per day	Weeks treated
Darifenacin 30 mg once a day						
Hill et al. ¹⁰⁰ 2006	96 (86.0)	54.0	10.4	8.2	2.2	12
Zinner et al. ⁸² 2005	15 (93.4)	59.9	10.4	8.9	1.6	2
Placebo						
Zinner et al. ¹⁰¹ 2006	188 (88.0)	59.1	11.2	9.4	1.8	12
Hill et al. ¹⁰⁰ 2006	101 (83.0)	53.7	10.1	9.0	1.1	12
Zinner et al. ⁸² 2005	15 (93.4)	59.9	10.4	9.6	0.8	2

Treatment with trospium chloride.

Content of the literature. Five RCTs evaluated trospium for reduction of symptoms of OAB. Four trials compared trospium to placebo, and one compared trospium to oxybutynin (5 mg twice daily) (Table 16). Four were conducted in the United States,^{103, 104, 105, 106} and the fifth at multiple centers in Europe and Asia (Tables 16 and 17).⁹⁰ These studies included a total of 1,309 women, with a mean age of 59.8 years in the trospium treatment arms, and a total of 1,130 women with a mean age of 60.1 in the placebo arms.¹⁰³⁻¹⁰⁶ One trial included a comparison to oxybutynin (5 mg twice a day).⁹⁰ The trospium arm is included in Tables 16 and 17; and the drug to drug comparison is noted in KQ3.

Outcomes. Reductions in numbers of urge urinary incontinence episodes per day at 12 weeks ranged from 1.8 to 2.5 relative to reductions in the placebo arms of 1.3 to 1.9. Reduction in voids per day on treatment ranged from 2.4 to 3.5 at 12 weeks, compared to placebo at 1.3 to 2.1

Meta-analysis could be used to estimate effects of trospium extended release only. This formulation reduced UUI episodes per day by 2.45 (95 percent CI: 2.19, 2.70) and voids per day by 2.68 (95 percent CI: 2.38, 2.98) episodes. The aggregate effect of placebo across all available study arms was a decrease in UUI episodes per day of 1.08 (95 percent CI: 0.86, 1.30) and voids by 1.48 (95 percent CI: 1.19, 1.71) per day.

Three studies evaluated urgency severity using the Indevus Urgency Severity Scale (IUSS). All three studies found statistically significant improvements in severity of urgency at trial end in the trospium arms versus placebo: $p=0.0004$, $p\leq 0.001$, and $p<0.0001$ respectively.¹⁰³⁻¹⁰⁵

Table 16. RCT arms for trospium chloride effect on urge incontinence

Author Year	N (% Women)	Mean age	Episodes per day baseline	Episodes per day on treatment	Decrease in episodes per day	Weeks treated
Trospium 60 mg once a day						
Dmochowski et al. ¹⁰⁵ 2008	267 (82.1)	61.2	4.0	1.6	2.4	12
Trospium 60 mg once a day						
Staskin et al. ¹⁰³ 2007	263 (85.2)	59.6	4.1	1.6	2.5	12
Staskin et al. ¹⁰³ 2007	263 (85.2)	59.6	4.1	1.8	2.4	4
Trospium 20 mg twice a day						
Rudy et al. ^{104, 242} 2006	323 (81.8)	61.1	2.9	1.0	1.9	12
Halaska et al. ⁹⁰ 2003	200 (85.0)	54.2	1.5	0.5	1.0	52
Zinner et al. ¹⁰⁶ 2004	256 (77.9)	63.0	3.0	1.2	1.8	12
Placebo						
Dmochowski et al. ¹⁰⁵ 2008	276 (87.7)	58.4	4.0	2.4	1.6	12
Staskin et al. ¹⁰³ 2007	273 (84.5)	59.3	4.1	2.2	1.9	12
Rudy et al. ^{104, 242} 2006	325 (81.2)	61.0	2.9	1.6	1.3	12
Zinner et al. ¹⁰⁶ 2004	256 (71.6)	61.5	4.3	2.4	1.9	12
Staskin et al. ¹⁰³ 2007	273 (84.5)	59.3	4.1	2.4	1.8	4

Table 17. RCT arms for trospium chloride effect on voids per day

Author Year	N (% Women)	Mean age	Voids per day baseline	Voids per day on treatment	Decrease in voids per day	Weeks treated
Trospium 60 mg once a day						
Dmochowski et al. ¹⁰⁵ 2008	267 (82.1)	61.2	12.8	10.3	2.5	12
Staskin et al. ¹⁰³ 2007	263 (85.2)	59.6	11.9	8.8	3.1	12
Staskin et al. ¹⁰³ 2007	263 (85.2)	59.6	11.9	9.2	2.7	4

Table 17. RCT arms for trospium chloride effect on voids per day (continued)

Author Year	N (% Women)	Mean age	Voids per day baseline	Voids per day on treatment	Decrease in voids per day	Weeks treated
Trospium 20 mg twice a day						
Rudy et al. ^{104, 242} 2006	323 (81.8)	61.1	12.9	10.3	2.7	12
Trospium 20 mg twice a day						
Zinner et al. ¹⁰⁶ 2004	256 (77.9)	63.0	12.7	10.3	2.4	12
Halaska et al. ⁹⁰ 2003	200 (85.0)	54.2	11.4	7.9	3.5	52
Placebo						
Dmochowski et al. ¹⁰⁵ 2008	276 (87.7)	58.4	12.9	11.1	1.8	12
Staskin et al. ¹⁰³ 2007	273 (84.5)	59.3	11.8	9.7	2.1	12
Staskin et al. ¹⁰³ 2007	273 (84.5)	59.3	11.8	10.1	1.7	4
Rudy et al. ^{104, 242} 2006	325 (81.2)	61.0	13.2	11.4	1.8	12
Zinner et al. ¹⁰⁶ 2004	256 (71.6)	61.5	12.9	11.6	1.3	12

Quality of life was addressed in three studies using the OAB-SCS validated questionnaire¹⁰³,¹⁰⁴ and the IIQ in one.¹⁰⁶ Only one of the studies using the OAB-SCS reported statistically significant improvement in quality of life compared to placebo, $p < 0.05$.¹⁰³

Treatment with estrogen.

Content of the literature. Three studies assessed the role of hormonal therapy in different doses and formulations in the alleviation of OAB symptoms.¹⁰⁷⁻¹⁰⁹ All studies were performed in Europe and Scandinavia. They included a total of 508 women with a mean age of 62.4 years. Three were RCTs and one was a case series.

Outcomes. One RCT compared oral estriol 3 mg per day to placebo at academic centers in England. They found that estriol was not significantly superior to placebo at improving symptoms or objective measures for all patients at one or three months.¹⁰⁷

The second RCT compared an estradiol releasing vaginal ring to an estriol vaginal pessary in a community population in Denmark, and found both equal in improvement of UUI episodes, urgency, and nocturia. In each arm, 58 percent of participants had decreased UUI at 24 weeks; 51 and 56 percent had decreased urgency, respectively.¹⁰⁸ This is difficult to interpret in the absence of a placebo comparison.

A recently published RCT compared tolterodine 2 mg twice daily with and without CEE (conjugated equine estrogens) vaginal cream 0.625 mg twice weekly. Voids per day decreased significantly in the CEE arm ($p = 0.001$), but did not show significant difference in UUI events per day. Subjective assessment with UDI-6 and IIQ-7 showed significant improvement in the tolterodine with CEE arm ($p = 0.001$).²⁴⁷

One case series found oral HRT significantly improved frequency at six-month followup where dosage was not associated with outcome. Hormone replacement therapy in the form of

estriol significantly decreased frequency in the OAB arm at six weeks.¹⁰⁹ Outcomes were assessed with a five-point unvalidated rating system and no helpful comparison group.

Table 18. Effect on quality of life and satisfaction for pharmacologic treatment

Author Year	Comparison Groups, N	Outcomes
Oxybutynin		
Dmochowski et al. ⁸⁵ 2002	G1: Oxybutynin TDS 1.3 mg G2: Oxybutynin TDS 2.6 mg G3: Oxybutynin TDS 3.9 mg G4: Placebo	<ul style="list-style-type: none"> • Participants in the treatment arm had significantly improved scores on the IIQ relative to placebo (p=0.033) • UDI scores showed greater improvement for treated patients compared to those on placebo (p=0.027)
Dmochowski et al. ⁸⁸ 2003	G1: Oxybutynin TDS 3.9 mg G2: Tolterodine ER 4 mg G3: Placebo	<ul style="list-style-type: none"> • Compared to those receiving placebo, participants in both the oxybutynin and tolterodine groups had significantly greater improvement in the travel domain of the IIQ (p<0.05 for both) and on the irritative symptoms domain of the UDI (p<0.02 for both)
Homma et al. ⁹² 2006	G1: Oxybutynin TDS 26 cm ² G2: Oxybutynin TDS 39 cm ² G3: Oxybutynin TDS 52 cm ² G4: Placebo	<ul style="list-style-type: none"> • Participants in the treatment arm had statistically significant improvements across all domains of the KHQ compared to placebo (p<0.05 for all comparisons)
Sand et al. ^{245, 246} 2006, 2007	G1: Oxybutynin TDS G2: Educational intervention	<ul style="list-style-type: none"> • Participants in the oxybutynin arm showed improvement in nine of ten domains on the KHQ and the PPBC vs. baseline (p<0.001), • Participants in the oxybutynin arm also showed improvement in embarrassment scores, effect on sex life, and relationships with partners (p<0.001) on BDI-II and KHQ • Oxybutynin also improved in interest in sex from baseline on BDI-II (p<0.001)
Sussman et al. ⁸⁶ 2002	G1: Tolterodine ER 2 mg G2: Tolterodine ER 4 mg G3: Oxybutynin ER 5 mg G4: Oxybutynin ER 10 mg	<ul style="list-style-type: none"> • Patients on tolterodine ER reported greater improvements after 8 weeks of treatment in bladder condition compared to tolterodine IR, oxybutynin ER or oxybutynin IR, on a validated 6-point Likert scale (all comparisons p<0.01).
Wang et al. ⁹³ 2006	G1: Electrical Stimulation G2: Oxybutynin 2.5 mg G3: Placebo	<ul style="list-style-type: none"> • Statistically significant differences were seen on the overall score and the incontinence impact domain of the KHQ between the ES group and oxybutynin (p<0.001 and p=0.038, respectively) and between the ES group and placebo (p=0.006 and p=0.012) • The difference between oxybutynin and placebo was also significant for the overall score only (p<0.001)

Table 18. Effect on quality of life and satisfaction for pharmacologic treatment (continued)

Author Year	Comparison Groups, N	Outcomes
Tolterodine		
Abrams et al. ⁹¹ 1998	G1: Tolterodine 2 mg G2: Oxybutynin 5 mg G3: Placebo	<ul style="list-style-type: none"> • After 12 weeks, 50% of patients on tolterodine, 49% on oxybutynin and 47% on placebo reported a perceived improvement in symptoms
Chapple et al. ¹⁴² 2008	G1: Tolterodine ER 4 mg G2: Fesoterodine 8 mg G3: Placebo	<ul style="list-style-type: none"> • Significant improvement in HRQoL as assessed by KHQ and ICIQ-SF in both fesoterodine 8 mg arm and the tolterodine ER 4 mg arm vs. placebo, but no significant difference between arms • Fesoterodine arm showed significant improvement in eight of nine domains (non significant in General Health domain) • Subset of patients who reported incontinence at baseline showed similar improvements versus baseline; however, no significant difference was seen between arms as well as between subsets of those reporting incontinence at baseline versus those without
Elinoff et al. ¹⁶⁰ 2006	G1: Tolterodine ER 4 mg	<ul style="list-style-type: none"> • After 12 weeks of treatment, improvement in bladder condition was noted on the PPBC by 78.8% of the intent to treat population, 86.2% of those bothered by urgency, 78.5% with nocturnal frequency and 74.6% of those with urge urinary incontinence • All patients reported significant decreases in OAB symptoms. Median percentage change for those with urge urinary incontinence was -86.1 (95% CI: -91.7, -80.0)
Freeman et al. ¹⁷⁴ 2003	G1: Tolterodine ER 4 mg G2: Placebo	<ul style="list-style-type: none"> • Significantly more participants in the tolterodine group reported “much benefit” and bladder symptom improvement than in the placebo group (43% vs. 24%; p<0.001 and 62% vs. 48%; p<0.001, respectively) • Participants in the tolterodine ER group had greater improvement in self reported urgency compared with placebo (46.6% vs. 26.6%; p=0.001)
Kelleher et al. ²²³ 2001	G1: Tolterodine ER 4 mg G2: Placebo	<ul style="list-style-type: none"> • Participants on tolterodine had significant improvements in several domains of the KHQ relative to placebo: incontinence impact, role limitations, physical limitations, sleep and energy, severity (coping), symptom severity, all at p<0.05 • No significant differences were observed for social limitations, personal relationships, etc. • A significantly higher proportion of patients on receiving placebo compared to tolterodine reported improvement in their bladder condition (58% vs. 43%; p=0.001)

Table 18. Effect on quality of life and satisfaction for pharmacologic treatment (continued)

Author Year	Comparison Groups, N	Outcomes
Tolterodine		
Robinson et al. ²⁴⁰ 2007	G1: Tamsulosin OCAS 0.25 mg G2: Tamsulosin OCAS 0.5 mg G3: Tamsulosin OCAS 1.0 mg G4: Tamsulosin OCAS 1.5 mg G5: Tolterodine ER G6: Placebo	<ul style="list-style-type: none"> No significant improvements were seen on the KHQ in either the tolterodine or tamsulosin groups
Rogers et al. ²⁴¹ 2008	G1: Tolterodine ER 4 mg G2: Placebo	<ul style="list-style-type: none"> Treatment group had improvements in sexual and emotional health versus placebo. SQoL-F; PISQ-total score, and HAD anxiety; p=0.004, p=0.009, p=0.03 respectively
Tseng et al. ²⁴⁷ 2009	G1: Tolterodine 2 mg G2: Tolterodine 2 mg + vaginal CEE	<ul style="list-style-type: none"> Both groups had improvements over baseline on the UDI-6 and IIQ-7
Zinner et al. ¹⁵⁸ 2002	G1: Tolterodine ER 4 mg G3: Placebo G3: Tolterodine ER 4 mg G4: Placebo	<ul style="list-style-type: none"> After 12 weeks, 69.8% of older (≥65) participants considered the treatment beneficial, compared to 46.9% on placebo (p<0.001) In the younger group, 78.3% of treated patients considered the treatment beneficial, compared to 58.3% placebo (p<0.001)
Solifenacin		
Chapple et al. ^{99, 211} 2007	G1: Solifenacin 5 mg G2: Tolterodine ER 4 mg	<ul style="list-style-type: none"> Significant change was seen on PPBC in solifenacin group compared to tolterodine group at 4 weeks (-1.51 vs. -1.33; p=0.006)
Garely et al. ²¹⁸ 2006	G1: Solifenacin 5 mg or 10 mg (flexible dosing)	<ul style="list-style-type: none"> Significant change was seen on the PPBC scale from baseline to study end (4.4 vs. 2.9; p<0.001) Participants reported improvement on all subscales of the OAB-q (mean changes, 14.7 to 29.6; all p<0.001) Using the VAS, participants reported significant reductions in degree of bother associated with urgency, urge urinary incontinence, frequency and/or nocturia (all p<0.001)
Kelleher et al. ²²⁵ 2005	G1: Solifenacin 5 mg G2: Solifenacin 10 mg G3: Placebo	<ul style="list-style-type: none"> Patients in the solifenacin arms had significantly greater improvement in nine of ten domains of the KHQ (except personal relationships) after 12 weeks (p<0.05)
Yamaguchi et al. ⁹⁸ 2007	G1: Solifenacin 5 mg G2: Solifenacin 10 mg G3: Propiverine 20 mg G4: Placebo	<ul style="list-style-type: none"> Solifenacin and propiverine were both associated with significant improvements in QoL as measured by the KHQ when compared to placebo (p<0.05 on all subscales) Greater improvements were reported on the severity domain in the solifenacin 10 mg group compared to placebo (p<0.05)

Table 18. Effect on quality of life and satisfaction for pharmacologic treatment (continued)

Author Year	Comparison Groups, N	Outcomes
Trospium		
Rudy et al. ¹⁰⁴ 2006	G1: Trospium 20 mg G2: Placebo	<ul style="list-style-type: none"> Participants in the trospium arm had significantly greater reduction in the OAB-SCS score compared to placebo (-8.4 vs. -4.6; p<0.0001)
Staskin et al. ¹⁰³ 2007	G1: Trospium 60 mg G2: Placebo	<ul style="list-style-type: none"> Using the OAB-SCS, the study demonstrated significant improvement of quality of life in the treatment group relative to placebo at 12 weeks (-11.2 vs. -7.8; p<0.001)
Zinner et al. ¹⁰⁶ 2004	G1: Trospium 20 mg G2: Placebo	<ul style="list-style-type: none"> Participants in the trospium arm had a reduction of their IIQ score of 59 compared to 36 in the placebo group (p≤0.05)
Darifenacin		
Hill et al. ¹⁰⁰ 2006	G1: Darifenacin 7.5 mg G2: Darifenacin 15 mg G3: Darifenacin 30 mg G4: Placebo	<ul style="list-style-type: none"> VAS was used to assess changes in severity of urgency before and after treatment Although changes were significant in pre-post measures in G2 and G3, no statistical comparisons are reported between groups
Chapple et al. ²¹⁰ 2007	G1: Darifenacin 7.5/15 mg G2: Placebo	<ul style="list-style-type: none"> Participants in darifenacin arm showed statistically significant improvement in all domains of the OAB-q versus placebo (p<0.001)
Dwyer et al. ²¹⁷	G1: Darifenacin 7.5/15 mg	<ul style="list-style-type: none"> Improvement in patient reported outcomes from baseline as measured by KHQ (eight of nine domains, p<0.001) at 24 months
Steers et al. ¹⁰² 2005	G1: Darifenacin 7.5 mg G2: Placebo	<ul style="list-style-type: none"> Participants in the darifenacin arm reported significantly greater reduction in the severity of urgency episodes (p<0.05), by validated 100 mm VAS
Zinner et al. ¹⁰¹ 2006	G1: Darifenacin 15 mg G2: Placebo	<ul style="list-style-type: none"> Participants in the darifenacin arm had a significantly greater change in OAB-q score after 12 weeks of treatment (26.4 vs. 19.1; p<0.001) Changes were also significantly greater in the treatment group for incontinence impact (-24.7 vs. -17.8; p=0.022) and severity measures (-24.3 vs. -15.6; p<0.001) by ICIQ KHQ

Harms of Pharmacologic Treatments

Proportions of individuals reporting harms in RCTs of pharmacologic treatments ranged from 9.7 to 63.6 percent of study participants; however, harms were generally mild in nature, and withdrawals due to adverse events did not exceed 17 percent in any study (Table 19). The risk of occurrence of harms reported in treatment arms often overlapped those observed with placebo.

Dry mouth was the most commonly reported harm, ranging from 5.9 percent to 88 percent in studies of oxybutynin IR, compared to 1.6 to 21 percent in placebo arms. Studies of transdermal oxybutynin had the lowest reported estimates of dry mouth (2.6 to 9.6 percent). Impaired urination, not defined by the authors, was reported in studies of oxybutynin and tolterodine. It

was highest in two studies of oxybutynin IR (14 to 29 percent), compared to 3.2 to 4.0 percent in two studies of oxybutynin ER, no events to 9.0 percent in six studies of tolterodine IR and 1.0 percent in one study of tolterodine ER. Urinary tract infections were reported by up to 11 percent of participants in eight placebo arms. Immediate release formulations of oxybutynin and tolterodine both had reports of up to about 18 percent of participants experiencing a UTI, compared to up to 12 percent in studies of oxybutynin ER and 4.1 percent in studies of tolterodine ER. Between 0 and 32 percent of participants in treatment arms reported constipation; again, the highest rate was reported in an oxybutynin IR arm. Darifenacin had the second highest proportion of participants reporting constipation (18.5 to 27.8 percent). Up to seven percent of participants in placebo arms also reported constipation.

Cardiac events, including new abnormalities on EKG, were very rare and reported in only a few studies. The highest reported rate of cardiac events was five percent in tolterodine ER. Generally, however, less than one percent of participants experienced any cardiac event, mostly tachycardia and arrhythmias in treatment arms, with events also occurring in placebo arms (0 to 0.9 percent).

Table 19. Side effects and harms of pharmacologic treatment

Range (Number of studies reporting)	Placebo	Oxybutynin IR	Oxybutynin ER	Oxybutynin TDS	Tolterodine IR	Tolterodine ER	Trospium	Darifenacin	Solifenacin	Fesoterodine
Cardiac events*	0-0.9 (3)	0.2-1.2 (3)	NR	0.8 (1)	0 (1)	0-5 (2)	0.1 (1)	0.5 (1)	0.3 (1)	3.3-3.9 (1)
Constipation	0-7 (19)	0-32 (10)	6.4-8.6 (5)	3.3-5.4 (2)	2.6-10.4 (12)	2.5-10.2 (11)	7-10.9 (5)	18.5- 27.8 (5)	6.4-18.9 (9)	3.3-14 (2)
Diarrhea	2-5.4 (4)	2-5 (2)	7.9-14 (3)	NR	3-3.4 (2)	2-6.8 (4)	1-3.1 (3)	NR	NR	NR
Dizziness	0-3.8 (5)	1.6-38 (6)	3.8-11 (5)	4 (1)	1.7-4.3 (4)	1.4-2.5 (5)	NR	0 (1)	1.2 (1)	1-1.5 (1)
Dry mouth	1.6-21 (23)	5.9-88 (15)	14-68 (9)	2.6-9.6 (3)	10-50 (15)	7.3-39 (13)	8.7-33 (5)	20.4- 59.1 (5)	17.9- 34.1 (9)	21.7-99 (2)
Dyspepsia	0.9-5 (4)	3-27 (5)	5.3-11 (3)	NR	3-9 (7)	2.7-3 (3)	5 (1)	5.2-8.7 (2)	NR	NR
Fatigue	<1-1.6 (3)	15 (1)	1.6-18 (2)	NR	1-3.6 (2)	2-3.4 (2)	NR	NR	NR	<1 (1)
Headache	0-5 (12)	1.2-22 (7)	5.6-12 (6)	NR	3-10.4 (10)	3-7 (9)	1-5.5 (4)	3.8-8.1 (4)	3.4-3.6 (4)	2.4-12 (2)
Impaired urination	0 (2)	14-29 (2)	3.2-4 (2)	NR	0-9 (6)	1 (1)	NR	NR	NR	NR
Insomnia	2-2.2 (2)	2 (1)	0.5-1.8 (3)	NR	0.5-1.8 (4)	0.8-1.7 (3)	4 (1)	NR	0.8 (1)	NR
Nausea	2-11 (4)	2-17 (7)	3.2-5 (3)	4.6 (1)	1.6-7 (6)	1-2.7 (4)	2 (1)	NR	1.8 (1)	<1-1.4 (1)
Respiratory events	0-14 (5)	3-13 (2)	6 (1)	NR	10-16 (2)	4 (1)	6.4 (1)	0.3-5.6 (2)	3.1-4.6 (3)	NR
Somnolence	0-2 (4)	3-40 (3)	1-4.3 (4)	1.6 (1)	1.6-2.7 (4)	2.3-3 (3)	NR	NR	NR	NR
Urinary tract infections	0-11 (8)	<1-18 (3)	5.1-12 (2)	2.4 (1)	3-19 (4)	2.7-4.1 (6)	4.9-12 (2)	1.1-4.8 (3)	3.4-3.6 (2)	10-15 (1)
Vision changes	0-7.7 (12)	1.2-22 (8)	2.2-3.3 (4)	2.3 (1)	0.6-7.5 (6)	<1-6 (6)	3 (1)	0-3.5 (2)	0.7-6.9 (9)	2.2-4.2 (1)
Any adverse event	17.5- 48.9 (6)	57 (1)	51 (1)	NR	53 (1)	9.7-74 (6)	26.8- 59.6 (2)	47.9- 63.6 (2)	59.4 (2)	50-69 (2)
Withdrawals due to adverse events	0-6 (11)	16-17 (2)	6.2-13 (4)	10.7 (1)	1.9-15 (6)	2.8-6.3 (5)	7.3-8.8 (1)	3.2 (1)	3.7-9.7 (4)	6-9 (1)

* Includes new abnormal EKG, tachycardia, other arrhythmias, palpitations, and other cardiac events as grouped together in the literature.

Procedural and Surgical Treatments of OAB

We reviewed 18 studies, of which 11 were fair quality and 7 poor. This section presents the results of our literature search and findings about outcomes of procedural and surgical treatments for OAB. These treatments include sacral neuromodulation, peripheral neuromodulation, electromagnetic nerve stimulation, injection or instillation of drugs into the bladder, bladder distention and bladder transection. No studies regarding augmentation cystoplasty or detrusor myomectomy met our search criteria. Detailed information on all studies relating to surgical management of OAB can be found in evidence tables in Appendix C.

Sacral neuromodulation. Stimulation of the sacral nerve roots is a technique in which an electrical stimulus directly stimulates the S3 sacral nerve root.²⁵⁶ The technique has evolved over time, but typically it is performed as a staged procedure. The first stage involves a “test” stimulation using a percutaneous needle to stimulate the S3 nerve root. If there is a favorable response during the trial period, then long-term stimulation can be provided by implanting an implantable pulse generator surgically. The implantable pulse generator is usually placed in the fatty tissues overlying the buttocks, although abdominal placement was used with some of the earlier studies. Recent evolutions in this technique now permit a permanent lead to be used for the test stimulation. If the test is unsuccessful, the lead can be removed, but if it is successful, this lead is attached to the permanent implantable pulse generator. This has the advantage of ensuring that stimulation is provided in the exact location as during the test period. (Previously, a new lead was placed at the time of the implantable pulse generator placement.) The mechanism by which neuromodulation acts to improve symptoms is not completely understood. The technique is used for urinary urgency, frequency, and urge incontinence refractory to other treatment modalities.²⁵⁶ It is also used for urinary retention. Given these seemingly contradictory applications, it is thought that the electrical stimulation affects the afferent nerves (which perceive bladder sensation), thus allowing them to appropriately transmit bladder sensations.

Peripheral neuromodulation. Other techniques for neuromodulation involve stimulating the S3 nerve fibers more peripherally, at the posterior tibial nerve or cutaneous stimulation of the pudendal nerve via an anal or vaginal probe.²⁵⁷ For the posterior tibial nerve stimulation, a needle is placed percutaneously near the ankle and is attached to an external electrical device. Instead of implanting an implantable pulse generator, the patient returns for periodic sessions, often weekly for a series of treatments. Small case series suggest that posterior tibial nerve stimulation may improve OAB symptoms.²⁵⁸⁻²⁶⁰ There were no studies involving this technique which met our search criteria. One study evaluating neuromodulation of the pudendal nerve with anal and/or vaginal probes met our search criteria. Similarly, this is performed on an outpatient basis with weekly treatment sessions.

For the purposes of this report, sacral neuromodulation will refer to techniques that directly stimulate the S3 nerve root. Peripheral neuromodulation will refer to nerve stimulation peripherally, such as the use of an anal or vaginal probe to stimulate the pudendal nerve. These approaches are reviewed in the same section of this text.

Electromagnetic sacral nerve stimulation. Electromagnetic stimulation is yet another modality to modulate the neurologic control of the bladder. Most treatments involve large, powerful magnets which require a dedicated facility as the magnets are not portable.²⁶¹ The study included in this review evaluated the use of a smaller, portable electromagnetic system.

Bladder instillation/injection of a drug. Another approach to treating urinary frequency, urgency, and urge incontinence is to instill or inject a drug into the bladder. Numerous drugs

have been administered using this approach. Two categories of intravesical drugs are included in this review, antimuscarinic agents and neurotoxins. By instilling or injecting the drug directly into the bladder, systemic adverse effects are theoretically avoided.

Two neurotoxins discussed in this review are resiniferatoxin and botulinum toxin. Botulinum toxin is a neuromuscular blocking agent which prevents nerve conduction. Typically botulinum toxin-A is used and can be injected directly into the wall of the bladder under cystoscopic guidance as a treatment for refractory OAB.²⁶²⁻²⁶⁴ It is not FDA approved for this indication at the time of the writing of this document. Concerns with this approach are the risk of urinary tract infections and urinary retention and that the ideal dosing has not yet been determined. Additionally, the effects of botulinum toxin are temporary and multiple courses of treatment would be anticipated. An RCT comparing botulinum toxin A to placebo for women with refractory urge incontinence found that incontinence episodes decreased from over 20 episodes per three day diary to less than 5 episodes per three day diary among those receiving botulinum toxin A, while those taking placebo had no difference in the number of incontinence episodes. Using a Patient Global Impression of Improvement score, approximately 60% of the women who received botulinum toxin A had a clinical response, and this response lasted six times longer than that achieved with placebo. The study was stopped early due to increased postvoid residuals in 43% of women receiving botulinum toxin A and a high rate of urinary tract infections among those with elevated postvoid residuals.²⁶⁵ Resiniferatoxin is a neurotoxin in the same category as capsaicin; these do not have FDA approval for the treatment of OAB. These agents block transmission along the C-fibers, nerve fibers involved in transmitting noxious stimuli.¹²⁷ It has been hypothesized that inhibition of these fibers may be a treatment for overactive bladder.

A review of RCTs evaluating the use of intravesical botulinum toxin for OAB was published by the Cochrane Collaboration in 2007.¹²⁸ Eight studies met their search criteria: five were published abstracts and three were full papers. Only one of the eight studies exclusively dealt with idiopathic OAB, the definition we used for our literature search. The remaining seven studies in the Cochrane review included subjects with neurogenic OAB.

The findings from the Cochrane review were that botulinum toxin injections were more effective than placebo, with fewer incontinence (unspecified type) episodes per day at 2 to 24 weeks and fewer incontinence episodes compared to baseline. Improvements following treatment were also seen in incontinence specific and overall quality of life. Urodynamic changes were also seen, including decreased pressure during a detrusor contraction and increased bladder capacity following treatment with botulinum toxin. The postvoid residual, the amount of urine left in the bladder after voiding, was also elevated. The numerical data were not provided by the author. Adverse events included cases of urinary retention requiring intermittent self catheterization following treatment. Of the patients requiring catheterization, 25 percent had a lower urinary tract infection.

In the Cochrane Collaboration, the limited number of studies, their small size and heterogeneous population highlight the need for more research regarding treatment with botulinum toxin.¹²⁸ One study compared botulinum toxin to bladder instillations with resiniferatoxin and found lower rates of incontinence, increased bladder capacity and lower detrusor pressure during uninhibited bladder contractions at 6 to 18 months with the botulinum toxin treatment. The optimal dose or long term effects of elevated postvoid residuals has not been determined.

Bladder distention and bladder transection. Two treatments that are no longer in common practice, prolonged bladder distention and bladder transection, are also included in this review. The bladder distention study describes distending the bladder to maximum capacity at a pressure equal to systolic blood pressure for four hours.¹¹⁸ Bladder transection involves cutting the

bladder wall and detrusor muscle. Both have significant morbidity and have been abandoned by most practitioners.

Outcomes of procedural and surgical treatments.

Content of the literature. We identified 18 studies reporting on surgical treatments and procedures for OAB (clinical trials reported in Table 20).¹¹⁰⁻¹²⁷

Eleven were of sacral neuromodulation, one of peripheral neuromodulation and one of electromagnetic sacral nerve stimulation. Three studied bladder instillation or injection of drugs, one was of bladder distention and one was of bladder transection.

Six of the 11 studies on sacral neuromodulation come from a family of papers financially supported by one company.^{110, 111, 114, 115, 123, 124} The patient population for this family of studies included an RCT to evaluate sacral neuromodulation versus medical therapy for six months.¹²⁴ The other studies involved similar inclusion and exclusion criteria, but varied in regards to the number of centers involved in recruitment (between 12 and 17 centers) and timing for enrollment. Given this, it is not possible to determine the degree of subject duplication.

Table 20. Outcomes of clinical trials of procedures

Author Year Design	Groups	N	Per day baseline	Per day on treatment	Decrease in episodes per day	Weeks at evaluation
Episodes of urge incontinence per day						
Rios et al. ¹²⁷ 2007	G1: Resiniferatoxin	34	3.1	2.7	0.4*	4
	G2: Placebo	24	5.8	4.2	1.6*	4
Schmidt et al. ¹²⁴ 1999	G1: SNM	34	9.7	2.6	7.1^	24
	G2: Usual care	42	9.3	11.3	+2.0^	24
Voids per day						
O'Reilly et al. ¹²⁵ 2008	G1: Trans-sacral magnetic stimulation	33	10.0	9.0	1.0*	12
	G2: Sham treatment	30	9.0	9.0	0.0*	12
Rios et al. ¹²⁷ 2007	G1: Resiniferatoxin	34	9.7	9.0	0.7*	4
	G2: Placebo	24	9.9	9.2	0.7*	4
Enzelsberger et al. ¹²⁶ 1995	G1: Oxybutynin instillation	26	12.6	5.8	6.8^	4
	G2: Sterile water instillation	26	12.8	10.4	2.4^	4

* Not significant differences between groups; ^ = p <0.01

This literature included 13 case series studies, which we operationally defined as descriptive analyses of a sequence of participants having the same type of procedure without a comparison to another type of surgery or treatment. Three of these studies are *retrospective* case series of a particular surgical treatment: two report on sacral neuromodulation^{119, 120} and one on bladder transection.¹²¹ Nine studies are *prospective* case series: six report on sacral neuromodulation,¹¹⁰⁻¹¹⁵ one on peripheral nerve stimulation with anal and/or vaginal probes,¹¹⁶ one on botulinum-A toxin injections¹¹⁷ and one on prolonged bladder distention.¹¹⁸ One case series had both a retrospective arm as well as a prospective arm; this study looked at sacral neuromodulation.¹²²

One study is a prospective cohort that compared outcomes among subjects receiving sacral neuromodulation and subjects receiving a control surgery for sacral neuromodulation, but without active electrical stimulation.¹²³

Four studies were RCTs: one looked at sacral neuromodulation versus medical therapy,¹²⁴ one evaluated transcutaneous electromagnetic stimulation versus sham,¹²⁵ and two evaluated instillation of a drug into the bladder versus placebo – one using oxybutynin¹²⁶ and one using resiniferatoxin.¹²⁷

The majority of the studies were conducted in Europe, six of which included involvement from Canada and the United States (there is subject duplication among these six studies). One study was conducted in the United States only. Two studies were performed in Australia and one in Brazil. Two of the studies involved national registries, one from Switzerland and one from Italy. Five studies specified that they occurred in an academic setting, three in a specialty setting and the remainder did not specify the clinical setting for the study.

The quality of the studies varied widely. Of the four randomized controlled trials, three had an appropriate control group, the fourth compared sacral neuromodulation to continued medical treatment among subjects who had already failed medical management for refractory OAB.¹²⁴ This does not adequately control for the placebo effect, which is quite prominent among these treatments. It is challenging to create an appropriate control group when evaluating sacral neuromodulation. This is illustrated by a cohort study in which surgical controls received sacral neuromodulation, but no electrical stimulation was applied. Given that subjects feel the electrical stimulation, they would be aware that this has been de-activated following the test stimulation.¹²³ Of the 11 studies on sacral neuromodulation, seven did not restrict the study population to OAB. Although sacral neuromodulation may be used to treat urinary retention, grouping these studies in such a fashion severely limited our ability to analyze the results. We restricted our discussion to the OAB findings in these papers. Case series studies are limited by the lack of a control group.

Outcomes assessed. Half of the studies reported on urge urinary incontinence outcomes (usually incontinence episodes per day as measured by a bladder diary, but also pad counts, pad weight, severity of incontinence or other measures). Nearly two-thirds of the studies reported some measure of urinary frequency, such as voids per day. Nearly all of the studies reported on a subjective measure of symptoms (such as percent cured, perceived severity, QoL). A third of the studies reported on urodynamic outcomes, such as the bladder volume at which there was a normal desire to void, and bladder capacity.

The majority (78 percent) of the studies, reported data on adverse effects or harms. Six studies reported on problems with constipation or gastrointestinal symptoms,^{111, 114, 116, 119, 120, 126} eight reported the presence of pain, six reported on infection and nearly half of the sacral neuromodulation studies reported the presence of lead migration. Other adverse effects were reported in 72 percent of studies.

Outcomes of sacral neuromodulation and peripheral neuromodulation.

The one RCT comparing sacral neuromodulation to medical therapy found a reduction in daily incontinence episodes from 9.7 to 2.6 in the intervention group, compared to an increase of 9.3 to 11.3 in the medical management group at six months ($p < 0.01$).¹²⁴ Of note, all subjects receiving medical therapy had already failed medical management; no benefit from continued medical therapy would be expected. The remaining six case series that reported on change in UUI had decreases in mean incontinence episodes per day of 51 percent to 80 percent^{111, 114, 115, 119, 122} and from a median of five down to zero incontinence episodes a day.¹¹² Length of followup in these studies ranged for six months to five years.

Pad use per day also decreased with sacral neuromodulation. Most studies started with a baseline of five or more pads used daily. One RCT reported an 82 percent decrease in pad use from 6.2 to 1.1 pads daily, six months following initiation of sacral neuromodulation.¹²⁴ Three case series evaluating sacral neuromodulation also found significant decreases in pad use ranging from 49 to 84 percent fewer mean pads^{111, 114, 115} and a 75 percent decrease in median pad use.¹¹² Length of followup on these studies ranged from six months to five years.

Some of the studies tried to characterize the severity of incontinence episodes. One RCT and two case series found a 64 percent to 92 percent decrease in the number of moderate to heavy urge urinary incontinence episodes at six months to five years of followup.^{111, 115, 124} The RCT reported the highest rate of decreased heavy incontinence episodes with a mean baseline in the neuromodulation group of 3.4 per day, reduced to 0.3 per day six months after treatment. In comparison, those with refractory OAB receiving usual therapy experienced an increase in mean heavy episodes per day from 2.6 to 3.9.¹²⁴

Improvement in episodes of urinary urgency without incontinence is difficult to measure as it is a more elusive symptom. Nonetheless, on a 3 point scale, (1=mild, 2=moderate, 3=severe), 69 percent of participants in one study reported improvement,¹¹⁴ with a second study showing no change in experience of urgency.¹¹⁵

Reduction in urinary frequency of between 31 and 45 percent is seen consistently across studies of sacral neuromodulation, regardless of study design. The majority of studies of sacral neuromodulation reported urinary frequency as the mean number of voids in 24 hours as recorded by a bladder diary. Six studies reported mean voids per day, one of which was a prospective cohort study comparing subjects with sacral neuromodulation and controls who had sacral neuromodulation placed, but did not receive active electrical stimulation.¹²³ At six months, those receiving sacral neuromodulation had a 45 percent decrease in the number of voids per day; no change was seen in the controls. Similar results were seen in the other studies, which found 31 to 40 percent fewer voids per day with sacral neuromodulation, regardless of whether they were prospective or retrospective case series studies.^{114, 115, 119} One study reported its results as a 40 percent decrease in the median number of voids per day.¹¹² Another study reported the results from an Italian national registry; they did not have baseline information regarding the number of voids in the registry prior to treatment, but 42 percent of subjects had fewer than 8 voids daily after treatment.¹²² The 31 to 45 percent decrease in mean (and median) voids per day seen across the studies was present at six months and up to two years following initiation of the sacral neuromodulation.^{112, 114, 115, 119, 123} The longest followup data was available from a prospective case series which found a 33 percent decrease in mean voids per day at one year which was reduced to a 23 percent decrease in mean voids per day at five years.¹¹⁵

Some studies also looked at the mean voided volume as a measure of treating urinary frequency. One cohort study and two case series found that sacral neuromodulation increased the mean voided volume between 1.7 to 1.9 fold, an increase of 78 mL to 108 mL per void.^{114, 115, 123}

Peripheral neuromodulation and electromagnetic stimulation were clinically ineffective in changing voiding frequency. One prospective case series found a 12 percent decrease in the mean voids per 24 hours was seen six weeks following 12 sessions of peripheral neuromodulation with an anal and/or vaginal probe.¹¹⁶ No decrease in mean voids per day was seen with electromagnetic stimulation of the sacral nerves.¹²⁵

Neither sacral neuromodulation nor peripheral neuromodulation with an anal and/or vaginal probe had a clinically relevant impact on nocturia rates, which were very low in these studies at baseline. Both treatment approaches reduced already low rates by approximately 30 percent.^{116, 119}

Several of the studies comment on clinical “cures” or “improvements”, but lack definitions for these criteria. Moreover, several of the case series on sacral neuromodulation include diverse patient populations which may include urinary retention or pelvic pain in addition to patients with symptoms of overactive bladder. The clinical endpoints for improvement in urinary retention are very different from those for urinary urgency and frequency.

Studies evaluating improvements following sacral neuromodulation describe cure rates of 26 to 65 percent (cure was defined as “completely dry” at six to twelve months).^{111, 122, 124} One study looked at improvements following the initial test stimulation and found 39 percent had greater than a 90 percent improvement in urinary frequency and/or urgency.¹¹³ As would be expected, higher rates are seen for classifications of “success” or “improvement” as compared to “cure.”

Compared to these rates, peripheral neuromodulation only had an 8.1 percent cure rate, with 31 percent reporting no change.¹¹⁶

Several studies evaluated QoL, either with the KHQ, a VAS or other validated QoL questionnaire (Table 21). Two studies evaluated the impact of sacral neuromodulation on QoL and found the treatment beneficial.^{122, 123} There was no improvement in QoL with electromagnetic stimulation.

Table 21. Effect on quality of life and satisfaction of procedural treatments

Author, Year Study Type	Comparison Groups, N	Outcomes
Hassouna et al. ¹²³ 2000 Cohort	G1: Neuromodulation (25) G2: Control (25)	<ul style="list-style-type: none"> At six months, the neuromodulation group reported greater improvement in quality of life compared to controls (p<0.0001) SF-36 scores²⁶⁶ were significantly higher (p<0.01) for sacral neuromodulation participants relative to controls on physical function, role physical, bodily pain, vitality, social function and mental health domains
O’Reilly et al. ¹²⁵ 2008 RCT	G1: Electromagnetic sacral nerve stimulation (33) G2: Sham (30)	<ul style="list-style-type: none"> No significant differences were observed by group for the KHQ domain(s) or the Australian Quality of Life Questionnaire domain(s) after Bonferroni correction for multiple testing
Rios et al. ¹²⁷ 2007 RCT	G1: Single dose 100 ml 50 nM resiniferatoxin (34) G2: Placebo (24)	<ul style="list-style-type: none"> No differences were reported from baseline to post-treatment QoL scores by the KHQ for general health perception, social limitations or personal relationships. Incontinence impact decreased a similar amount in both groups following intervention (p<0.05 pre-post). Emotions, sleep/energy and symptom severity scores decreased more for G1 following the intervention (no between group testing) Role limitations decreased for G2 (p<0.05) but not G1 following the intervention
Schmid et al. ¹¹⁷ 2006 Prospective case series	Botulinum Toxin (100)	<ul style="list-style-type: none"> 90% of patients experienced improvement in at least one category of the KHQ at three months, with a waning of benefit by nine months.
Spinelli et al. ¹²² 2001 Case series	G1: Sacral neuromodulation retrospective cases (93) G2: SNM prospective cases (103)	<ul style="list-style-type: none"> In a validated QoL questionnaire²⁶⁷ completed by 54% of prospective participants, all reported significant increases in life quality at all time points.

The evaluation of overactive bladder with urodynamics provides objective data, but the clinical relevance of this information is not always clear. Urodynamic parameters may indicate changes in bladder function, but unless symptoms were also evaluated, they do not necessarily carry over into improvements in clinical outcomes.

During bladder filling, known as cystometry, the first sensation of bladder filling can be measured. This is known to be one of the more variable urodynamic measurements.²⁶⁸ The bladder volume at first sensation increased by 80 mL after sacral neuromodulation in one case series¹¹⁰ and by 36 mL after peripheral neuromodulation with an anal/vaginal probe.¹¹⁶ Both of these studies also looked for changes in bladder capacity and found this increased 1.4 fold for sacral neuromodulation and found no difference for peripheral neuromodulation.^{110, 116}

By their very nature, surgical and procedural treatments are likely to have a higher incidence of adverse events than conservative and medical treatments for OAB. In the early studies of sacral neuromodulation, there was an average of 1.1 to 1.7 adverse events per participant.^{115, 124} Advances in technology, such as the use of tined leads, have decrease this rate and the more recent studies report 0.1 to 0.5 events per participant.^{112, 119} None of the studies exclusively used the newer technology. Of the adverse events, pain, lead migration or problems with the lead, infection and explantation of the device were the most common adverse events. From the test stimulation phase, pain at the needle site was 0.5 percent in a study employing newer technology¹¹² and 7 percent in a study with older technology.¹¹³ When the implantation phase is included, pain rates and uncomfortable stimulation responses were seen in 3.9 to 43 percent of subjects, with studies employing new techniques at the lower end of this spectrum.^{112, 114, 115, 120, 122, 124} The one RCT found a pain rate of 19.1 percent, but this was compared to medical management, not a sham procedure.¹²⁴ Pain at the implantable pulse generator implantation site was typically reported separately and occurred 15.4 to 27 percent of the time.^{114, 115, 120, 124} Problems with the lead or lead migration occurred between 3.3 and 11 percent of the time.^{114, 115, 120, 122, 124} This may be less common with the new tined leads, as one study had lead migration only with the older non-tined leads¹¹² and another found a lower loss of efficacy, 12.3 percent, with the tined leads compared to the non-tined leads, 31.7 percent.¹¹⁹ Infection occurred in 1.9 to 6.1 percent of participants, sometimes requiring hospitalization for intravenous antibiotics or removal of the device.^{114, 119, 120, 122, 124} Two papers noted a 0.5 to 1.7 percent risk of neuropraxia or nerve injury.^{114, 120} There was a high rate of needing surgical revision, with most studies showing 33 to 48.3 percent of subjects required a return to the operating room.^{114, 120, 124} Lower rates were documented in a study using newer technology, 7 percent,¹¹² and in a national registry, 9.7 percent.¹²² One study looked at return to the operating room rates at five years and found that that there was a 67 percent risk of return (numbers based on the population enrolled).¹¹⁵ However, at five years, many of the subjects were returning for new implantable pulse generator batteries, an expected development over time, thus this is not truly an adverse event. Unfortunately, the number of surgeries which were to replace implantable pulse generator batteries was not reported. Of the surgical adverse events, a significant proportion included explantations of the device. This occurred in 3.9 percent of subjects in the national registry;¹²² higher rates were seen in the remaining studies, 9.8 to 14 percent.^{111, 115, 120}

Adverse events following peripheral neuromodulation with an anal and/or vaginal probe led to a 19 percent dropout rate, with 29 percent of the dropouts attributed to pain. The other most common complaint was bowel irritation.¹¹⁶ Currently, techniques of peripheral neuromodulation with an anal/vaginal probe are not routinely used.

Outcomes of bladder instillation/injection of drugs. An RCT of resiniferatoxin bladder instillations versus placebo found no improvement in the number of urge urinary incontinence episodes per day with either arm.¹²⁷ However, bladder instillations with oxybutynin did decrease

mean voids per day. In an RCT evaluating bladder instillations with oxybutynin versus a placebo with sterile water, those receiving oxybutynin had a 47 percent decrease compared to a 16 percent decrease in mean voids per day with the placebo at two weeks following completion of treatment.¹²⁶ This decrease is similar to that seen with sacral neuromodulation. In a prospective case series evaluating the use of botulinum-A toxin injected into the detrusor muscle of the bladder wall, 74 percent of the subjects had 8 or fewer voids per day four weeks following treatment. All subjects had greater than 8 voids per day at the start of the study.¹¹⁷ In an RCT of bladder instillations with resiniferatoxin versus placebo, there was no difference in the mean voids per day four weeks following treatment.¹²⁷ Bladder instillations with oxybutynin decreased nocturia 65 percent from 5 to 1.8 episodes per night.¹²⁶ Following injection of botulinum-A toxin into the bladder, 66 percent had no urgency at 12 weeks and 80 percent reported no urge incontinence.¹¹⁷ No difference was seen compared to baseline or placebo in an RCT evaluating resiniferatoxin bladder instillations.¹²⁷

Two studies looked at what volume the normal desire to void occurred, finding this increased 1.6 fold (55 mL increase) for instillation of oxybutynin into the bladder¹²⁶ and 1.7 fold (83 mL) for botulinum-A toxin bladder injections.¹¹⁷ Theoretically these increases would result in less urinary frequency, and possibly less urgency. Similar changes were also seen in terms of bladder capacity after treatment, with bladder instillations of oxybutynin increasing capacity 1.5 fold (105 mL)¹²⁶ and 1.6 fold (135 mL) with botulinum-A toxin injections.¹¹⁷

A potential adverse effect of intravesical oxybutynin or botulinum-A toxin is impairment of the bladder's ability to empty.²⁶⁵ The goal of treatment with these agents is to provide a dose large enough to decrease the symptoms of OAB, without impairing the ability to void when physiologically necessary. An objective measurement that can serve as a proxy for this parameter is a postvoid residual, the volume of urine that remains in the bladder after voiding; however the clinical importance of an increased postvoid residual in the absence of symptomatic urinary retention is not clear. The mean postvoid residual following treatment with intravesical oxybutynin increased twofold to a mean of 40 mL (range 10 to 50 mL) and for botulinum-A toxin, increased fourfold to a mean of 75 ± 10 mL.^{117, 126} Both treatments seem to have similar effects on bladder capacity, but the risk of urinary retention may be higher with use of botulinum-A toxin.^{117, 126} Currently, oxybutynin and resiniferatoxin instillations are not routinely used for the treatment of OAB.

For instilled/injected bladder drugs, there were small benefits seen in the emotions, sleep/energy and symptoms severity subscales of the KHQ for subjects receiving the treatment compared to placebo. Incontinence impact scores improved a similar amount for both resiniferatoxin and placebo ($p < 0.05$). Emotions, sleep/energy and symptom severity scores decreased more for resiniferatoxin following the intervention ($p < 0.05$).¹²⁷ Ninety percent of subjects receiving botulinum A toxin had an improvement in QoL scores at 3 months; this effect was waning at 9 months.¹¹⁷

Outcomes of bladder distention and transection. Bladder distention and transection are treatment methods no longer in routine use. Following bladder distention, 18 percent were symptom free at 18 months with 52 percent unchanged.¹¹⁸ The study on bladder transection reports 65 percent were cured at two to five years and 16 percent were unchanged, but provides no information about the criteria for these categories.¹²¹ Prolonged bladder distention had a 4 percent rate of bladder rupture.¹¹⁸ Bladder transection was associated with a 14 percent rate of vesicoureteral reflux on urodynamics, the clinical significance of which was not determined, and 1 percent rate of a persistent urine leak requiring reoperation. The authors also noted “minor chest and urinary tract infections” but the rates of these adverse events were not reported.¹²¹

Behavioral Treatments

This section presents the results of our literature search and findings about outcomes of behavioral techniques to reduce overactive bladder in women. Behavioral treatment options for OAB have been used for managing urinary incontinence for more than 50 years, although large-scale and well-designed studies on them are fairly uncommon. We reviewed 27 studies, of which 14 were fair quality and 13 poor.

Behavioral techniques include the use of bladder training, pelvic floor muscle exercises (PME), biofeedback, dietary changes, and multicomponent approaches that combine bladder training with PME and/or biofeedback. Detailed information on all studies related to behavioral techniques for OAB can be found in evidence tables in Appendix C.

Bladder training. Bladder training was introduced in the 1960s (Jeffcoate and Francis), and modified by Frewen in the 1970s. It involves education, a strict schedule of daytime voiding with progressive increases in time between voids, urgency suppression techniques, and positive reinforcement. Frewen recommended that women be treated initially on an inpatient basis, and the training was often combined with antimuscarinic medication or sedatives to manage extreme urgency. Inpatient bladder training is no longer standard practice, and the technique has been modified to be administered on an outpatient basis, with the use of patient education and bladder diaries. Although the underlying mechanism for bladder training in OAB is not well understood, it is thought to reverse dysfunctional habits, increase bladder capacity and provide techniques for handling feelings of urgency.

Pelvic muscle exercises. Training the pelvic floor muscles were originally suggested for patients with stress incontinence, the idea being that patients could learn to contract the periurethral muscles to occlude the urethra during activities that caused leakage. However, it may also be useful in inhibiting detrusor contractions. PME can be implemented alone or with additional techniques such as biofeedback to help patients identify and contract the appropriate muscles.

Multicomponent approaches. Although both bladder training and PME can and are administered alone, they may also be combined with or without biofeedback for a multicomponent approach to reducing incontinence.

Tools for behavioral training. Behavioral training can be administered with written materials, verbal feedback, coaching in person or on the phone, in groups, or using other strategies, such as cognitive approaches or biofeedback for increasing the potential for success. Behavioral approaches can also be combined with medications such as antimuscarinics. Various combinations of approaches are examined in the literature and are described in this review.

For the purposes of this report, we will use the term “bladder training” to refer to bladder training alone (i.e., without PME without biofeedback). “Behavioral training” will refer to a multicomponent approach that includes bladder training. We indicate when biofeedback is used in conjunction with other behavioral techniques.

Behavioral treatments.

Content of the literature. We identified 29 papers from 27 studies that included arms with outcomes of behavioral interventions. We have divided the studies into three primary categories: those that compare only behavioral approaches, those that compare behavioral approaches to pharmaceutical ones directly and those that measure the effect of adding a behavioral approach to a pharmaceutical one (combination approaches). The first category is presented here and the second two are covered in KQ3.

Nine studies met criteria and included only behavioral arms. Among these, two studies focused on comparing delivery mechanisms or approaches and provide insight into techniques for teaching and encouraging participants in behavioral management.^{132, 135}

Four additional studies (represented in six papers) with multiple arms, including pharmacologic ones, provided data that allowed the comparison of behavioral management to placebo^{93, 143-145} or to another behavioral intervention.^{148, 201} These are counted in the direct comparisons section, and therefore are not represented in the counts below, but relevant outcomes are described here as well as the direct comparisons section, and they appear in tables in both sections. All three of these are RCTs.

The literature base of studies that had only behavioral arms included three retrospective case series, which we have operationally defined as a sequence of participants having the same intervention without a comparison to another type of intervention. One examined bladder training alone,¹²⁹ one examined pelvic muscle exercises¹³⁰ and the third reported on a series of participants who were provided either bladder training or biofeedback, but presented results only for the two groups combined.¹³¹ All three were conducted in community-based clinical settings.

One was a prospective cohort study comparing three bladder training approaches: self-administered, coaching and cognitive strategies.¹³²

Five studies were randomized controlled trials. One compared bladder training to a “control” condition.¹³³ One compared bladder training to pelvic muscle exercises.¹³⁴ One included the following three arms: pelvic floor muscle training, pelvic floor muscle training assisted with biofeedback, and electrical stimulation.⁹³ One compared three different approaches to multicomponent behavioral training: biofeedback, verbal feedback and self-administered using an instruction booklet.¹³⁵ One compared bladder training to bladder training with an additional caffeine reduction component.¹³⁶

Three of the studies were conducted in the United States, four in Europe, one in Australia and one in Taiwan. Four were conducted at academic medical centers; five were in community settings, of which two included an inpatient component.

Outcomes measured. For each type of intervention, we combed the publications for the outcomes and complications summarized in the analytic framework presented in Chapter 1.

Five studies reported a change in numbers of episodes of incontinence,^{93, 134-136, 143} although the time period varied, and three of these calculated a percent reduction in incontinence episodes.^{135, 136, 143} Three of the studies measured incontinence episodes over the course of a week,^{134, 135, 143} two did so over 24 hours.^{93, 136} One study reported that a significant decrease in frequency of voiding was observed, but did not provide data that could be included in this table.¹³²

Five studies presented the outcome of cure or improvement in incontinence (using various definitions)^{129-131, 133, 269} with three defining cure as complete resolution of incontinence.^{129, 131, 133}

One study presented data on episodes of urgency separate from incontinence.¹³⁶

Changes in frequency were reported in three studies.^{131, 136, 145} Two specified voids per day as an outcome.^{136, 145} In addition, Jarvis and colleagues¹³³ reported on the numbers of women indicating that they had nocturnal and diurnal frequency before and after treatment. One study

recorded frequency over a three-day period¹³² and reported a statistically significant effect, but did not report numbers of episodes, and one reported time between voids.¹³¹ One reported that cure included achieving a minimum of three to four hours between voids¹²⁹ and was therefore related to frequency but not summarized as effect on frequency.

Other patient-reported outcomes in this literature included reports of resolution or improvement and changes in severity, quality of life, and satisfaction.

Five of the studies provided some patient-reported outcome, with four reporting on perceived improvement.^{130, 131, 134, 135} Three provided some report on severity;^{130, 135, 143} four measured bother;^{132, 134, 135, 143} two measured impact or interference with daily activity;^{134, 135} one assessed changes in quality of life overall;¹³⁴ and three reported on patient satisfaction.^{134, 135, 143}

Outcomes of behavioral treatment.

UUI episodes. Lack of consistency in study design, interventions or comparison groups makes it impossible to provide consistent summary results across studies. Although each intervention was associated with reductions in incontinence episodes, no behavioral approach performed better than any other in any study in this category on incontinence outcomes over any time period greater than 12 weeks. Trials are summarized in Table 22.

Table 22. Outcomes of behavioral treatment trials

Author Year	Groups	N	Per day baseline	Per day on treatment	Decrease in episodes	Weeks at evaluation
Episodes of incontinence per day						
Arruda et al. ²⁰¹ 2008	G1: Oxybutynin	22	2.0	1.0	1.0	12
	G2: Functional electrostimulation	21	1.9	1.1	0.8	12
	G3: Pelvic floor training	21	2.3	1.1	1.2	12
Wang et al. ⁹³ 2006	G1: Electrical stimulation	25	1.0	0.5	0.5	12
	G2: Oxybutynin 2.5 mg 3x/day	26	0.0	0.0	0.0	12
	G3: Placebo	23	1.0	1.0	0.0	12
Bryant et al. ¹³⁶ 2002	G1: Bladder training + caffeine reduction	48	2.8	1.2	1.6*	4
	G2: Bladder training	47	3.1	1.4	1.7*	4
Burgio et al. ¹³⁵ 2002	G1: Multicomponent via biofeedback	73	2.2	0.9	1.3*	10
	G2: Multicomponent via verbal	74	2.5	0.9	1.6*	10
	G3: Self- administered	75	2.2	1.0	1.2*	10
Burgio et al. ¹⁴³ 1998	G1: Multicomponent	65	2.3	0.4	1.9^	10
	G2: Oxybutynin (range of doses)	67	2.3	0.8	1.5^	10
	G3: Placebo	65	2.2	1.2	1.0^	10

Table 22. Outcomes of behavioral treatment trials (continued)

Author Year	Groups	N	Per day baseline	Per day on treatment	Decrease in episodes	Weeks at evaluation
Episodes of incontinence per day (continued)						
Wyman et al. ¹³⁴ 1998	G1: Bladder Training	68	2.0	0.9	1.1†	12
	G2: Pelvic muscle exercise	64	3.0	1.7	1.3†	12
	G3: Combination therapy	61	2.3	0.8	1.5†	12
Voids per day						
Arruda et al. ²⁰¹ 2008	G1: Oxybutynin	22	7.7	6.4	1.3	12*
	G2: Functional electrostimulation	21	8.6	7.9	0.7	12*
	G3: Pelvic floor training	21	6.8	7.1	0.3^	12*
Wang et al. ⁹³ 2006	G1: Electrical Stimulation	25	12.8	7.8	5.0^	12
	G2: Oxybutynin 2.5 mg 3x/day	26	11.5	7.4	4.1^	12
	G3: Placebo	23	11.5	10.0	1.5^	12
Bryant et al. ¹³⁶ 2002	G1: Bladder training + caffeine reduction	48	11.1	6.8	4.3^	4
	G2: Bladder Training	47	11.2	7.9	3.3^	4
Goode et al. ¹⁴⁵ 2002	G1: Multicomponent behavioral ± biofeedback	65	10.0	8.2	1.8	10
	G2: Oxybutynin (range of doses)	67	10.9	8.8	2.1	10
	G3: Placebo + bladder diary	65	10.0	9.7	0.3	10

* Not significant differences between groups. ^p<0.05 † Significant comparisons BT vs. CT and PME vs. CT

One study evaluated the ability of biofeedback to improve outcomes associated with multicomponent behavioral therapy,¹³⁵ relative to providing the training with verbal feedback. The intervention included multicomponent behavioral training with either biofeedback or verbal feedback, compared to a self-help booklet. Multicomponent behavioral training combined bladder training techniques (e.g., relaxation approaches in the presence of urge, extending time between voids) with pelvic muscle exercises. The interventions took place over an eight-week period with four visits, followed by a two-week bladder diary. Reductions (mean: 58.6 to 69.4 percent) in episodes of incontinence were seen in all groups, with median percentage reductions ranging from 70.4 (IQR: -29.4, 100) to 82.8 (IQR: 0, 100). The wide IQR makes this finding somewhat difficult to interpret, and overall there were no by-group differences (p=0.23). Nonetheless, patients' perceptions of treatment benefit differed (see section below on patient-reported outcomes) with patients in the self-administered training significantly less satisfied (p=0.001).

Burgio et al¹³⁵ also compared their multicomponent behavioral approach to oxybutynin and placebo (study described in KQ3) and observed reductions in incontinence episodes of 81 percent among those in the behavioral group relative to 39 percent in the placebo arm.

Wyman and colleagues attempted to separate the role of pelvic muscle strengthening from bladder drill, and assess the potential for a combined impact. Combining the approaches provided the greatest reduction in incontinence episodes immediately after the 12 week intervention ($p=0.050$), but the difference did not persist at three months ($p=0.587$).¹³⁴

Only one study in this body of literature included vaginal electrical stimulation.²⁶⁹ In this study of pelvic floor muscle training with or without biofeedback compared to electrical stimulation, about half of women reported subjective improvement or cure in OAB when treated with electrical stimulation or biofeedback assisted pelvic floor muscle exercises, compared to 38 percent of women instructed in pelvic floor muscle exercises and told to perform them at home.²⁶⁹

Frequency outcomes. Number of voids per day was a common, objective measure of treatment effectiveness for behavioral interventions – particularly as a key element of the training is generally encouraging patients to extend time between voiding progressively, with a goal of reaching three to four hours between voids. Bryant found greater reduction in frequency when a caffeine reduction component was added to bladder training (reduction of 4.3 versus 3.3; $p=0.037$).¹³⁶ Women who simultaneously reduced caffeine intake had a 35 percent decrease in number of voids per day compared to 25 percent for those using bladder training alone. In the study described above of multicomponent behavioral intervention compared to placebo (and oxybutynin), frequency of micturition was reduced significantly in the behavioral arm, but not the placebo arm with a reduction of 1.8 micturitions per day.

Urodynamic outcomes. Three studies examined urodynamic measures pre- and post-treatment.^{131, 133, 135} In all of the studies, increased bladder capacity was seen along with changes in incontinence and frequency measures.

Pelvic muscle strength. Wang and colleagues²⁶⁹ found that biofeedback associated pelvic floor muscle exercises resulted in greater change in muscle strength than electrical stimulation, but the clinical significance of the change was not examined.

Patient reported outcomes. The nature of OAB is such that while specific morbidities and mortality are not primary outcomes, the interference that OAB creates in patients' lives, along with embarrassment and stigma, is often the concern that results in treatment seeking. Therefore, characteristics that make up and affect quality of life are of concern for studies of women with OAB, and are the focus of the results presented below. Four studies presented data on patient-reported outcomes other than urgency and frequency or cure/improvement.^{130, 132, 134, 135}

Wyman¹³⁴ examined the potential impact of combining pelvic muscle exercises with bladder training in an RCT of 204 women, of whom 59 had detrusor instability (results for the 145 women with SUI are not reported here as they are not relevant to this report). Although both impact (measured by the IIQ-R) and quality of life (measured by the UDI) were most improved in the combination group, relative to pelvic muscle exercises alone immediately after treatment, those effects were not sustained three months later. The combination group also had greater perceived improvement immediately after treatment, but again, no differences by group were sustained.

In the Burgio study¹³⁵ comparing biofeedback to verbal feedback and a pamphlet to teach multicomponent behavioral training, both biofeedback and verbal feedback performed equally well on all measures of patient perceptions of improvement, with verbal feedback better than the self-help booklet on five measures (accidents are smaller, $p<0.006$; comfortable with treatment, $p=0.01$; description of progress, $p<0.001$; satisfaction with progress, $p<0.001$; and restriction of

activities, $p=0.002$). The biofeedback group performed better on three measures (description of progress, $p<0.001$; satisfaction with progress, $p=0.03$ and restriction of activities, $p=0.002$). The biofeedback and verbal feedback groups saw no significant differences. All three groups had significantly improved quality of life scores (per Hopkins Symptom Checklist and Incontinence Impact Questionnaire), with no by-group differences. Similarly, Wang and colleagues²⁶⁹ found that biofeedback produced greater change in the overall Kings Health Questionnaire score among the group with biofeedback assisted pelvic floor muscle training relative to those who received PFMT without biofeedback or those who received electrical stimulation. Quality of life and satisfaction outcomes are summarized in Table 23.

Table 23. Effects on quality of life and satisfaction of behavioral treatments

Author Year	Comparison Groups	Outcomes
Burgio et al. ¹³⁵ 2002 RCT	G1: Behavioral training with biofeedback G2: Behavioral training with verbal feedback G3: Self-administered behavioral training	<ul style="list-style-type: none"> • 86% of those who received behavioral training with verbal feedback were completed satisfied relative to 75% of those receiving biofeedback and 56% in the self-administered group • Significantly more participants in the verbal feedback (89%) and biofeedback (79%) groups reported smaller accidents than in the self-administered group (68%) ($p=0.02$) • Almost all participants in the verbal (100%) and biofeedback (98%) groups were “comfortable enough” to continue treatment indefinitely, relative to self-administered (89%) ($p=0.009$)
Dorey et al. ¹³⁰ 2006 Case Series	G1: Pelvic muscle exercises	<ul style="list-style-type: none"> • 64 of 75 reported pelvic muscle exercises were useful • Severity of UUI at discharge was reported to be severe in four (6%) and moderate in 12 (18%) participants, relative to 32 (37%) and 40 (46%) at baseline, respectively
Dowd et al. ¹³² 2003 Cohort	G1: Bladder health information G2: Bladder health information + cognitive strategies (CS) G3: Bladder health information + CS + coaching	<ul style="list-style-type: none"> • Persons in G1 and G3 saw modest gains in UFICQ scores over time; G2 did not. Significant group-by-type of UI interaction: G1 and G3 with urge had more improvement than participants with stress or other UI ($F=3.61$; $p=0.037$)
Wang et al. ²⁶⁹ 2004 RCT	G1: Pelvic floor muscle training (PFMT) G2: Biofeedback assisted PFMT G3: Electrical Stimulation (ES)	<ul style="list-style-type: none"> • Significantly greater changes on the KHQ were observed for both biofeedback assisted PFMT and ES compared to PFMT alone ($p=0.003$), but not between biofeedback assisted PFMT and ES • Significantly greater changes were observed for ES compared to both other interventions on two specific domains of the KHQ: emotions and severity
Wyman et al. ¹³⁴ 1998 RCT	G1: Bladder training (BT) G2: Pelvic muscle exercise (PME) G3: Combination therapy (CT)	<ul style="list-style-type: none"> • Immediately after treatment, the combination group showed significantly more improvement in quality of life measures on the UDI ($p=0.054$) but the improvement was not sustained at three months ($p=0.126$) • Changes in life impact as measured by the IIQ-R was greatest in the combination group immediately after treatment ($p=0.03$), but not at three months ($p=0.85$) • Satisfaction levels were highest among the combination group immediately after treatment (82% very satisfied, relative to 73% in the PME group and 64% in the BT group); however, differences were not statistically significant ($p=0.096$)

Complementary and Alternative Therapies

Complementary and alternative therapies span a broad range of which only a small subset of modalities were used in studies of OAB. Acupuncture is an ancient Chinese medical system based on the balance of subtle energy flows (chi) in which imbalance of energy flows can result in disease. Acupuncture therapy aims to manipulate these energies through the insertion of fine needles at key, highly specific points related to chi flow to specific organs for varying periods of time. Foot reflexology is a variation of acupressure that postulates all body organs have corresponding external “reflex points on the foot” and the manipulation of these points can enhance the flow of energy to the reference organ. Specific areas of the sole of the foot are treated for specific medical conditions or symptoms. Hypnotherapy involves direct suggestion of symptom removal through therapeutic relaxation. Treatment of OAB is aimed at reduction of the component symptoms of urgency, frequency, and UII.

We identified three publications that used complementary and alternative medicine therapies to treat OAB: a fair quality trial of acupuncture,¹³⁷ a fair quality trial of foot reflexology,¹³⁸ and a poor quality prospective case series of hypnotherapy.¹³⁹ See complete evidence tables in Appendix C.

Acupuncture. The acupuncture trial was conducted in Oregon at an academic teaching center and was notable for incorporating sham acupuncture treatment as the comparison group.¹³⁷ Among 85 women randomized, 74 (87 percent) completed all four weekly treatment sessions and had complete outcome data at two to four weeks after treatment. Outcomes included comparison of baseline and post-treatment three-day voiding diaries as well as cystometrics, measurements of post-void residuals, the Urinary Distress Inventory, and the Incontinence Impact Questionnaire.

Episodes of urge urinary incontinence were statistically equivalent across groups at completion of four weeks of treatment. Number of voids per day were reduced 14 percent in the acupuncture group compared to 4 percent in the sham treatment group ($p=0.03$). This equated to a reduction of 1.4 voids per day among those receiving acupuncture. The experience of symptoms of urgency was reduced 30 versus 3 percent, with those treated having 1.6 fewer distinct episodes of awareness of urgency per day ($p<0.02$).

Some measures, including functional bladder capacity and cystometric maximum capacity, were modestly improved in the treatment group ($p<0.05$); others, including volume at urge to void and detrusor contractions during cystometry, were comparable across groups with no apparent trends. Scores improved meaningfully on both validated instruments that evaluate distress and impact on quality of life, with statistical significance.¹³⁷ No subjects withdrew for adverse events and treatment was well tolerated.

Reflexology. The study of foot reflexology was conducted in an academic center in Hong Kong. They incorporated sham reflexology in the form of a nonspecific foot massage without deep pressure. Among 120 women randomized, 97 (81 percent) completed all treatments and the assessment at three weeks. The reflexology group had three of 60 drop out; the sham group had six of 60 drop out. Losses were related to various competing demands, including four individuals who reported fear of SARS which was a threat during the study period in Hong Kong, with five losses to followup for other personal or medical reasons. There were no withdrawals because of discomfort or complications of treatment. Outcomes included comparisons of baseline and followup 24-hour voiding diaries and the King’s Health Questionnaire.

At completion, number of urge incontinence episodes, urgency episodes, and nocturnal voids were equivalent across groups. Daytime voids were reduced by 1.9 voids in the reflexology treatment group compared to 0.55 in the sham massage group ($p=0.03$). Quality of life measures did not differ. The authors note that their participants may have been unmasked by their familiarity with what to expect from reflexology treatments: as 88.9 percent of those in the reflexology group and 67.4 percent of those in the sham massage group believed that they had received “true” reflexology. This difference in unblinding could bias the findings.

Hypnotherapy. The hypnotherapy study was conducted in a UK academic setting and followed 63 women who were prospectively enrolled, had urodynamics, received 12 weeks of weekly hypnotherapy, and had followup urodynamic evaluation. Descriptive information is provided with little statistical analysis.¹³⁹

Ten of 63 participants discontinued hypnotherapy before completing all sessions. Of those completing all sessions, 29 were reported to be entirely free of OAB symptoms with 14 “considerably improved”. Of 44 women who had repeated urodynamics, 22 initially classified as having unstable bladder “converted to stability”, other improvements in cystometrics were also reported. This case series lacks masking of assessors and does not provide key patient reported outcomes.

In summary, a well-conducted small trial of acupuncture has intriguing results related to decreased frequency of voiding and reduced symptoms of urgency which are associated with changes in cystometrics related to improved bladder capacity that are logical intermediates of the improvement in symptoms. Women in the study felt they were improved as measured by scales that capture bother and quality of life. This evidence is insufficient to support definitive choice of acupuncture but offers preliminary information that promises modest improvements that are similar to those reported in many pharmacologic trials.

Reflexology is represented by a small trial with unmasking of participants that could have biased the results. There is not evidence to support choice of this modality. Likewise, hypnotherapy is not supported by the scant information provided by one case series with little detail, patient reported outcomes, or statistical assessment. Given the scope of placebo effects demonstrated in other well-conducted studies of OAB treatment, it is difficult to know whether to attribute any effect to hypnotherapy.

KQ 3: Comparisons of Treatments

Direct comparisons of treatments are made between entire approaches to management of OAB (e.g., pharmacologic to behavioral) or within approaches (e.g., drug to drug). In this section we present the results of any direct comparisons of either type, beginning with comparisons within the pharmacologic approach, followed by procedures compared to medical therapy or one another, and finally any comparison involving behavioral approaches. This third group comprises three distinct subgroups: behavioral compared directly to pharmacologic; combination pharmacologic plus behavioral compared to pharmacologic alone, and combination behavioral plus pharmacologic compared to behavioral alone.

Comparisons between pharmacologic treatments

All trial arms for RCTs of pharmacologic treatment for OAB are presented in evidence tables in Appendix C. Specific comparisons have been made in the literature for the following pairs of drugs that describe differences in reduction in urge urinary incontinence or voids per day:

- Oxybutynin ER to Tolterodine ER⁸⁴
- Oxybutynin ER to Tolterodine IR⁸³

- Oxybutynin IR to Tolterodine IR^{89, 91, 94}
- Oxybutynin IR to Darifenacin⁸²
- Oxybutynin IR to Trospium IR⁹⁰
- Oxybutynin TDS to Tolterodine ER⁸⁸
- Tolterodine ER to Tolterodine IR^{140, 141}
- Tolterodine ER to Solifenacin⁹⁹
- Tolterodine ER to Fesoterodine^{96, 142}
- Tolterodine IR to Solifenacin⁹⁷

These studies are generally powered simplistically only to assess non-inferiority; studies would need to be much larger for full assessment of comparability at robust power for small differences between pharmacologic agents. Nonetheless, in the majority of comparisons, neither drug was reported more effective at reducing either urge urinary incontinence episodes or voids per day with a few exceptions (Tables 24 and 25).

Both oxybutynin and tolterodine in their extended release forms demonstrated superiority in reducing incontinence episodes over tolterodine immediate release.^{83, 140} In the OBJECT trial, oxybutynin 10 mg ER was compared to tolterodine 2 mg IR twice a day.⁸³ At the end of 12 weeks, women taking oxybutynin reduced their episodes per week of urge urinary incontinence from 25.2 to 6.2, compared to a change from 25.1 to 8.5 in the tolterodine arm. The difference of 2.4 episodes per week between groups was statistically significant. However, upon stratifying by age group, the difference was maintained only among those age 64 and younger. Two studies compared the effectiveness of tolterodine 4 mg once per day to tolterodine 2 mg, taken twice per day, and found that the extended release formulation resulted in significantly greater reductions in incontinence episodes.^{140, 141}

Sand and colleagues found that voids per week diminished from 91.7 to 68.0 in the oxybutynin 10 mg extended release arm, compared to 91.6 to 71.2 in the tolterodine 2 mg immediate release twice a day arm ($p=0.024$).⁸³ However, as with the difference observed for incontinence episodes, upon stratifying by age, the difference was maintained only among those 64 years and younger. Diokno and colleagues (2003)⁸⁴ observed greater reductions in voids per week ($p=0.05$) in women taking oxybutynin 10 mg ER compared to those taking tolterodine 4 mg ER in the OPERA trial. Both studies provided treatment for 12 weeks, and data were obtained via bladder diaries. Harms were rare in both studies, although Diokno and colleagues report significantly higher rates of dry mouth with oxybutynin ($p=0.02$).

No other comparisons yielded statistically significant differences in terms of our primary outcomes.

Table 24. Direct comparisons of pharmaceutical treatments on urge incontinence

	Oxybutynin, ER	Oxybutynin, IR	Tolterodine, ER	Tolterodine, IR	Solifenacin	Darifenacin	Trospium, ER	Trospium, IR	Fesoterodine	Oxybutynin TDS
Oxybutynin, ER										
Oxybutynin, IR										
Tolterodine, ER	=									
Tolterodine, IR	O>T^	=	ER>IR*							
Solifenacin			=	=						
Darifenacin		=								
Trospium, ER										
Trospium, IR		=								
Fesoterodine, IR			=							
Oxybutynin, TDS			=							

Equivalence assigned if shown to be statistically insignificant or outcomes similar and no significance testing.

^Oxybutynin ER superior for this outcome to Tolterodine IR⁸³

*Tolterodine ER superior for this outcome to Tolterodine IR¹⁴⁰

Table 25. Direct comparisons of pharmaceutical treatments on voids per day

	Oxybutynin, ER	Oxybutynin, IR	Tolterodine, ER	Tolterodine, IR	Solifenacin	Darifenacin	Trospium, ER	Trospium, IR	Fesoterodine	Oxybutynin TDS
Oxybutynin, ER										
Oxybutynin, IR										
Tolterodine, ER	O>T*									
Tolterodine, IR	O>T^	=	=	=						
Solifenacin			=	=						
Darifenacin		=								
Trospium, ER										
Trospium, IR		=								
Fesoterodine			=							
Oxybutynin TDS			=							

Equivalence assigned if shown to be statistically insignificant or outcomes similar and no significance testing.

*Oxybutynin ER superior for this outcome to Tolterodine ER⁸⁴

^Oxybutynin ER superior for this outcome to Tolterodine IR⁸³

Comparisons between procedural and pharmacologic treatments

The only procedure to be compared to another treatment modality was sacral neuromodulation, which was compared to medical therapy in one RCT. In this study, 98 participants refractory to medical therapy were randomized to immediate sacral nerve stimulation or delayed sacral nerve stimulation. The delay group continued unspecified medical management for a six month period before having the procedure. The study found a reduction in daily urge urinary incontinence episodes from 9.7 to 2.6 in the sacral neuromodulation group, compared to an increase from 9.3 to 11.3 in the medical management group at six months ($p < 0.01$).¹²⁴ At 18 months, 76 percent of patients reported that they were completely dry or had experienced a reduction in symptoms of 50 percent or greater. It is important to note that those receiving medical therapy knew they were awaiting treatment with a modality that they were invested in believing was superior to their current level of symptom management. The differences in risk between sacral neuromodulation and medical management are important. Six patients had permanent explantation: three for pain, two for infection, and three for change in bowel function.

Comparisons between behavioral and pharmacologic treatments

Nine studies, with 11 publications, included behavioral and pharmaceutical arms in direct comparison to one another.^{93, 143-150, 201, 254}

The literature base included one prospective cohort study,¹⁴⁷ and eight RCTs.^{93, 133, 143-146, 148, 150}

Three of the studies were conducted in Europe, two were in the United States, two were in Asia (Taiwan and Korea), one in Brazil, and one in New Zealand. Seven were conducted at academic medical centers; two were in community settings.

The behavioral approaches examined included bladder training,^{146-150, 254} multicomponent behavioral approaches,¹⁴³⁻¹⁴⁵ pelvic floor training,²⁰¹ and electrical stimulation (Table 26).⁹³

Table 26. Direct comparisons between pharmacologic and behavioral interventions

Author Year Design	Groups	N	Episodes per day baseline	Episodes per day on treatment	Decrease in episodes per day	Weeks treated
Urge incontinence per day						
Arruda et al. ²⁰¹ 2008	G1: Oxybutynin 5 mg b.i.d.	22	2.0	1.0	1.0	12*
	G2: Electrical stimulation	21	1.9	1.1	0.8	12*
	G3: Pelvic floor training	21	2.3	1.1	1.2	12*
Lauti et al. ²⁵⁴ 2008	G1: Oxybutynin	21	2.2	0.8	1.4	12*
	G2: Bladder retraining	16	1.0	0.1	0.9	12*
	G3: Combination	19	1.8	0.6	1.2	12*

Table 26. Direct comparisons between pharmacologic and behavioral interventions (continued)

Author Year Design	Groups	N	Episodes per day baseline	Episodes per day on treatment	Decrease in episodes per day	Weeks treated
Urge incontinence per day						
Burgio et al. ¹⁴³ 1998	G1: Multicomponent behavioral ± biofeedback	65	2.3	0.4	1.9 [^]	10
	G2: Oxybutynin (range of doses)	67	2.3	0.8	1.5 [^]	10
	G3: Placebo	65	2.2	1.2	1.0 [^]	10
Wang et al. ⁹³ 2006	G1: Electrical stimulation	25	1.0	0.5	0.5 [*]	12
	G2: Oxybutynin (2.5 mg 3x/day)	26	0.0	0.0	0.0 [*]	12
	G3: Placebo	23	1.0	1.0	0.0 [*]	12
Voids per day						
Arruda et al. ²⁰¹ 2008	G1: Oxybutynin (5 mg b.i.d.)	22	7.7	6.4	1.3	12 [*]
	G2: Electrical stimulation	21	8.6	7.9	0.7	12 [*]
	G3: Pelvic floor training	21	6.8	7.1	0.3 [^]	12
Lauti et al. ²⁵⁴ 2008	G1: Oxybutynin	21	7.8	6.7	1.1	12 [*]
	G2: Bladder retraining	16	8.0	6.3	1.3	12 [*]
	G3: Combination	19	8.4	6.7	1.7	12 [*]
Song et al. ¹⁵⁰ 2006 RCT	G1: Bladder training	46	10.9	8.1	2.8 ^{*^}	12
	G2: Tolterodine (2 mg qd)	47	11.6	8.1	3.5 [*]	12
	G3: Tolterodine + bladder training	46	11.9	7.9	4.0 [^]	12
Wang et al. ⁹³ 2006	G1: Electrical stimulation	25	12.8	7.8	5.0 [^]	12
	G2: Oxybutynin (2.5 mg 3x/day)	26	11.5	7.4	4.1 [^]	12
	G3: Placebo	23	11.5	10.0	1.5 [^]	12
Goode et al. ¹⁴⁵ 2002 RCT	G1: Multicomponent behavioral ± biofeedback	65	10.0	8.2	1.8 [†]	10
	G2: Oxybutynin (range of doses)	67	10.9	8.8	2.1 [†]	10
	G3: Placebo	65	10.0	9.7	0.3 [†]	10

* Not significant differences between groups; [^]p<0.05; [†]significance not reported.

Episodes of Incontinence. Two RCTs considered urge urinary incontinence episodes as a primary outcome.^{93, 143} One compared multicomponent behavioral interventions to pharmacologic interventions¹⁴³ and reported on changes in incontinence episodes. This study was a three-arm study in which multicomponent behavioral treatment was compared to pharmacologic treatment and to placebo. The behavioral arm had a significantly higher percent reduction in episodes of incontinence at the ten-week followup ($p < 0.001$) (80.7 percent compared to 68.5 percent for pharmacologic and 39.4 percent for placebo). In a pilot study comparing oxybutynin to bladder retraining, no difference was observed in effectiveness by group.²⁵⁴ The authors calculated that to observe a difference between these arms in a full-scale study would require 165 women in each arm, rather than the approximately 20 in the pilot. This study also had a combination therapy arm. Two studies examined the effects of electrical stimulation with different comparison groups.^{93, 201} One⁹³ found no difference in reduction of incontinence comparing electrical stimulation to oxybutynin or to placebo either within or by groups. The other²⁰¹ showed no difference in effectiveness between three groups: electrical stimulation, oxybutynin (5 mg b.i.d.) or pelvic floor exercises.

Cure. Three studies reported on numbers of patients who achieved “cure” or resolution of UUI, without further definition.^{146, 147, 149} The results were inconsistent, the studies were of poor quality, and all were conducted prior to the ICS definition of OAB. The behavioral intervention in Jarvis, 1981¹⁴⁹, was inpatient bladder drill, which is not a current treatment approach.¹⁴⁹ Diokno (1995) is a report on a series of patients who chose their own treatment modality.¹⁴⁷ Only Colombo (1995) in this series was an RCT.¹⁴⁶ Bladder training was provided over a six-week period on an outpatient basis; although cure rates at the end of treatment were essentially the same, after six months, those who had received bladder training maintained a higher cure rate (96 percent of those initially cured in the bladder training group versus 57 percent in the oxybutynin group). However, the numbers reported in this study were quite small (only 53 in all groups at six months followup).

Frequency. Of the six studies that provided data on voids per day, all found that both pharmacologic and behavioral approaches could reduce frequency, but that there was no difference between the two approaches.^{93, 145, 148, 150, 201, 254} Goode and colleagues conducted a secondary analysis of the original Burgio and colleagues 1998 study. They used structural equation modeling to determine that changes in voiding frequency were not mediating factors associated with decreases in incontinence.¹⁴⁵

Similarly, both bladder training approaches (either multicomponent or bladder drill) and pharmacologic treatment were associated with urodynamic changes, including increased maximum cystometric capacity overall and at first and strong desire to void. These changes were statistically significant, and large enough to be clinically relevant as well. However, Goode and colleagues¹⁴⁵ examined the potential role of urodynamic changes to mediate the perceived effects of behavioral and pharmacologic treatment on incontinence and found, once again, that they did not seem to be associated with observed reductions in incontinence.¹⁴⁵

Patient reported outcomes. Burgio and colleagues used the Hopkins Symptom checklist to consider psychological changes potentially associated with improvement in continence.¹⁴⁴ Behavioral management and individually titrated oxybutynin were both associated with improvement in psychological status overall (including the placebo condition) and on a range of subscales, but psychological changes measured on the Hopkins Symptom checklist did not correlate with changes in rates of incontinence episodes. Satisfaction was highest among patients who received behavioral management (77.6 versus 54.7 percent with oxybutynin); interestingly, satisfaction on placebo was not substantially different than that on drug (43.1 percent). A very high proportion of the women receiving multicomponent behavioral training felt that they were

“comfortable enough” to continue with the approach (96.5 percent) relative to those on drug (54.7 percent) or placebo (43.1 percent).

Comparisons of combined behavioral and pharmacologic treatment to pharmacologic treatment alone

Seven studies examined the effect of adding a behavioral intervention to drug compared to drug alone, one of which was a feasibility study.^{150-155, 254} In all but two studies, the drug was tolterodine. The literature included six RCTs^{143, 150, 151, 154, 155} and two randomized open-label trial (Table 27).^{153, 254}

Two of the studies were conducted in the United States, one was in Canada, one was in multiple Scandinavian countries, one in Korea, one in New Zealand, and one did not specify, but indicated that it took place internationally in multiple sites. The two United States studies, the Korean one, and the one in New Zealand were conducted at academic medical centers.

Burgio and colleagues (2008) provided multicomponent behavioral treatment as an adjunct to pharmacologic treatment compared to pharmacologic treatment alone¹⁵² to examine the potential for behavioral management to aid patients in ceasing medication use and staying off. The primary endpoint of interest was a combined effect of a 70 percent reduction in incontinence plus no medication use or other therapy for incontinence. This study examined outcomes at 10 weeks, immediately after treatment, and at 8 months after a period of no treatment to examine persistent effects. During the post-treatment period, participants could request a return to medication. The participants in the behavioral group reported a greater reduction in episodes of incontinence than those in the oxybutynin group at 10 weeks; however the difference did not persist to 8 months (ability to discontinue drugs was 41 percent in both groups at 8 months). The two groups also experienced similar percentage reductions in episodes of urge incontinence (20.4 percent in the behavioral group versus 18.5 percent in tolterodine alone). The behavioral group did report statistically significantly greater satisfaction or quality of life.

Table 27. Comparisons of pharmaceuticals with and without behavioral interventions

Author Year	Groups	N	Episodes per day baseline	Episodes per day on treatment	Decrease in episodes per day	Weeks treated
Episodes of urge incontinence per day						
Burgio et al. ¹⁵² 2008	G1: Tolterodine ER 4 mg+ behavioral	153	3.3	0.4	2.9†	10
	G2: Tolterodine	154	3.3	0.7	2.6†	10
Chancellor et al. ¹⁵³ 2008	G1: Darifenacin (ranges 7-15 mg qd) + behavioral	190	2.8	1.5	1.3*	12
	G2: Darifenacin	205	3.0	1.0	1.0*	12

Table 27. Comparisons of pharmaceuticals with and without behavioral interventions (continued)

Author Year	Groups	N	Episodes per day baseline	Episodes per day on treatment	Decrease in episodes per day	Weeks treated
Episodes of urge incontinence per day						
Lauti et al. ²⁵⁴ 2008	G1: Oxybutynin	21	7.8	6.7	1.1	12*
	G2: Bladder retraining	16	8.0	6.3	1.3	12*
	G3: Combination	19	8.4	6.7	1.7	12*
Millard et al. ¹⁵⁴ 2004	G1: Tolterodine 2 mg b.i.d. with PFME	227	3.4	1.3	2.1*	12
	G2: Tolterodine	253	3.2	1.1	2.1*	12
Mattiasson et al. ¹⁵¹ 2003	G1: Tolterodine 2 mg b.i.d. + BT	244	2.0	0.3	1.7*	24
	G2: Tolterodine	257	2.3	0.3	2.0*	24
Voids per day						
Burgio et al. ¹⁵² 2008	G1: Tolterodine ER 4 mg + behavioral	153	NR	NR	0.5†	10
	G2: Tolterodine	154	NR	NR	0.04†	10
Song et al. ¹⁵⁰ 2006	G1: Tolterodine 2 mg b.i.d. + BT	46	11.9	7.9	4.0 [^] *	12
	G2: BT	46	10.9	8.1	2.8 [^]	12
	G3: Tolterodine	47	11.6	8.1	3.5*	12
Herschorn et al. ¹⁵⁵ 2004	G1: Tolterodine + behavioral information	39	NR	NR	1.8†	16
	G2: Tolterodine	45	NR	NR	2.2†	16
Millard et al. ¹⁵⁴ 2004	G1: Tolterodine 2 mg b.i.d. + PFME	227	11.9	9.2	2.7†	12
	G2: Tolterodine	253	12.8	9.4	3.4†	12
Mattiasson et al. ¹⁵¹ 2003	G1: Tolterodine 2 mg b.i.d.+ BT	244	10.3	6.9	3.4†	24
	G2: Tolterodine	257	10.6	8.0	2.6†	24

* Not significant differences between groups. [^]p<0.05; † significance not reported; ↑ increase in episodes per day.

Three additional studies compared the effectiveness of combinations of drug and behavioral approaches to drug alone in changing episodes of incontinence as well as episodes of urgency per day.^{151, 153, 154} Differences between the groups were small and non-significant, although there were significant decreases within all groups. Episodes of urgency decreased in all groups as well, with decreases ranging from approximately 1.9 to 2.7 episodes per day, but again there were no differences between the study groups.

Two studies found that adding behavioral training to tolterodine was associated with further reductions in frequency compared to tolterodine alone;^{151, 152} two found no significant effect of

adding behavioral training.^{150, 155} However, the type of the training provided differed dramatically – for example, the intervention provided by Herschorn¹⁵⁵ was purely informational, while Burgio¹⁵² provided a multicomponent system that included biofeedback and pelvic floor muscle exercises. Per one study,¹⁵⁴ addition of pelvic floor exercises alone in addition to tolterodine immediate release (2 mg b.i.d.) did not confer added reductions in frequency.

In those studies that measured quality of life and participant satisfaction, improvements were significantly greater among those patients receiving combination therapy compared to those receiving pharmacologic therapy alone (Table 28).

Table 28. Effect on quality of life and satisfaction of combination treatment

Author Year	Comparison Groups, N	Quality of Life and Patient Satisfaction Outcomes
Burgio et al. ¹⁵² 2008	G1: Tolterodine ER 4 mg plus behavioral training G2: Tolterodine ER 4 mg	<ul style="list-style-type: none"> • 53% of participants in the combination group were completely satisfied at 10 weeks, compared to 40% in the drug only group (difference of 13, range 1–25) • At 8 months, 33% of the combination group were completely satisfied compared to 30% in the drug only group • OAB-q bother scores decreased by 36.7 at 10 weeks and 30.4 at 8 months in the combination group relative to 30.9 and 20.4 in the drug only group. The difference was statistically significant (p<0.0001) • Health related quality of life improved in both groups with only small differences between groups
Chancellor et al. ¹⁵³ 2008	G1: Darifenacin (7.5-15 mg qd) G2: Darifenacin + behavioral modification	<ul style="list-style-type: none"> • Improvement in total OAB-q scores and OAB-SAT-q at 12 wks, no difference between groups (numbers not provided)
Song et al. ¹⁵⁰ 2006	G1: BT G2: Tolterodine 2 mg b.i.d. G3: Tolterodine 2 mg b.i.d. + BT	<ul style="list-style-type: none"> • More participants in combination treatment group (71%) had improved satisfaction scores at completion compared to the drug group (63%) or the bladder training group (54%) (difference not significant) • Urgency scores were reduced by 60% in the combination group, relative to 62% in the drug group and 45% in the bladder training group. The difference between drug and combination was not statistically significant, but differences between bladder training and combination and between bladder training and drug were at p<0.05

Comparisons of combined behavioral and pharmacologic treatment to behavioral treatment alone

Three studies measured the effect of adding a pharmaceutical approach to a behavioral one. All three of these studies used bladder training as the behavioral technique.

Ghei and colleagues²⁵³ describe a series of cases in which patients chose management approaches that could include either bladder training alone, or bladder training in addition to an antimuscarinic agent. Only 52 of 708 patients chose the bladder training alone, and although they experienced greater reduction in frequency (p<0.0001), those in the combination group had greater reductions in incontinence episodes (p=0.024).

A retrospective chart review of 92 patients treated with bladder retraining drill, among whom 36 also received antimuscarinics, was reported by Fantl et al.²⁵² The outcome of cure, in this

clinical populations with 6 months to 6 years of followup, was defined as no further episodes of incontinence and voiding every 3 to 5 hours with no associated symptoms. Cure was achieved in 83.3 percent of patients using bladder training with antimuscarinics and 78.6 percent of patients using bladder training alone; this difference was not significant ($p>0.6$).

In a double-blind, placebo-controlled RCT, patients were randomized to receive placebo or oxybutynin 2.5 mg twice a day in addition to bladder training.²⁵⁵ The patients taking oxybutynin had a greater reduction in daytime frequency when compared to the patients taking placebo ($p<0.05$). There were no differences between groups in the change in incontinence episodes.

Finally, Burgio and colleagues²⁵¹ provided the opportunity to patients in their trial of biofeedback-assisted therapy versus oxybutynin¹⁴³ whose treatment was not completely successful from the patient perspective to receive combined pharmacologic and behavioral management. Of the 35 individuals who met criteria and agreed to move onto combined treatment, eight crossed from behavioral alone to combination, and 27 went from pharmacologic alone to combined. Both groups experienced significant reductions in incontinence over the effect of the initial treatment. The behavioral to combined group went from 58 percent reduction at the end of single therapy to 89 percent reduction after combined therapy for an additional eight weeks. The pharmacologic to combined group also improved from 73 percent reduction to 84 percent reduction in incontinence episodes.

KQ 4. Modifiers of Treatment Outcomes

This section includes information related to how individual characteristics may influence likelihood of responding to treatment and outcomes of treatment. We included publications that explicitly presented stratification by a baseline characteristic that can be determined in a clinical setting and that presented statistical analysis related to interpreting the influence of the characteristic on treatment effects as they relate to outcomes. Detailed analyses of personality characteristics or psychometrics that require specialized expertise or survey instruments unlikely to be used by those providing care for women with OAB are not reviewed here.

Identified Modifiers

Age. Eight publications examined the relationship of age to response to pharmacologic treatment.^{83, 102, 156-161}

Tolterodine was the focus of four of these studies;^{156-158, 160} and one compared oxybutynin to tolterodine.⁸³ The largest study of symptom-related outcomes was an open label clinical cohort with 2,250 patients from 462 urology practices in Germany. The mean dose received was 3.8 ± 1.2 mg, with a median of 2 mg. Average age of those treated was 61 ± 14 years with range not provided. Increasing age was associated with being more likely to have incontinence episodes which were not strictly required to be urge incontinence. In multivariable regression models, with age as a continuous variable, each year of increasing age was associated with small absolute reductions in global efficacy (OR for global efficacy 0.986; 95 percent CI: 0.98, 0.99 per year of age). Global efficacy was defined as eight or fewer voids a day, fewer than two urge episodes, and no incontinence episodes per day.¹⁵⁷

The largest RCT evaluated age effects among 1,015 participants and found that tolterodine drug was superior to placebo among patients aged younger than 65 compared to those 65 and older. They reported that treatment effects on incontinence episodes per week, voids per day, and subjective reports of experience of urgency and the ability to hold urine, were comparable across age groups at 12 weeks. A side-light of interest was that placebo effect was more pronounced for

reduction of voids per day in the younger group ($p < 0.045$); however change from baseline in the treatment arm was comparable regardless of age.¹⁵⁸

The IMPACT trial enrolled 896 individuals with urge incontinence from primary care settings and conducted an open-label evaluation of tolterodine extended release 4 mg once daily.^{160, 239} All parameters, including UUI, urgency, frequency, and nocturnal frequency, were improved at 12 weeks for both those younger than 65 and those older. However, decreases in frequency were less pronounced among the older group who experienced on average a 22.2 percent reduction in daytime frequency (95 percent CI: -26.7, -15.2) and 28.6 percent decrease in nighttime frequency (95 percent CI: -35.7, -20.0); while younger participants had a 33 percent decrease in daytime frequency (95 percent CI: -36.0, -30.3) and a 50 percent decrease in nighttime frequency (95 percent CI: -53.8, 40.0). The most common treatment-related adverse events among those younger than 65 were dry mouth (11.4 percent), constipation (2.7 percent), and dry eyes (1.0 percent); among those who were older dry mouth (6.6 percent), constipation (4.4 percent) and headache (2.2 percent) were most common. Retention (<1 percent) occurred only among those older than 75 and none of four individuals with this complication required catheterization. Statistical comparisons for harms by age were not provided.¹⁶⁰

A prescription-event monitoring study, conducted in the United Kingdom to assess population impact of tolterodine (range 1 to 4 mg) entering the prescription drug market, monitored more than 14,500 patients who filled prescriptions over a minimum of six months.¹⁵⁶ Average age was 63 ± 16 . Analysis of risks of rare adverse events (fewer than 40 events in the full population) found the upper quartile of age, those over 74, had greater risk of rare events including hallucination, heart palpitations, and tachycardia. Those under 50 had the lowest risk of cardiac events. No note was made of whether more common side effects varied with age.

A small comparative study of oxybutynin (10 mg once daily) and tolterodine (2 mg twice daily) ($n=315$) found an advantage for oxybutynin extended release over tolterodine twice daily for decreasing urge incontinence, urgency, and frequency, among those 64 and younger. This effect was not apparent in older age groups in which both were comparably effective.⁸³

Other pharmacologic treatments. A single placebo controlled RCT of trospium (20 mg twice a day) evaluated whether CNS adverse effects, specifically daytime drowsiness, varied with age. Using the Stanford Sleepiness Scale, they found fewer than 1.5 percent of those on trospium (and 2.5 percent of those on placebo) experienced a clinically relevant three or more point increase. Using continuous scores, neither age grouping as <65 and older or <75 and older revealed meaningful differences. Average changes in scores across groups were most often improvements of less than half a point, generally less than a quarter point.¹⁶¹ Darifenacin has been studied among those age 65 and older in a two-year, open-label extension of 716 participants, that documented comparable effectiveness among older and younger participants with respect to sustained or improved treatment response over time as defined by a global response score, and individual measures that included incontinence episodes, voiding frequency, urgency, and OAB-related nocturnal waking.¹⁵⁹ All statistical testing for these outcomes across time points through 24 months had $p < 0.088$ for the comparisons with baseline status.¹⁵⁹ A trial that allowed dose adjustment of darifenacin over the course of the study found the mean age of those requiring dose adjustments “for additional efficacy” was equal to those who did not change dose.¹⁰²

Though some studies reported reduced efficacy for specific endpoints among older participants in their study populations, none reported complete lack of benefit among older populations.

Prior treatment. Seven publications investigated whether prior treatment with antimuscarinics predicted treatment response.^{85, 86, 102, 162-165} In three placebo-controlled trials of oxybutynin patch,⁸⁵ , and tolterodine IR,¹⁶² participants who had previously been on

antimuscarinics had comparable outcomes to those who were treatment naïve. The tolterodine study specifically commented on prior treatment failures, noting improvements in those who had failed prior treatments that were above placebo but not statistically significant; few participants were in this category.¹⁶²

In two drug-to-drug comparison trials, outcomes, including UUI, total incontinence episodes, and “perceived improvement of bladder condition,” were likewise reported to be comparable for treatment naïve and previously treated participants. These studies investigated oxybutynin ER (5 mg and 10 mg) and tolterodine ER (2 mg and 4 mg);⁸⁶ and oxybutynin ER (10 mg) and tolterodine ER (4 mg).¹⁶⁴ A nine-month open label study of tolterodine 2 mg twice daily found that 89 percent of individuals previously unable to tolerate oxybutynin, tolerated tolterodine well.¹⁶³ One study included participants who had not been on medications as well as those switching from oxybutynin immediate release to extended release. In a subanalysis of those switching from immediate to extended release, the total proportions continent at 12 weeks is similar to that presented for the whole study population; no statistical test related to this comparison is provided. Some who had been on IR dosing regimens had worsening of symptoms on comparable total doses of ER.¹⁶⁵ This study was small (n=256 spread across 16 centers without placebo comparisons) which hinders interpretation. In a trial that allowed dose adjustments, prior treatment was associated with higher rates of dose increase.¹⁰²

Baseline severity.

Presence, type, and severity of incontinence. Two studies contrasted those with UUI at baseline to participants without UUI. In an open-label study of tolterodine 3,824 participants with nine months of treatment, urge, frequency, nocturia, and OAB scales were similarly improved regardless of UUI baseline status.¹⁶⁶ The VOLT study of solifenacin was an open label study of flexible dosing.¹⁶⁷ Participants who had UUI at baseline and reported it was their most bothersome symptom, reported improvements in urgency, UUI, frequency, and nocturia, from “moderate to severe” to “very minor to some minor.” With the exception of nocturia, the point estimates for improvement in individual measures were better in the group with UUI at baseline, although not always statistically significantly better.¹⁶⁷ Severity of UUI was not significantly associated with improvement in UUI. Subjects with severe UUI (defined as >20 episodes per week) had 67.6 percent decrease in UUI episodes, compared with 71.4 percent in subjects with less severe UUI.¹⁶⁸

The response of those with UUI (n=552) was compared to those with urge-predominant mixed urinary incontinence (n=171) in a 16-week single blind trial of tolterodine that allowed dose adjustments from 1 mg twice daily to 2 mg twice daily. The authors reported cure rates were comparable: 44 percent of those with UUI achieved “dryness” and 39 percent of those with mixed urinary incontinence; 24.0 and 23.5 percent respectively had normalized voiding frequency (<8 voids per 24 hours).¹⁶⁹ Other studies included participants with urge-dominant mixed urinary incontinence, or failed to specifically address inclusion or exclusion of those with mixed urinary incontinence, and did not report assessing trial data for differential treatment effects.

Specific symptoms and outcomes. A single study of transdermal oxybutynin and tolterodine ER 4 mg reported on ability to improve frequency in the lowest quartile of frequency, which was those with <10 voids a day at baseline, and found that neither drug had a significant effect in that subgroup compared to placebo.⁸⁸ The IMPACT trial, an open label trial of tolterodine ER in primary care practices, documented improvements among all groups of participants whether the most bothersome symptom was daytime frequency, nocturnal frequency, urge urinary incontinence, or urgency. Statistical testing was not provided to determine if specific groups were more likely to benefit; the data presented suggests similar improvements in self-reported

improvement, symptom bother, coping, health-related quality of life, and AUA symptom index scores across all groups.²³⁹

Other authors used severity scales or single questionnaire items to group participants and to conduct sub-analyses by severity. In the German open-label, observational study of tolterodine 2 mg twice daily, patients with greater baseline symptom severity experienced greater magnitude of improvements. This trend was for global efficacy; individual component symptoms varied across groups and were not statistically significant. Overall, those with severe baseline symptoms were less likely than others to be symptom free at 12 weeks.¹⁵⁷ In the ACET study among those whose self-reported symptom scores were graded as moderate to severe, those on tolterodine ER achieved greater magnitude of improvements on a global improvement scale than those on oxybutynin ER; comparison across moderate-to-severe to less affected participants was not provided.⁸⁶

Urodynamic findings. Four publications related baseline urodynamic findings to outcomes of treatment.^{89, 170, 171, 173} Three of these studies did not identify urodynamic findings that predicted poor response or non-response to treatment. The finding of detrusor overactivity compared to its absence was not associated with outcomes in a case series of 365 women treated with bladder retraining and oxybutynin 2.5 mg twice a day. Both women with and without detrusor instability had comparable benefits from treatment, as measured by voids per day and incontinence episodes.¹⁷¹ Likewise in an RCT of duloxetine (an SSRI) versus placebo which showed treatment benefits for OAB with and without UI, the classification of participants as having detrusor overactivity or sensory urgency based on urodynamic findings was not statistically significant as an effect modifier or predictor of treatment outcomes.¹⁷⁰ The third study to group participants by urodynamic finding had group sizes (n = 6, 25, 36, 40) too small to make definitive assessments but suggested in the two larger groups that those with low volume and high pressure profiles had comparable results to those with low volume and low pressure profiles. The last case series, reporting on 111 women with OAB based on symptoms and urodynamic diagnosis of detrusor overactivity, found that those women who had involuntary detrusor contractions with provocative maneuvers like coughing, washing hands in cold water, and the sound of running water, were less likely to respond to treatment, which was not operationally defined by the authors, with tolterodine 4 mg a day than those who did not respond to provocation (p=0.0008).¹⁷²

A last study conducted urodynamics among 1,133 women of whom 132 met the criteria for detrusor overactivity “conforming” to the definitions of the ICS; the researchers then compared treatment outcomes among women with and without coital incontinence. Among women with DO treated with tolterodine ER 4 mg daily for 12 weeks, those women with coital incontinence at orgasm were more likely to be non-responders to treatment (“unimproved by self-report”) for their OAB (41.2 percent) than those without coital incontinence (17.0 percent), (p=0.23).¹⁷³

Other candidate modifiers.

Race and ethnicity. The VOLT study, an open label trial of solifenacin has resulted in two publications addressing race and ethnicity. The authors report these analyses were motivated by lack of literature that explicitly addresses whether minority groups achieve comparable outcomes to the overall study group; the publications focused on Hispanic and black participants in VOLT. They provide detailed symptom and outcome profiles and report that outcomes were similar to the larger cohort. However direct statistical comparison is not provided by race/ethnicity.^{205, 229}

A single three-arm trial of tolterodine ER 4 mg, tolterodine IR 2 mg twice daily, and placebo (n=1,235) reported that women who were above the mean for BMI (> 27kg/m²) were more likely to have UI at baseline but achieve comparable reductions in number of incontinence episodes, to those with lower BMI after 12 weeks of treatment.¹⁴⁰

Anterior vaginal wall prolapse. Anterior vaginal wall prolapse was the sole pelvic organ prolapse measure that we identified in the literature about treatment outcomes. Anterior vaginal wall prolapse was defined as descent to at least 1 cm proximal to the hymen (POP-Q measurement \geq stage IIa). Women with such prolapse in this clinical case series of women treated with 4 mg of tolterodine extended release formula once daily were more likely to show no improvement at 12 weeks (39.2 percent) compared to those without prolapse (14.1 percent) using a three-point scale of no change, improved, and cured ($p=0.0002$).²⁴³

Gender. Authors frequently reported that men, especially older men, fared less well in resolution of symptoms of OAB.^{85, 106, 156, 157, 174} This evidence review was focused on outcomes of treatment among women. However, in order to retain landmark studies we included a number of studies that enrolled men as long as the proportion of women in the study was 75 percent or more. This means that treatment effects may be attenuated when men are included.

KQ5. Costs of Overactive Bladder Treatment

This section presents the results of our literature search and findings about financial costs associated with treatment for overactive bladder. Direct medical costs, indirect medical costs, and lost productivity for individuals with OAB are significant issues, in part because these women may be at greater risk for comorbidities such as falls and fractures, urinary tract infections, depression, and skin conditions.^{188, 189} OAB symptoms can interfere substantially with work and other activities,²⁵⁰ affecting productivity. Nonetheless, the proportion of individuals with OAB who seek medical treatment for OAB has been estimated to be extremely low with possibly as few as 4 percent of individuals with symptoms seeking treatment.²⁴ Therefore, estimates of costs of care for OAB are likely a reflection of care provided to only a small proportion of a group that may be able to benefit from effective treatment.

Although several studies have estimated the cost of illness of urinary incontinence, few studies in the United States have estimated the cost of illness of OAB, and even fewer have focused on treatment costs, as specified for this review. In particular it is difficult to estimate costs related to OAB because it is a symptom-based syndrome with significant inter- and intra-individual variation in etiology and in symptoms. To answer the question of costs related to the treatments reviewed in this systematic review, we restricted our analysis to studies that included direct costs of treatment for OAB in the United States.

Detailed information on all studies related to financial costs associated with management of OAB can be found in Appendix C. Studies analyzed costs for at least one of the following cost categories:

- **Direct costs.** Depending on the bundle of treatment for OAB, costs may include PCP services, specialist care, skilled nursing facilities, home health care, and/or surgery. Direct costs include diagnostic costs, radiology costs, laboratory costs, treatment costs, and costs related to the consequences of OAB.
- **Indirect Costs.** Indirect medical costs include lost productivity and an informal caregiver's time.

Financial Costs

Content of the literature. We identified five studies on financial costs related to OAB that met criteria for inclusion.¹⁷⁵⁻¹⁷⁹ All studies included assessment of direct medical costs related to OAB, and two included costs due to lost productivity. One study additionally assessed financial implications for pain and suffering. Two additional studies do not meet criteria for measuring direct costs of treatment, but provide context related to health care utilization.^{24, 250}

The literature base included three analyses of administrative-claims databases: one using Medicare data; one using a large, private health plan affiliated with Ingenix; one using seven plans administered by UnitedHealth Group. These studies analyzed claims for ICD-9 codes they selected as indicative of OAB; some ICD-9 codes were unique to specific studies.

One study was a national telephone survey of community-dwelling adults¹⁷⁹ that was designed to measure prevalence in the community. A followup survey was sent to a selected set of age and sex-matched cases (with OAB) and controls to estimate treatment use, medication, routine care, OAB related consequences and work productivity.

One focused on community-dwelling adults, one on persons younger than 65 years of age who filled prescriptions for drug treatment of OAB, one on persons of any age with prescription-drug benefit who filled prescriptions for drug treatment of OAB, and one on persons who had failed tolterodine ER and sought alternative drug therapy. Each of the studies used a different definition of OAB (Table 29). Because Jensen’s study¹⁷⁷ represents the Medicare population in 1994 to 1995 and is therefore likely substantially out of date, we do not summarize the data here. Data are available in the evidence table (Appendix C).

Table 29. Study definitions used for cost determination

Author Year	Population	Definition of Costs	Definition of OAB (ICD-9 Codes)
Hu et al. ¹⁷⁹ 2003	National telephone survey of community-dwelling adults	Direct: Routine care Diagnostic costs Treatment costs Consequences of OAB Indirect: Lost productivity Informal caregiver’s time	Self report in the NOBLE study
Hall et al. ¹⁷⁶ 2001	Claims database of seven plans affiliated with UnitedHealth Group	Physician office visits Pharmacy claims Laboratory claims Radiology claims Outpatient hospitalizations Inpatient hospitalizations ER visits	(788.30, 788.31, 788.33, 788.34, 788.35, 788.36, 788.37, 788.39, 788.40, 788.41, 788.42, 788.43)
Nitz et al. ¹⁷⁸ 2005	Claims database associated with Ingenix	Hospital ER Physician outpatient visits Laboratory Other non-pharmacy	(595.3, 596.51, 597.81, 788.30, 788.31, 788.33, 788.34, 788.36, 788.39, 788.4, 788.41, 788.43)
Jensen et al. ¹⁷⁷ 2003	Medicare-claims database	Hospital inpatient care Outpatient care Physician services Laboratory and x-ray services Skilled nursing facility care Home health care services	(596.51, 596.59, 788.30, 788.31, 788.33, 788.41, 788.43)
Varadharajan et al. ¹⁷⁵ 2005	PharMetrics Patient-Centric database	Pharmacy claims Outpatient medical care Laboratory/diagnostic costs Inpatient costs and days in hospital	(596.5x [excluding 596.53, 596.54], 788.3x, 788.41, 788.43)

Costs of treatment. Only Varadharajan and colleagues provided expenditures on medications for OAB, comparing sets of matched patients who used either tolterodine or oxybutynin in their extended release form, or who used tolterodine ER compared to oxybutynin in its immediate release formulation. They did not provide estimates by gender. Costs of drug ranged from an average low of \$56 over 12 months for oxybutynin immediate release to a high of \$360 for extended release tolterodine. The extended release formulation of oxybutynin had intermediate costs at \$317. Tolterodine extended release was significantly more expensive over the course of treatment than both oxybutynin ER and oxybutynin IR ($p < 0.001$). These differences in part reflect higher number of prescriptions filled in the extended release groups over the study period compared to the immediate release group ($p < 0.0001$). Nonetheless, total healthcare costs, including those specifically related to OAB, were highest for users of oxybutynin immediate release compared to the extended release formulations ($p < 0.0001$).

Three studies examined total medical costs (not just treatment costs) for persons who filled prescriptions for OAB drug treatment (Table 30). These studies calculated the total healthcare costs for persons with OAB, per person per year by drug (PPPY). No studies did so by surgical or behavioral approach and only Varadharajan presented data for women only.¹⁷⁵

Table 30. Total cost differences in annual medical care among persons filling prescriptions for OAB drug treatment

Author Year	Groups	Cost Outcomes			
		Average PPPY Total Costs before treatment (\$), Unadjusted, Mean \$ cost \pm SD	Average PPPY Total Costs after treatment initiation (\$), Unadjusted, Mean \$ \pm SD	Statistical significance	Reference standard
Nitz et al. ¹⁷⁸ 2005	G1: Oxybutynin IR G2: Oxybutynin ER G3: Tolterodine ER	G1: 4956 \pm 14396 G2: 4146 \pm 8695 G3: 3349 \pm 6715	G1: 7083 \pm 39420 G2: 5980 \pm 13263 G3: 5074 \pm 11007	G1/G3: $P < 0.05$ G1/G2: NS G2/G3: NS	NR
Hall et al. ¹⁷⁶ 2001	G1: Tolterodine G2: Oxybutynin G3: Flavoxate or other OAB medication G4: No drug treatment for OAB	G1: 5004 G2: 5688 G3: 5352 G4: 2928	G1: 7020 G2: 7116 G3: 7380 G4: 5040	NR	NR
Varadharajan et al. ¹⁷⁵ 2005 (women only)	G1: Tolterodine ER* G2: Oxybutynin ER G3: Tolterodine ER* G4: Oxybutynin IR	NR	G1: 8303 \pm 18802 G2: 8862 \pm 18684 G3: 9975 \pm 42860 G4: 10521 \pm 22602	G1vG2: $p = 0.0109$ G3vG4: $p = 0.3612$	2004 US dollars

PPPY = per person per year; *Separate groups matched according to G2 and G4

In their followup survey of individuals with OAB, Hu and colleagues¹⁷⁹ collected data on a range of health care utilization measures, including pharmacologic and surgical treatment for OAB. They used several sources of cost data, including the Red Book, to assign the annual costs of treatment for OAB in the United States in 2000. Pharmacologic treatment costs for women in 2000 were estimated at approximately \$1.2 billion for women overall, with approximately equal

costs across age groups. OAB surgical costs among women in 2000 were estimated at approximately \$550 million.¹⁷⁹ Overall health care costs, including lost productivity for the nation were \$7.4 billion for women (approximately \$3.1 billion for those under 65).

This body of literature is particularly challenged by the varying methodology for identifying and defining patients with OAB. In part, this variability is a reflection of the overall literature on OAB, which also uses varying definitions. Most problematic for definitional purposes is whether and when authors included incontinence in any of its forms.

None of the cost papers qualifying for inclusion used data beyond 2002, when the ICS definition of OAB was put into place.

Chapter 4. Discussion

This chapter summarizes the strength of the evidence to address our key questions and then presents methodologic considerations and a discussion of the findings for each of our five key questions. We conclude with a discussion of the status of research, limitations of the current literature, and our recommendations for future research priorities.

Strength of Evidence

We have summarized the quality of individual studies in categories of good, fair, or poor (with grading explained in Chapter 2) for each key question or sub-question within the summaries below and the information is included on the evidence table for each study in Appendix C. The strength of the evidence for each question or sub-question is a broad assessment of the totality of the literature available to provide evidence on a specific question or sub-question.

To reiterate the strength grades, the levels of strength of evidence are as follows:

- I. **Strong:** The evidence is from studies of strong design; results are both clinically important and consistent with minor exceptions at most; results are free from serious doubts about generalizability, bias, or flaws in research design. Studies with negative results have sufficiently large samples to have adequate statistical power.
- II. **Moderate:** The evidence is from studies of strong design, but some uncertainty remains because of inconsistencies or concern about generalizability, bias, research design flaws, or adequate sample size. Alternatively, the evidence is consistent but derives from studies of weaker design.
- III. **Weak:** The evidence is from a limited number of studies of weaker design. Studies with strong design either have not been done or are inconclusive.
- IV. **No evidence:** No published literature.

As a global assessment of this literature, no treatments reach the level of strong evidence. For the majority of interventions, strength hovers between moderate and weak because of crucial study design and reporting flaws. When there is no evidence we have noted that.

Principal Findings and Considerations

KQ1: Prevalence and Incidence of OAB

Methodologic issues. Data related to prevalence and incidence of OAB and urge urinary incontinence is notably coherent given the immense technical challenges to comparability. Response rates vary widely across studies with no clear patterns relating type of questionnaire (phone, mailed, administered) or population to completeness of response. Most research teams documented exhausting conventional options for improving response rates and many presented thoughtful analyses of non-response. While estimates, especially of self-reported symptom complexes, are very sensitive to the operational definitions used, these measures were able to be compared when the authors provided information about details like wording of survey items and the required frequency of an event to meet criterion definitions. With some exceptions, such information provided sufficient context to provide a global and United States picture of OAB and urge incontinence.

Appendixes and Evidence Tables for this report are provided electronically at <http://www.ahrq.gov/downloads/pub/evidence/pdf/bladder/bladder.pdf>

Publications that appeared after the ICS consensus definitions had more similar results, even when they used operational definitions that differed from ICS. In general, greater detail was provided in the publications after the ICS standardization to support comparisons across studies. We found no evidence that studies supported by pharmaceutical companies returned higher estimates for similar measures, with the exception of a single United States publication reporting on a cluster of consumer surveys done in malls. The estimates from those surveys were high enough to be considered outliers and the operational definitions used to derive the estimates were not provided in order to determine how their results compared to others.³⁶ There was no readily apparent relationship between method of administration (e.g., mailed survey, administered, or other methods) and findings which is reassuring.

Summary of prevalence and incidence findings. Overactive bladder and urge urinary incontinence occur in women of all ages around the world. The type of healthcare system appears not to clearly relate to the proportion of women affected at any point in time, suggesting that universal access to care such as in the United Kingdom and Canada versus more socioeconomically correlated access in the United States, is not the driver of whether or not women currently report symptoms. Several factors work in synergy to determine whether a woman with OAB symptoms is counted among prevalent OAB cases: degree of bother of symptoms, which varies widely; individual decision to seek care, which is shown to be a low proportion of those affected (13 percent of nurses in the Nurses Health Study); availability of care; and the degree to which an individual woman is distressed by the symptoms or interprets them as a normal part of aging.

Conservatively estimated more than 10 to 15 percent of adult, community-dwelling women in the United States meet criteria for OAB and as many as 5 to 10 percent have urinary incontinence associated with urgency. OAB and continence status are not static; both OAB and urge incontinence can resolve, though the data suggest most women are affected for multiple years at minimum if they develop symptoms.^{64, 77} How often resolution is related to treatment, individual behavioral change, or the natural history of the condition is completely undocumented. Evidence to describe the relationship of UI or OAB with age is varied. The preponderance of the literature suggests that OAB risk is more directly associated with increasing age than urge incontinence. For both conditions the relationship may be complex, such that rates of increase in the proportion of women who are symptomatic are relatively consistent until a threshold age. The most probable threshold ages at which rates plateau are well past the average age of menopause and reported to fall between 60 and 75 years of age.

Because of the population-based design of these reports, often details like recent urinary tract infections or childbirth were not able to be taken into account, though future researchers could incorporate self-reported status in inclusion and exclusion for the studies as some of the strongest publications in this literature did. Likewise, future research on incidence and regression of symptoms would do well to attempt to account for intercurrent diagnosis and whether or not an individual has received treatment.

The epidemiology of OAB and urge incontinence lead us to conclude, as have others, that the conditions are common, occur across the lifespan, and that providers of all types – primary care, specialists, and advanced practice nurses and health educators – will be called on to advise patients and provide care.¹⁹⁶ Opportunities for detection and treatment in clinical settings should be frequent and attention should be addressed to the degree to which patients find the symptoms distressing including impact of self-image, quality of life, and sexual function.

KQ2: Outcomes of Treatments for OAB

Pharmacologic treatments.

Methodologic consideration for pharmacologic treatment studies. The content of this literature is predominantly of fair to poor quality and strength of the evidence is at best moderate, and improved only by the consistency with which medications for OAB are shown to have modest advantage over placebo. The duration of followup during the masked portion of trials is short, rarely longer than 12 weeks. Given the longevity of treatment likely to be used by patients this is concerning and limits the generalizability to clinical practice. No investigators were able to determine whether or not there is a time period after which individuals can discontinue treatment and still retain some or all of the benefits of treatment reduction. Statistical and methodologic concerns are addressed in detail in the Future Research Section of this chapter. After review and analysis of 110 studies, of which four were good quality, 75 fair and 31 poor, with 68 RCTs, the strength of the evidence for managing OAB with pharmacologic treatment is weak to moderate for short term outcomes and weak for long term outcomes and harms.

Findings from pharmacologic treatment studies. All pharmacologic treatments were effective at improving one or more OAB symptoms when compared to placebo. Reductions ranged from 0.9 to 4.6 in urge urinary incontinence episodes per day across all drug treatments and from 0.7 to 4.2 in voids per day. Study by study, extended release formulations achieved better effects than immediate release, although statistical significance varied. No one drug was definitively superior to others by preponderance of evidence, including comparison of newer selective agents to older antimuscarinics. As estimated by meta-analysis extended release forms (taken once a day) reduce UII by 1.78 (95 percent CI: 1.61, 1.94) episodes per day, and voids by 2.24 (95 percent CI: 2.03, 2.46) per day. Immediate release forms (taken twice or more a day) reduce UII by 1.46 (95 percent CI: 1.28, 1.64), and voids by 2.17 (95 percent CI: 1.81, 2.54). Of note, placebo reduces UII by 1.08 (95 percent CI: 0.86, 1.30), and voids by 1.48 (95 percent CI: 1.19, 1.71). Even in the context of small to moderate affect on symptoms, pharmacologic treatments were generally associated with increased quality of life and reductions in measures of impact or distress, compared to baseline and to placebo.

Findings reported in this review are consistent with three prior reviews.^{196, 270, 271} The most recent found: (1) antimuscarinics are efficacious compared to placebo, (2) mean decrease in UII episodes per day ranged from 0.4 to 1.1, (3) mean decrease in the number of voids per day ranged from 0.5 to 1.3, (4) every treatment, with two exceptions, was associated with greater risk of adverse events compared to placebo, and (5) improvements were seen in quality of life.²⁷¹ This review added an additional 28 studies and incorporated evidence from study designs other than randomized clinical trials.

Table 31 below provides estimates of treatment effects for pharmacologic treatments represented by more than one trial arm. Some drugs and doses of drugs are not reported because the publications with trial arms for that treatment did not provide sufficient information to estimate variance in meta-analysis models. The models required that we have some estimate of the variance of the effect size such as standard deviation, standard error, or confidence bound, in order to achieve appropriate estimates.

Table 31. Estimates of mean reductions in incontinent episodes and voids per day

Drug	Decrease in Incontinent Episodes per Day			Decrease in Voids per Day		
	Estimate	95% CI		Estimate	95% CI	
Single drug estimates						
Placebo	1.08	0.86	1.30	1.48	1.19	1.71
Oxybutynin IR	1.49	1.18	1.80	2.18	1.75	2.61
Oxybutynin ER	*	*	*	*	*	*
Tolterodine IR	1.45	1.24	1.66	2.19	1.76	2.61
Tolterodine ER	1.75	1.65	1.85	2.48	1.94	3.02
Fesoterodine	2.03	1.74	2.31	1.84	1.64	2.03
Darifenacin	*	*	*	*	*	*
Solifenacin	1.46	1.32	1.59	2.19	1.94	3.02
Trospium IR	*	*	*	*	*	*
Trospium ER	2.45	2.19	2.70	2.68	2.38	2.98
Combined comparison of extended versus immediate release formulations						
Placebo	1.08	0.86	1.30	1.48	1.19	1.71
Extended Release	1.78	1.61	1.94	2.24	2.03	2.46
Immediate Release	1.46	1.28	1.64	2.17	1.81	2.54

*Estimates could not be calculated for these formulations because authors did not provide adequate data on variance for weighting of the raw data

Since baseline episodes of UUI per day ranged from 1.6 to 5.3, and voids per day from 7.2 to 13.7, these reductions (Table 31) reflect modest margins of benefit from baseline above placebo. Data was not consistently provided across studies to estimate the proportion of women who became symptom free.

Procedural and surgical treatments.

Methodologic consideration for procedural and surgical treatment studies. Studies in this treatment domain are of limited quality and predominantly case series in specialized treatment settings. Sacral neuromodulation has not had properly masked randomized clinical trials,¹⁹³ and botulinum toxin injections are promising but based on a small number of studies that identified urinary retention as a distinct risk factor that is self-resolving but troublesome.¹⁹³ Other procedures found no benefits or are no longer used in practice. Given consideration of 18 studies, of which 11 were fair quality and seven poor, with five RCTs, the strength of the evidence for managing OAB with procedural and surgical treatment is weak for all aspects of understanding outcomes of care.

Findings from procedural and surgical treatment studies. Among the trials of procedures and surgery, one demonstrated a statistically significant benefit of sacral neuromodulation over usual care for the reduction of episodes of urge urinary incontinence per day (average reduction of 7.1 compared to 2.1 increase with usual care) among subjects with OAB known to be refractory to medical therapy.¹²⁴ Enthusiasm is tempered primarily by reports from multiple case series that harms are not rare with these treatment approaches. Data that reflects newer sacral neuromodulation techniques in this refractory population are lacking.

Surgical and procedural treatments for OAB are not first line management due to the cost and risks of these types of procedures. Sacral neuromodulation decreases the number of urge incontinence episodes by at least 50 percent among patients refractory to conservative therapies. Moreover, the number of moderate-heavy incontinence episodes also decreases. These results persist even at five years. Sacral neuromodulation seems to have less of a benefit for urinary urgency (mixed results found in our review) and frequency (31 to 45 percent decrease), with benefits in frequency tapering off over time (23 percent reduction in daily voids at 5 years). These benefits come with a relatively high rate of adverse events. Early studies using older

technology had more than one adverse event per subject, on average;^{115, 124} studies employing newer technology report lower rate, with events in 11 to 53 percent of subjects.^{112, 119} Nearly 40 percent describe pain or an unpleasant stimulation, 7 to 48 percent returned to the operating room (this increased to 67 percent at five years, but includes the need to change the implantable pulse generator battery) and between 2 to 6 percent have an infection (often requiring hospitalization, intravenous antibiotics or explantation). The overall explantation rate hovers around 10 percent.

Our review included only one study on peripheral neuromodulation, using an anal and/or vaginal probe. Benefits in frequency were unlikely to be clinically significant (decrease from 9 to 8 voids daily) and there was a high dropout rate due to pain and effects on the bowels. Other forms of peripheral neuromodulation such as posterior tibial stimulation were not reviewed.

Electromagnetic stimulation with a portable unit was not found to be beneficial for OAB.

Of the drugs injected or instilled into the bladder, botulinum toxin and oxybutynin had the greatest benefit. One trial demonstrated benefit of instillation of oxybutynin compared to sterile water in the reduction of voids per day (average reduction of 6.8 compared to 2.4).¹²⁶ A trial we identified and the recent review by the Cochrane Collaboration found that small trials suggest benefit though researchers continue to evaluate means to decrease the risk of undesired side effects like urinary retention with botulinum toxin. Both botulinum toxin and instilled oxybutynin increase the postvoid residuals and the long term effects of higher residuals in terms of bladder infection and other risks is not known. Although evidence for these approaches is promising, the strength of the literature is inadequate to recommend any of these approaches for broader use in general practice. As of the date of the report, neither treatment is FDA approved for OAB.

Resiniferatoxin injection was not beneficial in the study reviewed. Older treatment modalities such as prolonged bladder distention or bladder transection are no longer commonly used due to the morbidity of these procedures. The reviewed studies also lacked rigorous methods for evaluating treatment benefit.

Behavioral treatments.

Methodologic considerations in behavioral treatment studies. Most of the literature addressing behavioral interventions (with or without comparison to pharmacologic intervention) was of fair or poor quality. In general, studies of behavioral approaches rarely included a true and comparable placebo arm. Although it is well recognized that there are inherent challenges to developing placebos for behavioral techniques, a reasonably strong evidence base on means of doing so suggests that it is possible. Burgio and colleagues¹⁴³ worked to mitigate this issue by maintaining similar visit schedules and through the use of bladder diaries in all groups, which was considered adequate masking for quality grading purposes. Particularly in older studies (prior to 2002), the behavioral approach often is not fully described; and inconsistency in the language used to identify different approaches requires the reader to examine the manuscripts very carefully – multiple studies may have called their approach by the same name, but in fact be studying quite different interventions.

To mitigate against such confusion, we have attempted in this report to always describe the intervention along with the first description of results from a given study. Studies conducted prior to the ICS definition of OAB in 2002 tended to examine more limited approaches to bladder training, while later studies focused more on multicomponent approaches and delivery of the training in different ways. Prior treatment attempts are rarely documented in this work, which may make it difficult to compare treatment groups across studies. Finally, this body of literature includes very few studies that included similar combinations of intervention and comparator making it almost impossible to summarize across them. After review and analysis of 29 studies, of which 14 were fair quality and 15 poor, including 17 RCTs, the strength of the evidence for

managing OAB with behavioral approaches treatment is moderate to weak for short term outcomes and weak for long term outcomes and harms.

Findings from behavioral treatment studies. Conclusions about the effectiveness of behavioral techniques for addressing the symptoms of overactive bladder are based on a total of 29 papers (27 studies) that encompass behavioral to behavioral comparisons as well as studies of combining behavioral approaches with pharmacologic treatment, and the reverse, combining pharmacologic approaches with behavioral ones. No two studies could be combined to produce summary data. Overall, behavioral approaches can be effective in reducing episodes of incontinence and daily voids. Multicomponent approaches are most effective, and they perform relatively equivalent to pharmacologic treatment. Generally speaking, reductions in symptoms were modest, with potential decreases in incontinence episodes of up to 1.9 per day, and reductions of up to about four voids per day. The addition of caffeine reduction to behavioral modification reduced frequency, but made no difference in reduction of incontinence episodes. There is no evidence the behavioral approaches enhance the effectiveness of pharmacologic therapy to reduce episodes of incontinence; and like pharmacologic approaches, there is no evidence for long term effectiveness beyond the period during which the intervention is being provided in the health care setting.

Complementary and alternative medicine treatments.

Findings from complementary and alternative therapy studies. We identified three studies that used complementary and alternative medicine therapies to treat OAB: a fair quality trial of acupuncture,¹³⁷ a fair quality trial of foot reflexology,¹³⁸ and a poor quality prospective case series of hypnotherapy. There is weak to no evidence for complementary and alternative approaches to managing OAB.

Outcomes of treatment. The small trial of acupuncture has intriguing results related to decreased frequency of voiding and reduced symptoms of urgency which are associated with changes in cystometrics related to improved bladder capacity that are logical intermediates of the improvement in symptoms. Women felt they were improved as measured by scales that capture bother and quality of life. Evidence is insufficient to support definitive choice of acupuncture but offers preliminary information that promises modest improvements similar to those reported in many pharmacologic trials.

Reflexology is represented by a small trial with unmasking of participants that could have biased the results. No evidence supports choice of this modality. Likewise, hypnotherapy is not supported by the scant information provided by one case series with little detail, patient reported outcomes, or statistical assessment. Given the scope of placebo effects demonstrated in other well-conducted studies of OAB treatment, it is difficult to know whether to attribute any effect to hypnotherapy.

KQ3: Comparisons of Treatments

Methodologic issues in studies that compare treatments. Evidence for direct comparisons of treatments was based on 19 studies: 12 were fair and 7 poor. Of these, 14 were RCTS. Evidence is currently weak to absent for choosing one therapy over another.

Pharmacologic. A number of pharmaceutical agents have been studied in direct comparison. Nine of these comparisons explicitly examined outcomes for urge urinary incontinence episodes and voids per day. Fewer report on the symptom of urgency. Five of these studies are comparability studies in which a new drug aimed to establish equivalence to oxybutynin, the first drug FDA approved for OAB. One study tested the hypothesis of whether the ER formulation of tolterodine was superior to the IR formulation and the remaining studies were “challenges” of newer drugs to tolterodine, which was the second drug to receive FDA approval. In this context

lack of statistical differences between active drugs is somewhat uninformative as trials were often powered for the comparison of the drugs individually to placebo and in many cases statistical testing of the outcomes of drug-to-drug comparison are not provided. Where studies reported being powered to detect differences between drugs, the postulated differences were unlikely and/or the withdrawals from protocol prevented meaningful effectiveness analysis beyond ITT. In a health services context, ideal comparisons would be of extended durations, since OAB is a chronic condition, and would report on all primary outcomes of relevance including differences in medication adherence, satisfaction with treatment, and quality of life.

Procedures. Participants in these studies were often not exclusively those with OAB. Strict application of inclusion criteria for this review would have eliminated virtually all studies of procedures for this reason. Studies of OAB included those with urinary retention and at times neurologic conditions as the indication. Masking, though challenging, was approximated by insertion of leads for sacral neuromodulation without activation; however given that a test period is done to establish efficacy before implantation of the permanent device, individuals may have been aware of their status and assessment of unmasking is not provided. Developing sham procedure methods may be of special importance in this area.

Behavioral. Variations in the behavioral approaches used and methods for teaching them, as well as differences in the duration and intensity of treatment make comparisons challenging. As a category the methods were generally strong with the continued challenge of developing attention control comparison methods and documenting testing of the degree to which individuals believed they knew their treatment status.

Findings of direct comparisons of treatments.

Pharmacologic. Among 14 pharmacologic RCTs that made ten unique comparisons among pharmacologic agents, the majority did not report statistical tests that showed one drug to be superior to another. The exceptions were from three RCTs. Both oxybutynin and tolterodine in extended release form were superior to tolterodine in immediate release forms.^{83, 140} One trial reported oxybutynin ER superior to tolterodine ER for reducing voids per day.⁸⁴ Given heterogeneity of participant populations and study designs, this limited number of studies is insufficient for any drug to be considered definitively superior.

Procedures. For procedures, sacral neuromodulation was compared to wait list participants on medications. It is important to note that failure of prior medical management was a criteria for entry into the study; those waiting had worsening of many symptoms. This is in contradistinction to improvements noted in virtually all other comparison groups and likely reflects bias from the knowledge that they were awaiting what they considered more definitive treatment for severe OAB. No conclusions can be reached with this data and future research should address the differences in risk profile of sacral neuromodulation versus medications.¹²⁴

Behavioral. Seven studies compared behavioral to pharmacologic treatments. One study in this group reported significant reductions in incontinence episodes with a multi-component intervention;¹⁴³ no study found differences in reductions in voids per day. Participants in one study who were queried reported a preference for behavioral treatment over pharmacologic.¹⁴³ Adding behavioral treatments to pharmacologic treatments did not improve outcomes for incontinence episodes or voids per day above pharmacologic treatment alone.

KQ4: Modifiers of Treatment Outcomes

Methodologic issues for study of modifiers. Individual characteristics of participants in OAB studies were highly varied on characteristics that are plausibly related to treatment outcomes such as: age, menopausal status, prior treatment, prior refractory symptoms during treatment, prior incontinence or gynecologic surgeries, severity of OAB at baseline, and presence

and type of incontinence. Higher quality studies reported on these characteristics and either found them comparable across trial arms or adjusted for baseline differences in their analyses. However, few studies indicated *a priori* goals of conducting sub-analyses in order to better understand treatment response. Among publications that did report on predictors of treatment response, the majority were under-powered to detect differences so the resulting claims of comparability are of limited value from an individual study. We found cross-cutting similarities for several of these characteristics such as age, severity, and prior treatment and have compiled those to present the limited picture that is coming into view. Overall this treatment literature is at an early stage of development in which the primary objective has been documenting superiority of the treatment to placebo. Population-based cohorts, such as the few provided by national and payor databases and larger clinical trials designed explicitly to more closely examine treatment response patterns and long-term effectiveness and tolerability will be required to have definitive information that can be used clinically with confidence.

Findings about modifiers of treatment outcomes. Advancing age was associated with more severe symptoms at baseline and with potentially observed attenuated treatment effects. Nonetheless, the majority of studies that reported on the effect of age in relationship to treatment outcomes found significant improvements in the older groups when active treatment was compared to placebo. No studies reported lack of efficacy among older participants. Older individuals do benefit from treatment. Race and ethnicity were not associated with outcomes in two analyses addressing this topic.

Presence of UUI, or urge-dominant mixed urinary incontinence, was not associated with treatment failure. Neither was severity of symptoms; in some cases those with the most severe symptoms, including more severe UUI, achieved the greatest treatment gains but were less likely to be symptom free.

Urodynamic findings do not provide consistent information about likelihood of treatment benefit or failure. A single study noted, among women who all had documented detrusor overactivity at enrollment, that those who had detrusor response to provocative maneuvers such as running water and washing hands were more likely to fail treatment. Further investigation of this finding from a small study would be intriguing as would examining the overall self-reported relationship between cues, urgency, UUI and treatment response. Another small study reported that anterior vaginal wall prolapse was a strong predictor of non-response to treatment. It is important to note that while these factors were associated with lower likelihood of treatment response, the majority of those treated with these characteristics did see improvement in symptoms.

Gender, while not a focus of this report, is important in interpreting findings with caution. Multiple research teams noted outcomes were not as favorable in men as in women. This likely reflects different pathophysiology behind the symptoms. Because some studies in this report included men (up to 25 percent of participants), results should be viewed in light of this potential bias.

KQ5: Costs of OAB Treatment

Methodologic considerations about cost studies. Studies that use administrative data are limited in their ability to adjust by clinical comorbidities and concomitant conditions, although they benefit from large enough numbers to provide a reasonable global estimate. Conversely studies that survey patients on their own health and health care may obtain more detailed information, but suffer from recall bias on the part of the respondents. None of the studies reports on their funding source; nor do they provide any information on investigator conflict of interest.

Findings about costs of treatment. Total direct health care costs for women with OAB in 2000 were estimated at \$6.9 billion, of which \$1.1 billion was for pharmacologic treatment and \$550 million for surgical treatment. The rest was estimated for “consequences” costs, which would include things like falls, longer hospital LOS and skin conditions. Medication costs for OAB with the two most commonly used drugs (oxybutynin and tolterodine) range from \$56 to \$360 over a twelve month period for newly diagnosed patients. However, overall health care costs were highest for patients who take oxybutynin, relative to tolterodine in any formulation, with costs lowest for patients on tolterodine ER. No study adequately determines why the observed differences exist, in particular whether they are actually a reflection of the differences in the populations who are prescribed the various medications. The one study that measures baseline costs found that patients whose incident OAB prescription was for tolterodine ER had lower costs in the six months prior to prescription than did patients whose incident prescription was either oxybutynin ER or oxybutynin IR. None of the studies conducted a cost effectiveness analysis, although cost-utility analyses have been conducted in Europe (which would be difficult given the low effectiveness of any of the drugs, and short followup of almost all efficacy studies). Nor does any study specifically assess the effect of medication on peripheral outcomes of OAB such as falls.

In studies of adherence and persistence, most of which are conducted in managed care populations, fewer than half of patients ever refill their prescriptions for OAB medications. Average quit time is about a month; among those who persist, adherence is best among patients taking extended release versus immediate release formulations. Even in this group, adherence is low; the highest medication possession ratio noted in the studies we identified was about 36 percent.²⁷²⁻²⁷⁵

Future Research

State of the Literature

The study of OAB as a syndrome is entering its second decade. As is typical of advancing areas of research, publications based on case series are giving way to observational cohorts. Trials, beyond those required for FDA approval of indication for OAB, are appearing in the literature and health services researchers are investigating population-level factors such as cost of care and risk of rare but serious side effects of treatments.

The 2002 ICS standardization of terminology¹⁸⁰ was associated with a productive trend toward greater attention to and clarity of operational definitions in research. Documentation of inclusion and exclusion criteria, baseline characteristics, and change in symptom profiles have become more detailed and nuanced in the last five to seven years. Improved clarity about research definitions for conducting the study and analyzing data was the case even when authors departed from ICS definitions. Simultaneously important research gains have been made in crafting, refining, and validating questionnaire and interview instruments for classifying symptoms, assessing severity of symptoms, describing impact of OAB on quality of life, and measuring satisfaction with outcomes.

Concerning deficits. As a body of literature, a number of concerning deficits were common. Fewer than seven percent of included studies met criteria for good quality. For example, for clinical trials, this meant that publications lacked one or more of the following:

- Any description of randomization method
- Masking of participants *and* assessors to treatment assignment
- Description of participant and selection process sufficient to understand generalizability

- Details of intervention provided to subjects sufficient to replicate
- Followup of treatment effects for 12 weeks or longer
- Loss to followup less than 10 percent
- Drop out less than 10 percent
- Power calculations (preferably for two-sided tests)
- Use of appropriate statistical comparisons and tests
- Clear description of methods used to measure outcomes
- Description of validity or reliability of outcome measures for primary outcomes

Each of these criteria is fundamental to the conduct and reporting of research of sufficient quality to build knowledge and inform care. The treatment literature is currently hindered by critical flaws that must be eliminated. These include use of data from only those who completed the whole treatment course and not intention-to-treat analyses. Likewise authors frequently noted covariates that were associated with either baseline severity or outcomes and did not adjust for these factors in analysis or conduct stratified analyses. This was often the case because the size of the study would not support modeling or sub-analyses. Conclusions often over-reached findings in ways that in some instances were blatantly biased in favor of a newer treatment.

The large magnitude of placebo effect in OAB studies deserves special note. The fact of robust placebo response implies that uncontrolled studies will be notably biased. Indeed in this literature observational studies, with rare exceptions, overestimated treatment benefits when compared to trials. High quality trials and innovations in masking treatment group (especially for procedural and behavioral studies) will be essential to firmly establishing treatment effects.

Conflict of Interest. Trends in increasing transparency about sources of funding and potential conflicts of interest have been steadily positive over the past two decades (Table 32). However, a fundamental mismatch exists between the initial research needed to obtain regulatory approval and broader, longer term research needs to assess a wider range of questions about outcomes of care in typical practice settings. For a condition as common as OAB, the funding and conflict of interest picture that emerges suggests a research area that is urgently in need of additional sources of independent funding for the next wave of clinical effectiveness research.

Table 32. Funding sources and conflict of interest by decade of publication

Study Focus	Source of funding for the research						Reporting of conflict of conflict of interest					
	Publications reporting research funding source, (%)			Industry funded among those reporting funding source, (%)			Publications reporting on author conflict of interest, (%)			Authors with COI for publications reporting COI, (%)*		
	1980s	1990s	2000s	1980s	1990s	2000s	1980s	1990s	2000s	1980s	1990s	2000s
Surgical or procedural treatments (n =18)	0/2 (0)	1/4 (25)	8/12 (75)	---	1/1 (100)	5/8 (63)	0/2 (0)	1/4 (25)	6/12 (50)	---	5/7 (71)	32/64 (50)
Medications (n=119)	1/3 (33)	11/17 (65)	78/99 (79)	0/1 (0)	9/11 (82)	73/78 (94)	0/3 (0)	0/17 (0)	50/99 (50)	---	---	228/329 (69)
Behavioral interventions (n = 25)	1/6 (17)	3/5 (60)	10/14 (71)	0/1 (0)	1/3 (33)	3/10 (30)	0/6 (0)	0/5 (0)	1/14 (7)	---	---	7/7 (100)
Complementary and alternative medical treatments (n=3)	0/1 (0)	---	1/2 (50)	---	---	0/1 (0)	0/1 (0)	---	0/2 (0)	---	---	---

* Data presented is the total number of authors reporting they had a conflict of interest (numerator) over the total number of authors in those publications that reported on their individual conflict of interest status (denominator). All other data in the table is the number of publications with the characteristic over the number of publications of that type in the respective decades.

Future Research Directions

Momentum in the direction of higher quality and more informative research will follow from attention to:

- Reporting greater information about key characteristics of populations studied in order to allow assessment of comparability of study populations and applicability of findings. This also facilitates understanding of candidate confounders and variations in findings across studies and study types.
- Conducting studies of sufficient size to conduct hypothesis testing or assess treatment effects. Small studies preclude meaningful descriptive analysis of modifiers and appropriate adjustment of confounders. Inclusion of small numbers of men in much larger studies was a recurrent example of a modifier of treatment outcomes that was noted or ignored without sufficient study size to meaningfully interpret.
- Continuing the expansion of standardized nomenclature and use of validated measures. The current literature is challenging to synthesize and interpret because outcomes measured are varied, not cross-cutting, and measured on different time scales (e.g., episodes per day, per week or per other unit of time). Use of validated measures is improving but measures are perhaps too numerous to help bring results into focus.

Networks of researchers or those working in common areas would benefit from a rigorous, evidence-based consensus process to prioritize tools to use across studies to improve comparability of measures of outcomes like severity and quality of life.

Content priorities. Well-conducted larger studies with study populations that reflect the severity of conditions seen in both primary care and specialty practice settings are critical. It is imperative that the time window of followup be extended. The literature suggests that treatment effects can be achieved in the early weeks of treatment with pharmacologic, procedural, behavioral, and complementary and alternative therapies. However this does not mean that the duration of study should be truncated. To the degree that long-term efficacy and effectiveness is poorly documented, the resolution or worsening of side effects is poorly characterized, and satisfaction and continuation of treatment is not assessed over extended periods of time, the literature is not relevant for informing care for this chronic condition.

Each area that was a focus of this report would benefit from continued study of:

- Etiology and natural history of disease, risk factors and potential preventive measures
- Novel treatments
- Properly powered investigations of direct comparisons of existing treatments
- Longer term investigation of benefits and side effects of treatment
- Continued study of combinations of treatments
- Further investigation of complementary and alternative treatments
- Investigation of predictors of both good and poor treatment responses across treatment modalities.
- Selection of treatment options after prior treatment failure

Conclusions

We find a concerning lack of high-quality evidence to inform clinical decision-making for millions of women in the United States. Both medical and behavioral interventions can provide symptom relief which is often not complete, but valued by women who struggle with OAB. Well-conducted trials of greater duration and sophistication, separate from drug development and marketing efforts, are crucial. Because benefits of current treatments are modest, because drug side effects can be bothersome, opportunities exist to study how to gain synergy from combinations of types of treatments. We must note that lack of evidence of benefit is not equivalent to evidence of no benefit. A number of treatments that are potentially promising warrant continued investigation. Cross-cutting concerns about the quality of research must be addressed to achieve literature that can be meaningfully synthesized. Current literature does not permit definitive conclusions about relative benefit, harm, or costs to achieve similar results. Given how common and concerning OAB is, a priority on promoting high-quality research in the United States is imperative. Women and their care providers deserve better information to guide their choices.

References and Included Studies

1. U. S. Census Bureau. Table 2. Annual Estimates of the Population by Sex and Selected Age Groups for the United States: April 1, 2000 to July 1, 2007 (NC-EST2007-02). May 1, 2008.
2. Lawrence JM, Lukacz ES, Nager CW, et al. Prevalence and co-occurrence of pelvic floor disorders in community-dwelling women. *Obstet Gynecol.* 2008 Mar;111(3):678-85.
3. Lukacz ES, Lawrence JM, Contreras R, et al. Parity, mode of delivery, and pelvic floor disorders. *Obstet Gynecol.* 2006 Jun;107(6):1253-60.
4. Wagner TH, Hu TW, Bentkover J, et al. Health-related consequences of overactive bladder. *Am J Manag Care.* 2002 Dec;8(19 Suppl):S598-607.
5. Hozo SP, Djulbegovic B and Hozo I. Estimating the mean and variance from the median, range, and the size of a sample. *BMC Med Res Methodol.* 2005;5(1):13.
6. DerSimonian R and Laird N. Meta-analysis in clinical trials. *Control Clin Trials.* 1986 Sep;7(3):177-88.
7. Sismondo S. How pharmaceutical industry funding affects trial outcomes: causal structures and responses. *Soc Sci Med.* 2008 May;66(9):1909-14.
8. Sismondo S. Pharmaceutical company funding and its consequences: a qualitative systematic review. *Contemp Clin Trials.* 2008 Mar;29(2):109-13.
9. Lexchin J, Bero LA, Djulbegovic B, et al. Pharmaceutical industry sponsorship and research outcome and quality: systematic review. *BMJ.* 2003 May 31;326(7400):1167-70.
10. Fenner DE, Trowbridge ER, Patel DL, et al. Establishing the prevalence of incontinence study: racial differences in women's patterns of urinary incontinence. *J Urol.* 2008 Apr;179(4):1455-60.
11. Dooley Y, Kenton K, Cao G, et al. Urinary incontinence prevalence: results from the National Health and Nutrition Examination Survey. *J Urol.* 2008 Feb;179(2):656-61.
12. Herschorn S, Gajewski J, Schulz J, et al. A population-based study of urinary symptoms and incontinence: the Canadian Urinary Bladder Survey. *BJU Int.* 2008 Jan;101(1):52-8.
13. Lawrence JM, Lukacz ES, Liu IL, et al. Pelvic floor disorders, diabetes, and obesity in women: findings from the Kaiser Permanente Continence Associated Risk Epidemiology Study. *Diabetes Care.* 2007 Oct;30(10):2536-41.
14. Choo MS, Ku JH, Lee JB, et al. Cross-cultural differences for adapting overactive bladder symptoms: results of an epidemiologic survey in Korea. *World J Urol.* 2007 Oct;25(5):505-11.
15. Townsend MK, Danforth KN, Lifford KL, et al. Incidence and remission of urinary incontinence in middle-aged women. *Am J Obstet Gynecol.* 2007 Aug;197(2):167 e1-5.
16. Link CL, Lutfey KE, Steers WD, et al. Is abuse causally related to urologic symptoms? Results from the Boston Area Community Health (BACH) Survey. *Eur Urol.* 2007 Aug;52(2):397-406.
17. Waetjen LE, Liao S, Johnson WO, et al. Factors associated with prevalent and incident urinary incontinence in a cohort of midlife women: a longitudinal analysis of data: study of women's health across the nation. *Am J Epidemiol.* 2007 Feb 1;165(3):309-18.
18. Irwin DE, Milsom I, Hunskaar S, et al. Population-based survey of urinary incontinence, overactive bladder, and other lower urinary tract symptoms in five countries: results of the EPIC study. *Eur Urol.* 2006 Dec;50(6):1306-14; discussion 1314-5.
19. Huang AJ, Thom DH, Kanaya AM, et al. Urinary incontinence and pelvic floor dysfunction in Asian-American women. *Am J Obstet Gynecol.* 2006 Nov;195(5):1331-7.
20. Hsieh CH, Chen HY, Hsu CS, et al. Prevalence of urinary frequency in Taiwanese women aged 20-59 years. *Taiwan J Obstet Gynecol.* 2006 Sep;45(3):244-6.
21. Homma Y, Yamaguchi O and Hayashi K. Epidemiologic survey of lower urinary tract symptoms in Japan. *Urology.* 2006 Sep;68(3):560-4.
22. Odeyemi IA, Dakin HA, O'Donnell RA, et al. Epidemiology, prescribing patterns and resource use associated with overactive bladder in UK primary care. *Int J Clin Pract.* 2006 Aug;60(8):949-58.
23. Teloken C, Caraver F, Weber FA, et al. Overactive bladder: prevalence and implications in Brazil. *Eur Urol.* 2006 Jun;49(6):1087-92.
24. Kim SH, Boye M, Bhattacharyya SK, et al. Medical visits among adults with symptoms commonly associated with an overactive bladder. *BJU Int.* 2006 Mar;97(3):551-4.
25. Fitzgerald MP, Thom DH, Wassel-Fyr C, et al. Childhood urinary symptoms predict adult overactive bladder symptoms. *J Urol.* 2006 Mar;175(3 Pt 1):989-93.
26. McGrother CW, Donaldson MM, Hayward T, et al. Urinary storage symptoms and comorbidities: a prospective population cohort study in middle-aged and older women. *Age Ageing.* 2006 Jan;35(1):16-24.
27. Thom DH, van den Eeden SK, Ragins AI, et al. Differences in prevalence of urinary incontinence by race/ethnicity. *J Urol.* 2006 Jan;175(1):259-64.
28. Donaldson MM, Thompson JR, Matthews RJ, et al. The natural history of overactive bladder and stress urinary incontinence in older women in the community: a 3-year prospective cohort study. *Neurourol Urodyn.* 2006;25(7):709-16.
29. Zhang W, Song Y, He X, et al. Prevalence and risk factors of overactive bladder syndrome in Fuzhou Chinese women. *Neurourol Urodyn.* 2006;25(7):717-21.
30. Yu HJ, Liu CY, Lee KL, et al. Overactive bladder syndrome among community-dwelling adults in Taiwan: prevalence, correlates, perception, and treatment seeking. *Urol Int.* 2006;77(4):327-33.
31. Homma Y, Yamaguchi O and Hayashi K. An epidemiological survey of overactive bladder symptoms in Japan. *BJU Int.* 2005 Dec;96(9):1314-8.
32. MacDiarmid S and Rosenberg M. Overactive bladder in women: symptom impact and treatment expectations. *Curr Med Res Opin.* 2005 Sep;21(9):1413-21.

33. Zhang W, Song Y, He X, et al. Prevalence and risk factors of lower urinary tract symptoms in Fuzhou Chinese women. *Eur Urol*. 2005 Aug;48(2):309-13.
34. Song YF, Zhang WJ, Song J, et al. Prevalence and risk factors of urinary incontinence in Fuzhou Chinese women. *Chin Med J (Engl)*. 2005 Jun 5;118(11):887-92.
35. Shakhathreh FM. Epidemiology of urinary incontinence in Jordanian women. *Saudi Med J*. 2005 May;26(5):830-5.
36. Muller N. What Americans understand and how they are affected by bladder control problems: highlights of recent nationwide consumer research. *Urol Nurs*. 2005 Apr;25(2):109-15.
37. Rohr G, Christensen K, Ulstrup K, et al. Reproducibility and validity of simple questions to identify urinary incontinence in elderly women. *Acta Obstet Gynecol Scand*. 2004 Oct;83(10):969-72.
38. Corcos J and Schick E. Prevalence of overactive bladder and incontinence in Canada. *Can J Urol*. 2004 Jun;11(3):2278-84.
39. Dallosso HM, McGrother CW, Matthews RJ, et al. Nutrient composition of the diet and the development of overactive bladder: a longitudinal study in women. *Neurourol Urodyn*. 2004;23(3):204-10.
40. Espino DV, Palmer RF, Miles TP, et al. Prevalence and severity of urinary incontinence in elderly Mexican-American women. *J Am Geriatr Soc*. 2003 Nov;51(11):1580-6.
41. Chen YC, Chen GD, Hu SW, et al. Is the occurrence of storage and voiding dysfunction affected by menopausal transition or associated with the normal aging process? *Menopause*. 2003 May-Jun;10(3):203-8.
42. Stewart WF, Van Rooyen JB, Cundiff GW, et al. Prevalence and burden of overactive bladder in the United States. *World J Urol*. 2003 May;20(6):327-36.
43. Siracusano S, Pregazzi R, d'Aloia G, et al. Prevalence of urinary incontinence in young and middle-aged women in an Italian urban area. *Eur J Obstet Gynecol Reprod Biol*. 2003 Apr 25;107(2):201-4.
44. Rortveit G, Daltveit AK, Hannestad YS, et al. Urinary incontinence after vaginal delivery or cesarean section. *N Engl J Med*. 2003 Mar 6;348(10):900-7.
45. Eva UF, Gun W and Preben K. Prevalence of urinary and fecal incontinence and symptoms of genital prolapse in women. *Acta Obstet Gynecol Scand*. 2003 Mar;82(3):280-6.
46. Yip SK and Chung TK. Treatment-seeking behavior in Hong Kong Chinese women with urinary symptoms. *Int Urogynecol J Pelvic Floor Dysfunct*. 2003 Feb;14(1):27-30; discussion 30.
47. Chen GD, Lin TL, Hu SW, et al. Prevalence and correlation of urinary incontinence and overactive bladder in Taiwanese women. *Neurourol Urodyn*. 2003;22(2):109-17.
48. van der Vaart CH, de Leeuw JR, Roovers JP, et al. The effect of urinary incontinence and overactive bladder symptoms on quality of life in young women. *BJU Int*. 2002 Oct;90(6):544-9.
49. Liu C and Andrews GR. Prevalence and incidence of urinary incontinence in the elderly: a longitudinal study in South Australia. *Chin Med J (Engl)*. 2002 Jan;115(1):119-22.
50. Aaron R, Muliylil J and Abraham S. Medico-social dimensions of menopause: a cross-sectional study from rural south India. *Natl Med J India*. 2002 Jan-Feb;15(1):14-7.
51. Milsom I, Abrams P, Cardozo L, et al. How widespread are the symptoms of an overactive bladder and how are they managed? A population-based prevalence study. *BJU Int*. 2001 Jun;87(9):760-6.
52. Tseng JJ, Chen YT, Chen MT, et al. Prevalence of urinary incontinence and intention to seek treatment in the elderly. *J Formos Med Assoc*. 2000 Oct;99(10):753-8.
53. Moller LA, Lose G and Jorgensen T. The prevalence and bothersomeness of lower urinary tract symptoms in women 40-60 years of age. *Acta Obstet Gynecol Scand*. 2000 Apr;79(4):298-305.
54. Ueda T, Tamaki M, Kageyama S, et al. Urinary incontinence among community-dwelling people aged 40 years or older in Japan: prevalence, risk factors, knowledge and self-perception. *Int J Urol*. 2000 Mar;7(3):95-103.
55. Bortolotti A, Bernardini B, Colli E, et al. Prevalence and risk factors for urinary incontinence in Italy. *Eur Urol*. 2000 Jan;37(1):30-5.
56. Kuh D, Cardozo L and Hardy R. Urinary incontinence in middle aged women: childhood enuresis and other lifetime risk factors in a British prospective cohort. *J Epidemiol Community Health*. 1999 Aug;53(8):453-8.
57. Simeonova Z, Milsom I, Kullendorff AM, et al. The prevalence of urinary incontinence and its influence on the quality of life in women from an urban Swedish population. *Acta Obstet Gynecol Scand*. 1999 Jul;78(6):546-51.
58. Kay L, Stigsby B, Brasso K, et al. Lower urinary tract symptoms--a population survey using the Danish Prostatic Symptom Score (DAN-PSS) questionnaire. *Scand J Urol Nephrol*. 1999 Apr;33(2):94-9.
59. Koyama W, Koyanagi A, Mihara S, et al. Prevalence and conditions of urinary incontinence among the elderly. *Methods Inf Med*. 1998 Jun;37(2):151-5.
60. Schulman C, Claes H and Matthijs J. Urinary incontinence in Belgium: a population-based epidemiological survey. *Eur Urol*. 1997;32(3):315-20.
61. Brieger GM, Mongelli M, Hin LY, et al. The epidemiology of urinary dysfunction in Chinese women. *Int Urogynecol J Pelvic Floor Dysfunct*. 1997;8(4):191-5.
62. Bogren MA, Hvarfwen E and Fridlund B. Urinary incontinence among a 65-year old Swedish population: medical history and psychosocial consequences. *Vard Nord Utveckl Forsk*. 1997 Winter;17(4):14-7.
63. Brieger GM, Yip SK, Hin LY, et al. The prevalence of urinary dysfunction in Hong Kong Chinese women. *Obstet Gynecol*. 1996 Dec;88(6):1041-4.
64. Nygaard IE and Lemke JH. Urinary incontinence in rural older women: prevalence, incidence and remission. *J Am Geriatr Soc*. 1996 Sep;44(9):1049-54.
65. Lara C and Nacey J. Ethnic differences between Maori, Pacific Island and European New Zealand women in prevalence and attitudes to urinary incontinence. *N Z Med J*. 1994 Sep 28;107(986 Pt 1):374-6.
66. Ju CC, Swan LK, Merriman A, et al. Urinary incontinence among the elderly people of Singapore. *Age Ageing*. 1991 Jul;20(4):262-6.

67. Hording U, Pedersen KH, Sidenius K, et al. Urinary incontinence in 45-year-old women. An epidemiological survey. *Scand J Urol Nephrol*. 1986;20(3):183-6.
68. Iosif CS and Bekassy Z. Prevalence of genito-urinary symptoms in the late menopause. *Acta Obstet Gynecol Scand*. 1984;63(3):257-60.
69. Danforth KN, Shah AD, Townsend MK, et al. Physical activity and urinary incontinence among healthy, older women. *Obstet Gynecol*. 2007 Mar;109(3):721-7.
70. Ege E, Akin B, Altuntug K, et al. Prevalence of urinary incontinence in the 12-month postpartum period and related risk factors in Turkey. *Urol Int*. 2008;80(4):355-61.
71. Hall SA, Cinar A, Link CL, et al. Do urological symptoms cluster among women? Results from the Boston Area Community Health Survey. *BJU Int*. 2008 May;101(10):1257-66.
72. Minassian VA, Stewart WF and Wood GC. Urinary incontinence in women: variation in prevalence estimates and risk factors. *Obstet Gynecol*. 2008 Feb;111(2 Pt 1):324-31.
73. Tikkinen KA, Auvinen A, Tiitinen A, et al. Reproductive factors associated with nocturia and urinary urgency in women: a population-based study in Finland. *Am J Obstet Gynecol*. 2008 Aug;199(2):153 e1-12.
74. Townsend MK, Curhan GC, Resnick NM, et al. BMI, waist circumference, and incident urinary incontinence in older women. *Obesity (Silver Spring)*. 2008 Apr;16(4):881-6.
75. Wesnes SL, Rortveit G, Bo K, et al. Urinary incontinence during pregnancy. *Obstet Gynecol*. 2007 Apr;109(4):922-8.
76. Hannestad YS, Rortveit G, Sandvik H, et al. A community-based epidemiological survey of female urinary incontinence: the Norwegian EPINCONT study. Epidemiology of Incontinence in the County of Nord-Trøndelag. *J Clin Epidemiol*. 2000 Nov;53(11):1150-7.
77. Herzog AR, Diokno AC, Brown MB, et al. Two-year incidence, remission, and change patterns of urinary incontinence in noninstitutionalized older adults. *J Gerontol*. 1990 Mar;45(2):M67-74.
78. Perry S, Shaw C, Assassa P, et al. An epidemiological study to establish the prevalence of urinary symptoms and felt need in the community: the Leicestershire MRC Incontinence Study. Leicestershire MRC Incontinence Study Team. *J Public Health Med*. 2000 Sep;22(3):427-34.
79. Yarnell JW, Voyle GJ, Richards CJ, et al. The prevalence and severity of urinary incontinence in women. *J Epidemiol Community Health*. 1981 Mar;35(1):71-4.
80. Zhu L, Lang J, Wang H, et al. The prevalence of and potential risk factors for female urinary incontinence in Beijing, China. *Menopause*. 2008 May-Jun;15(3):566-9.
81. Nuotio M, Tammela TL, Luukkaala T, et al. Urgency and urge incontinence in an older population: ten-year changes and their association with mortality. *Aging Clin Exp Res*. 2002 Oct;14(5):412-9.
82. Zinner N, Tuttle J and Marks L. Efficacy and tolerability of darifenacin, a muscarinic M3 selective receptor antagonist (M3 SRA), compared with oxybutynin in the treatment of patients with overactive bladder. *World J Urol*. 2005 Sep;23(4):248-52.
83. Sand PK, Miklos J, Ritter H, et al. A comparison of extended-release oxybutynin and tolterodine for treatment of overactive bladder in women. *Int Urogynecol J Pelvic Floor Dysfunct*. 2004 Jul-Aug;15(4):243-8.
84. Diokno AC, Appell RA, Sand PK, et al. Prospective, randomized, double-blind study of the efficacy and tolerability of the extended-release formulations of oxybutynin and tolterodine for overactive bladder: results of the OPERA trial. *Mayo Clin Proc*. 2003 Jun;78(6):687-95.
85. Dmochowski RR, Davila GW, Zinner NR, et al. Efficacy and safety of transdermal oxybutynin in patients with urge and mixed urinary incontinence. *J Urol*. 2002 Aug;168(2):580-6.
86. Sussman D and Garely A. Treatment of overactive bladder with once-daily extended-release tolterodine or oxybutynin: the antimuscarinic clinical effectiveness trial (ACET). *Curr Med Res Opin*. 2002;18(4):177-84.
87. Davila GW, Daugherty CA and Sanders SW. A short-term, multicenter, randomized double-blind dose titration study of the efficacy and anticholinergic side effects of transdermal compared to immediate release oral oxybutynin treatment of patients with urge urinary incontinence. *J Urol*. 2001 Jul;166(1):140-5.
88. Dmochowski RR, Sand PK, Zinner NR, et al. Comparative efficacy and safety of transdermal oxybutynin and oral tolterodine versus placebo in previously treated patients with urge and mixed urinary incontinence. *Urology*. 2003 Aug;62(2):237-42.
89. Giannitsas K, Perimenis P, Athanasopoulos A, et al. Comparison of the efficacy of tolterodine and oxybutynin in different urodynamic severity grades of idiopathic detrusor overactivity. *Eur Urol*. 2004 Dec;46(6):776-82; discussion 782-3.
90. Halaska M, Ralph G, Wiedemann A, et al. Controlled, double-blind, multicentre clinical trial to investigate long-term tolerability and efficacy of trospium chloride in patients with detrusor instability. *World J Urol*. 2003 May;20(6):392-9.
91. Abrams P, Freeman R, Anderstrom C, et al. Tolterodine, a new antimuscarinic agent: as effective but better tolerated than oxybutynin in patients with an overactive bladder. *Br J Urol*. 1998 Jun;81(6):801-10.
92. Homma Y and Koyama N. Minimal clinically important change in urinary incontinence detected by a quality of life assessment tool in overactive bladder syndrome with urge incontinence. *Neurourol Urodyn*. 2006;25(3):228-35.
93. Wang AC, Chih SY and Chen MC. Comparison of electric stimulation and oxybutynin chloride in management of overactive bladder with special reference to urinary urgency: a randomized placebo-controlled trial. *Urology*. 2006 Nov;68(5):999-1004.
94. Lee JG, Hong JY, Choo MS, et al. Tolterodine: as effective but better tolerated than oxybutynin in Asian patients with symptoms of overactive bladder. *Int J Urol*. 2002 May;9(5):247-52.
95. Nitti VW, Dmochowski R, Sand PK, et al. Efficacy, safety and tolerability of fesoterodine for overactive bladder syndrome. *J Urol*. 2007 Dec;178(6):2488-94.
96. Chapple C, Van Kerrebroeck P, Tubaro A, et al. Clinical efficacy, safety, and tolerability of once-daily fesoterodine in subjects with overactive bladder. *Eur Urol*. 2007 Oct;52(4):1204-12.

97. Chapple CR, Rechberger T, Al-Shukri S, et al. Randomized, double-blind placebo- and tolterodine-controlled trial of the once-daily antimuscarinic agent solifenacin in patients with symptomatic overactive bladder. *BJU Int.* 2004 Feb;93(3):303-10.
98. Yamaguchi O, Marui E, Kakizaki H, et al. Randomized, double-blind, placebo- and propiverine-controlled trial of the once-daily antimuscarinic agent solifenacin in Japanese patients with overactive bladder. *BJU Int.* 2007 Sep;100(3):579-87.
99. Chapple CR, Fianu-Jonsson A, Indig M, et al. Treatment outcomes in the STAR study: a subanalysis of solifenacin 5 mg and tolterodine ER 4 mg. *Eur Urol.* 2007 Oct;52(4):1195-203.
100. Hill S, Khullar V, Wyndaele JJ, et al. Dose response with darifenacin, a novel once-daily M3 selective receptor antagonist for the treatment of overactive bladder: results of a fixed dose study. *Int Urogynecol J Pelvic Floor Dysfunct.* 2006 May;17(3):239-47.
101. Zinner N, Susset J, Gittelman M, et al. Efficacy, tolerability and safety of darifenacin, an M(3) selective receptor antagonist: an investigation of warning time in patients with OAB. *Int J Clin Pract.* 2006 Jan;60(1):119-26.
102. Steers W, Corcos J, Foote J, et al. An investigation of dose titration with darifenacin, an M3-selective receptor antagonist. *BJU Int.* 2005 Mar;95(4):580-6.
103. Staskin D, Sand P, Zinner N, et al. Once daily trospium chloride is effective and well tolerated for the treatment of overactive bladder: results from a multicenter phase III trial. *J Urol.* 2007 Sep;178(3 Pt 1):978-83; discussion 983-4.
104. Rudy D, Cline K, Harris R, et al. Time to onset of improvement in symptoms of overactive bladder using antimuscarinic treatment. *BJU Int.* 2006 Mar;97(3):540-6.
105. Dmochowski RR, Sand PK, Zinner NR, et al. Trospium 60 mg once daily (QD) for overactive bladder syndrome: results from a placebo-controlled interventional study. *Urology.* 2008 Mar;71(3):449-54.
106. Zinner N, Gittelman M, Harris R, et al. Trospium chloride improves overactive bladder symptoms: a multicenter phase III trial. *J Urol.* 2004 Jun;171(6 Pt 1):2311-5, quiz 2435.
107. Cardozo L, Rekers H, Tapp A, et al. Oestriol in the treatment of postmenopausal urgency: a multicentre study. *Maturitas.* 1993 Dec;18(1):47-53.
108. Lose G and Englev E. Oestradiol-releasing vaginal ring versus oestriol vaginal pessaries in the treatment of bothersome lower urinary tract symptoms. *BJOG.* 2000 Aug;107(8):1029-34.
109. Molander U, Milsom I, Ekelund P, et al. A health care program for the investigation and treatment of elderly women with urinary incontinence and related urogenital symptoms. *Acta Obstet Gynecol Scand.* 1991;70(2):137-42.
110. Groenendijk PM, Lycklama Nyeholt AA, Heesakkers JP, et al. Urodynamic evaluation of sacral neuromodulation for urge urinary incontinence. *BJU Int.* 2008 Feb;101(3):325-9.
111. Janknegt RA, Hassouna MM, Siegel SW, et al. Long-term effectiveness of sacral nerve stimulation for refractory urge incontinence. *Eur Urol.* 2001 Jan;39(1):101-6.
112. Kessler TM, Buchser E, Meyer S, et al. Sacral neuromodulation for refractory lower urinary tract dysfunction: results of a nationwide registry in Switzerland. *Eur Urol.* 2007 May;51(5):1357-63.
113. Koldewijn EL, Rosier PF, Meuleman EJ, et al. Predictors of success with neuromodulation in lower urinary tract dysfunction: results of trial stimulation in 100 patients. *J Urol.* 1994 Dec;152(6 Pt 1):2071-5.
114. Siegel SW, Catanzaro F, Dijkema HE, et al. Long-term results of a multicenter study on sacral nerve stimulation for treatment of urinary urge incontinence, urgency-frequency, and retention. *Urology.* 2000 Dec 4;56(6 Suppl 1):87-91.
115. van Kerrebroeck PE, van Voskuilen AC, Heesakkers JP, et al. Results of sacral neuromodulation therapy for urinary voiding dysfunction: outcomes of a prospective, worldwide clinical study. *J Urol.* 2007 Nov;178(5):2029-34.
116. Fossberg E, Sorensen S, Ruutu M, et al. Maximal electrical stimulation in the treatment of unstable detrusor and urge incontinence. *Eur Urol.* 1990;18(2):120-3.
117. Schmid DM, Sauermann P, Werner M, et al. Experience with 100 cases treated with botulinum-A toxin injections in the detrusor muscle for idiopathic overactive bladder syndrome refractory to anticholinergics. *J Urol.* 2006 Jul;176(1):177-85.
118. Korda A, Krieger M and Hunter P. The use of prolonged bladder distension in the treatment of intractable urinary incontinence in females with detrusor instability. *Aust N Z J Obstet Gynaecol.* 1987 May;27(2):155-7.
119. Sutherland SE, Lavers A, Carlson A, et al. Sacral nerve stimulation for voiding dysfunction: One institution's 11-year experience. *Neurourol Urodyn.* 2007;26(1):19-28; discussion 36.
120. van Voskuilen AC, Oerlemans DJ, Weil EH, et al. Long term results of neuromodulation by sacral nerve stimulation for lower urinary tract symptoms: a retrospective single center study. *Eur Urol.* 2006 Feb;49(2):366-72.
121. Mundy AR. Long-term results of bladder transection for urge incontinence. *Br J Urol.* 1983 Dec;55(6):642-4.
122. Spinelli M, Bertapelle P, Cappellano F, et al. Chronic sacral neuromodulation in patients with lower urinary tract symptoms: results from a national register. *J Urol.* 2001 Aug;166(2):541-5.
123. Hassouna MM, Siegel SW, Nyeholt AA, et al. Sacral neuromodulation in the treatment of urgency-frequency symptoms: a multicenter study on efficacy and safety. *J Urol.* 2000 Jun;163(6):1849-54.
124. Schmidt RA, Jonas U, Oleson KA, et al. Sacral nerve stimulation for treatment of refractory urinary urge incontinence. *Sacral Nerve Stimulation Study Group. J Urol.* 1999 Aug;162(2):352-7.
125. O'Reilly BA, Fynes M, Ahtari C, et al. A prospective randomised double-blind controlled trial evaluating the effect of trans-sacral magnetic stimulation in women with overactive bladder. *Int Urogynecol J Pelvic Floor Dysfunct.* 2008 Apr;19(4):497-502.
126. Enzelsberger H, Helmer H and Kurz C. Intravesical instillation of oxybutynin in women with idiopathic detrusor instability: a randomised trial. *Br J Obstet Gynaecol.* 1995 Nov;102(11):929-30.
127. Rios LA, Panhoca R, Mattos D, Jr., et al. Intravesical resiniferatoxin for the treatment of women with idiopathic

- detrusor overactivity and urgency incontinence: A single dose, 4 weeks, double-blind, randomized, placebo controlled trial. *Neurourol Urodyn.* 2007;26(6):773-8.
128. Duthie J, Wilson DI, Herbison GP, et al. Botulinum toxin injections for adults with overactive bladder syndrome. *Cochrane Database of Systematic Reviews.* 2007;2007(3).
 129. Frewen WK. A reassessment of bladder training in detrusor dysfunction in the female. *Br J Urol.* 1982 Aug;54(4):372-3.
 130. Dorey G, Sultan H, Eckford SD, et al. Pelvic floor muscle exercises and urge suppression techniques in women with urge urinary incontinence: a retrospective study from a dedicated urogynaecology clinic. *Journal of the Association of Chartered Physiotherapists in Women's Health.* 2006;99(Autumn):43-8.
 131. Millard RJ and Oldenburg BF. The symptomatic, urodynamic and psychodynamic results of bladder re-education programs. *J Urol.* 1983 Oct;130(4):715-9.
 132. Dowd T, Kolcaba K and Steiner R. The addition of coaching to cognitive strategies: interventions for persons with compromised urinary bladder syndrome. *J Wound Ostomy Continence Nurs.* 2003 Mar;30(2):90-9.
 133. Jarvis GJ and Millar DR. Controlled trial of bladder drill for detrusor instability. *Br Med J.* 1980 Nov 15;281(6251):1322-3.
 134. Wyman JF, Fantl JA, McClish DK, et al. Comparative efficacy of behavioral interventions in the management of female urinary incontinence. *Continence Program for Women Research Group. Am J Obstet Gynecol.* 1998 Oct;179(4):999-1007.
 135. Burgio KL, Goode PS, Locher JL, et al. Behavioral training with and without biofeedback in the treatment of urge incontinence in older women: a randomized controlled trial. *JAMA.* 2002 Nov 13;288(18):2293-9.
 136. Bryant CM, Dowell CJ and Fairbrother G. Caffeine reduction education to improve urinary symptoms. *Br J Nurs.* 2002 Apr 25-May 8;11(8):560-5.
 137. Emmons SL and Otto L. Acupuncture for overactive bladder: a randomized controlled trial. *Obstet Gynecol.* 2005 Jul;106(1):138-43.
 138. Mak HL, Cheon WC, Wong T, et al. Randomized controlled trial of foot reflexology for patients with symptomatic idiopathic detrusor overactivity. *Int Urogynecol J Pelvic Floor Dysfunct.* 2007 Jun;18(6):653-8.
 139. Freeman RM and Baxby K. Hypnotherapy for incontinence caused by the unstable detrusor. *Br Med J (Clin Res Ed).* 1982 Jun 19;284(6332):1831-4.
 140. Swift S, Garely A, Dimpfl T, et al. A new once-daily formulation of tolterodine provides superior efficacy and is well tolerated in women with overactive bladder. *Int Urogynecol J Pelvic Floor Dysfunct.* 2003 Feb;14(1):50-4; discussion 54-5.
 141. Van Kerrebroeck P, Kreder K, Jonas U, et al. Tolterodine once-daily: superior efficacy and tolerability in the treatment of the overactive bladder. *Urology.* 2001 Mar;57(3):414-21.
 142. Chapple CR, Van Kerrebroeck PE, Junemann KP, et al. Comparison of fesoterodine and tolterodine in patients with overactive bladder. *BJU Int.* 2008 Nov;102(9):1128-32.
 143. Burgio KL, Locher JL, Goode PS, et al. Behavioral vs drug treatment for urge urinary incontinence in older women: a randomized controlled trial. *JAMA.* 1998 Dec 16;280(23):1995-2000.
 144. Burgio KL, Locher JL, Roth DL, et al. Psychological improvements associated with behavioral and drug treatment of urge incontinence in older women. *J Gerontol B Psychol Sci Soc Sci.* 2001 Jan;56(1):P46-51.
 145. Goode PS, Burgio KL, Locher JL, et al. Urodynamic changes associated with behavioral and drug treatment of urge incontinence in older women. *J Am Geriatr Soc.* 2002 May;50(5):808-16.
 146. Colombo M, Zanetta G, Scalabrino S, et al. Oxybutynin and bladder training in the management of female urge urinary incontinence: a randomized study. *Int Urogynecol J.* 1995;6(2):63-67.
 147. Diokno A and Yuhico M, Jr. Preference, compliance and initial outcome of therapeutic options chosen by female patients with urinary incontinence. *J Urol.* 1995 Nov;154(5):1727-30; discussion 1731.
 148. Macaulay AJ, Stern RS, Holmes DM, et al. Micturition and the mind: psychological factors in the aetiology and treatment of urinary symptoms in women. *Br Med J (Clin Res Ed).* 1987 Feb 28;294(6571):540-3.
 149. Jarvis GJ. A controlled trial of bladder drill and drug therapy in the management of detrusor instability. *Br J Urol.* 1981 Dec;53(6):565-6.
 150. Song C, Park JT, Heo KO, et al. Effects of bladder training and/or tolterodine in female patients with overactive bladder syndrome: a prospective, randomized study. *J Korean Med Sci.* 2006 Dec;21(6):1060-3.
 151. Mattiasson A, Blaakaer J, Hoye K, et al. Simplified bladder training augments the effectiveness of tolterodine in patients with an overactive bladder. *BJU Int.* 2003 Jan;91(1):54-60.
 152. Burgio KL, Kraus SR, Menefee S, et al. Behavioral therapy to enable women with urge incontinence to discontinue drug treatment: a randomized trial. *Ann Intern Med.* 2008 Aug 5;149(3):161-9.
 153. Chancellor MB, Kianifard F, Beamer E, et al. A comparison of the efficacy of darifenacin alone vs. darifenacin plus a Behavioural Modification Programme upon the symptoms of overactive bladder. *Int J Clin Pract.* 2008 Apr;62(4):606-13.
 154. Millard RJ. Clinical efficacy of tolterodine with or without a simplified pelvic floor exercise regimen. *Neurourol Urodyn.* 2004;23(1):48-53.
 155. Herschorn S, Becker D, Miller E, et al. Impact of a health education intervention in overactive bladder patients. *Can J Urol.* 2004 Dec;11(6):2430-7.
 156. Layton D, Pearce GL and Shakir SA. Safety profile of tolterodine as used in general practice in England: results of prescription-event monitoring. *Drug Saf.* 2001;24(9):703-13.
 157. Michel MC, Schneider T, Krege S, et al. Does gender or age affect the efficacy and safety of tolterodine? *J Urol.* 2002 Sep;168(3):1027-31.
 158. Zinner NR, Mattiasson A and Stanton SL. Efficacy, safety, and tolerability of extended-release once-daily tolterodine treatment for overactive bladder in older versus younger patients. *J Am Geriatr Soc.* 2002 May;50(5):799-807.

159. Hill S, Elhilali M, Millard RJ, et al. Long-term darifenacin treatment for overactive bladder in patients aged 65 years and older: analysis of results from a 2-year, open-label extension study. *Curr Med Res Opin.* 2007 Nov;23(11):2697-704.
160. Elinoff V, Bavendam T, Glasser DB, et al. Symptom-specific efficacy of tolterodine extended release in patients with overactive bladder: the IMPACT trial. *Int J Clin Pract.* 2006 Jun;60(6):745-51.
161. Staskin DR and Harnett MD. Effect of trospium chloride on somnolence and sleepiness in patients with overactive bladder. *Curr Urol Rep.* 2004 Dec;5(6):423-6.
162. Jacquetin B and Wyndaele J. Tolterodine reduces the number of urge incontinence episodes in patients with an overactive bladder. *Eur J Obstet Gynecol Reprod Biol.* 2001 Sep;98(1):97-102.
163. Appell RA, Abrams P, Drutz HP, et al. Treatment of overactive bladder: long-term tolerability and efficacy of tolterodine. *World J Urol.* 2001 Apr;19(2):141-7.
164. Anderson RU, MacDiarmid S, Kell S, et al. Effectiveness and tolerability of extended-release oxybutynin vs extended-release tolterodine in women with or without prior anticholinergic treatment for overactive bladder. *Int Urogynecol J Pelvic Floor Dysfunct.* 2006 Sep;17(5):502-11.
165. Gleason DM, Susset J, White C, et al. Evaluation of a new once-daily formulation of oxybutynin for the treatment of urinary urge incontinence. Ditropan XL Study Group. *Urology.* 1999 Sep;54(3):420-3.
166. Michel MC, de la Rosette JJ, Piro M, et al. Comparison of symptom severity and treatment response in patients with incontinent and continent overactive bladder. *Eur Urol.* 2005 Jul;48(1):110-5.
167. Garely AD, Lucente V, Vapnek J, et al. Solifenacin for overactive bladder with incontinence: symptom bother and health-related quality of life outcomes. *Ann Pharmacother.* 2007 Mar;41(3):391-8.
168. Landis JR, Kaplan S, Swift S, et al. Efficacy of antimuscarinic therapy for overactive bladder with varying degrees of incontinence severity. *J Urol.* 2004 Feb;171(2 Pt 1):752-6.
169. Kreder KJ, Jr., Brubaker L and Mainprize T. Tolterodine is equally effective in patients with mixed incontinence and those with urge incontinence alone. *BJU Int.* 2003 Sep;92(4):418-21.
170. Steers WD, Herschorn S, Kreder KJ, et al. Duloxetine compared with placebo for treating women with symptoms of overactive bladder. *BJU Int.* 2007 Aug;100(2):337-45.
171. Malone-Lee J, Henshaw DJ and Cummings K. Urodynamic verification of an overactive bladder is not a prerequisite for antimuscarinic treatment response. *BJU Int.* 2003 Sep;92(4):415-7.
172. Salvatore S, Serati M, Digesu GA, et al. Efficacy of tolterodine in relation to different urodynamic findings of detrusor overactivity. *Int Urogynecol J Pelvic Floor Dysfunct.* 2008 May;19(5):701-4.
173. Serati M, Salvatore S, Uccella S, et al. Urinary incontinence at orgasm: relation to detrusor overactivity and treatment efficacy. *Eur Urol.* 2008 Oct;54(4):911-5.
174. Freeman R, Hill S, Millard R, et al. Reduced perception of urgency in treatment of overactive bladder with extended-release tolterodine. *Obstet Gynecol.* 2003 Sep;102(3):605-11.
175. Varadharajan S, Jumadilova Z, Girase P, et al. Economic impact of extended-release tolterodine versus immediate- and extended-release oxybutynin among commercially insured persons with overactive bladder. *Am J Manag Care.* 2005 Jul;11(4 Suppl):S140-9.
176. Hall JA, Nelson MA, Meyer JW, et al. Costs and resources associated with the treatment of overactive bladder using retrospective medical care claims data. *Manag Care Interface.* 2001 Aug;14(8):69-75.
177. Jensen GA, Zhou Z and Torigoe Y. Medical care of overactive bladder in elderly patients. *Journal of Aging & Pharmacotherapy.* 2003;13(1):13-27.
178. Nitz NM, Jumadilova Z, Darkow T, et al. Medical costs after initiation of drug treatment for overactive bladder: effects of selection bias on cost estimates. *Am J Manag Care.* 2005 Jul;11(4 Suppl):S130-9.
179. Hu TW, Wagner TH, Bentkover JD, et al. Estimated economic costs of overactive bladder in the United States. *Urology.* 2003 Jun;61(6):1123-8.
180. Abrams P, Cardozo L, Fall M, et al. The standardisation of terminology of lower urinary tract function: report from the Standardisation Sub-committee of the International Continence Society. *Neurourol Urodyn.* 2002;21(2):167-78.
181. Gardyn R. Detrol. *Advertising Age.* 1999 June 28, 1999.
182. Fisher JA. *Medical Research for Hire: The Political Economy of Pharmaceutical Clinical Trials.* New Brunswick, NJ: Rutgers University Press; 2009.
183. Kassirer JP. *On the Take: How America's Complicity with Big Business Can Endanger Your Health.* New York: Oxford University Press; 2005.
184. Ronald L. *Empowered to Consume: Direct-to-consumer advertising and pharmaceutical governance.* Troy, NY: Rensselaer Polytechnic Institute; 2006.
185. Petersen M. *Our Daily Meds: How the Pharmaceutical Companies Transformed Themselves into Slick Marketing Machines and Hooked the Nation on Prescription Drugs* New York: Farrar, Straus and Giroux; 2008.
186. Pfizer. *Pfizer Achieves Key 2006 Financial Targetes.* 2007 [updated 2007; cited 2009 January 8]; Available from: <http://www.prnewswire.com/cgi-bin/stories.pl?ACCT=104&STORY=/www/story/01-22-2007/0004509853&EDATE=>.
187. Bren L. *Controlling Urinary Incontinence.* FDA Consumer Magazine. 2005 Sept-Oct.
188. Brown JS, Vittinghoff E, Wyman JF, et al. Urinary incontinence: does it increase risk for falls and fractures? Study of Osteoporotic Fractures Research Group. *J Am Geriatr Soc.* 2000 Jul;48(7):721-5.
189. Darkow T, Fontes CL and Williamson TE. Costs associated with the management of overactive bladder and related comorbidities. *Pharmacotherapy.* 2005 Apr;25(4):511-9.
190. Matchar DB, Myers ER, Barber MW, et al. Management of uterine fibroids. *Evid Rep Technol Assess (Summ).* 2001;Jan(34):1-6.
191. Schulz KF, Chalmers I, Grimes DA, et al. Assessing the quality of randomization from reports of controlled

- trials published in obstetrics and gynecology journals. *JAMA*. 1994 Jul 13;272(2):125-8.
192. Schulz KF, Chalmers I, Hayes RJ, et al. Empirical evidence of bias. Dimensions of methodological quality associated with estimates of treatment effects in controlled trials. *JAMA*. 1995 Feb 1;273(5):408-12.
 193. Altman DG, Schulz KF, Moher D, et al. The revised CONSORT statement for reporting randomized trials: explanation and elaboration. *Ann Intern Med*. 2001 Apr 17;134(8):663-94.
 194. West S, King V, Carey TS, et al. Systems to rate the strength of scientific evidence. *Evid Rep Technol Assess (Summ)*. 2002;Mar(47):1-11.
 195. Viswanathan M, Hartmann K, McKoy N, et al. Management of Uterine Fibroids: An Update of the Evidence. Evidence Report/Technology Assessment No. 154 (Prepared by RTI International-University of North Carolina Evidence-based Practice Center under Contract No. 290-02-0016. AHRQ Publication No. 07-E011. Rockville, MD: Agency for Healthcare Research and Quality; 2007.
 196. Shamlivan T, Wyman J, Bliss DZ, et al. Prevention of urinary and fecal incontinence in adults. *Evid Rep Technol Assess (Full Rep)*. 2007 Dec(161):1-379.
 197. Wilkinson E. Urinary incontinence in women. *Nurs Times*. 2006 Nov 28-Dec 4;102(48):23-4.
 198. Abrams P, Cardozo L, Chapple C, et al. Comparison of the efficacy, safety, and tolerability of propiverine and oxybutynin for the treatment of overactive bladder syndrome. *Int J Urol*. 2006 Jun;13(6):692-8.
 199. Anderson RU, Mobley D, Blank B, et al. Once daily controlled versus immediate release oxybutynin chloride for urge urinary incontinence. OROS Oxybutynin Study Group. *J Urol*. 1999 Jun;161(6):1809-12.
 200. Appell RA, Sand P, Dmochowski R, et al. Prospective randomized controlled trial of extended-release oxybutynin chloride and tolterodine tartrate in the treatment of overactive bladder: results of the OBJECT Study. *Mayo Clin Proc*. 2001 Apr;76(4):358-63.
 201. Arruda RM, Castro RA, Sousa GC, et al. Prospective randomized comparison of oxybutynin, functional electrostimulation, and pelvic floor training for treatment of detrusor overactivity in women. *Int Urogynecol J Pelvic Floor Dysfunct*. 2008 Aug;19(8):1055-61.
 202. Barkin J, Corcos J, Radomski S, et al. A randomized, double-blind, parallel-group comparison of controlled- and immediate-release oxybutynin chloride in urge urinary incontinence. *Clin Ther*. 2004 Jul;26(7):1026-36.
 203. Bemelmans BL, Kiemeny LA and Debruyne FM. Low-dose oxybutynin for the treatment of urge incontinence: good efficacy and few side effects. *Eur Urol*. 2000 Jun;37(6):709-13.
 204. Boone TB, Kusek JW, Nyberg LM, et al. Treatment patterns and associated symptom improvement during six months of care for overactive bladder: a prospective, observational study. *Clin Ther*. 2002 Mar;24(3):397-408.
 205. Capo JP, Jr., Laramée C, Lucente V, et al. Solifenacin treatment for overactive bladder in Hispanic patients: patient-reported symptom bother and quality of life outcomes from the VESiCare Open-Label Trial. *Int J Clin Pract*. 2008 Jan;62(1):39-46.
 206. Cardozo L, Hessdorfer E, Milani R, et al. Solifenacin in the treatment of urgency and other symptoms of overactive bladder: results from a randomized, double-blind, placebo-controlled, rising-dose trial. *BJU Int*. 2008 Nov;102(9):1120-7.
 207. Cardozo L, Lisek M, Millard R, et al. Randomized, double-blind placebo controlled trial of the once daily antimuscarinic agent solifenacin succinate in patients with overactive bladder. *J Urol*. 2004 Nov;172(5 Pt 1):1919-24.
 208. Chancellor MB, Zinner N, Whitmore K, et al. Efficacy of solifenacin in patients previously treated with tolterodine extended release 4 mg: results of a 12-week, multicenter, open-label, flexible-dose study. *Clin Ther*. 2008 Oct;30(10):1766-81.
 209. Chancellor M, Freedman S, Mitcheson HD, et al. Tolterodine, an effective and well tolerated treatment for urge incontinence and other overactive bladder symptoms. *Clinical Drug Investigation*. 2000;19(2):83-91.
 210. Chapple C, DuBeau C, Ebinger U, et al. Darifenacin treatment of patients >or= 65 years with overactive bladder: results of a randomized, controlled, 12-week trial. *Curr Med Res Opin*. 2007 Oct;23(10):2347-58.
 211. Chapple CR, Martinez-Garcia R, Selvaggi L, et al. A comparison of the efficacy and tolerability of solifenacin succinate and extended release tolterodine at treating overactive bladder syndrome: results of the STAR trial. *Eur Urol*. 2005 Sep;48(3):464-70.
 212. Choo MS, Doo CK and Lee KS. Satisfaction with tolterodine: assessing symptom-specific patient-reported goal achievement in the treatment of overactive bladder in female patients (STARGATE study). *Int J Clin Pract*. 2008 Feb;62(2):191-6.
 213. Chu FM, Dmochowski RR, Lama DJ, et al. Extended-release formulations of oxybutynin and tolterodine exhibit similar central nervous system tolerability profiles: a subanalysis of data from the OPERA trial. *Am J Obstet Gynecol*. 2005 Jun;192(6):1849-54; discussion 1854-5.
 214. Diokno A, Sand P, Labasky R, et al. Long-term safety of extended-release oxybutynin chloride in a community-dwelling population of participants with overactive bladder: a one-year study. *Int Urol Nephrol*. 2002;34(1):43-9.
 215. Dmochowski R, Kreder K, MacDiarmid S, et al. The clinical efficacy of tolterodine extended-release is maintained for 24 h in patients with overactive bladder. *BJU Int*. 2007 Jul;100(1):107-10.
 216. Drutz HP, Appell RA, Gleason D, et al. Clinical efficacy and safety of tolterodine compared to oxybutynin and placebo in patients with overactive bladder. *Int Urogynecol J Pelvic Floor Dysfunct*. 1999;10(5):283-9.
 217. Dwyer P, Kelleher C, Young J, et al. Long-term benefits of darifenacin treatment for patient quality of life: results from a 2-year extension study. *Neurourol Urodyn*. 2008;27(6):540-7.
 218. Garely AD, Kaufman JM, Sand PK, et al. Symptom bother and health-related quality of life outcomes following solifenacin treatment for overactive bladder: the VESiCare Open-Label Trial (VOLT). *Clin Ther*. 2006 Nov;28(11):1935-46.
 219. Haab F, Corcos J, Siami P, et al. Long-term treatment with darifenacin for overactive bladder: results of a 2-year, open-label extension study. *BJU Int*. 2006 Nov;98(5):1025-32.

220. Haab F, Cardozo L, Chapple C, et al. Long-term open-label solifenacin treatment associated with persistence with therapy in patients with overactive bladder syndrome. *Eur Urol*. 2005 Mar;47(3):376-84.
221. Jonas U, Hofner K, Madersbacher H, et al. Efficacy and safety of two doses of tolterodine versus placebo in patients with detrusor overactivity and symptoms of frequency, urge incontinence, and urgency: urodynamic evaluation. The International Study Group. *World J Urol*. 1997;15(2):144-51.
222. Karram MM, Togli MR, Serels SR, et al. Treatment with solifenacin increases warning time and improves symptoms of overactive bladder: results from VENUS, a randomized, double-blind, placebo-controlled trial. *Urology*. 2009 Jan;73(1):14-8.
223. Kelleher CJ, Kreder KJ, Pleil AM, et al. Long-term health-related quality of life of patients receiving extended-release tolterodine for overactive bladder. *Am J Manag Care*. 2002 Dec;8(19 Suppl):S616-30.
224. Kelleher CJ, Reese PR, Pleil AM, et al. Health-related quality of life of patients receiving extended-release tolterodine for overactive bladder. *Am J Manag Care*. 2002 Dec;8(19 Suppl):S608-15.
225. Kelleher CJ, Cardozo L, Chapple CR, et al. Improved quality of life in patients with overactive bladder symptoms treated with solifenacin. *BJU Int*. 2005 Jan;95(1):81-5.
226. Koonings PP and Bergman A. Urethral pressure changes in women with detrusor instability. Bladder or urethral pathologic process? *Urology*. 1991 Jun;37(6):540-2.
227. Kreder K, Mayne C and Jonas U. Long-term safety, tolerability and efficacy of extended-release tolterodine in the treatment of overactive bladder. *Eur Urol*. 2002 Jun;41(6):588-95.
228. Leung HY, Yip SK, Cheon C, et al. A randomized controlled trial of tolterodine and oxybutynin on tolerability and clinical efficacy for treating Chinese women with an overactive bladder. *BJU Int*. 2002 Sep;90(4):375-80.
229. Mallett V, Burks D, Garely AD, et al. Solifenacin treatment for overactive bladder in black patients: patient-reported symptom bother and health-related quality of life outcomes. *Curr Med Res Opin*. 2007 Apr;23(4):821-31.
230. Messelink EJ. Treatment of the overactive bladder with tolterodine, a new muscarinic receptor antagonist. *BJU Int*. 1999 Mar;83 Suppl 2:48-52.
231. Michel MC, Wetterauer U, Vogel M, et al. Cardiovascular safety and overall tolerability of solifenacin in routine clinical use: a 12-week, open-label, post-marketing surveillance study. *Drug Saf*. 2008;31(6):505-14.
232. Michel MC, Oelke M, Goepel M, et al. Relationships among symptoms, bother, and treatment satisfaction in overactive bladder patients. *Neurourol Urodyn*. 2007;26(2):190-5.
233. Millard R, Tuttle J, Moore K, et al. Clinical efficacy and safety of tolterodine compared to placebo in detrusor overactivity. *J Urol*. 1999 May;161(5):1551-5.
234. Minassian VA, Ross S, Sumabat O, et al. Randomized trial of oxybutynin extended versus immediate release for women aged 65 and older with overactive bladder: lessons learned from conducting a trial. *J Obstet Gynaecol Can*. 2007 Sep;29(9):726-32.
235. Moore KH and Sutherst JR. Response to treatment of detrusor instability in relation to psychoneurotic status. *Br J Urol*. 1990 Nov;66(5):486-90.
236. Moore KH, Hay DM, Imrie AE, et al. Oxybutynin hydrochloride (3 mg) in the treatment of women with idiopathic detrusor instability. *Br J Urol*. 1990 Nov;66(5):479-85.
237. Preik M, Albrecht D, O'Connell M, et al. Effect of controlled-release delivery on the pharmacokinetics of oxybutynin at different dosages: severity-dependent treatment of the overactive bladder. *BJU Int*. 2004 Oct;94(6):821-7.
238. Rentzhog L, Stanton SL, Cardozo L, et al. Efficacy and safety of tolterodine in patients with detrusor instability: a dose-ranging study. *Br J Urol*. 1998 Jan;81(1):42-8.
239. Roberts R, Bavendam T, Glasser DB, et al. Tolterodine extended release improves patient-reported outcomes in overactive bladder: results from the IMPACT trial. *Int J Clin Pract*. 2006 Jun;60(6):752-8.
240. Robinson D, Cardozo L, Terpstra G, et al. A randomized double-blind placebo-controlled multicentre study to explore the efficacy and safety of tamsulosin and tolterodine in women with overactive bladder syndrome. *BJU Int*. 2007 Oct;100(4):840-5.
241. Rogers R, Bachmann G, Jumadilova Z, et al. Efficacy of tolterodine on overactive bladder symptoms and sexual and emotional quality of life in sexually active women. *Int Urogynecol J Pelvic Floor Dysfunct*. 2008 Nov;19(11):1551-7.
242. Rudy D, Cline K, Harris R, et al. Multicenter phase III trial studying trospium chloride in patients with overactive bladder. *Urology*. 2006 Feb;67(2):275-80.
243. Salvatore S, Serati M, Ghezzi F, et al. Efficacy of tolterodine in women with detrusor overactivity and anterior vaginal wall prolapse: is it the same? *BJOG*. 2007 Nov;114(11):1436-8.
244. Salvatore S, Khullar V, Cardozo L, et al. Long-term prospective randomized study comparing two different regimens of oxybutynin as a treatment for detrusor overactivity. *Eur J Obstet Gynecol Reprod Biol*. 2005 Apr 1;119(2):237-41.
245. Sand P, Zinner N, Newman D, et al. Oxybutynin transdermal system improves the quality of life in adults with overactive bladder: a multicentre, community-based, randomized study. *BJU Int*. 2007 Apr;99(4):836-44.
246. Sand PK, Goldberg RP, Dmochowski RR, et al. The impact of the overactive bladder syndrome on sexual function: a preliminary report from the Multicenter Assessment of Transdermal Therapy in Overactive Bladder with Oxybutynin trial. *Am J Obstet Gynecol*. 2006 Dec;195(6):1730-5.
247. Tseng LH, Wang AC, Chang YL, et al. Randomized comparison of tolterodine with vaginal estrogen cream versus tolterodine alone for the treatment of postmenopausal women with overactive bladder syndrome. *Neurourol Urodyn*. 2009;28(1):47-51.
248. Versi E, Appell R, Mobley D, et al. Dry mouth with conventional and controlled-release oxybutynin in urinary incontinence. The Ditropan XL Study Group. *Obstet Gynecol*. 2000 May;95(5):718-21.

249. Wang PS, Levin R, Zhao SZ, et al. Urinary antispasmodic use and the risks of ventricular arrhythmia and sudden death in older patients. *J Am Geriatr Soc.* 2002 Jan;50(1):117-24.
250. Zinner N, Noe L, Rasouliyan L, et al. Impact of solifenacin on resource utilization, work productivity and health utility in overactive bladder patients switching from tolterodine ER. *Curr Med Res Opin.* 2008 Jun;24(6):1583-91.
251. Burgio KL, Locher JL and Goode PS. Combined behavioral and drug therapy for urge incontinence in older women. *J Am Geriatr Soc.* 2000 Apr;48(4):370-4.
252. Fantl JA, Hurt WG and Dunn LJ. Detrusor instability syndrome: the use of bladder retraining drills with and without anticholinergics. *Am J Obstet Gynecol.* 1981 Aug 15;140(8):885-90.
253. Ghei M, Miller R and Malone-Lee J. Case series data to encourage randomized trials of bladder retraining compared to antimuscarinic agents. *J Urol.* 2006 Apr;175(4):1411-5; discussion 1415-6.
254. Lauti M, Herbison P, Hay-Smith J, et al. Anticholinergic drugs, bladder retraining and their combination for urge urinary incontinence: a pilot randomised trial. *Int Urogynecol J Pelvic Floor Dysfunct.* 2008 Nov;19(11):1533-43.
255. Szonyi G, Collas DM, Ding YY, et al. Oxybutynin with bladder retraining for detrusor instability in elderly people: a randomized controlled trial. *Age Ageing.* 1995 Jul;24(4):287-91.
256. Oerlemans DJ and van Kerrebroeck PE. Sacral nerve stimulation for neuromodulation of the lower urinary tract. *Neurourol Urodyn.* 2008;27(1):28-33.
257. Cooperberg MR and Stoller ML. Percutaneous neuromodulation. *Urol Clin North Am.* 2005 Feb;32(1):71-8, vii.
258. van Balken MR, Vandoninck V, Gisolf KW, et al. Posterior tibial nerve stimulation as neuromodulative treatment of lower urinary tract dysfunction. *J Urol.* 2001 Sep;166(3):914-8.
259. Vandoninck V, Van Balken MR, Finazzi Agro E, et al. Posterior tibial nerve stimulation in the treatment of urge incontinence. *Neurourol Urodyn.* 2003;22(1):17-23.
260. McGuire EJ, Zhang SC, Horwinski ER, et al. Treatment of motor and sensory detrusor instability by electrical stimulation. *J Urol.* 1983 Jan;129(1):78-9.
261. Goldberg RP and Sand PK. Electromagnetic pelvic floor stimulation: applications for the gynecologist. *Obstet Gynecol Surv.* 2000 Nov;55(11):715-20.
262. Giannantoni A, Di Stasi SM, Stephen RL, et al. Intravesical resiniferatoxin versus botulinum-A toxin injections for neurogenic detrusor overactivity: a prospective randomized study. *J Urol.* 2004 Jul;172(1):240-3.
263. Karsenty G, Reitz A, Lindemann G, et al. Persistence of therapeutic effect after repeated injections of botulinum toxin type A to treat incontinence due to neurogenic detrusor overactivity. *Urology.* 2006 Dec;68(6):1193-7.
264. Kuo HC. Urodynamic evidence of effectiveness of botulinum A toxin injection in treatment of detrusor overactivity refractory to anticholinergic agents. *Urology.* 2004 May;63(5):868-72.
265. Brubaker L, Richter HE, Visco A, et al. Refractory idiopathic urge urinary incontinence and botulinum A injection. *J Urol.* 2008 Jul;180(1):217-22.
266. McHorney CA, Ware JE, Jr. and Raczek AE. The MOS 36-Item Short-Form Health Survey (SF-36): II. Psychometric and clinical tests of validity in measuring physical and mental health constructs. *Med Care.* 1993 Mar;31(3):247-63.
267. Wagner TH, Patrick DL, Bavendam TG, et al. Quality of life of persons with urinary incontinence: development of a new measure. *Urology.* 1996 Jan;47(1):67-71; discussion 71-2.
268. Sorensen SS, Nielsen JB, Norgaard JP, et al. Changes in bladder volumes with repetition of water cystometry. *Urol Res.* 1984;12(4):205-8.
269. Wang AC, Wang YY and Chen MC. Single-blind, randomized trial of pelvic floor muscle training, biofeedback-assisted pelvic floor muscle training, and electrical stimulation in the management of overactive bladder. *Urology.* 2004 Jan;63(1):61-6.
270. Nabi G, Cody JD, Ellis G, et al. Anticholinergic drugs versus placebo for overactive bladder syndrome in adults. *Cochrane Database Syst Rev.* 2006(4):CD003781.
271. Chapple CR, Khullar V, Gabriel Z, et al. The effects of antimuscarinic treatments in overactive bladder: an update of a systematic review and meta-analysis. *Eur Urol.* 2008 Sep;54(3):543-62.
272. D'Souza AO, Smith MJ, Miller LA, et al. Persistence, adherence, and switch rates among extended-release and immediate-release overactive bladder medications in a regional managed care plan. *J Manag Care Pharm.* 2008 Apr;14(3):291-301.
273. Balkrishnan R, Bhosle MJ, Camacho FT, et al. Predictors of medication adherence and associated health care costs in an older population with overactive bladder syndrome: a longitudinal cohort study. *J Urol.* 2006 Mar;175(3 Pt 1):1067-71; discussion 1071-2.
274. Lawrence M, Guay DR, Benson SR, et al. Immediate-release oxybutynin versus tolterodine in detrusor overactivity: a population analysis. *Pharmacotherapy.* 2000 Apr;20(4):470-5.
275. Yu YF, Nichol MB, Yu AP, et al. Persistence and adherence of medications for chronic overactive bladder/urinary incontinence in the california medicaid program. *Value Health.* 2005 Jul-Aug;8(4):495-505.

List of Acronyms/Abbreviations

±	plus or minus
≤	less than or equal to
≥	greater than or equal to
AE	adverse events
AHCPR	Agency for Health Care Policy and Research
AHRQ	Agency for Healthcare Research and Quality
AUC	area under the curve
AUM	ambulatory urodynamic monitoring
avg.	average
BAPFMT	biofeedback-assisted pelvic floor muscle training
BFQ	Bladder Function Questionnaire
b.i.d.	twice a day
BL	baseline
BM	bowel movements
BMI	body mass index
BOO	bladder outlet obstruction
BPH	benign prostatic hyperplasia
bpm	beats per minute
BRD	bladder retraining drill
BT	bladder training
CEE	Conjugated equine estrogens
CHF	Congestive heart failure
CI	confidence interval(s)
cm	centimeter
cmH ₂ O	centimeters of water
CR	controlled release
CS	cognitive strategies
CT	combination therapy
CUBS	Compromised urinary bladder syndrome
d	day
d/t	drug treatment
DI	Detrusor instability
dL	deciliter
DM	Diabetes mellitus
DO	detrusor overactivity
Dx	diagnosis
EKG	electrocardiogram
ER	extended release
ES	electrical stimulation (electrostimulation)
etc.	et cetera
EtOH	Ethanol (alcohol)
F	F-distribution
G	group
GI	gastrointestinal
GII	global impression of improvement
GSI	genuine stress incontinence
GU	genitourinary
HAD	Hospital Anxiety and Depression Scale
H ₂ O	water
hr(s)	hour(s)
HRQoL	Health related quality of life
HRT	hormone replacement therapy

Hx	history
Hz	hertz
IBD	irritable bowel disease
IC	interstitial cystitis
ICIQ	International Consultation on Incontinence Modular Questionnaire
ICIQ-SF	International Consultation on Incontinence Questionnaire-Short Form
IIQ	Incontinence Impact Questionnaire
IIQ-R	Incontinence Impact Questionnaire-Revised
IMPACT	Improvement in Patients: Assessing symptomatic control with tolterodine
IQR	interquartile range
IR	immediate release
ITT	intention to treat
IUSS	Indevus Urgency Severity Scale
KQ	key question
kg/m ²	kilograms per meter squared
KHQ	King's Health Questionnaire
L	liter
LCB	low-compliance bladder
LOCF	last observation carried forward
LOS	length of stay
LS	least square
Ltd	limited
LTFU	Loss to followup
LUT	Lower urinary tract
LUTS	Lower urinary tract syndrome
mg	milligram
min	minute(s)
mL	milliliter
mL/s	milliliters per second
mm	millimeter
MMSE	Mini Mental Status Exam
mo(s)	month(s)
MUI	Mixed urinary incontinence
MVV	Mean volume voided
n, N	number
ng	nanogram
NIH	National Institutes of Health
NR	not reported
NS	Not significant
OAB	Overactive bladder
OAB-q	OAB questionnaire
OAB-SCS	OAB-Symptom Composite Score
OCAS	oral controlled absorption system
P, p	p value
PFME	pelvic floor muscle exercises
PFMT	pelvic floor muscle training
PGA	patient-reported goal achievement
PISQ	Pelvic Organ Prolapse/Urinary Incontinence Sexual Questionnaire
PME	pelvic muscle exercises
PNN50	measure of heart rate variability
P.O.	per oral (by mouth)
PPBC	Patient perception of bladder condition

PPBCS	Patient perception of bladder condition scale
PPIUS	Patient Perception of Intensity of Urgency Scale
PPPY	per person per year
PRO	Patient reported outcome(s)
Pt	patient
PVR	post-void residual
qAM	every morning
qd	every day
qid	four times per day
QoL	quality of life
QTcB	QT interval correct for heart rate using Bazett's formula
RCT	randomized controlled trial
RTX	resiniferatoxin
s	second
SCL-90-R	Symptom Checklist-90-Revised
SD	standard deviation
SE	standard error
SF	Sexual function
sec	second
SNM	sacroneuromodulation
SQoL-F	Sexual Quality of Life Questionnaire-Female
subj.	subjects
SUI	Stress urinary incontinence
Sx	symptoms
t.i.d.	three times a day
TDS	transdermal delivery system
UDI	Urogenital Distress Inventory
UDS	urodynamics
UFICQ	Urinary Frequency and Incontinence Questionnaire
UI	urinary incontinence
US	United States
UTI	urinary tract infection
UUDI	Urge Urogenital Distress Inventory
UUI	Urge urinary incontinence
VAS	visual analog scale
vs., v	versus
w/	with
wk(s)	week(s)
x	times
yr(s)	year(s)

APPENDIX A. Exact Search Strings

PubMed search strategies (last updated October 1, 2008)

	Search terms	Search results
#1	("Urinary Bladder, Overactive"[Mh] OR "overactive bladder" OR "urge incontinence" OR urinary incontinence, urge[mh] OR "detrusor instability" OR "overactive detrusor" OR "urinary urgency" OR "urinary frequency" OR "irritable bladder" OR "detrusor overactivity") AND "female"[MeSH Terms] AND "humans"[MeSH Terms] AND English[lang]	2886
#2	#1 AND editorial[pt]	10
#3	#1 AND letter[pt]	30
#4	#1 AND case reports[pt]	164
#5	#1 AND review[pt]	299
#6	#1 AND practice guideline[pt]	2
#7	#1 NOT (#2 OR #3 OR #4 OR #5 OR #6)	2400*†

* Approximately 250 of these citations represent pediatric literature (due to variability in indexing for this topic, we were unable to exclude pediatric literature at the search strategy level).

† Numbers do not total due to exclusions in more than one category; 5 items were indexed as both letters and case reports and 14 items were indexed as both reviews and case reports

EMBASE search (OVID) (last updated October 1, 2008)

	Search Terms	Search Results
#1	*overactive bladder/ or *urinary urgency/ or *urge incontinence/ or *urinary frequency/ or *detrusor dyssynergia/ or *bladder irritation/	1624
#2	limit 1 to (human and female and english language and (adult <18 to 64 years> or aged <65+ years>))	363
#3	#2 and review.pt.	12
#4	#2 and conference paper.pt.	4
#5	#2 and editorial.pt.	1
#6	#2 and letter.pt.	0
#7	#2 and note.pt.	3
#8	#2 and short survey.pt.	4
#9	#2 and case report/	18
#10	#2 and practice guideline/	4
#11	#2 and "systematic review"/	1
#12	#2 and meta analysis/	1
#13	#2 not (3 or 4 or 5 or 6 or 7 or 8 or 9 or 10 or 11 or 12)	318*†

* Overlap with PubMed: 310 citations; 8 new citations retrieved for inclusion.

† Numbers do not total due to exclusions in more than one category: 1 item was indexed as both a case report and review; 1 item was indexed as both a case report and a note; and 1 item was indexed as both a review and a systematic review.

CINAHL search (EBSCO) (last updated October 1, 2008)

	Search Terms	Search Results
#1	(MH "Urge Incontinence") or (MH "Overactive Bladder") or "overactive bladder" or "urge incontinence" or "urge urinary incontinence" or "detrusor instability" or "overactive detrusor" or "urinary urgency" or "urinary frequency" or "detrusor overactivity") and (MH "Adult+") and (ZL "ENGLISH") and (PT "Journal Article")	305
#2	#1 and case reports	18
#3	#1 and review	6
#4	#1 and CE material	5
#5	#1 and abstract/commentary	7
#6	#1 and consumer literature	5
#7	1 not (2 or 3 or 4 or 5 or 6)	264*

* Overlap with PubMed: 240 citations; 24 new citations retrieved for inclusion.

APPENDIX B. Sample Data Abstract Forms

Systematic Review of the Etiology and Treatment of Overactive Bladder in Women Abstract Review Form

First Author, Year: _____

Reference # _____

Abstractor Initials: ____ _

Primary Inclusion/Exclusion Criteria			
1. Applies to SER topic (If not, select at least one of the following reasons): a. ___ Not OAB (including post-operative/iatrogenic) b. ___ Stress or mixed incontinence c. ___ Isolated nocturia d. ___ Interstitial cystitis/painful bladder syndrome e. ___ Pelvic organ prolapse f. ___ Neurogenic conditions g. ___ Basic science or anatomy only h. ___ Imaging/diagnostic study only i. ___ Other _____	Yes	No	Cannot Determine
2. Original research (exclude editorials, commentaries, letters to editor, reviews, etc)	Yes	No	Cannot Determine
3. Study published in English	Yes	No	
4. Adult female study population (or includes women)	Yes	No	Cannot Determine
5. Ambulatory population (exclude if exclusively institutionalized or home-bound)	Yes	No	
6. Eligible Study type a. ___ RCT b. ___ Cohorts with comparison c. ___ Case-control d. ___ Case series e. ___ Incidence/prevalence in representative populations f. ___ Cost of treatment in US populations (monetary & non-monetary)	Yes	No	Cannot Determine
7. Eligible study size Record N if < 50 relevant subjects enrolled: _____	Yes	No	Cannot Determine

OAB is operationalized as idiopathic urinary urgency and frequency

Retain for:

___ **BACKGROUND/DISCUSSION**

___ **REVIEW OF REFERENCES**

___ **Other** _____

COMMENTS :

**Systematic Review of the Treatment Alternatives of Overactive Bladder in Women
Full-text Review Form**

First Author, Year: _____

Reference # _____

Abstractor Initials: _____

OAB is operationalized as idiopathic urinary urgency and frequency

Primary Inclusion/Exclusion Criteria		
<p>8. Applies to SER topic (If not, select at least one of the following reasons):</p> <ul style="list-style-type: none"> a. ___ Not OAB (including post-operative/iatrogenic) b. ___ Stress or mixed incontinence c. ___ Isolated nocturia d. ___ Interstitial cystitis/painful bladder syndrome e. ___ Pelvic organ prolapse f. ___ Neurogenic conditions g. ___ Basic science or anatomy only h. ___ Imaging/diagnostic study only i. ___ Other _____ 	Yes	No
<p>9. Original research (exclude editorials, commentaries, letters to editor, reviews, etc)</p>	Yes	No
<p>10. Study published in English</p>	Yes	No
<p>11. Adult female study population (or reports data by gender) If No, % female _____</p>	Yes	No
<p>12. Ambulatory population (exclude if exclusively institutionalized or home-bound)</p>	Yes	No
<p>13. Eligible Study type</p> <ul style="list-style-type: none"> g. ___ RCT /CCT h. ___ Cohorts with comparison i. ___ Case-control j. ___ Case series k. ___ Incidence/Prevalence study (survey-based) l. ___ Cost benefit/utility/effectiveness study 	Yes	No
<p>14. Eligible study size Record N if < 50 relevant subjects enrolled: _____</p>	Yes	No
<p>15. Does study address one of the following:</p> <ul style="list-style-type: none"> a. ___ Treatment of OAB b. ___ Incidence/prevalence of OAB c. ___ Monetary costs of treatment d. ___ Non-monetary costs/harms of treatment 	Yes	No

EXCLUDE IF AN ITEM IN A GRAY BOX IS SELECTED

Content Inventory

1. _____ Treatment of women with symptoms of OAB

- a. _____ Pharmacologic
- b. _____ Surgical
 - i. _____ Botox
 - ii. _____ Central neuromodulation
 _____ Sacral
 - iii. _____ Peripheral neuromodulation
 _____ Tibial
 _____ Pudendal
 - iv. _____ Augmentation cystoplasty
 - v. _____ Other _____
- c. _____ Behavioral/Physical Therapy
- d. _____ Complementary and alternative therapies
- e. _____ Other _____

2. Modification of outcomes by:

- a. _____ Age
- b. _____ Body habitus/BMI
- c. _____ Clinical presentation, physical exam findings, urodynamic findings, symptom cluster
- d. _____ Diabetes
- e. _____ Functional status
- f. _____ Hormone replacement therapy
- g. _____ Menopausal status
- h. _____ Parity/post-partum/route-of-delivery
- i. _____ Prior treatment
- j. _____ Race/ethnicity
- k. _____ Smoking
- l. _____ Hysterectomy
- m. _____ Other factors _____

Length of follow-up: _____

Retain for:

- _____ **BACKGROUND/DISCUSSION**
- _____ **REVIEW OF REFERENCES**
- _____ **Other** _____

COMMENTS :

Evidence Table

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Author:	Design:	Inclusion criteria:	Incontinence:	Outcomes:	
Country and setting:	Intervention:	Exclusion criteria:	Urgency:	Modifiers	
Enrollment period:	Groups:		Frequency:		
Funding:	N at enrollment:				
Author industry relationship disclosures:	N at follow-up:				
	Age, yrs ± SD:				
	Race/ethnicity, mean ± SD:				
	Women, N (%):				
	Parity mean ± SD:				

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence	Quality Rating
<p>Author: Aaron et al., 2002</p> <p>Country: India</p> <p>Study period: NR</p> <p>Funding: Department of Community Health and Development, Christian Medical College</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Cross-sectional administered questionnaire</p> <p>Base population: Permanent residents of the Kaniambadi rural development block of Vellore district in Tamil Nadu</p> <p>Sampling frame: Residents of 7 representative villages as enumerated by census; menopausal participant selected then matched to premenopausal control in same age strata</p> <p>N sampled: NR</p> <p>N screened: NR</p> <p>N eligible: Menopausal: 100 Pre-menopausal: 100</p> <p>N respondents: NR</p> <p>N included: Menopausal: 100 Pre-menopausal: 100</p> <p>Age, mean ± SD: Menopausal: 46.6 ± 2.2 Pre-menopausal: 45.4 ± 2.3</p> <p>Race/ethnicity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> women age 40 to 49 <p>Exclusion criteria:</p> <ul style="list-style-type: none"> hysterectomy <p>Effective response: NR</p>	<p>Urgency: NR</p> <p>Frequency: NR</p>	<p>Prevalence of urgency, %: Menopausal: 18 Pre-menopausal: 8</p> <p>Prevalence of frequency, %: Menopausal: 17 Pre-menopausal: 11</p>	<p>Overall quality: Poor</p> <p>Internal validity score: 2, -</p> <p>External Validity Score: 2, -</p> <p>Sampling Method Described: +</p> <p>N sampled provided: -</p> <p>N eligible provided: -</p> <p>N included respondents: +</p> <p>Response Rate:* NR</p> <p>Inclusion (Exclusion) Specified: +</p> <p>Age of population described: +</p> <p>Operational definition provided*: -</p> <p>Required frequency defined*: -</p>

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence[^]	Quality Rating
Author: Bogren et al., 1997 Country: Sweden Study period: NR Funding: NR Author industry relationship disclosures: NR	Design: Cross sectional mailed questionnaire Base population: Primary healthcare district in southwest Sweden Sampling frame: All residents N screened: Total: 458 Women: 225 N eligible: Total: 458 Women: 225 N respondents: Total: 419 Women: 216 N included: Total: 419 Women: 216 Age, %: 65: 100 Race/ethnicity: NR	Inclusion criteria: UUI: • adults • 65 years old Exclusion criteria: NR Effective response, %: 96 [^]	UUI: NR	Prevalence UUI, %: 18.5	Overall quality: Poor Internal validity score: 5, ++ External Validity Score: 1,- Sampling Method Described: + N sampled provided: + N eligible provided: + N included respondents: + Response Rate:* 96 Inclusion (Exclusion) Specified: + Age of population described: 0? Operational definition provided*: - Required frequency defined*: -

[^]Data presented for women only.

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence[^]	Quality Rating
<p>Author: Bortolotti et al., 2000</p> <p>Country: Italy</p> <p>Study period: March 1997 to October 1997</p> <p>Funding: Pharmacia & Upjohn Italia</p> <p>Author industry relationship disclosures: At least 2 of 12 Pharmacia & Upjohn (2)</p>	<p>Design: Cross-sectional telephone questionnaire</p> <p>Base population: Registered participants in primary care networks in six areas of Italy</p> <p>Sampling frame: NR</p> <p>N sampled: NR</p> <p>N screened: NR</p> <p>N eligible: NR</p> <p>N respondents: Women: 2,767</p> <p>N included: Women: 2,767</p> <p>Age, %:[^] 40-50: 22.6 51-60: 30.4 61-70: 19.8 > 70: 27.2</p> <p>Race/ethnicity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • adults • women age ≥ 40 • men age ≥ 50 <p>Exclusion criteria: NR</p> <p>Effective response, %: NR (“practically 100%”)</p>	<p>Incontinence: Loss of urine at least once in prior year</p> <p>UII: NR</p> <p>SUI: NR</p> <p>MUI: NR</p>	<p>Prevalence of UII, %:[*] 1.4</p> <p>Prevalence of MUI, %:[*] 2.7</p>	<p>Overall quality: Poor</p> <p>Internal validity score: 2, -</p> <p>External Validity Score: 3, +</p> <p>Sampling Method Described: +</p> <p>N sampled provided: -</p> <p>N eligible provided: -</p> <p>N included respondents: +</p> <p>Response Rate:[*] NR</p> <p>Inclusion (Exclusion) Specified: +</p> <p>Age of population described: +</p> <p>Operational definition provided[*]: -</p> <p>Required frequency defined[*]: +</p>

[^]Data presented for women only.

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence[^]	Quality Rating
<p>Author: Brieger et al., 1997 Yip and Chung, 2003 Country: Territory of Hong Kong Study period: May 1996 to November 1996 Funding: NR Author industry relationship disclosures: NR</p>	<p>Design: Cross-sectional telephone questionnaire Base population: Chinese women in Hong Kong Sampling frame: Multistage random sample of more than 1.7 million residential telephone listings in Hong Kong N screened: 3,509 N eligible: NR N respondents: 1,500 N included: 1,500 Age, mean ± SD: 45 ± 15 Race/ethnicity, %: Chinese: 100 BMI, mean ± SD: 22.4 ± 2.8 Parity, %: 83</p>	<p>Inclusion criteria: • women • oldest of eligibles in household • ethnic Chinese Exclusion criteria: NR Effective response, %: 43</p>	<p>UUI: NR MUI: NR Detrusor dysfunction: One or more of the following: UUI, urgency, frequency, or nocturia, in the absence of SUI Urgency: NR Frequency: NR</p>	<p>Prevalence of UUI, %: 0.7 Prevalence of MUI, %: 4.7 Prevalence of detrusor dysfunction, %: 2.4 Prevalence of urgency, %: 4.3 Prevalence of frequency, %: 4.2</p>	<p>Overall quality: Poor Internal validity score: 4, + External Validity Score: 1, - Sampling Method Described: + N sampled provided: + N eligible provided: - N included respondents: + Response Rate:* 43 Inclusion (Exclusion) Specified: + Age of population described: - Operational definition provided*:- Required frequency defined*:-</p>

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence [^]	Quality Rating
<p>Author: Brieger et al., 1996</p> <p>Country: Territory of Hong Kong</p> <p>Study period: NR</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Cross sectional telephone questionnaire (Kings College Urodynamics Questionnaire)</p> <p>Base population: Chinese women in Hong Kong</p> <p>Sampling frame: Multistage random sample of more than 1.7 million residential telephone listings in Hong Kong</p> <p>N screened: 3,248</p> <p>N eligible: NR</p> <p>N respondents: NR</p> <p>N included: 819</p> <p>Age, mean ± SD: 41.5 ± 16</p> <p>Race/ethnicity: NR</p> <p>Parity, %: 66.9</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> women age between 10 and 90 oldest of eligibles in household <p>Exclusion criteria: NR</p> <p>Effective response, %: 25.2</p>	<p>UI: Involuntary loss of urine that is socially or hygienically unacceptable</p> <p>UUI: NR</p> <p>Frequency: NR</p>	<p>Prevalence of urgency and/or UUI, %: 14.7</p> <p>Prevalence of frequency, %: 18.8</p>	<p>Overall quality: Poor</p> <p>Internal validity score: 3, -</p> <p>External Validity Score: 2, -</p> <p>Sampling Method Described: +</p> <p>N sampled provided: +</p> <p>N eligible provided: -</p> <p>N included respondents: +</p> <p>Response Rate:* 25.2</p> <p>Inclusion (Exclusion) Specified: +</p> <p>Age of population described: +</p> <p>Operational definition provided*: -</p> <p>Required frequency defined*: -</p>

[^]Data presented for women only. Prevalence also reported by age.

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence	Quality Rating
<p>Author: Chen, Lin, Hu et al., 2003</p> <p>Chen, Chen, Hu et al., 2003</p> <p>Country: Taiwan</p> <p>Study period: 1999</p> <p>Funding: National Science Council in Taiwan</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Cross-sectional administered questionnaire (Bristol)</p> <p>Base population: Female residents of Dali</p> <p>Sampling frame: 3% random sample based on national census records to achieve nationally representative age strata</p> <p>N sampled: 1,584</p> <p>N screened: NR</p> <p>N eligible: NR</p> <p>N respondents: 1,253</p> <p>N included: 1,247</p> <p>Age, mean ± SD: 43.2 ± 15.1</p> <p>Race/ethnicity: NR</p> <p>BMI, mean ± SD: 23.1 ± 3.3</p> <p>Parity, % 86.5</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> women age ≥ 20 <p>Exclusion criteria: NR</p> <p>Effective response, %: 78.7</p>	<p>UII: Involuntary loss of urine preceded by the urge to void or relatively uncontrollable voiding with little or no warning, without requirement for “bother” or perceived social or hygienic problem.</p> <p>OAB: Frequency and urgency, or nocturia, with or without UII</p> <p>Urgency: Per ICS 2002</p> <p>Frequency: Per ICS 2002</p> <p>Strict definitions:* Above with requirement for “bother” and/or perceived social or hygienic problem</p>	<p>Prevalence of UII, %: 9.1</p> <p>Prevalence of UII-strict, %: 1.5</p> <p>Prevalence of MUI, %: 17.1</p> <p>Prevalence of MUI by age, %: 20-30: 10.0 >30-40: 16.2 >40-50: 23.1 >50-65: 20.2 > 65: 16.6 <i>P</i> < 0.05</p> <p>Prevalence of MUI-strict, %: 1.8</p> <p>Prevalence of MUI-strict by age, %: 20-30: 0.8 >30-40: 0.5 >40-50: 3.5 >50-65: 3.7 > 65: 2.1 <i>P</i> < 0.05</p> <p>Prevalence of OAB, % 18.6</p> <p>Prevalence of OAB by age, % 20-30: 11.7 >30-40: 13.3 >40-50: 20.7 >50-65: 20.2 > 65: 39.3 <i>P</i> < 0.05</p> <p>Prevalence of OAB-strict, % 2.5</p> <p>Prevalence of OAB-strict by age, % 20-30: 0.9 >30-40: 0.5 >40-50: 1.5 >50-65: 4.8 > 65: 9.7 <i>P</i> < 0.05</p>	<p>Overall quality: Fair</p> <p>Internal validity score: 4+</p> <p>External Validity score: 3, +</p> <p>Sampling Method Described: +</p> <p>N sampled provided: +</p> <p>N eligible provided: -</p> <p>N included respondents: +</p> <p>Response Rate:* 78.7</p> <p>Inclusion (Exclusion) Specified: +</p> <p>Age of population described: +</p> <p>Operational definition provided*: +</p> <p>Required frequency defined*: -</p>

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence	Quality Rating
Chen, Lin, Hu et al., 2003				Prevalence of urgency, %: 12.6	
Chen, Chen, Hu et al., 2003 (continued)				Prevalence of urgency-strict, %: 1.7	
				Prevalence of frequency, %: 21.1	
				Prevalence of frequency-strict, %: 2.3	

* Authors provide the strict definition and term it "meeting the criteria of the ICS" in the publication.

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence	Quality Rating
<p>Author: Choo et al., 2007</p> <p>Country: Korea</p> <p>Study period: 2000</p> <p>Funding: Korean Continenence Society-Johnson & Johnson Medical</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Cross-sectional telephone questionnaire</p> <p>Base population: Representative cross section of Korean population stratified by age, sex, and region</p> <p>Sampling frame: Random selection of telephone numbers</p> <p>N sampled/ screened: 14,559</p> <p>N eligible: Total: 2,005 Women: 1,005</p> <p>Age, mean ± SD:[^] 59.4 ± 11.6</p> <p>Age, median:[^] 59.0</p> <p>Age, n (%):[^] 40-49: 254 (25.3) 50-59: 252 (25.1) 60-69: 246 (24.5) ≥ 70: 253 (25.2)</p> <p>Race/ethnicity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> adults age 40 to 89 community-dwelling <p>Exclusion criteria: None</p> <p>Effective response: NA</p>	<p>Urgency: “Do you have to rush to the toilet?”</p> <p>Frequency: “How often do you pass urine on average?”</p> <p>UUI: “When you have an urge to urinate, do you loose urine before you could reach the bathroom?”</p> <p>OAB: Per ICS</p> <p>OAB-dry: Per ICS urge + freq; or urge + nocturia; or urge + freq + nocturia, without UUI</p> <p>OAB-wet: Any combination with UUI</p>	<p>Prevalence of UUI by age, %: All: 19.2 40-49: 13.0 50-59: 15.1 60-69: 24.4 ≥ 70: 24.5</p> <p>Prevalence of OAB-wet by age, %: All: 15.0 40-49: 10.2 50-59: 11.9 60-69: 19.5 ≥ 70: 18.6</p> <p>Prevalence of OAB-dry by age, %: All: 16.3 40-49: 16.9 50-59: 15.5 60-69: 14.2 ≥ 70: 18.6</p> <p>Prevalence of urgency by age, %: All: 32.5 40-49: 29.1 50-59: 28.6 60-69: 35.0 ≥ 70: 37.5</p> <p>Prevalence of frequency by age, %: All: 17.7 40-49: 16.5 50-59: 14.7 60-69: 17.1 ≥ 70: 22.5</p>	<p>Overall quality: Fair</p> <p>Internal validity score: 3,-</p> <p>External Validity Score: 3,+</p> <p>Sampling Method Described: +</p> <p>N sampled provided: +</p> <p>N eligible provided: +</p> <p>N included respondents: -</p> <p>Response Rate:* NR</p> <p>Inclusion (Exclusion) Specified: +</p> <p>Age of population described: +</p> <p>Operational definition provided*: +</p> <p>Required frequency defined*: -</p>

[^]Data presented for women only.

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence	Quality Rating
<p>Author: Corcos and Schick, 2004</p> <p>Country: Canada</p> <p>Study period: Spring 2002</p> <p>Funding: Pfizer Novartis</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Cross-sectional telephone questionnaire</p> <p>Base population: Canada</p> <p>Sampling frame: Stratified random sample of census areas in four census metropolitan areas by gender from the telephone registry</p> <p>N sampled: Total: 7,487</p> <p>N screened: NR</p> <p>N eligible: NR</p> <p>N respondents: Total: 3,249 Women: 1,683</p> <p>N included: Total: 3,249 Women: 1,683</p> <p>Age, %:[^] 35-44: 35.6 45-54: 28.1 55-64: 17.0 65-74: 11.9 ≥ 75: 6.8</p> <p>Race/ethnicity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • adults • age ≥ 35 <p>Exclusion criteria: NR</p> <p>Effective response, %: 53.7</p>	<p>OAB-wet (UUI): Urgency with urinary leaks</p> <p>OAB-dry (urgency): “Urgency with need to urinate that runs the risk of urine loss” at least weekly; or half the time or more it is “difficult to postpone urination” and they engage in bathroom seeking behavior.</p> <p>OAB: NR</p> <p>Frequency: ≥ 9 voids per day</p>	<p>Prevalence of OAB-wet, %: 6.5</p> <p>Prevalence of OAB, %: 21.3</p> <p>Prevalence of OAB by age, %: 35-44: 17.9 45-54: 21.1 55-64: 26.8 65-74: 24.0 ≥ 75: 21.4</p> <p>Prevalence of urgency, %: 23.3</p> <p>Prevalence of frequency, %: 15.5</p>	<p>Overall quality: Fair</p> <p>Internal validity score: 4, +</p> <p>External Validity Score: 4, ++</p> <p>Sampling Method Described: +</p> <p>N sampled provided: +</p> <p>N eligible provided: -</p> <p>N included respondents: +</p> <p>Response Rate:* 53.7</p> <p>Inclusion (Exclusion) Specified: +</p> <p>Age of population described: +</p> <p>Operational definition provided*: +</p> <p>Required frequency defined*: -</p>

[^]Data presented for women only.

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence	Quality Rating
<p>Author: Dallosso et al., 2004</p> <p>Country: UK</p> <p>Study period: October 1998 to December 1998* Follow-up 1999</p> <p>Funding: Medical Research Council</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Cross-sectional mailed questionnaire with repeated measures mailed food frequency (FFQ) questionnaire</p> <p>Base population: Patients registered in any of 108 general practices in Leicestershire</p> <p>Sampling frame: Random sample Leicestershire Health Authority registry</p> <p>N sampled: 20,247</p> <p>N screened: NR</p> <p>N eligible: 19,241</p> <p>N respondents: Baseline: 12,568 FFQ: 7,046 Follow-up FFQ: 5,816</p> <p>N included: 5,816</p> <p>Age, median: 61</p> <p>Race/ethnicity: NR</p>	<p>Inclusion criteria: OAB:</p> <ul style="list-style-type: none"> women age ≥ 40 community dwelling <p>Exclusion criteria: NR</p> <p>Effective response, %: 30.2</p>	<p>Study Definitions Urge leakage at least several times a month and/or very strong or overwhelming urgency</p>	<p>Annual incidence of OAB, %: 8.8</p> <p>Incidence of OAB by age, %: 40-49: 7.8 50-59: 6.5 60-69: 10.0 70-79: 11.7 ≥ 80: 14.3</p> <p>Baseline prevalence of OAB, %: 15.9</p> <p>Baseline prevalence of OAB by age, %: 40-49: 12.1 50-59: 15.4 60-69: 16.8 70-79: 18.5 ≥ 80: 26.1</p>	<p>Overall quality: Fair</p> <p>Internal validity score: 4, +</p> <p>External Validity Score: 4, ++</p> <p>Sampling Method Described: +</p> <p>N sampled provided: +</p> <p>N eligible provided: +</p> <p>N included respondents: +</p> <p>Response Rate:* 30.2</p> <p>Inclusion (Exclusion) Specified: +</p> <p>Age of population described: +</p> <p>Operational definition provided*: +</p> <p>Required frequency defined*: +</p>

*Initial mailing in October with two reminders sent in four week intervals

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence	Quality Rating
<p>Author: Dooley et al., 2008*</p> <p>Minassian et al., 2008</p> <p>Country: US</p> <p>Study period: 2001 to 2004* 2001 to 2002</p> <p>Funding: NR*, NR</p> <p>Author industry relationship disclosures: NR*, NR</p>	<p>Design: Cross-sectional administered questionnaire</p> <p>Base Population: NHANES a probability sample of the US non-institutionalized civilian population</p> <p>Sampling frame: NHANES participants</p> <p>N sampled: 21,161*</p> <p>N screened: NA</p> <p>N eligible: 4,541* 2,875</p> <p>N participants: 4,229* 2,577</p> <p>Age, %:* 20-39: 36.3 40-59: 28.0 ≥ 60: 35.7</p> <p>Age, mean ± SD: 50.8 ± 20.1</p> <p>Race/ethnicity, %:* White: 57.9 Black: 20.5 Hispanic: 21.5</p> <p>Race/ethnicity, %: White: 55 Black: 18 Hispanic: 24 Other: 3</p> <p>BMI, kg/m² n:* ≤ 25.0: 1,361 25.0-29.9: 1,320 30.0-39.9: 1,238 ≥ 40.0: 310</p> <p>BMI, mean ± SD: 28.6 ± 6.8</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> women age 20 or older completed the standardized interview and examination answered the questions about incontinence <p>Exclusion criteria:</p> <ul style="list-style-type: none"> ethnicity classified as "other"* <p>Effective response: NA</p>	<p>UII: in past 12 months, "leaked or lost control of even a small amount of urine with an urge or pressure to urinate and you could not get to a toilet fast enough"</p> <p>SUI: In past 12 months, "leaked or lost control of even a small amount of urine with an activity like coughing, lifting, or exercise."</p> <p>MUI: affirmative response to both SUI and UII.</p> <p>Severity of incontinence: Mild: few times a year Moderate: few times a month Severe: daily or few times a week</p>	<p>Prevalence of UII, %:* 7.9</p> <p>Prevalence of UII, %: 8.8</p> <p>Prevalence of UII by Age, %:* 20-39: 4.6 40-59: 8.7 ≥ 60: 11.7</p> <p>Severity of UII, % Mild: 41 Moderate: 31 Severe: 57</p> <p>Prevalence of MUI, %:* 17.0</p> <p>Prevalence of MUI, %: 13.0</p> <p>Prevalence of MUI by Age, %:* 20-39: 7.7 40-59: 18.6 ≥ 60: 28.7</p> <p>Severity of MUI, % Mild: 13 Moderate: 31 Severe: 57</p>	<p>Overall quality: Fair</p> <p>Internal validity score: 4, +</p> <p>External Validity Score: 4, ++</p> <p>Sampling Method Described: +</p> <p>N sampled provided: +</p> <p>N eligible provided: +</p> <p>N included respondents: +</p> <p>Response Rate:* NR</p> <p>Inclusion (Exclusion) Specified: +</p> <p>Age of population described: +</p> <p>Operational definition provided*: +</p> <p>Required frequency defined*: +</p>

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence	Quality Rating
<p>Author: Ege et al., 2008</p> <p>Country: Turkey</p> <p>Study period: May 2006</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Cross-sectional administered questionnaire</p> <p>Base population: Female residents of Konya</p> <p>Sampling frame: All postpartum women registered in seven health centers in the metropolitan area</p> <p>N sampled: 2,200</p> <p>N screened: NR</p> <p>N eligible: NR</p> <p>N respondents: 1,749</p> <p>N included: 1,749</p> <p>Age, mean ± SD: 26.8 ± 5.1</p> <p>Race/ethnicity: NR</p> <p>BMI, mean ± SD: 25.9 ± 3.8</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> women within 12 months postpartum <p>Exclusion criteria:</p> <ul style="list-style-type: none"> pregnant again at time of interview <p>Effective response, %: 79.5</p>	<p>UII: NR</p> <p>SUI: NR</p> <p>MUI: NR</p>	<p>Prevalence of UII postpartum, %: 2.0</p> <p>Prevalence of MUI postpartum, %: 9.3</p>	<p>Overall quality: Poor</p> <p>Internal validity score: 4, +</p> <p>External Validity Score: 1, -</p> <p>Sampling Method Described: +</p> <p>N sampled provided: +</p> <p>N eligible provided: -</p> <p>N included respondents: +</p> <p>Response Rate:* 79.5</p> <p>Inclusion (Exclusion) Specified: +</p> <p>Age of population described: -</p> <p>Operational definition provided*: -</p> <p>Required frequency defined*: -</p>

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence	Quality Rating
<p>Author: Espino et al., 2003</p> <p>Country: US</p> <p>Study period: Fall 1993 to Spring 1994</p> <p>Funding: NIH</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Cross-sectional administered in-home interview</p> <p>Base population: Mexican-Americans, age ≥ 65 residing in southwestern states (AZ, CA, CO, NM, TX)</p> <p>Sampling frame: Representative probability sample from the Hispanic EPESE*</p> <p>N screened: NR</p> <p>N eligible: NR</p> <p>N respondents: 1,755</p> <p>N included: 1,589</p> <p>Age, mean ± SD: 73.3 ± 6.2</p> <p>Race/ethnicity, %: Hispanic: 100</p> <p>BMI: NR for total N</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> women age ≥ 65 community dwelling Mexican-American <p>Exclusion criteria:</p> <ul style="list-style-type: none"> individuals with indwelling catheters <p>Effective response, %: 90.5</p>	<p>UII: Feeling need to urinate before losing urine</p> <p>MUI: UUI and leaking urine during sneezing, coughing, vomiting, lifting, laughing or straining</p>	<p>Prevalence of UII, %: 5.0</p> <p>Frequency of UII, %: Hardly ever: 17.7 Some of the time: 59.5 Most of the time: 13.9 All of the time: 8.9</p> <p>Volume of urine loss UII, %: Small: 50.0 Moderate: 39.5 Large: 10.5</p> <p>Wear protection all the time UII, %: 10.1</p> <p>UII Inhibits social activity, %: 16.7</p> <p>Prevalence of MUI, %: 6.3</p> <p>Frequency of MUI, %: Hardly ever: 12.0 Some of the time: 60.0 Most of the time: 18.0 All of the time: 10.0</p> <p>Volume of urine loss MUI, %: Small: 59.2 Moderate: 31.6 Large: 9.2</p> <p>Wear protection all the time MUI, %: 16.0</p> <p>MUI Inhibits social activity, %: 21.0</p>	<p>Overall quality: Fair</p> <p>Internal validity score: 3, +</p> <p>External Validity Score: 3, +</p> <p>Sampling Method Described: +</p> <p>N sampled provided: -</p> <p>N eligible provided: -</p> <p>N included respondents: +</p> <p>Response Rate:* 90.5</p> <p>Inclusion (Exclusion) Specified: +</p> <p>Age of population described: +</p> <p>Operational definition provided*: +</p> <p>Required frequency defined*: -</p>

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence	Quality Rating
<p>Author: Eva et al., 2003</p> <p>Country: Sweden</p> <p>Study period: 1997</p> <p>Funding: County of Östergötland (Folkshälsoanslag et); Linköping University Hospital</p> <p>Author -industry relationship disclosures: NR</p>	<p>Design: Cross-sectional mailed questionnaire</p> <p>Base population: Women in Östergötland born in 1937 or 1957</p> <p>Sampling frame: 39% random sample of the base</p> <p>N screened: 2,000</p> <p>N eligible: NR</p> <p>N respondents: 1,336</p> <p>N included: 1,317</p> <p>Age, n (%): 40-yr olds: 643 (48.8) 60-yr olds: 674 (51.2)</p> <p>Race/ethnicity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> women born in designated years <p>Exclusion criteria: NR</p> <p>Effective response, %: 65.9</p>	<p>UII: Urge to void before leaking</p>	<p>Prevalence of UII, %: 40-yr olds: Sometimes: 11.3 Mostly: 2.1 60-yr olds: Sometimes: 15.1 Mostly: 5.2</p> <p>Prevalence of daytime voids, Number %: 40-yr olds: 8-10: 4.2 > 10: 2.1 60-yr olds: 8-10: 7.8 > 10: 2.1</p>	<p>Overall quality: Fair</p> <p>Internal validity score: 4, +</p> <p>External Validity Score: 3, +</p> <p>Sampling Method Described: +</p> <p>N sampled provided: +</p> <p>N eligible provided: -</p> <p>N included respondents: +</p> <p>Response Rate:* 65.9</p> <p>Inclusion (Exclusion) Specified: +</p> <p>Age of population described: +</p> <p>Operational definition provided*: +</p> <p>Required frequency defined*: -</p>

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence	Quality Rating
<p>Author: Fenner et al., 2008</p> <p>Country: US</p> <p>Study period: Summer 2002 to Fall 2004</p> <p>Funding: NIH</p> <p>Author industry relationship disclosures: 2 of 7: Johnson & Johnson (1) Novartis (1) Novasys (1)</p>	<p>Design: Cross-sectional telephone questionnaire</p> <p>Base population: Residents of 3 Michigan counties</p> <p>Sampling frame: Telephone listings from commercial survey sampling group</p> <p>N sampled: 12,541</p> <p>N screened: 9,199</p> <p>N eligible: 3,692</p> <p>N respondents: 2,814</p> <p>Age, %: 35-44: 40.6 45-54: 37.7 55-64: 21.7</p> <p>Race/ethnicity, n (%): Black: 1,922 (68.3) White: 892 (31.7)</p> <p>Vaginally Parous, %: 69.3</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • black or white women • age 35 to 64 • community dwelling <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • pregnancy in last 12 months <p>Effective response, %: 69</p>	<p>Incontinence: Losing urine ≥ 12 times in 12 months</p> <p>UUI: based on at least one factor from urge component of factor analysis and no stress factors</p> <p>SUI: based on at least one factor from stress component of factor analysis</p>	<p>Prevalence of UUI, %: 3.6</p> <p>Prevalence of UUI by race/ethnicity, %: Black: 3.5 White: 3.6</p> <p>Prevalence of MUI, %: 6.0</p> <p>Prevalence of MUI by race/ethnicity, %: Black: 4.1 White: 7.1</p>	<p>Overall quality: Good</p> <p>Internal validity score: 5, ++</p> <p>External Validity Score: 4, ++</p> <p>Sampling Method Described: +</p> <p>N sampled provided: +</p> <p>N eligible provided: +</p> <p>N included respondents: +</p> <p>Response Rate:* 69</p> <p>Inclusion (Exclusion) Specified: +</p> <p>Age of population described: +</p> <p>Operational definition provided*: +</p> <p>Required frequency defined*: +</p>

Estimates of proportions of population with different types of UI use weights “constructed based on age, race and geographic location to adjust for the oversampling and for survey nonresponse.” Estimates of the prevalence of UUI and MUI are computed by multiplying the overall prevalence of UI (reported at the top of page 1457) by the proportions of types of UI (reported in Table 4 on page 1459). For example, overall prevalence of UUI is $0.265 \times 13.6\% = 3.6\%$

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence	Quality Rating
<p>Author: Fitzgerald et al., 2006[^] Thom et al., 2006* Huang et al., 2006[†]</p> <p>Country: US</p> <p>Study period: NR</p> <p>Funding: NIH</p> <p>Author industry relationship disclosures: 4 of 7[^] Allergan (2) Eli Lilly (1) Novartis (1) Pfizer (3) Q-med(1) Watson (1) Yamanouchi (4)</p>	<p>Design: Cross-sectional Written questionnaire and in-person interview follow-up</p> <p>Base population: Members of Kaiser Permanente Northern Care California</p> <p>Sampling frame: Random sampling of women by age and race strata to achieve equal strata size for age and fixed ratios by race/ethnicity</p> <p>N screened: 10,230*</p> <p>N screened: 8,835*</p> <p>N eligible: 3,240 estimated</p> <p>N respondents: 2,109*</p> <p>N included: 2,109 1,348[†]</p> <p>Age, mean ± SD: 56 ± 9 Asian: 53.2 ± 7.4[†] White: 56.0 ± 9.1[†]</p> <p>Race/ethnicity, %: White: 48 Black: 18 Latina: 17 Asian: 16</p> <p>Race/ethnicity, %:[†] White: 74 Asian: 26</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> women age 40 to 69 members since age 18 ≥ 50% births with Kaiser <p>Exclusion criteria:</p> <ul style="list-style-type: none"> race/ethnicity other than White or Asian[†] <p>Effective response, %: 65.1</p>	<p>Urgency: Strong urge or pressure to urinate without actually leaking, at least monthly</p> <p>Frequency: ≥ 7 voids/day; NR[†]</p> <p>UUI: NR, at least weekly; At least weekly incontinence with only or predominantly urge episodes in the last 7 days[†]</p> <p>UUI only:* NR, at least weekly</p> <p>MUI: NR, at least weekly</p> <p>MUI, predominantly urge:* NR, at least weekly</p> <p>MUI, equal stress and urge:* NR, at least weekly</p>	<p>Prevalence of UUI, %: 10</p> <p>Prevalence of UUI, %:[†] Asian: 7.3 White: 9.5 P=.04</p> <p>Prevalence of UUI only, age-adjusted, by race/ethnicity, % (95%CI):* White: 4.8 (3.9, 5.7) Hispanic: 5.8 (4.8, 6.8) Black: 7.6 (6.5, 8.8) Asian-American: 3.0 (2.3, 3.8) P = 0.027</p> <p>Prevalence of MUI, %: 5</p> <p>Prevalence of MUI, predominantly urge, by race/ethnicity, % (95%CI):* White: 4.0 (3.1, 4.8) Hispanic: 4.2 (3.3, 5.0) Black: 6.0 (5.0, 7.1) Asian-American: 4.4 (3.5, 5.2) P = NS</p> <p>Prevalence of MUI, predominantly urge, by race/ethnicity, % (95%CI):* White: 3.3 (2.5, 4.1) Hispanic: 5.3 (4.3, 6.3) Black: 1.9 (1.3, 2.5) Asian-American: 3.2 (2.5, 4.0) P = NS</p>	<p>Overall quality: Good</p> <p>Internal validity score: 5, ++</p> <p>External Validity Score: 4, ++</p> <p>Sampling Method Described: +</p> <p>N sampled provided: +</p> <p>N eligible provided: +</p> <p>N included respondents: +</p> <p>Response Rate: * 65.1</p> <p>Inclusion (Exclusion) Specified: +</p> <p>Age of population described: +</p> <p>Operational definition provided*: +</p> <p>Required frequency defined*: +</p>

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/ Exclusion Criteria	Study Definitions	Incidence/ Prevalence	Quality Rating
Fitzgerald et al., 2006 [^]	BMI, mean ± SD: NR for total N Asian: 25.8 ± 4.8 [†] White: 28.0 ± 6.7 [†]			Prevalence of urgency, %: 34	
Thom et al., 2006 [*]				Prevalence of frequency, %: 24%	
Huang et al., 2006 [†] (continued)	Parity, %: Total: 80 White: 80.8 [†] Asian: 79.3 [†]			Prevalence of daily frequency, %: Asian: 7.8 White: 13.0	
				Prevalence of weekly frequency, %: Asian: 10.6 White: 17.8 <i>P</i> <.01	

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence	Quality Rating
<p>Author: Hannestad et al., 2000</p> <p>Rortveit et al., 2003*</p> <p>Country: Norway</p> <p>Study period: 1995 to 1997</p> <p>Funding: National Health Screening Service of Norway; National Institute of Public Health; Norwegian University of Science and Technology; Research Council of Norway</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: cross-sectional mailed questionnaire (EPINCONT)</p> <p>Base population: Nord-Trøndelag County</p> <p>Sampling frame: All female residents of county; All women with vital records linkage*</p> <p>N sampled: 47,313</p> <p>N screened: NR</p> <p>N eligible:[^] 34,755</p> <p>Birth substudy:* 15,307</p> <p>N respondents:[^] 27,936</p> <p>Birth substudy:* 15,307</p> <p>N included: 27,936</p> <p>Birth substudy:* 15,307</p> <p>Age, %: 20-24: 6.7 25-29: 7.6 30-34: 8.9 35-39: 9.7 40-44: 10.5 45-49: 10.7 50-54: 9.9 55-59: 7.3 60-64: 6.6 65-69: 6.6 70-74: 6.4 75-79: 5.1 80-84: 2.7 85-89: 1.0 ≥ 90: 0.2</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> women age ≥ 20 community dwelling <p>Birth substudy: • no births, only cesareans, or only vaginal births of singletons*</p> <p>Exclusion criteria: NR; Birth substudy:*</p> <ul style="list-style-type: none"> more than four children (no women had more than four cesareans) births prior to 1967 (start of compulsory birth registration) age ≥ 65 (birth records not consistently accurate) <p>Effective response, %: 80.4%</p>	<p>UII: Any loss of urine with sudden and strong urge to go to the toilet</p> <p>SUI: Any loss of urine with coughing, sneezing, laughing, lifting, etc.</p> <p>MUI: Both UII and SUI</p>	<p>Prevalence of UII, %:^{**} 2.7</p> <p>Prevalence of UII by age, %:^{**} 20-24: 1.3 25-29: 1.9 30-34: 1.8 35-39: 1.5 40-44: 1.9 45-49: 2.0 50-54: 2.1 55-59: 2.5 60-64: 2.6 65-69: 4.4 70-74: 4.8 75-79: 6.4 80-84: 7.4 85-89: 8.0 ≥ 90: 4.8</p> <p>Prevalence of UII, by birth type, %:[*] None: 1.6 Cesarean: 2.2 Vaginal: 1.8 P = NS</p> <p>Prevalence of MUI, %:^{**} 8.8</p> <p>Prevalence of MUI by age, %:^{**} 20-24: 3.4 25-29: 4.0 30-34: 4.9 35-39: 6.1 40-44: 6.9 45-49: 7.7 50-54: 10.9 55-59: 10.2 60-64: 12.1 65-69: 70-74: 75-79: 80-84: 85-89: ≥ 90:</p> <p>Prevalence of MUI, by birth type, %:[*] None: 3.1 Cesarean: 5.5 Vaginal: 6.8</p>	<p>Overall quality: Fair</p> <p>Internal validity score: 5, ++</p> <p>External Validity Score: 3, +</p> <p>Sampling Method Described: +</p> <p>N sampled provided: +</p> <p>N eligible provided: +</p> <p>N included respondents: +</p> <p>Response Rate:[*] 80.4</p> <p>Inclusion (Exclusion) Specified: +</p> <p>Age of population described: +</p> <p>Operational definition provided*: +</p> <p>Required frequency defined*: -</p>

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/ Exclusion Criteria	Study Definitions	Incidence/ Prevalence	Quality Rating
Hannestad et al., 2000	Age, birth sub-study, %:* 20-24: 12.1			<i>P</i> vaginal compared to none:*	
Rortveit et al., 2003* (continued)	25-29: 13.0 30-34: 14.0 35-39: 15.4 40-44: 16.9 45-49: 16.5 50-54: 8.7 55-59: 2.2 60-64: 1.2			<i>P</i> < 0.05	
	Race/ethnicity: NR			<i>P</i> vaginal compared to cesarean:*	
	BMI birth sub-study kg/m², %:* < 25.0: 54.2 25.5-29.0: 32.9 ≥ 30: 12.9			<i>P</i> = NS	

[^] Data presented for women only; for each type of incontinence severity increased with age.

^{**} Prevalence is calculated by multiplying the prevalence of incontinence by the reported proportion of each type of incontinence

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence	Quality Rating
<p>Author: Herschorn et al., 2008</p> <p>Country: Canada</p> <p>Study period: October 2002</p> <p>Funding: Janssen-Ortho Canada</p> <p>Author industry relationship disclosures: 3 of 4 Astellas (1) Gynecare (1) Janssen-Ortho (3) Pfizer (2) Purdue (1) Triton (2)</p>	<p>Design: Cross-sectional telephone questionnaire in English or French</p> <p>Base population: Representative cross section of Canadian population stratified by age, sex, province, and census division</p> <p>Sampling frame: Modified random-digit dialing</p> <p>N sampled: NA</p> <p>N screened: 2,500</p> <p>N eligible: Total: 1,000 Women: 518</p> <p>Age, mean ± SD:[^] 44.5 ± 17.2</p> <p>Age, %:[^] 18-40: 42.8 41-64: 45.8 ≥ 65: 11.4</p> <p>Race/ethnicity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> adults age 18 or older community dwelling <p>Exclusion criteria:</p> <ul style="list-style-type: none"> institutionalized men and women <p>Effective response: NR</p>	<p>OAB: Per ICS</p> <p>Urgency: Per ICS</p> <p>Frequency: Per ICS</p> <p>UUI: Per ICS</p>	<p>Prevalence of UUI, %: Total (any): 9.3 ≤ 1x/wk: 5.0 2-3x/wk: 1.4 ~1x/day: 1.0 Several x/day: 0.8 All the time: 0.6</p> <p>Prevalence of UUI by age, %: 18-40: 4.6 41-64: 10.7 ≥ 65: 22.4</p> <p>Prevalence of OAB, %: 14.7</p> <p>Prevalence of OAB by age, %: 18-40: 12.8 41-64: 13.7 ≥ 65: 27.5</p> <p>Prevalence of urgency, %: Total (any): 14.1 ≤ 1x/wk: 4.4 2-3x/wk: 2.9 ~1x/day: 3.7 Several x/day: 2.0 All the time: 1.2</p> <p>Prevalence of urgency by age, %: 18-40: 13.3 41-64: 14.2 ≥ 65: 19.0</p> <p>Prevalence of frequency, %: 14.9</p> <p>Prevalence of frequency by age, %: 18-40: 15.1 41-64: 13.3 ≥ 65: 22.4</p>	<p>Overall quality: Poor</p> <p>Internal validity score: 2, -</p> <p>External Validity Score: 3, +</p> <p>Sampling Method Described: +</p> <p>N sampled provided: -</p> <p>N eligible provided: +</p> <p>N included respondents: -</p> <p>Response Rate:* NR</p> <p>Inclusion (Exclusion) Specified: +</p> <p>Age of population described: +</p> <p>Operational definition provided*: +</p> <p>Required frequency defined*: -</p>

[^]Data presented for women only.

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence	Quality Rating
<p>Author: Herzog et al., 1990</p> <p>Country: US</p> <p>Study period: 1983 to 1986*</p> <p>Funding: NIH</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Baseline with follow-up telephone questionnaire</p> <p>Base population: Residents of Washtenaw County</p> <p>Sampling frame: Multistage stratified random sample of households</p> <p>N screened: Total: 13,912</p> <p>N eligible: Total: 2,968 Women: NR</p> <p>N respondents: 1,956</p> <p>N included:[^] Baseline: 1,154 One-Yr: 1,056 Two-Yrs: 776</p> <p>Age, %: NR</p> <p>Race/ethnicity: NR</p>	<p>Inclusion criteria: UUI: • adults • age ≥ 60</p> <p>Exclusion criteria: NR</p> <p>Effective response, %: Total: 65.9</p>	<p>UUI: Urine loss preceded by urge to void or uncontrollable voiding with little or no warning</p> <p>SUI: Urine loss at times of exertion such as sneezing, lifting, bending</p> <p>MUI: Both UUI and SUI</p>	<p>Annual incidence of UUI, %: Baseline to Yr1: 1.7 Yr1 to Yr2: 1.5</p> <p>Annual incidence of MUI, %: Baseline to Yr1: 9.8 Yr1 to Yr2: 5.4</p> <p>Annual remission of UUI, %: Baseline to Yr1: 22.7 Yr1 to Yr2: 24.0</p> <p>Annual remission of MUI, %: Baseline to Yr1: 4.8 Yr1 to Yr2: 8.6</p> <p>Baseline prevalence of UUI, %: 2.8</p> <p>Baseline prevalence of MUI, %: 21.2</p>	<p>Overall quality: Fair</p> <p>Internal validity score: 5, ++</p> <p>External Validity Score: 3, +</p> <p>Sampling Method Described: +</p> <p>N sampled provided: +</p> <p>N eligible provided: +</p> <p>N included respondents: +</p> <p>Response Rate:* 65.9</p> <p>Inclusion (Exclusion) Specified: +</p> <p>Age of population described: +</p> <p>Operational definition provided*: +</p> <p>Required frequency defined*: -</p>

*Baseline was in 1983 and 1984 with one year and two year follow-up from those dates.

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence	Quality Rating
<p>Author: Homma et al., 2005</p> <p>Homma et al., 2006</p> <p>Country: Japan</p> <p>Study period: November 2002 to March 2003</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: None</p>	<p>Design: Cross-sectional mailed questionnaire</p> <p>Base population: Population of Japan age ≥ 40</p> <p>Sampling frame: Two-stage random sample of households</p> <p>N sampled: 10,096</p> <p>N screened: NA</p> <p>N returned: 4,605</p> <p>N participants: Total: 4,570 Women: 2,380</p> <p>Age, mean (range):^ 61 (41,100)</p> <p>Race/ethnicity: NR</p>	<p>Inclusion criteria: OAB:</p> <ul style="list-style-type: none"> • adults • age ≥ 40 <p>Exclusion criteria: NR</p> <p>Effective response, %: Total: 45 Women: NR</p>	<p>OAB: ≥8 voids per day and ≥1 episode of urgency/week</p> <p>UII: ICS definition</p> <p>Frequency: ≥ 8 voids per day</p> <p>Frequency: ≥ 11 voids per day</p> <p>Urgency: ICS definition per day or week</p>	<p>Prevalence of UII , %: 7</p> <p>Prevalence of OAB, %: 11</p> <p>Prevalence of UII and OAB by age: Prevalence of both types increase with age. P=NS</p> <p>Prevalence of urgency ≥ 1/day, %: 7</p> <p>Prevalence of urgency ≥ 1/wk, %: 13</p> <p>Prevalence of frequency ≥ 8/day, %: 49</p> <p>Prevalence of frequency ≥ 11/day, %: 10</p>	<p>Overall quality: Fair</p> <p>Internal validity score: 4, +</p> <p>External Validity Score: 4, ++</p> <p>Sampling Method Described: +</p> <p>N sampled provided: +</p> <p>N eligible provided: -</p> <p>N included respondents: +</p> <p>Response Rate:* 45</p> <p>Inclusion (Exclusion) Specified: +</p> <p>Age of population described:</p> <p>Operational definition provided*: +</p> <p>Required frequency defined*: +</p>

^Data presented for women only.

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence	Quality Rating
<p>Author: Hording et al., 1986</p> <p>Country: Denmark</p> <p>Study period: April 1981 to July 1982</p> <p>Funding: The Legacy of Emmy Lange, née Kramp, Rahbek Foundation</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Cross-sectional administered questionnaire</p> <p>Base population: Female residents of the Glostrup area of Copenhagen county born in 1936</p> <p>Sampling frame: All women born in 1936 and living in the Glostrup area</p> <p>N sampled: 613</p> <p>N screened: NR</p> <p>N respondents: 528</p> <p>N included: 515</p> <p>Age: 45</p> <p>Race/ethnicity: NR</p> <p>Parity, %: 93</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Women • Age 45 <p>Exclusion criteria: NR</p> <p>Effective response, %: 85</p>	<p>UII: Involuntary loss of urine following a pathologically strong desire to void</p> <p>SUI: Urine loss accompanying coughing, laughing, sneezing, running, or jumping with a full bladder or during sexual intercourse</p> <p>MUI: Mixture of UII and SUI</p>	<p>Prevalence of UII, %: 5.4</p> <p>Prevalence of MUI, %: 3.1</p>	<p>Overall quality: Fair</p> <p>Internal validity score: 4, +</p> <p>External Validity Score: 4, ++</p> <p>Sampling Method Described: +</p> <p>N sampled provided: 1</p> <p>N eligible provided: +-N included respondents: +</p> <p>Response Rate:* 85</p> <p>Inclusion (Exclusion) Specified: +</p> <p>Age of population described: +</p> <p>Operational definition provided*:</p> <p>Required frequency defined*: -</p>

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence	Quality Rating
Author: Hsieh et al., 2006 Country: Taiwan Study period: NR Funding: NR Author industry relationship disclosures: NR	Design: Cross-sectional administered questionnaire Base population: Female residents of Taiwan Sampling frame: Three stage random sample: township, block, individual to obtain nationally representative sample N sampled: NR N screened: NR N eligible: 4,546 N respondents: 3,537 N included: 3,519 Age, %: 20-29: 27.1 30-39: 29.8 40-49: 26.5 50-59: 16.6 Race/ethnicity: NR	Inclusion criteria: <ul style="list-style-type: none"> women age 20 to 59 Exclusion criteria: NR Effective response, %: 77.4	Frequency: “void too often during the day”	Prevalence of frequency, %: 5.2 Prevalence of frequency, voids per day, %: 8-15: 2.4 16-23: 1.9 24-31: 0.05 ≥ 32: 0.7 Prevalence of frequency by age, %: 20-29: 4.7 30-39: 5.8 40-49: 5.2 50-59: 5.9 $P = 0.33$	Overall quality: Poor Internal validity score: 4, + External Validity Score: 2, - Sampling Method Described: + N sampled provided: - N eligible provided: + N included respondents: + Response Rate:* 77.4 Inclusion (Exclusion) Specified: + Age of population described: + Operational definition provided*: - Required frequency defined*: -

3537 was described by authors as 77.8% response rate. Therefore ‘N’ whom they considered eligible would be 4546, so effective response rate is $3519/4546 = 77.4\%$

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence	Quality Rating
<p>Author: Iosif and Bekassy, 1984</p> <p>Country: Sweden</p> <p>Study period: NR</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Cross-sectional mailed questionnaire</p> <p>Base population: Malmöhus County, Sweden</p> <p>Sampling frame: Approximately 3,000 female residents born in 1921</p> <p>N sampled: 1,200</p> <p>N screened: NA</p> <p>N eligible: NA</p> <p>N respondents: 902</p> <p>Age: NR</p> <p>Race/ethnicity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> women born in 1921 <p>Exclusion criteria: NR</p> <p>Effective response, %: 75</p>	<p>UUI: “do you usually get such a very strong urge that you cannot hold back until you reach a toilet?”</p> <p>MUI: Combination of urge and stress (“involuntary loss of urine when cough, laugh,” etc) incontinence</p>	<p>Prevalence UUI, %: 8.0</p> <p>Prevalence MUI, %: 9.5</p>	<p>Overall quality: Fair</p> <p>Internal validity score: 4, +</p> <p>External Validity Score: 2 -</p> <p>Sampling Method Described: +</p> <p>N sampled provided: +</p> <p>N eligible provided: -</p> <p>N included respondents: +</p> <p>Response Rate:* 75</p> <p>Inclusion (Exclusion) Specified: +</p> <p>Age of population described: -</p> <p>Operational definition provided*: +</p> <p>Required frequency defined*: -</p>

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence [^]	Quality Rating
<p>Author: Irwin et al., 2006</p> <p>Country: Canada, Germany, Italy, Sweden, UK</p> <p>Study period: April 2005 to December 2005</p> <p>Funding: Pfizer</p> <p>Author industry relationship disclosures: 10 of 11 Astellas (5) Bayer (1) Boehringer-Ingelheim (1) Diagnostic Ultrasound (1) Ferring (1) Janssen-Ortho (1) Lilly (2) Novartis (3) Paladin (1) Pfizer (10) Plethora (1) Schwarz-Pharma (1) Tena (1) UCB (1) Yamanouchi (1)</p>	<p>Design: Cross-sectional telephone questionnaire (EPIC)</p> <p>Base population: Residents of Canada, Germany, Italy, Sweden, UK</p> <p>Sampling frame: Two-step random sampling of households and individuals via residential telephone</p> <p>N screened: 58,139</p> <p>N eligible: NR</p> <p>N respondents: 19,165</p> <p>N included: Total: 19,165 Women: NR</p> <p>Age, %:[^] 18-29: 13.6 30-34: 7.5 35-39: 10.2 40-44: 11.7 45-49: 10.5 50-54: 9.7 55-59: 9.2 60-64: 7.0 ≥ 70: 12.5</p> <p>Race/ethnicity, %: White: 95.6</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> adults age ≥ 18 most recent birthday in household <p>Exclusion criteria: NR</p> <p>Effective response, %: 33</p>	<p>Urgency: Per ICS 2002</p> <p>Frequency: Per ICS 2002</p> <p>UII: Per ICS 2002</p> <p>MUI: Per ICS 2002</p>	<p>Prevalence of UII, % (95% CI): 1.5 (1.2, 1.7)</p> <p>Prevalence of UII by age, % (95% CI): ≤ 39: 1.0 (0.6, 1.3) 40-59: 1.1 (0.7, 1.5) ≥ 60: 2.5 (1.9, 3.0)</p> <p>Prevalence of MUI, % (95% CI): 2.4 (2.1, 2.7)</p> <p>Prevalence of MUI by age, % (95% CI): ≤ 39: 1.0 (0.6, 1.3) 40-59: 2.4 (1.9, 3.0) ≥ 60: 4.1 (3.4, 4.8)</p> <p>Prevalence of urgency, % (95% CI): 12.8 (12.2, 13.5)</p> <p>Prevalence of urgency by age, % (95% CI): ≤ 39: 9.7 (8.8, 10.7) 40-59: 11.2 (10.1, 12.3) ≥ 60: 18.3 (16.9, 19.6)</p> <p>Prevalence of frequency, % (95% CI): 7.4 (6.9, 7.9)</p> <p>Prevalence of frequency by age, % (95% CI): ≤ 39: 7.9 (7.0, 8.8) 40-59: 5.8 (5.0, 6.6) ≥ 60: 8.4 (7.5, 9.4)</p>	<p>Overall quality: Fair</p> <p>Internal validity score: 4, +</p> <p>External Validity Score: 4, ++</p> <p>Sampling Method Described: +</p> <p>N sampled provided: +</p> <p>N eligible provided: -</p> <p>N included respondents: +</p> <p>Response Rate:* 33</p> <p>Inclusion (Exclusion) Specified: +</p> <p>Age of population described: +</p> <p>Operational definition provided*: -</p> <p>Required frequency defined*: -</p>

[^]Data presented for women only.

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence [^]	Quality Rating
<p>Author: Ju, et al. 1991</p> <p>Country: Singapore</p> <p>Study period: July 1989 to November 1989</p> <p>Funding: National University of Singapore</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Cross-sectional administered questionnaire, with follow-up interview if incontinent</p> <p>Base population: Elderly residents of a postal district in Telok Blangah Housing Estate</p> <p>Sampling frame: All elderly people in Ministry of Home Affairs registry for study district</p> <p>N sampled: 1,511</p> <p>N eligible: 1,143</p> <p>N respondents: 919</p> <p>N included: Total: 919 Women: 484</p> <p>Age: NR by gender</p> <p>Race/ethnicity: NR by gender</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Adults • Age ≥ 65 <p>Exclusion criteria: NR</p> <p>Effective response, %: 80.4</p>	<p>Incontinence: Leakage of urine two or more times in the past month</p> <p>UUI: Urine loss because of inability to delay voiding following an urge to micturition</p> <p>SUI: Urine loss associated with increases in intra-abdominal pressure (physical exertion, straining, coughing, or laughing)</p> <p>MUI: NR</p>	<p>Prevalence of UUI, %: 3.3</p> <p>Prevalence of UUI, %: 0.8</p>	<p>Overall quality: Fair</p> <p>Internal validity score: 4, +</p> <p>External Validity Score: 4,++</p> <p>Sampling Method Described: +</p> <p>N sampled provided: +</p> <p>N eligible provided: +</p> <p>N included respondents: +</p> <p>Response Rate:* 82</p> <p>Inclusion (Exclusion) Specified: +</p> <p>Age of population described: +</p> <p>Operational definition provided*: +</p> <p>Required frequency defined*: -</p>

[^]Data presented is for women only.

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence [^]	Quality Rating
<p>Author: Kay et al., 1999</p> <p>Country: Denmark</p> <p>Study period: NR</p> <p>Funding: Gammelgårds Grant, Synthelabo Scandinavia</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Cross-sectional mailed questionnaire (DAN-PPS)</p> <p>Base population: Population of Herlev municipality</p> <p>Sampling frame: Gender and age stratified random sample of adults in age range</p> <p>N sampled: Total: 500 Women: 250</p> <p>N screened: NA</p> <p>N eligible: NR</p> <p>N respondents: All: 368 Women: 158</p> <p>Age, n:[^] 40-49: 33 50-59: 40 60-69: 28 70-79: 36 80-89: 21</p> <p>Age, mean:[^] 62.5</p> <p>Age, median (range):[^] 61 (41, 89)</p> <p>Race/ethnicity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • adults • age 40 to 89 <p>Exclusion criteria: NR</p> <p>Effective response, %: All: 74 Women: 63</p>	<p>All measures are for two week window</p> <p>UUI: NR</p> <p>Urgency: NR</p> <p>Frequency: NR</p>	<p>Prevalence of UUI, % (95% CI): 25.3 (18.9, 33.0)</p> <p>Prevalence of urgency, % (95% CI): 67.1 (59.1, 74.2)</p> <p>Prevalence of frequency, % (95% CI): 42.2 (34.7, 50.5)</p>	<p>Overall quality: Fair</p> <p>Internal validity score: 4, +</p> <p>External Validity Score: 3,+</p> <p>Sampling Method Described: +</p> <p>N sampled provided: +</p> <p>N eligible provided: -</p> <p>N included respondents: +</p> <p>Response Rate:* 63</p> <p>Inclusion (Exclusion) Specified: +</p> <p>Age of population described: +</p> <p>Operational definition provided*: -</p> <p>Required frequency defined*: +</p>

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence [^]	Quality Rating
<p>Author: Kim et al., 2006</p> <p>Country: US</p> <p>Study period: 2000</p> <p>Funding: Pfizer</p> <p>Author industry relationship disclosures: 5 of 5 Pfizer (5)</p>	<p>Design: Database linkage study</p> <p>Base population: US population</p> <p>Sampling frame: National Ambulatory Medical Care Survey, National Hospital Ambulatory Medical Care Survey, National Discharge Survey data for 2000</p> <p>N sampled: NR</p> <p>N screened: NR</p> <p>N eligible: NR</p> <p>N respondents: NR</p> <p>N included: NR</p> <p>Age: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> adults age ≥ 18 first three coded diagnoses per care episode <p>Exclusion criteria: NR</p> <p>Effective response: NA</p>	<p>OAB visits: ICD-9: 788.31, 788.41 and 788.33</p> <p>Frequency: coded indication for visit.</p> <p>Urge incontinence: Coded indication for visit.</p> <p>Prevalence: expressed as number of adults with visit type per 10,000 adult population weighted for gender based on 2000 census.</p>	<p>National annual OAB visits: 1.4 million (1.1, 1.8)</p> <p>Prevalence of visits for UUI: 16 per 10,000 adults</p> <p>Prevalence of visits for MUI: 5 per 10,000 adults</p> <p>Prevalence of visits for any OAB symptom: Women: 81/10,000 Men: 56/10,000</p> <p>Prevalence of visits for frequency: 48 per 10,000 adults</p>	<p>Overall quality: Fair</p> <p>Internal validity score: (+)</p> <p>External Validity Score: 2, -</p> <p>Sampling Method Described: +</p> <p>N sampled provided: NA</p> <p>N eligible provided: NA</p> <p>N included respondents: NA</p> <p>Response Rate:* NA</p> <p>Inclusion (Exclusion) Specified: +</p> <p>Age of population described: -</p> <p>Operational definition provided*: +</p> <p>Required frequency defined*: -</p>

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence [^]	Quality Rating
<p>Author: Koyama et al., 1998</p> <p>Country: Japan</p> <p>Study period: NR</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Cross-sectional mailed questionnaire</p> <p>Base population: Residents of the selected farming village or suburban town</p> <p>Sampling frame: Distribution to all residents, 970 in village; 1,508 in suburb</p> <p>N sampled: Village: 970 Suburb: 1,508</p> <p>N screened: NA</p> <p>N eligible: NA</p> <p>N respondents: Village: 937 Suburb: 934 Women: 1,120</p> <p>Age, %: 60-69*: 31.7 70-79: 47.9 ≥ 80: 20.4</p> <p>Race/ethnicity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • adults • age > 65 • community dwelling <p>Exclusion criteria: NR</p> <p>Effective response, %: Village: 98.4 Suburb: 65.0</p>	<p>Incontinence: “even a small amount of urinary leakage at a time when there was no intention of urinating.”</p> <p>UUI: Not specified, however “urgency” subtype presented</p>	<p>Prevalence of UUI, %: Women: 5.3**</p>	<p>Overall quality: Poor</p> <p>Internal validity score: 4, +</p> <p>External Validity Score: 2, -</p> <p>Sampling Method Described: +</p> <p>N sampled provided: +</p> <p>N eligible provided: -</p> <p>N included respondents: +</p> <p>Response Rate:* >65</p> <p>Inclusion (Exclusion) Specified: +</p> <p>Age of population described: +</p> <p>Operational definition provided*: -</p> <p>Required frequency defined*: -</p>

*Age range as presented “60-69” however, inclusion says all participants >65 years old?!

**Denominator for total community dwelling in Table 1 = 1120 women, Table 3 = 59 with urgency as “nature of urinary incontinence”

[^]Data presented for women only.

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence [^]	Quality Rating
<p>Author: Kuh et al., 1999</p> <p>Country: UK</p> <p>Study period: 1993 to 1994</p> <p>Funding: Medical Research Council</p> <p>Author industry relationship disclosures: None</p>	<p>Design: Cross-sectional mailed questionnaire</p> <p>Base population: Women in the UK</p> <p>Sampling frame: Members of the Medical Research Council National Survey of Health and Development a nationally representative birth cohort begun in March 1946</p> <p>N sampled: 1,486</p> <p>N screened: NR</p> <p>N eligible: 1,486</p> <p>N respondents: 1,378</p> <p>N included: 1,333</p> <p>Age, %: 48: 100%</p> <p>Race/ethnicity: NR</p> <p>BMI kg/m², %: < 20: 7.2 20.1-25.0: 55.9 25.1-30.0: 25.0 > 30.0: 12.0</p> <p>Vaginally parous, %: 81.7</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> women born in March 1946 members of cohort <p>Exclusion criteria: NR</p> <p>Effective response, %: 89.7</p>	<p>UII: Urgent and strong desire to pass urine which is difficult to control and ever any loss of urine before getting to toilet</p> <p>SUI: Loss of urine when you cough, sneeze, laugh, run, or exercise</p> <p>MUI: Both UII and SUI</p>	<p>Prevalence of UII, %: 22</p> <p>Prevalence of MUI, %: 20</p>	<p>Overall quality: Fair</p> <p>Internal validity score: 5, ++</p> <p>External Validity Score: 3, +</p> <p>Sampling Method Described: +</p> <p>N sampled provided: +</p> <p>N eligible provided: +</p> <p>N included respondents: +</p> <p>Response Rate:* 89.7</p> <p>Inclusion (Exclusion) Specified: +</p> <p>Age of population described: +</p> <p>Operational definition provided*: +</p> <p>Required frequency defined*: -</p>

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence [^]	Quality Rating
<p>Author: Lara and Nacey, 1994</p> <p>Country: New Zealand</p> <p>Study period: NR</p> <p>Funding: Wellington Medical Research Foundation</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Cross-sectional mailed questionnaire</p> <p>Base population: Female residents of three areas of central Wellington</p> <p>Sampling frame: Random sample of the electoral roll</p> <p>N sampled: 1,028</p> <p>N screened: NA</p> <p>N eligible: NA</p> <p>N respondents: 556</p> <p>Age, mean (range): Maoris: 39.1 (18, 95) Pacific Islanders: 34.9 (18, 79) Europeans: 47 (18, 90)</p> <p>Race/ethnicity, %: Maoris: 25.7 Pacific Islanders: 21.9 Europeans: 52.3</p>	<p>Inclusion criteria: • women on electoral roll</p> <p>Exclusion criteria: NR</p> <p>Effective response, %: 54</p>	<p>Incontinence definition: “any involuntary loss of urine”</p> <p>UUI: “sudden involuntary loss”</p> <p>MUI: Elements of UUI and stress (“loss with increased intraabdominal pressure”)</p>	<p>Prevalence of UUI, %: 9.2</p> <p>Prevalence of MUI, %: 7.4</p>	<p>Overall quality: Fair</p> <p>Internal validity score: 4, +</p> <p>External Validity Score: 2, -</p> <p>Sampling Method Described: +</p> <p>N sampled provided: +</p> <p>N eligible provided: -</p> <p>N included respondents: +</p> <p>Response Rate:* 54</p> <p>Inclusion (Exclusion) Specified: -</p> <p>Age of population described: +</p> <p>Operational definition provided*.: +</p> <p>Required frequency defined*.: -</p>

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence [^]	Quality Rating
<p>Author: Link et al., 2007 Hall et al., 2008</p> <p>Country: US</p> <p>Study period: 2002 to 2005</p> <p>Funding: NIH</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Cross-sectional administered questionnaire</p> <p>Base population: Residents of 16/17 Boston residential planning districts*</p> <p>Sampling frame: Two-stage random sample stratified to achieve gender and race/ethnicity representation</p> <p>N screened: 24,063 households* 9,066 individuals*</p> <p>N eligible: 8,702*</p> <p>N respondents: 5,506*</p> <p>N included: Total: 5,506 Women: 3,205*</p> <p>Age, %:[^] 30-39: 24.7* 40-49: 26.2* 50-59: 24.3* 60-79: 24.7*</p> <p>Race/ethnicity, n (%):[^] Black: 1,070* (33.4) White: 1,024* (32.0) Hispanic: 1,111* (34.7)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • adults • age 30 to 79 • community dwelling <p>Exclusion criteria: NR</p> <p>Effective response, %: 57.3*</p>	<p>Frequency: Needed to urinate again < 2hrs after urinating (fairly often, usually, almost always) and/or frequent urination during the day (fairly often, usually, almost always) and/or on average 8 or more urinations per day</p> <p>Urgency: Difficulty postponing urination (fairly often, usually, almost always) and/or strong urge or pressure to urinate immediately with little or no warning (fairly often, usually, almost always) and/or strong urge to urinate immediately whether or not they urinated or leaked urine (several times, many times per day)</p> <p>† Cluster analysis groupings: Four symptomatic clusters emerged which cannot be readily mapped to clinically useful tool/descriptors.</p>	<p>Prevalence of urgency, %: 14.2</p> <p>Prevalence of frequency, %: 36.9</p>	<p>Overall quality: Fair</p> <p>Internal validity score: 5, ++</p> <p>External Validity Score: 3, +</p> <p>Sampling Method Described: +</p> <p>N sampled provided: +</p> <p>N eligible provided: +</p> <p>N included respondents: +</p> <p>Response Rate:* 57.3</p> <p>Inclusion (Exclusion) Specified: +</p> <p>Age of population described:+</p> <p>Operational definition provided*: +</p> <p>Required frequency defined*: -</p>

[^]Data presented for women only.

† Groupings may have etiologic or other research utility see publication for details.

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence [^]	Quality Rating
<p>Author: Liu and Andrews, 2002</p> <p>Country: Australia</p> <p>Study period: 1992 to 1994</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Repeated measures telephone questionnaire</p> <p>Base population: Residents of Southern Australia</p> <p>Sampling frame: Random sample, stratified by age from the State Electoral Data Base for South Australia</p> <p>N sampled: 4,184*</p> <p>N screened: NR</p> <p>N eligible: 2,272</p> <p>N respondents: 2,087</p> <p>N included: 2,087</p> <p>Age: "sample size of more than 190 in each gender and age group [70-74, 75-79,80-84,≥85]"</p> <p>Race/ethnicity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> adults age ≥ 70 <p>Exclusion criteria: NR</p> <p>Effective response, %: 49.9</p>	<p>UII: NR, "at least occasionally"</p> <p>SUI: NR, "at least occasionally"</p> <p>Mixed: Both UII and SUI, "at least occasionally"</p>	<p>Annual Incidence of UII, %: 22.6</p> <p>Two-Year Incidence of UII, %: 37.5</p> <p>Prevalence of UII by year, %:** Yr 1: 18.2 Yr 2: 16.1 Yr 3: 21.2</p> <p>Prevalence of MUI by year, %:** Yr 1: 23.2 Yr 2: 26.5 Yr 3: 26.8</p>	<p>Overall quality: Fair</p> <p>Internal validity score: 4, +</p> <p>External Validity Score: 2, -</p> <p>Sampling Method Described: +</p> <p>N sampled provided: +</p> <p>N eligible provided: +</p> <p>N included respondents: +</p> <p>Response Rate:* 49.9</p> <p>Inclusion (Exclusion) Specified: +</p> <p>Age of population described:-</p> <p>Operational definition provided*: +</p> <p>Required frequency defined*: -</p>

[^]Data presented for women only.

* Study also included 151 individuals identified within households sampled

** Prevalence for UII and MUI is calculated by multiplying the prevalence of the combined UII and MUI (reported in Table 1) by the relevant proportions from Table 3, e.g. 41.4 * 0.384/(0.384+0.495)=18.2

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence	Quality Rating
<p>Author: Lukacz et al., 2006</p> <p>Lawrence et al., 2007[^]</p> <p>Lawrence et al., 2008[†]</p> <p>Country: US</p> <p>Study period: April 2004 to January 2005</p> <p>Funding: NIH NIH; Kaiser Permanente Direct Community Benefit Funds^{^†}</p> <p>Author industry relationship disclosures: † 2 of 5 Astellas (1) Novartis (1) Pfizer (2) Watson (1)</p>	<p>Design: Cross-sectional mailed questionnaire (EPIQ[*])</p> <p>Base population: Kaiser Permanente Southern California members</p> <p>Sampling frame: Age-stratified random sample of 950,000 female members in age range in April 2004</p> <p>N sampled: 12,200</p> <p>N eligible: 12,200</p> <p>N respondents: 4,458</p> <p>N included: 4,103 3,962[^]</p> <p>Age, mean ± SD: 56.5 ± 15.8 56.6 ± 15.8[^]</p> <p>Race/ethnicity, %: White: 61.0 Hispanic: 19.4 Black: 9.7 Asian/Pacific: 7.9 Other: 2.1</p> <p>Race/ethnicity, %:[^] White: 61.7 Hispanic: 19.2 Black: 9.6 Asian/Pacific: 8.2 Other/unknown: 1.3</p> <p>BMI, mean ± SD: 27.4 ± 6.3 27.8 ± 6.2[^]</p> <p>Parity, %: 80.8</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> women age 25 to 84 address on record with HMO <p>Exclusion criteria:</p> <ul style="list-style-type: none"> insufficient data to categorize <p>Effective response, %: 33.6 32.5[^]</p>	<p>OAB: Per EPIQ[*]</p> <p>OAB-any: Includes urgency and frequency, with or without incontinence</p> <p>OAB-wet: Includes urgency and frequency, with incontinence</p> <p>SUI: Per EPIQ[*]</p>	<p>Prevalence of OAB-any, % (95% CI): 13.3 (12.2, 14.4)</p> <p>Prevalence of OAB-any by age, %:[†] 25-39: 5.9 40-54: 10.9 55-69:14.8 70-84: 19.0 <i>P</i><0.01</p> <p>Prevalence of OAB, %:[^] All: 13.4 Nondiabetic: 12.5 Diabetic: 21.4 <i>P</i><0.0001 for diabetic vs. non</p> <p>Prevalence of OAB by parity, % (95% CI): Nulliparous: 9 (7, 11) Cesarean: 9 (7, 13) Vaginal: 15 (14, 16) <i>P</i><0.05 both comparisons to vaginal</p> <p>Prevalence of OAB-wet, % (95% CI): † 12.7 (11.7, 13.9)</p> <p>Prevalence of mixed OAB-wet and SUI, % (95% CI): † 8.3 (7.5, 9.2)</p>	<p>Overall quality: Fair</p> <p>Internal validity score: 5, ++</p> <p>External Validity Score: 3, +</p> <p>Sampling Method Described: +</p> <p>N sampled provided: +</p> <p>N eligible provided: +</p> <p>N included respondents: +</p> <p>Response Rate:[*] 34</p> <p>Inclusion (Exclusion) Specified: +</p> <p>Age of population described: +</p> <p>Operational definition provided[*]: +</p> <p>Required frequency defined[*]: -</p>

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/ Exclusion Criteria	Study Definitions	Incidence/ Prevalence[^]	Quality Rating
Author: Lukacz et al., 2006	Vaginally Parous, %: [^] 71.6				
Lawrence et al., 2007 [^]					
Lawrence et al., 2008 [†] (continued)					

^{**}12,200 mailed, 4458 returned, 3962 evaluable = effective response of 3962/12,200 = 32.5%

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence [^]	Quality Rating
<p>Author: MacDiarmid and Rosenberg, 2005</p> <p>Country: US</p> <p>Study period: September 2004</p> <p>Funding: Ortho-McNeil</p> <p>Author industry relationship disclosures: 2 of 2 GlaxoSmithKline (1) Lilly (1) Novartis (1) Odyssey (1) Ortho-McNeil (2) Pfizer (2) Reliant (1) Watson (1) Yamanouchi (1)</p>	<p>Design: E-mailed invitation to online questionnaire</p> <p>Base population: Adult members of a multi-million-member online panel</p> <p>Sampling frame: Random electronic mailing</p> <p>N eligible: 2,951</p> <p>N respondents: 2,951</p> <p>Age: NR for total N</p> <p>Race/ethnicity: NR for total N</p> <p>BMI: NR for total N</p> <p>Parity: NR for total N</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • women • age ≥ 30 <p>Exclusion criteria: NR</p> <p>Effective response: NR</p>	<p>UII: ICS definition</p> <p>MUI: ICS definition</p>	<p>Prevalence of UII, %: 9.2*</p> <p>Prevalence of MUI, %: 13.2*</p>	<p>Overall quality: Poor</p> <p>Internal validity score: 2, -</p> <p>External Validity Score: 2, -</p> <p>Sampling Method Described: +</p> <p>N sampled provided: -</p> <p>N eligible provided: +</p> <p>N included respondents: +</p> <p>Response Rate:* NR</p> <p>Inclusion (Exclusion) Specified: +</p> <p>Age of population described: -</p> <p>Operational definition provided*: +</p> <p>Required frequency defined*: -</p>

*Calculated from reported frequencies: 271/2,951 and 389/2,951

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence	Quality Rating
<p>Author: McGrother et al., 2006[^]</p> <p>Donaldson et al., 2006*</p> <p>Country: UK</p> <p>Study period: Baseline questionnaires mailed in 1998*</p> <p>Funding: Medical Research Council</p> <p>Author industry relationship disclosures: None</p>	<p>Design: Repeated measures mailed questionnaires at baseline and 1, 2*, and 3*-year follow-up</p> <p>Base population: Leicestershire Health Authority (108 general practices)</p> <p>Sampling frame: Random sample from 20,247 women on registers at baseline</p> <p>N eligible: 19,241</p> <p>N respondents: Baseline: 12,750 Follow-up: NR Year 1: NR Year 2: NR Year 3: NR</p> <p>Age, mean ± SD: 59.5 ± 13.0*</p> <p>Race/ethnicity, %: White: 85.5*</p> <p>BMI kg/m², %:[^] < 20: 3.2 > 20-25: 38.7 > 25-30: 38.6 > 30: 19.5</p> <p>Parity, %: 87.2*</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> women age ≥ 40 community dwelling <p>Exclusion criteria: NR[^];</p> <ul style="list-style-type: none"> very elderly people in residential care;* <p>Effective Response, %: Baseline: 65.3 Follow-up: 79.7[^] Year 1: 76.4* Year 2: 71.2* Year 3: 67.0*</p>	<p>OAB: “a strong desire to pass urine resulting in leakage or urgency occurring monthly or more.” (exclusive of SUI)[^] Very strong or overwhelming urgency and UUI several times a month or more*</p> <p>UUI: strong desire to pass urine resulting in leakage before reaching toilet (exclusive of SUI)</p> <p>MUI:[^] Not explicitly defined</p>	<p>Annual incidence of MUI, %:[^] 4.5</p> <p>Annual incidence of OAB, %:[^] 5.4</p> <p>Year 1 Incidence of OAB, %:[*] 6.9</p> <p>Year 1 Incidence of OAB by age, %:[*] 40-49: 6.7 50-59: 6.5 60-69: 7.0 70-79: 6.9 80+: 9.5</p> <p>Year 1 Remission of OAB, %:[*] 38.8</p> <p>Year 1 Remission of OAB by age, %:[*] 40-49: 36.3 50-59: 32.9 60-69: 45.4 70-79: 39.7 80+: 45.9</p> <p>Year 2 Incidence of OAB, %:[*] 6.0</p> <p>Year 2 Remission of OAB, %:[*] 38.9</p> <p>Year 3 Incidence of OAB, %:[*] 6.8</p> <p>Year 3 Remission of OAB, %:[*] 36.9</p> <p>Baseline prevalence of MUI, %:[^] 12.7</p> <p>Baseline prevalence of OAB, %:[^] 7.7</p>	<p>Overall quality: Fair</p> <p>Internal validity score: 4, +</p> <p>External Validity Score: 4, ++</p> <p>Sampling Method Described: +</p> <p>N sampled provided: +</p> <p>N eligible provided: +</p> <p>N included respondents: -</p> <p>Response Rate:[*] +</p> <p>Inclusion (Exclusion) Specified: +</p> <p>Age of population described: +</p> <p>Operational definition provided[*]: +</p> <p>Required frequency defined[*]: +</p>

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/ Exclusion Criteria	Study Definitions	Incidence/ Prevalence[^]	Quality Rating
McGrother et al., 2006 [^]				Baseline prevalence of OAB by age, %:* 40-49: 11.4 50-59: 14.060-69: 12.5 70-79: 12.1 80+: 15.5	
Donaldson et al., 2006* (continued)					

Prevalence, year 1 incidence and remission are reported by age in Table 2 on page 712; urgency is defined in Table 1, but the threshold is only given for "When you need to pass urine, how strong is the urge usually?"

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence [^]	Quality Rating
<p>Author: Milsom et al., 2001</p> <p>Country: France, Germany, Italy, Spain, Sweden, UK</p> <p>Study period: NR</p> <p>Funding: Pharmacia Corporation</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Cross-sectional telephone questionnaire, except Spain where in-person interviews were done</p> <p>Base population: Residents of France, Germany, Italy, Spain, Sweden, UK</p> <p>Sampling frame: Stratified random sample to obtain nationally representative respondents from telephone listings (or voter roles in Spain)</p> <p>N sampled: NR</p> <p>N screened: NR</p> <p>N eligible: NR</p> <p>N respondents: Total: 16,776 Women: 9,728</p> <p>Age, %: 40-44: 17.4 45-49: 14.8 50-54: 13.2 55-59: 12.2 60-64: 11.1 65-69: 10.9 70-74: 8.9 ≥ 75: 11.5</p> <p>Race/ethnicity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> adults age ≥ 40 <p>Exclusion criteria:</p> <ul style="list-style-type: none"> voiding symptoms suggestive of stress incontinence, prostatic obstruction, or urinary tract infection <p>Effective response: NR</p>	<p>Frequency: > 8 voids/day</p> <p>Urgency: “Strong urge to go to toilet with no advance warning; have to hurry to toilet in time to urinate, have to keep running to toilet to urinate”</p> <p>UUI: “unable to get to bathroom in time; have wetting accidents at night when asleep; sudden and uncontrolled loss of urine; leak urine during the day without being able to control it”</p> <p>OAB: Frequency, urgency, and urge urinary incontinence alone or in any combination</p>	<p>Prevalence of OAB, %: 17.4</p> <p>Prevalence of OAB by age, %: 40-44: 8.7 45-49: 10.6 50-54: 11.9 55-59: 16.9 60-64: 16.9 65-69: 17.5 70-74: 22.1 75+: 31.3</p>	<p>Overall quality: Poor</p> <p>Internal validity score: 2, -</p> <p>External Validity Score: 3, +</p> <p>Sampling Method Described: +</p> <p>N sampled provided: -</p> <p>N eligible provided: -</p> <p>N included respondents: +</p> <p>Response Rate:* NR</p> <p>Inclusion (Exclusion) Specified: +</p> <p>Age of population described: +</p> <p>Operational definition provided*: +</p> <p>Required frequency defined*: -</p>

[^]Data presented for women only.

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence [^]	Quality Rating
<p>Author: Møller et al., 2000</p> <p>Country: Denmark</p> <p>Study period: June 1996</p> <p>Funding: Coloplast A/S Pharmacia A/S Four Danish foundations</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Cross sectional mailed questionnaire</p> <p>Base population: Residents of Denmark</p> <p>Sampling frame: Random sampling of Danish Civil Registry System in age and urban vs. rural strata</p> <p>N screened: 4,000</p> <p>N eligible: NR</p> <p>N respondents: 3,204</p> <p>N included: 2,860</p> <p>Age, %: 40: 21.5 45: 20.3 50: 20.0 55: 19.2 60: 18.9</p> <p>Race/ethnicity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> women age 40, 45, 50, 55, or 60 <p>Exclusion criteria: NR</p> <p>Effective response, %: 71.7</p>	<p>UUI: Do you leak urine if suddenly you need to void?</p> <p>Urgency: Do you rush to the toilet because of a sudden desire to void?</p> <p>Daytime frequency: Number of episodes exceeds 10 voids daily.*</p>	<p>Prevalence UUI, weekly or more, %: 7.2 (95% CI : 6.2,8.2)</p> <p>Prevalence of urgency, weekly or more, %: 7.1 (95% CI: 6.1,8.1)</p> <p>Prevalence of daytime frequency, %: 3.6 (95% CI: 2.9,4.3)</p>	<p>Overall quality: Fair</p> <p>Internal validity score: 4, +</p> <p>External Validity Score: 3, +</p> <p>Sampling Method Described: +</p> <p>N sampled provided: +</p> <p>N eligible provided: -</p> <p>N included respondents: +</p> <p>Response Rate:* 71.7</p> <p>Inclusion (Exclusion) Specified: +</p> <p>Age of population described: +</p> <p>Operational definition provided*:</p> <p>Required frequency defined*: -</p>

[^]Data presented for women only.

*More than 10 voids per day is categorized as "often" (and reported in Table III).

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence[^]	Quality Rating
Author: Muller, 2005 Country: US Study period: October 2000 to December 2000 Funding: Pfizer Author industry relationship disclosures: NR	Design: Cross sectional administered questionnaire (NS) Base population: US population Sampling frame: Unspecified sampling of shoppers in retail malls of 20 major urban areas N sampled: NR N screened: NR N eligible: NR N respondents: NR N included: Total: 1,001 Women: NR Age: NR Race/ethnicity: NR	Inclusion criteria: <ul style="list-style-type: none"> • adults • age 30 to 70 • community dwelling Exclusion criteria: NR Effective response: NR	Definitions: NR	Prevalence of “sudden urge to urinate”, %: 13 Prevalence of “both urge and leakage”, %: 22	Overall quality: Poor Internal validity score: 1, - External Validity Score: 1,- Sampling Method Described:+ N sampled provided: - N eligible provided: - N included respondents: - Response Rate:* NR Inclusion (Exclusion) Specified: + Age of population described: - Operational definition provided*:- Required frequency defined*: -

Note: This is a report from the Executive Director of The National Association for Continence presenting for academic audiences consumer research done as part of pre- or post-marketing research by for profit entities. The Association assisted with formation and scope of questions included in the questionnaire.

[^]Data presented for women only.

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence [^]	Quality Rating
<p>Author: Nuotio et al., 2002</p> <p>Country: Finland</p> <p>Study period: 1979 to 1989</p> <p>Funding: Medical Research Fund of Tampere University Hospital, the Academy of Finland, and the Uulo Arhio Foundation</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Cross-sectional administered questionnaire with similar follow-up questionnaire 10 years later</p> <p>Base population: Elderly residents of Tampere, Finland</p> <p>Sampling frame: Age and gender stratified random sample; part of the European Longitudinal Study on Aging (ELSA)</p> <p>N sampled: 1,494</p> <p>N screened: NR</p> <p>N eligible: Baseline: 1,309 Follow-up: 518</p> <p>N included: Total: 1,052 Women: 528 Follow-up: 435 Women: 260</p> <p>Age, mean:[^] 73.2</p> <p>Race/ethnicity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Adults • Age ≥ 60 <p>Exclusion criteria: NR</p> <p>Effective response, %: Baseline: 82 Follow-up: 84</p>	<p>Urgency: “Do you ever have trouble getting to the lavatory in time?”</p> <p>UUI: Urgency and urinary leakage, regardless of the frequency of urine loss</p>	<p>Prevalence of urgency, %: 8</p> <p>Prevalence of UUI, %: 6</p>	<p>Overall quality: Fair</p> <p>Internal validity score: 4, +</p> <p>External Validity Score: 4, ++</p> <p>Sampling Method Described: +</p> <p>N sampled provided: +</p> <p>N eligible provided: -</p> <p>N included respondents: +</p> <p>Response Rate:* 82</p> <p>Inclusion (Exclusion) Specified: +</p> <p>Age of population described: +</p> <p>Operational definition provided*: +</p> <p>Required frequency defined*:</p>

[^]Data presented is for women only.

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence [^]	Quality Rating
<p>Author: Nygaard and Lemke, 1996</p> <p>Country: US</p> <p>Study period: 1981 to 1988</p> <p>Funding: NIH</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Repeated measures, in-person interviews</p> <p>Base population: Residents of two counties in rural Iowa</p> <p>Sampling frame: Age stratified random sample of women with address on record</p> <p>N sampled: NR</p> <p>N screened: NA</p> <p>N eligible: NR</p> <p>N respondents: 2025</p> <p>Age, %: 65-69: 25.6 70-74: 25.9 75-79: 23.0 80-84: 15.9 85-89: 7.5 ≥ 90: 2.1</p> <p>Race/ethnicity, %: Non-Hispanic white: 61.7 Hispanic: 19.2 Black: 9.6 Asian-Pacific Islander: 8.2 Other/unknown: 1.3</p> <p>BMI, mean: 25.4</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> women age ≥ 65 living in community <p>Exclusion criteria: NR</p> <p>Effective response, %: 80</p>	<p>UII: “How often do you have difficulty holding your urine until you can get to a toilet?”</p> <p>For prevalence, incidence, and remission never and hardly ever = negative response</p>	<p>3-year incidence of UII (first follow-up), %: 20.4</p> <p>3-year remission of UII (first follow-up), %: 31.7</p> <p>3-year incidence of UII (second follow-up), %: 24.0*</p> <p>3-year remission of UII (second follow-up), %: 34.9*</p> <p>Baseline prevalence of UII, %: Total ever: 55 Positive response: 36.3</p> <p>Baseline prevalence of UII by how often, %: Hardly ever: 18.6 Some of the time: 27.8 Most of the time: 6.7 All of the time: 1.6</p>	<p>Overall quality: Fair</p> <p>Internal validity score: 3, +</p> <p>External Validity Score: 4, ++</p> <p>Sampling Method Described: +</p> <p>N sampled provided: -</p> <p>N eligible provided: -</p> <p>N included respondents: +</p> <p>Response Rate:* 80</p> <p>Inclusion (Exclusion) Specified: +</p> <p>Age of population described: +</p> <p>Operational definition provided*: +</p> <p>Required frequency defined*: +</p>

*3-year incidence and remission in the second follow-up are calculated from those women without UII at baseline. Effective response rate is reported on page 1050, but the number sampled is not reported.

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence [^]	Quality Rating
<p>Author: Odeyemi et al., 2006</p> <p>Country: UK</p> <p>Study period: 1987 to 2004</p> <p>Funding: Allergan</p> <p>Author industry relationship disclosures: 1 of 6: Allergan (1)</p>	<p>Design: Healthcare database</p> <p>Base population: Residents of the UK receiving care in one of 350 general practices selected to be representative of the National Health Service</p> <p>Sampling frame: General Practice Research Database (4.6% representative sample with longitudinal data)</p> <p>N sampled: NR</p> <p>N screened: NA</p> <p>N eligible: 68,910</p> <p>Age: NR for total N</p> <p>Race/ethnicity: NR for total N</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • adults • in GPRD sample <p>Exclusion criteria: UTI within two weeks before or after visit meeting OAB criteria, renal dysfunction, diuretics use</p> <p>Effective response: NR</p>	<p>OAB-related symptoms: Oxford Medical Index System and Read Clinical Classification codes for urgency, frequency, urge, urgency incontinence, nocturia, unstable bladder or irritable bladder</p>	<p>Annual incidence of OAB-related symptoms (95% CI): 2.60/1,000 (2.57, 2.63)</p> <p>Prevalence of OAB-related symptoms: 3.64/1,000</p>	<p>Overall quality: Fair</p> <p>Internal validity score: (+)</p> <p>External Validity Score: 2, -</p> <p>Sampling Method Described: +</p> <p>N sampled provided:</p> <p>NA</p> <p>N eligible provided: NA</p> <p>N included respondents: NA</p> <p>Response Rate:* NA</p> <p>Inclusion (Exclusion) Specified: +</p> <p>Age of population described: -</p> <p>Operational definition provided*.: +</p> <p>Required frequency defined*.: -</p>

[^]Data presented for women only.

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence [^]	Quality Rating
<p>Author: Perry et al., 2000</p> <p>Country: UK</p> <p>Study period: October 1996 to June 1997</p> <p>Funding: Medical Research Council</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Cross-sectional mailed questionnaire</p> <p>Base population: Patients registered in any of 108 general practices in Leicestershire</p> <p>Sampling frame: Random sample Leicestershire Health Authority registry</p> <p>N sampled: 15,904</p> <p>N screened: NR</p> <p>N eligible: Total: 14,600 Women: 7,659</p> <p>N respondents: Total: 10,226 Women: NR</p> <p>N included: Total: 10,116 Women: 5,544</p> <p>Age, %:[^] 40-49: 25.1 50-59: 24.8 60-69: 21.8 70-79: 17.9 ≥ 80: 10.4</p> <p>Race/ethnicity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • adults • age ≥ 40 • community dwelling <p>Exclusion criteria: NR</p> <p>Effective response, %:[^] 72.4</p>	<p>Incontinence: Loss of urine several times a month or more</p> <p>Urgency: Experienced an overwhelming urge to pass urine or had difficulty holding urine most of the time or overwhelming</p> <p>Frequency: Voiding hourly or more often</p>	<p>Prevalence of urgency, %: 8.8</p> <p>Prevalence of frequency, %: 9.1</p>	<p>Overall quality: Good</p> <p>Internal validity score: 5, ++</p> <p>External Validity Score: 4, ++</p> <p>Sampling Method Described: +</p> <p>N sampled provided: +</p> <p>N eligible provided: +</p> <p>N included respondents: +</p> <p>Response Rate:[*] 72.4</p> <p>Inclusion (Exclusion) Specified: +</p> <p>Age of population described: +</p> <p>Operational definition provided[*]: +</p> <p>Required frequency defined[*]: +</p>

[^]Data presented is for women only.

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence [^]	Quality Rating
<p>Author: Rohr et al., 2004</p> <p>Country: Denmark</p> <p>Study period: NR</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Cross-sectional telephone questionnaire</p> <p>Base population: Female residents of central Odense</p> <p>Sampling frame: Central Person Register listing of residents in postal district 5000</p> <p>N sampled: 498</p> <p>N screened: 498</p> <p>N eligible: 421</p> <p>N respondents: 223</p> <p>N included: 223</p> <p>Age, median (range): 82 (75, 96)</p> <p>Race/ethnicity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> women age ≥ 70 <p>Exclusion criteria:</p> <ul style="list-style-type: none"> need for proxy respondent <p>Effective response, %: 53</p>	<p>UII: Such a strong urge that it was impossible to get to the toilet in time</p> <p>SUI: Involuntary loss of urine in connection with physical exertion in the past month</p> <p>MUI: Both UII and SUI</p>	<p>Prevalence of UII, % (95% CI): 16 (10.9, 20.5)</p> <p>Prevalence of MUI, % (95% CI): 13 (8.2, 16.9)</p>	<p>Overall quality: Good</p> <p>Internal validity score: 5, ++</p> <p>External Validity Score: 4, ++</p> <p>Sampling Method Described: +</p> <p>N sampled provided: +</p> <p>N eligible provided: +</p> <p>N included respondents: +</p> <p>Response Rate:* 53</p> <p>Inclusion (Exclusion) Specified: +</p> <p>Age of population described: +</p> <p>Operational definition provided*: +</p> <p>Required frequency defined*: +</p>

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence [^]	Quality Rating
<p>Author: Schulman et al., 1997</p> <p>Country: Belgium</p> <p>Study period: November 1994 to February 1995</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: 1 of 3 Marrion Merrell Dow (1)</p>	<p>Design: Cross sectional written questionnaire in home</p> <p>Base population: Residents of Belgium</p> <p>Sampling frame: Two stage, random sampling, stratified by region, gender, age, and profession for nationally representative sample.</p> <p>N screened: 8,000</p> <p>N eligible: NR</p> <p>N respondents: 5,920</p> <p>N included: Total: 5,269 Women: 2,770</p> <p>Age, %: 30-34: 13.3 35-49: 34.0 50-64: 28.0 ≥ 65: 24.7</p> <p>Race/ethnicity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • adults • age 30 and older • community dwelling <p>Exclusion criteria: NR</p> <p>Effective response, %: NR</p>	<p>UUI: Arriving too late at the toilet</p>	<p>Prevalence of UUI, %: 2.7</p>	<p>Overall quality: Poor</p> <p>Internal validity score: 3, -</p> <p>External Validity Score: 2, -</p> <p>Sampling Method Described: +</p> <p>N sampled provided: +</p> <p>N eligible provided: -</p> <p>N included respondents: +</p> <p>Response Rate:* NR</p> <p>Inclusion (Exclusion) Specified: +</p> <p>Age of population described: +</p> <p>Operational definition provided*: -</p> <p>Required frequency defined*: -</p>

[^]Data presented for women only.

The total number of UUI cases is $0.026 \times 5,269 = 137$, of which 55% (see page 317) are women. The prevalence of UUI can then be calculated as $137 \times 0.55 / 2,770 = 75 / 2,770 = 2.7\%$

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence [^]	Quality Rating
<p>Author: Shakhatreh, 2005</p> <p>Country: Jordan</p> <p>Study period: July 2003</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Cross-sectional administered questionnaire</p> <p>Base population: Female residents of Southern Jordan in this age group</p> <p>Sampling frame: 68 census blocks selected from among 228 to achieve representative sample; household and then individuals randomly selected within blocks</p> <p>N sampled: Households: 810</p> <p>N screened: NR</p> <p>N eligible: Women: 190*</p> <p>N respondents: 182</p> <p>N included: 182</p> <p>Age, %: 50-54: 38.9 55-59: 23.9 60-65: 37.2</p> <p>Race/ethnicity: NR</p> <p>BMI kg/m², %: 18.5-24: 20 25-29: 32 ≥ 30: 48</p> <p>Parity, %: 95.8</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> women age 50 to 65 community dwelling <p>Exclusion criteria: NR</p> <p>Effective response, %: 96*</p>	<p>UII: Inability to postpone urination with strong urge to void</p> <p>SUI: Involuntary leakage of urine on exertion as in coughing, sneezing, laughing, bending over, or with exercise as in jumping</p> <p>MUI: Combination of UII and SUI</p>	<p>Prevalence of UII, %: 26.4</p> <p>Prevalence of MUI, %: 18.1</p>	<p>Overall quality: Fair</p> <p>Internal validity score: 5, ++</p> <p>External Validity Score: 3, +</p> <p>Sampling Method Described: +</p> <p>N sampled provided: +</p> <p>N eligible provided: +</p> <p>N included respondents: +</p> <p>Response Rate:* 96</p> <p>Inclusion (Exclusion) Specified: +</p> <p>Age of population described: +</p> <p>Operational definition provided*: +</p> <p>Required frequency defined*: -</p>

* Authors report 96% of eligible women participated and 182 = respondents.

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence [^]	Quality Rating
<p>Author: Simeonova et al., 1999</p> <p>Country: Sweden</p> <p>Study period: NR</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Cross-sectional mailed questionnaire</p> <p>Base population: Female residents of the Central [primary care] District of Göteborg</p> <p>Sampling frame: Every fourth female resident (other detail NR)</p> <p>N sampled: 2,911</p> <p>N screened: NR</p> <p>N eligible: NR</p> <p>N respondents: 2,248</p> <p>N included: 2,176</p> <p>Age, %: 20-29: 22.0 30-39: 18.2 40-49: 14.8 50-59: 12.6 60-69: 10.2 70-79: 12.0 ≥ 80: 10.2</p> <p>Race/ethnicity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> women age ≥ 20 <p>Exclusion criteria: NR</p> <p>Effective response, %: 74.8</p>	<p>Incontinence: involuntary loss of urine at least 1x/wk, considered by women to be hygienic or social problem</p> <p>UUI: incontinence preceded by urge to void or uncontrollable voiding with little or no warning</p> <p>SUI: Incontinence precipitated by coughing, sneezing, or physical exertion</p> <p>Mixed: Both UUI and SUI</p>	<p>Prevalence of UUI, %: 5.1</p> <p>Prevalence of UUI by age, %: 20-29: 0.8 30-39: 1.5 40-49: 4.4 50-59: 4.9 60-69: 7.7 70-79: 10.0 ≥ 80: 13.7 <i>P</i> < 0.01 for increase with age</p> <p>Prevalence of MUI, %: 3.9</p> <p>Prevalence of MUI by age, %: 20-29: 1.1 30-39: 1.5 40-49: 2.5 50-59: 4.9 60-69: 5.5 70-79: 6.7 ≥ 80: 10.0 <i>P</i> < 0.01 for increase with age</p>	<p>Overall quality: Fair</p> <p>Internal validity score: 4, +</p> <p>External Validity Score: 4, ++</p> <p>Sampling Method Described: +</p> <p>N sampled provided: +</p> <p>N eligible provided: -</p> <p>N included respondents: +</p> <p>Response Rate:* 74.8</p> <p>Inclusion (Exclusion) Specified: +</p> <p>Age of population described: +</p> <p>Operational definition provided*: +</p> <p>Required frequency defined*: +</p>

[^]Urge and mixed incontinence prevalence (overall and by age) computed from proportions of types of incontinence estimated from Figure 3.

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence[^]	Quality Rating
Author: Siracusano et al., 2003 Country: Italy Study period: NR Funding: Associazione Progetto Continenza Author industry relationship disclosures: NR	Design: Cross-sectional mailed questionnaire Base population: Female residents of Trieste Sampling frame: NR N sampled: 10,000 N screened: NR N eligible: NR N respondents: 3,557 N included: 2,900 Age: NR for respondents Race/ethnicity: NR	Inclusion criteria: <ul style="list-style-type: none"> • women • age 18 to 49 Exclusion criteria: NR Effective response, %: 29	UII: NR	Prevalence of UII, %: 8.7*	Overall quality: Poor Internal validity score: 2, - External Validity Score: 1, - Sampling Method Described: - N sampled provided: + N eligible provided: - N included respondents: + Response Rate:* 29% Inclusion (Exclusion) Specified: + Age of population described:- Operational definition provided*:- Required frequency defined*:-

* Calculated: 43.5% of 581 women with incontinence had urge incontinence (Table 2) and denominator is 2900.

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence [^]	Quality Rating
<p>Author: Song et al., 2005 Zhang et al., 2005 Zhang et al., 2006*</p> <p>Country: China</p> <p>Study period: April 2002</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Cross-sectional mailed questionnaire (BFLUTS)</p> <p>Base population: Residents of Fuzhou, China</p> <p>Sampling frame: Random sample (3%) of national census records for the city</p> <p>N sampled: 6,066</p> <p>N screened: NA</p> <p>N eligible: 4,745</p> <p>N respondents: 4,684</p> <p>Age, mean ± SD: 40.0 ± 11.1</p> <p>Race/ethnicity: Han Chinese: 100</p> <p>BMI, mean ± SD: 21.9 ± 3.0</p> <p>Vaginally Parous, %: 79.8</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> women age ≥ 20 <p>Exclusion criteria: NR</p> <p>Effective response, %: 77.2</p>	<p>Incontinence: Any involuntary loss of urine in last month</p> <p>UII: incontinence preceded by a sudden urge to void or uncontrollable voiding with little or no warning</p> <p>MUI: Symptoms of stress and urge incontinence</p> <p>OAB: OAB-wet or OAB-dry</p> <p>OAB-wet*: Urge incontinence with any combination of at least one of the following: urgency, frequency, nocturia, or mixed incontinence</p> <p>OAB-dry*: Urgency with either or both frequency and nocturia</p> <p>Frequency: ≥ 8 voids a day</p>	<p>Prevalence of UII, %: 10.0</p> <p>Prevalence of MUI, %: 7.7</p> <p>UII and MUI Prevalence increases with age <i>P</i><0.01</p> <p>Prevalence of OAB-wet*, %: 5.6</p> <p>Prevalence of OAB-dry*, %: 2.4</p> <p>Overall prevalence of OAB, %: 8.0</p> <p>Prevalence of urgency, %: 10.2</p> <p>Prevalence of frequency, %: 16.4</p>	<p>Overall quality: Good</p> <p>Internal validity score: 5, ++</p> <p>External Validity Score: 4, ++</p> <p>Sampling Method Described: +</p> <p>N sampled provided: +</p> <p>N eligible provided: +</p> <p>N included respondents: +</p> <p>Response Rate:* 77.2</p> <p>Inclusion (Exclusion) Specified: +</p> <p>Age of population described:+</p> <p>Operational definition provided*:+</p> <p>Required frequency defined*:+</p>

Note: differences between UII and OAB-wet prevalence result from differences in operational definitions used in Zhang et al., 2006.

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence [^]	Quality Rating
<p>Author: Stewart et al., 2003 Wagner et al., 2002</p> <p>Country: US</p> <p>Study period: November 2000 to January 2001</p> <p>Funding: Pharmacia</p> <p>Author industry relationship disclosures: 2 of 8 Pharmacia (2)</p>	<p>Design: Cross sectional telephone questionnaire</p> <p>Base population: US adults who are English speaking</p> <p>Sampling frame: Nationwide random sample by telephone to achieve representative sample by age, sex, and region</p> <p>N sampled: 17,231</p> <p>N screened: 11,740</p> <p>N eligible: 6,201</p> <p>N respondents: 6,201</p> <p>N included: Total: 5,204 Women: 2,735</p> <p>Age, %:[^] < 25: 11.2 25-34: 15.3 35-44: 22.2 45-54: 16.5 55-64: 14.1 65-75: 11.1 > 75: 8.6</p> <p>Race/ethnicity, %:[^] White: 81.7 Black: 9.1 Hispanic: 4.2 Other: 3.8</p> <p>BMI:[^] < 22.0: 25.3 22.0-23.9: 15.7 24.0-25.9: 15.9 26.0-29.9: 19.3 ≥ 30: 23.8</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> adults age 18 or older English speaking residential phone most recent birthday among eligibles <p>Exclusion criteria: NR</p> <p>Effective response, %: 83.9</p>	<p>OAB-wet: Per ICS 2002</p> <p>OAB-dry: Per ICS 2002</p>	<p>Prevalence of OAB-wet, %: 9.3</p> <p>Prevalence of OAB-dry, %: 7.6</p>	<p>Overall quality: Fair</p> <p>Internal validity score: 5, ++</p> <p>External Validity Score: 3, +</p> <p>Sampling Method Described: +</p> <p>N sampled provided: +</p> <p>N eligible provided: +</p> <p>N included respondents: +</p> <p>Response Rate:* 83.9</p> <p>Inclusion (Exclusion) Specified: +</p> <p>Age of population described: +</p> <p>Operational definition provided*: +</p> <p>Required frequency defined*: -</p>

[^]Data presented for women only.

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence [^]	Quality Rating
<p>Author: Teloken et al., 2006</p> <p>Country: Brazil</p> <p>Study period: November 2003 to August 2004</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Cross sectional, written questionnaire (King's Health Questionnaire)</p> <p>Base population: Residents of Porto Alegre, Brazil</p> <p>Sampling frame: NR ("population based")</p> <p>N screened: NR</p> <p>N eligible: NR</p> <p>N respondents: 913</p> <p>N included: Total: 848 Women: 449</p> <p>Age, %: < 25: 59.3 26-35: 21.4 36-45: 13.2 46-55: 6.1</p> <p>Race/ethnicity, %: White: 90.8 Black: 3.9 Other: 5.3</p> <p>Parity, %:[^] 39.9</p>	<p>Inclusion criteria: OAB:</p> <ul style="list-style-type: none"> • individuals • ages 15 to 55 <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • current pregnancy • UTI • Diabetes • SUI • diuretic use • urinary tract or gynecologic cancer • renal stones • previous urogenital tract surgery <p>Effective response, %: NR</p>	<p>Per ICS 2002</p>	<p>Prevalence of OAB, %: 23.2</p>	<p>Overall quality: Poor</p> <p>Internal validity score: 1, -</p> <p>External Validity Score: 2, -</p> <p>Sampling Method Described: -</p> <p>N sampled provided: -</p> <p>N eligible provided: -</p> <p>N included respondents: +</p> <p>Response Rate:* NR</p> <p>Inclusion (Exclusion) Specified: +</p> <p>Age of population described: +</p> <p>Operational definition provided*: +</p> <p>Required frequency defined*: -</p>

[^]Data presented for women only.

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence [^]	Quality Rating
<p>Author: Tikkinen et al., 2008</p> <p>Country: Finland</p> <p>Study period: November 2003 to February 2004</p> <p>Funding: Competitive Research Funding of the Pirkanmaa Hospital District; Pfizer</p> <p>Author industry relationship disclosures: 1 of 6 Astellas (1) Orion (1)</p>	<p>Design: Cross-sectional mailed questionnaire (DAN-PSS; AUA-SI)</p> <p>Base population: Women in Finland</p> <p>Sampling frame: Random sample with over-sampling in younger age strata from Finish Population Registry to achieve representative sample with good precision of estimates for all ages</p> <p>N sampled: 3,000</p> <p>N screened: NR</p> <p>N eligible: 2,989</p> <p>N respondents: 2,002</p> <p>N included: 1,728</p> <p>Age, mean ± SD: 42.3 ± 15.6</p> <p>Race/ethnicity: NR</p> <p>Parity, %: 64</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> women age 18 to 79 <p>Exclusion criteria:</p> <ul style="list-style-type: none"> pregnant or postpartum women those reporting current UTI <p>Effective response, %: 57.6</p>	<p>Urgency: An imperative (strong) urge to urinate, often or always</p>	<p>Age-standardized prevalence of urgency, % (95% CI): 10.2 (8.5, 11.9)</p>	<p>Overall quality: Good</p> <p>Internal validity score: 5, ++</p> <p>External Validity Score: 4, ++</p> <p>Sampling Method Described: +</p> <p>N sampled provided: +</p> <p>N eligible provided: +</p> <p>N included respondents: +</p> <p>Response Rate:* 57.6</p> <p>Inclusion (Exclusion) Specified: +</p> <p>Age of population described: +</p> <p>Operational definition provided*: +</p> <p>Required frequency defined*: +</p>

Prevalence of urgency by age in Figure 2 (no data).

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence [^]	Quality Rating
<p>Author: Danforth et al., 2007*</p> <p>Townsend et al., 2008</p> <p>Country: US</p> <p>Study period: Baseline 2000 Follow-up 2002</p> <p>Funding: NIH</p> <p>Author industry relationship disclosures: None</p>	<p>Design: Repeated measures mailed questionnaires at two time points, with supplement to some individuals with incident 1x/wk incontinence at follow-up</p> <p>Base population: Nurses Health Study (NHS) participants</p> <p>Sampling frame: Members of the NHS cohort</p> <p>N sampled/ screened: 121,700 Supplement: 1,939</p> <p>N respondents: 77,696 Supplement: 1,753</p> <p>N eligible: 35,754 Supplement: 1,601</p> <p>Age, mean: 65.3</p> <p>Race/ethnicity: White: 96.5 Black: 1.6 Hispanic: 0.9 Asian: 0.8 Other: 0.2</p> <p>BMI kg/m², %: < 21: 12.6 21-24: 36.3 25-29: 33.9 ≥ 30: 17.1</p> <p>Parity, %: 94.0</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> women NHS participants returning both baseline and follow-up community dwelling <p>Exclusion criteria:</p> <ul style="list-style-type: none"> missing data on BMI or UI prevalent UI 1x/mo at baseline history of major neurologic conditions significant functional limitations <p>Effective response, %: 46.0 Supplement: 82.6</p>	<p>Frequent incontinence: Leaking ≥ 1x/wk</p> <p>UII: Frequent incontinence with a sudden feeling of bladder fullness or when a toilet was inaccessible</p> <p>SUI: Frequent incontinence with coughing or sneezing, lifting things, laughing, or doing exercise</p> <p>MUI: Reporting UII and SUI symptoms equally</p>	<p>2-year incidence of UII, %:** 1.4</p> <p>2-year incidence of MUI, %:** 1.9</p>	<p>Overall quality: Fair</p> <p>Internal validity score: 3, +</p> <p>External Validity Score: 4, ++</p> <p>Sampling Method Described: +</p> <p>N sampled provided: +</p> <p>N eligible provided: +</p> <p>N included respondents: -</p> <p>Response Rate:* 46</p> <p>Inclusion (Exclusion) Specified: +</p> <p>Age of population described: +</p> <p>Operational definition provided*: +</p> <p>Required frequency defined*: +</p>

* Incidence of UII and MUI from Danforth et al., 2007 is 1.4% and 1.8%, respectively.

** Incidence calculated as incidence of frequent UI (1x/wk) times estimated proportion with each type of UI, e.g. (2,416/35,754)*(342/1,601) = 1.4%

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence [^]	Quality Rating
<p>Author: Townsend et al., 2007</p> <p>Country: US</p> <p>Study period: Baseline 2001 Follow-up 2003</p> <p>Funding: NIH</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Repeated measures mailed questionnaires at two time points, with supplemental questionnaire if \geq 1x/wk incontinence at follow-up</p> <p>Base population: Nurses Health Study (NHS) participants</p> <p>Sampling frame: Members of the NHSII cohort</p> <p>N sampled/ screened: 115,713</p> <p>N respondents: Baseline: 85,994 Follow-up: 70,712</p> <p>N eligible: Total: 64,650</p> <p>Age, mean: 46.4</p> <p>Age, %: 36-40: 13.2 41-45: 29.1 46-50: 34.8 51-55: 22.9</p> <p>Race/ethnicity: "Largely Caucasian"</p> <p>BMI kg/m², %: \geq 25: 50.3</p> <p>Parity, %: 79.0</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> women age 35 to 54 at baseline NHS participants returning both baseline and follow-up <p>Exclusion criteria:</p> <ul style="list-style-type: none"> missing data unclear continence status at baseline <p>Response, %: Baseline: 74 Follow-up: 61</p> <p>Effective response, %: Total: 57</p>	<p>Incontinence: "During last 12 months, how often have you leaked or lost control of your urine?" (required \geq 1x/wk)</p> <p>UII: Predominance of urge symptoms in the Medical Epidemiological and Social Aspects of Aging questionnaire, per Diokno (1995)</p> <p>SUI: Predominance of stress symptoms</p> <p>MUI: Equal predominance of UII and SUI symptoms</p>	<p>2-year incidence of UII, %: 0.4</p> <p>2-year incidence of UII by age, %: 36-40: 0.4 41-45: 0.4 46-50: 0.5 51-55: 0.4</p> <p>2-year incidence of MUI, %: 0.6</p> <p>2-year incidence of MUI by age, %: 36-40: 0.3 41-45: 0.6 46-50: 0.7 51-55: 0.9</p>	<p>Overall quality: Fair</p> <p>Internal validity score: 5, ++</p> <p>External Validity Score: 4, ++</p> <p>Sampling Method Described: +</p> <p>N sampled provided: +</p> <p>N eligible provided: +</p> <p>N included respondents: +</p> <p>Response Rate:* 61</p> <p>Inclusion (Exclusion) Specified: +</p> <p>Age of population described:+</p> <p>Operational definition provided*:+</p> <p>Required frequency defined*:+</p>

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence [^]	Quality Rating
<p>Author: Tseng et al., 2000</p> <p>Country: Taiwan</p> <p>Study period: March 1997 to April 1997</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Cross-sectional administered questionnaire</p> <p>Base population: Residents of Tungkang Town</p> <p>Sampling frame: Random sample stratified to obtain age and gender representation by city region</p> <p>N sampled: NR</p> <p>N screened: NR</p> <p>N eligible: NR</p> <p>N respondents: NR</p> <p>N included: Total: 504 Women: 256</p> <p>Age, %: 65-70: 43.1 71-75: 28.1 76-80: 18.4 > 80: 12.1</p> <p>Race/ethnicity: NR</p> <p>BMI, %: "Overweight": 47.3*</p>	<p>Inclusion criteria: UI:</p> <ul style="list-style-type: none"> • adults • age ≥ 65 <p>Exclusion criteria: NR</p> <p>Effective response, %: 80 (per authors cannot calculate)</p>	<p>UI: "ever experienced inappropriate loss of urine"</p> <p>UUI: Because of inability to delay voiding following micturition urge</p> <p>SUI: Loss associated with physical exertion (coughing, sneezing, lifting, or other physical activity)</p> <p>MUI: Features of UUI and SUI</p>	<p>Prevalence of UUI, % : 6.6</p> <p>Prevalence of MUI, %: 6.3</p>	<p>Overall quality: Poor</p> <p>Internal validity score: 2, -</p> <p>External Validity Score: 4, ++</p> <p>Sampling Method Described: +</p> <p>N sampled provided: -</p> <p>N eligible provided: -</p> <p>N included respondents: +</p> <p>Response Rate:* NR</p> <p>Inclusion (Exclusion) Specified: +</p> <p>Age of population described: +</p> <p>Operational definition provided*: +</p> <p>Required frequency defined*: +</p>

[^] Data presented for women only.

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence [^]	Quality Rating
<p>Author: Ueda et al., 2000</p> <p>Country: Japan</p> <p>Study period: NR</p> <p>Funding: LACADIA Health and Welfare Foundation; Social Department Foundation for Senior Citizens</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Cross-sectional mailed questionnaire</p> <p>Base population: Residents of Shiga Prefecture</p> <p>Sampling frame: Random sample of individuals in seven towns</p> <p>N sampled: 3,500</p> <p>N screened: NR</p> <p>N eligible: NR</p> <p>N respondents: 1,836</p> <p>N included: Total: 1,786 Women: 968</p> <p>Age, %: ^ 40-49: 26.4 50-59: 28.8 60-69: 29.4 ≥ 70: 15.4</p> <p>Race/ethnicity: NR</p> <p>BMI: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • adults • age 40 to 80 • community-dwelling <p>Exclusion criteria: NR</p> <p>Effective response, %: 51.0</p>	<p>Incontinence: involuntary loss of urine within prior year</p> <p>UUI: Often have difficulty holding urine until able to get to toilet</p> <p>SUI: Leak when cough, sneeze, or laugh?</p> <p>MUI: Both UUI and SUI</p>	<p>Prevalence of UUI, %: 6.9</p> <p>Prevalence of UUI by age, %: 40-49: 5.1 50-59: 4.3 60-69: 8.8 ≥ 70: 11.4</p> <p>Prevalence of MUI, %: 12.9</p> <p>Prevalence of MUI by age, %: 40-49: 14.5 50-59: 14.3 60-69: 6.3 ≥ 70: 20.1</p>	<p>Overall quality: Fair</p> <p>Internal validity score: 4, +</p> <p>External Validity Score: 4, ++</p> <p>Sampling Method Described: +</p> <p>N sampled provided: +</p> <p>N eligible provided: -</p> <p>N included respondents: +</p> <p>Response Rate:* 51</p> <p>Inclusion (Exclusion) Specified: +</p> <p>Age of population described: +</p> <p>Operational definition provided*: +</p> <p>Required frequency defined*: +</p>

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence [^]	Quality Rating
<p>Author: van der Vaart et al., 2002</p> <p>Country: Netherlands</p> <p>Study period: 1999</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Cross-sectional mailed questionnaire (UDI)</p> <p>Base population: Female residents of the Central Netherlands</p> <p>Sampling frame: Random sample of women from the population registry office of a suburban area</p> <p>N sampled: 1,393</p> <p>N screened: NR</p> <p>N eligible: NR</p> <p>N respondents: 1,029</p> <p>N included: 933</p> <p>Age, mean ± SD: 34.2 ± 3.2</p> <p>Race/ethnicity: NR</p> <p>BMI, mean ± SD: 23.6 ± 4.2</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • women • age 20 to 45 <p>Exclusion criteria: NR</p> <p>Effective response, %: 67</p>	<p>UUI: Urine loss related to feeling of urgency</p> <p>OAB: Per ICS</p> <p>Urgency: Per ICS</p> <p>Frequency: Per ICS</p>	<p>Prevalence of UUI, %: 15.3</p> <p>Prevalence of OAB, %: 11.9</p> <p>Prevalence of urgency, %: 45.4</p> <p>Prevalence of frequency, %: 34.0</p>	<p>Overall quality: Fair</p> <p>Internal validity score: 4, +</p> <p>External Validity Score: 3, +</p> <p>Sampling Method Described: +</p> <p>N sampled provided: +</p> <p>N eligible provided: -</p> <p>N included respondents: +</p> <p>Response Rate:* 67%</p> <p>Inclusion (Exclusion) Specified: +</p> <p>Age of population described:+</p> <p>Operational definition provided*:+</p> <p>Required frequency defined*:-</p>

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence [^]	Quality Rating
<p>Author: Waetjen et al., 2007</p> <p>Country: US</p> <p>Study period: 1995 to 2001</p> <p>Funding: NIH, Univ. of California Davis Health Systems Research Award</p> <p>Author industry relationship disclosures: None</p>	<p>Design: Repeated measures administered questionnaire and follow-up visits</p> <p>Base population: Women in six states near seven urban study centers</p> <p>Sampling frame: Random digit dialing, snowballing, and list sampling based on study center</p> <p>N sampled: 16,065</p> <p>N screened: 16,065</p> <p>N eligible: NR</p> <p>N respondents: 16,065 baseline 3,302 included in follow-up based on per center quotas</p> <p>Age, mean ± SD: 45.8 ± 2.7</p> <p>Race/ethnicity, %: Caucasian: 47.0 Black: 28.3 Chinese: 7.6 Japanese: 8.5 Hispanic: 8.7</p> <p>BMI, mean ± SD: 28.2 ± 7.2</p> <p>Parity, mean ± SD: 2.0 ± 1.4</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> women age 42 to 52 self-identified as member of one of five race/ethnic groups <p>Exclusion criteria:</p> <ul style="list-style-type: none"> inability to speak English, Spanish, Japanese, or Cantonese no menstrual period in prior 3 months hysterectomy and/or oophorectomy prior to study current use of hormones including birth control pills <p>Effective response, %: NR</p>	<p>UII: at least monthly leakage when “urge to void and can’t reach the toilet fast enough.”</p> <p>SUI: at least monthly leakage “with coughing, laughing, sneezing, jogging, jumping, or picking up an object from the floor.”</p> <p>MUI: Affirmative answers to both UII and SUI at least monthly</p>	<p>5-year incidence of UII, %: 15.9</p> <p>5-year incidence of MUI, %: 11.9</p> <p>Baseline prevalence of UII, %: 7.6</p> <p>Baseline prevalence of MUI, %: 12.4</p>	<p>Overall quality: Fair</p> <p>Internal validity score: 3, -</p> <p>External Validity Score: 3, ++</p> <p>Sampling Method Described: +</p> <p>N sampled provided: +</p> <p>N eligible provided: -</p> <p>N included respondents: +</p> <p>Response Rate:* NR</p> <p>Inclusion (Exclusion) Specified: +</p> <p>Age of population described: +</p> <p>Operational definition provided*.: +</p> <p>Required frequency defined*.: +</p>

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence [^]	Quality Rating
<p>Author: Wesnes et al., 2007</p> <p>Country: Norway</p> <p>Study period: 1999 to 2006</p> <p>Funding: Research Council of Norway</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Cross-sectional mailed questionnaire</p> <p>Base population: Pregnant women in Norway</p> <p>Sampling frame: Women receiving prenatal care in one of 52 maternity units</p> <p>N screened: NR</p> <p>N eligible: NR</p> <p>N respondents: 46,262</p> <p>N included: 43,279</p> <p>Age, mean (range): 29.5 (14, 47)</p> <p>Race/ethnicity: NR</p> <p>Pre-pregnancy BMI, mean (range): 24.1 (13, 59)</p> <p>Parity, %: 53.8</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • pregnant women • read and write Norwegian • completion of questionnaire by 30-week visit <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • participated in prior pregnancy <p>Effective response, %: 45</p>	<p>UII: Per ICS 2002</p> <p>MUI: Per ICS 2002</p>	<p>Prevalence of UII, before pregnancy, %: 3.8</p> <p>Prevalence of UII during pregnancy, %: 4.8</p> <p>Prevalence of MUI, before pregnancy, %: 5.5</p> <p>Prevalence of MUI during pregnancy, %: 16.4</p>	<p>Overall quality: Fair</p> <p>Internal validity score: 3, +</p> <p>External Validity Score: 3, +</p> <p>Sampling Method Described: +</p> <p>N sampled provided: -</p> <p>N eligible provided: -</p> <p>N included respondents: +</p> <p>Response Rate:* 45</p> <p>Inclusion (Exclusion) Specified: +</p> <p>Age of population described: +</p> <p>Operational definition provided*: +</p> <p>Required frequency defined*: -</p>

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence [^]	Quality Rating
<p>Author: Yarnell et al., 1981</p> <p>Country: UK</p> <p>Study period: NR</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Cross-sectional administered questionnaire</p> <p>Base population: Female residents of South Wales</p> <p>Sampling frame: Random sample of electoral role from base population of 38,000</p> <p>N sampled: 1,140</p> <p>N screened: NR</p> <p>N eligible: 1,060</p> <p>N respondents: 1,022</p> <p>N included: 1,000</p> <p>Age, %: 17-24: 12 23-34: 21 35-44: 15 45-54: 19 55-64: 15 65-74: 11 ≥ 75: 7</p> <p>Race/ethnicity: NR</p> <p>BMI: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> women age ≥ 18 <p>Exclusion criteria: NR</p> <p>Effective response, %: 94.3</p>	<p>UII: Ever (12 months) have to rush to the toilet to pass water and ever loose water before reaching toilet</p> <p>SUI: Ever (12 months) loose urine when you cough, laugh, sneeze, etc.</p> <p>MUI: Both UII and MUI</p>	<p>Prevalence of UII, %: 9.2</p> <p>Prevalence of UII by age, %: 17-24: 9 23-34: 5 35-44: 10 45-54: 9 55-64: 10 65-74: 13 ≥75: 14</p> <p>Prevalence of MUI, %: 13.5</p> <p>Prevalence of MUI by age, %: 17-24: 4 23-34: 8 35-44: 16 45-54: 16 55-64: 15 65-74: 13 ≥ 75: 28</p>	<p>Overall quality: Good</p> <p>Internal validity score: 5, ++</p> <p>External Validity Score: 4, ++</p> <p>Sampling Method Described: +</p> <p>N sampled provided: +</p> <p>N eligible provided: +</p> <p>N included respondents: +</p> <p>Response Rate:* 94.3</p> <p>Inclusion (Exclusion) Specified: +</p> <p>Age of population described: +</p> <p>Operational definition provided*: +</p> <p>Required frequency defined*: +</p>

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence [^]	Quality Rating
<p>Author: Yu et al., 2006</p> <p>Country: Taiwan</p> <p>Study period: February 2003 to February 2005</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Cross-sectional administered questionnaire</p> <p>Base population: Residents of Matsu</p> <p>Sampling frame: 100% of residents</p> <p>N sampled: 5,456</p> <p>N screened: NR</p> <p>N eligible: NR</p> <p>N respondents: 1,921</p> <p>N included: Total: 1,827 Women: 925</p> <p>Age, %:[^] 30-39: 22.6 40-49: 34.5 50-59: 19.4 60-69: 12.8 70-79: 10.8</p> <p>Race/ethnicity: NR</p> <p>BMI kg/m², %:[^] < 24.0: 70.9 24.0-26.9: 20.5 ≥ 27.0: 8.5</p> <p>Parity, %: 89.4</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> adults age ≥ 30 in 2000 <p>Exclusion criteria: NR</p> <p>Effective response, %: 33.5</p>	<p>UII: Urine loss associated with an urge to void</p> <p>OAB: Per ICS 2002</p>	<p>Prevalence of OAB, %: 18.3</p> <p>Prevalence of OAB by age, %: 30-39: 14.4 40-49: 14.7 50-59: 19.6 60-69: 25.4 70-79: 31.0 <i>P</i> = 0.001 for trend</p>	<p>Overall quality: Fair</p> <p>Internal validity score: 4, +</p> <p>External Validity Score: 3, +</p> <p>Sampling Method Described: +</p> <p>N sampled provided: +</p> <p>N eligible provided: -</p> <p>N included respondents: +</p> <p>Response Rate:* 33.5</p> <p>Inclusion (Exclusion) Specified: +</p> <p>Age of population described: +</p> <p>Operational definition provided*: +</p> <p>Required frequency defined*: -</p>

[^]Data presented for women only.

Evidence Table 1. KQ 1 Prevalence and Incidence of OAB (continued)

Study Description	Study Design and Sampling	Inclusion/Exclusion Criteria	Study Definitions	Incidence/Prevalence	Quality Rating
<p>Author: Zhu et al., 2008</p> <p>Country: China</p> <p>Study period: April 2005 to July 2005</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Cross-sectional administered questionnaire (Chinese ICIQ)</p> <p>Base population: Female residents of Beijing</p> <p>Sampling frame: 1.0% of female Beijing residents age ≥ 20 (2000 national census data)</p> <p>N sampled: 5,300</p> <p>N included: 5,221</p> <p>Age, mean: 46.4</p> <p>Race/ethnicity: NR</p>	<p>Inclusion criteria: • Age ≥ 20</p> <p>Exclusion criteria: NR</p> <p>Effective response, %: 98.5</p>	<p>Incontinence: Any leakage or involuntary loss of urine within the last month.</p> <p>SUI: A leak or loss of urine caused by sneezing, coughing, exercising, lifting, or physical activity.</p> <p>UUI: An urge to urinate but being unable to reach the toilet before leaking or having a strong sudden urge to go to the toilet to urinate with no advance warning.</p> <p>MUI: Experiencing at least one stress and one urge symptom.</p>	<p>Prevalence of UUI, %: 2.8</p> <p>Prevalence of UUI by menopausal status, %: Premenopause: 1.9 Menopause: 4.0</p> <p>Prevalence of MUI, %: 12.4</p> <p>Prevalence of MUI by menopausal status, %: Premenopause: 6.0 Menopause: 20.4</p>	<p>Overall quality: Good</p> <p>Internal validity score: 4, +</p> <p>External Validity Score: 4, ++</p> <p>Sampling Method Described: +</p> <p>N sampled provided: +</p> <p>N eligible provided: +</p> <p>N included respondents: +</p> <p>Response Rate:* 98.5</p> <p>Inclusion (Exclusion) Specified: +</p> <p>Age of population described: +</p> <p>Operational definition provided*: +</p> <p>Required frequency defined*: -</p>

Mean age and prevalence of UUI and MUI by menopausal status calculated from results reported. Prevalence of UUI and MUI by age groups in Figure 3 on page 567 (no results reported).

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Abrams et al., 1998</p> <p>Country and setting: UK, Ireland, Sweden, Academic medical center</p> <p>Enrollment period: July 1995 to July 1996</p> <p>Funding: Pharmacia Upjohn</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: Tolterodine vs. Oxybutynin vs. Placebo x 12 wks</p> <p>Groups: G1: Tolterodine 2 mg b.i.d. (dose could be reduced to 1.0 mg b.i.d.) G2: Oxybutynin 5 mg t.i.d. (dose could be reduced to 2.5 mg t.i.d.) G3: Placebo</p> <p>N at enrollment: G1: 118 G2: 118 G3: 57</p> <p>N at follow-up: G1: 118 G2: 117 G3: 56</p> <p>Women, n (%): G1: 91 (77.1) G2: 88 (74.5) G3: 43 (75.4)</p> <p>Age, mean (range): G1: 55 (19-80) G2: 58 (21-80) G3: 58 (26-78)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age \geq 18 • UDS confirmed bladder overactivity • \geq 8 voids/day • \geq 1 episode UUI/ day <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • SUI • Detrusor hyperreflexia • Hepatic, renal, hematological disorders • Symptomatic or recurrent UTI • BOO • Bladder retraining • Electrical stimulation therapy • Indwelling catheter • CIC • Pregnant/ nursing • Women without reliable BC 	<p>Incontinence episodes, n (%): G1: 93 (79) G2: 88 (75) G3: 40 (70)</p> <p>Incontinence episodes/day, mean (range): G1: 2.9 (0.1-24.0) G2: 2.6 (0.1-24.0) G3: 3.3 (0.1-23.5)</p> <p>Voids/day, mean (range): G1: 11.5 (6.3-37.0) G2: 10.7 (5.3-27.7) G3: 11.7 (6.9-34.4)</p> <p>\geq 8 voids/day, n (%): G1: 112 (95) G2: 109 (92) G3: 53 (93)</p> <p>Voided volume (mL), mean (range): G1: 166 (55-426) G2: 176 (57-423) G3: 157 (59-276)</p> <p>Previous OAB therapy, n (%): G1: 61 (52) G2: 71 (60) G3: 43 (75)</p>	<p>Incontinence episodes/day, mean change \pm SD: G1: -1.3 \pm 3.2 G2: -1.7 \pm 3.1 G3: -0.9 \pm 1.5 G1/G3: $P = 0.22$ G2/G3: $P = 0.023$</p> <p>Voids/day, mean change \pm SD: G1: -2.7 \pm 3.8 G2: -2.3 \pm 2.7 G3: -1.6 \pm 3.6 G1/G3: $P = 0.002$ G2/G3: $P = 0.068$</p> <p>Voided volume (mL), mean change \pm SD: G1: 38 \pm 54 G2: 47 \pm 58 G3: 6 \pm 42 G1/G3: $P < 0.001$ G2/G3: $P < 0.001$</p> <p>Subjective improvement in bladder symptoms, %: G1: 50 G2: 49 G3: 47</p> <p>Dose reductions, n (%): G1: 9 (8) G2: 38 (32) G3: 1 (2) G2/G1: $P < 0.001$ G2/G3: $P < 0.001$</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: fair</p> <p>Randomization: +</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: ++</p> <p>Drop-out rates: -</p> <p>Power calculation: +</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Abrams et al., 1998 (continued)				<p>Dry mouth, n (%): G1: 59 (50) G2: 102 (86) G3: 12 (21) G1/G3: $P < 0.001$ G2/G1: $P < 0.001$ G2/G3: $P < 0.001$</p> <p>Dyspepsia, n (%): G1: 11 (9) G2: 27 (23) G3: 3 (5)</p> <p>Nausea, n (%): G1: 4 (3) G2: 7 (6) G3: 6 (11)</p> <p>Upper respiratory infection, n (%): G1: 12 (10) G2: 3 (3) G3: 8 (14)</p> <p>Adverse events reported, N: G1: 302 G2: 412 G3: 117</p> <p>Patients with ≥ 1 AE, n (%): G1: 105 (89) G2: 114 (97) G3: 46 (81) G2/G1: $P = 0.023$ G2/G3: $P < 0.001$</p> <p>Discontinued due to AEs, n (%): G1: 10 (8) G2: 20 (17) G3: 1 (2)</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Abrams et al., 2006</p> <p>Country and setting: UK, 8 sites</p> <p>Enrollment period: NR</p> <p>Funding: Pfizer</p> <p>Author industry relationship disclosures: 2 of 6 Pfizer (2)</p>	<p>Design: Crossover RCT</p> <p>Intervention: 8 weeks consisting of 2-week run-in period followed by two-week treatment period, 2 week washout, and additional 2 week treatment.</p> <p>Groups: G1: Propiverine 20 mg qd G2: Propiverine 15 mg t.i.d. G3: Oxybutynin 5 mg t.i.d. G4: Placebo</p> <p>N at enrollment: Total: 77 G1: 38 G2: 42 G3: 41 G4: 24</p> <p>N at follow-up: Safety Analysis: 77 AUM parameter analysis: 69</p> <p>Women, n (%)*: 59 (76.6)</p> <p>Age, range: 47-56</p> <p>Race/ethnicity: NR</p> <p>Weight, range (kg): 74-81</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age \geq 18 • Clinical diagnosis of idiopathic OAB with detrusor overactivity • Two or more of the following during 2-week run-in period: urinary frequency (\geq 7 voids/ day), UUI (\geq 1 episode/week requiring change of clothing or pad), urinary urgency (\geq 7 episodes preceding week) <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Hepatic, renal or cardiac abnormalities • SUI • Untreated narrow-angle glaucoma • Urinary or gastric retention • BOO > 40 (Abrams-Griffiths number) • Indwelling catheter • Recent urogenital surgery • Use of investigational drugs in 30 days preceding the study 	<p>Duration of OAB (years), range: 5-15 f</p>	<p>IDC number, all activities, mean difference (SE): G2/G1: -1.3 (1.4)* G3/G1: -3.8 (1.4) $P \leq$ 0.01 G3/G2: -2.6 (1.4)*</p> <p>IDC number, with concurrent symptoms, mean difference (SE): G2/G1: -0.9 (0.7)* G3/G1: -1.7 (0.7) $P \leq$ 0.05 G3/G2: -0.8 (0.7)*</p> <p>IDC duration (sec), all activities, mean difference (SE): G2/G1: -133.6 (88.0)* G3/G1: -244 (91.3) $P \leq$ 0.01 G3/G2: -110.4 (88.9)*</p> <p>IDC duration (sec), with concurrent symptoms, mean difference (SE): G2/G1: -54.3 (48.0)* G3/G1: 108.1 (50.0) $P \leq$ 0.05 G3/G2: 53.7 (49.0)*</p> <p>AUC detrusor pressure vs time (cmH₂O * s), all activities, mean difference (SE): G2/G1: -0.1 (0.5)* G3/G1: -0.3 (0.5)* G3/G2: -0.2 (0.5)*</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: - Masking: + Pt selection criteria: + Loss to followup: + Drop-out rates: ++ Power calculation: - Statistical issues: +</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: -, NR Baseline OAB status: NR Baseline characteristics: - Length of followup: - Measurement methods: + Measurement reliability: + Intervention description: +</p>

^ gender accounted for in mixed-effects statistical model; * $P = NS$

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Abrams et al., 2006 (continued)				<p>AUC detrusor pressure vs. time (cmH₂O * s), with concurrent symptoms, mean difference (SE): G2/G1: -0.2 (0.8)* G3/G1: -1.3 (0.8)* G3/G2: -1.1 (0.8)*</p> <p>IDC max amplitude (cmH₂O), all activities, mean difference (SE): G2/G1: -3.8 (8.2)* G3/G1: -16.8 (8.5)* G3/G2: 12.7 (8.3)*</p> <p>IDC max amplitude (cmH₂O), with concurrent symptoms, mean difference (SE): G2/G1: -5.0 (7.6)* G3/G1: -15.8 (7.9) <i>P</i> ≤ 0.05 G3/G2: -10.7 (7.7)*</p> <p>Salivary flow rate, mean difference (SE): G2/G1: 0.0 (0.7)* G3/G1: -0.6 (0.1) <i>P</i> < 0.0001 G3/G2: -0.6 (0.1) <i>P</i> < 0.0001</p> <p>Visual near point, mean difference (SE): G2/G1: 1.6 (1.2)* G3/G1: 0.6 (1.2)* G3/G2: -1.0 (1.2)*</p> <p>Heart rate, mean difference (SE): G2/G1: 2.7 (1.4) <i>P</i> < 0.05 G3/G1: -5.1 (1.4) <i>P</i> < 0.0001 G3/G2: -7.8 (1.4) <i>P</i> < 0.0001</p>	

^ gender accounted for in mixed-effects statistical model; * *P* = NS

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Abrams et al., 2006 (continued)				<p>PNN50, mean difference (SE): G2/G1: -3.6 (1.8) <i>P</i> < 0.05 G3/G1: 6.2 (1.8) <i>P</i> < 0.0001 G3/G2: 9.9 (1.8) <i>P</i> < 0.0001</p> <p>St. George's index, mean difference (SE): G2/G1: -3.5 (1.9) <i>P</i> < 0.05 G3/G1: 3.4 (1.9) <i>P</i> < 0.05 G3/G2: 6.9 (1.9) <i>P</i> < 0.0001</p> <p>Dry mouth, n (%): G1: 13 (34) G2: 22 (52) G3: 34 (83) G4: 4 (17)</p> <p>Abnormal vision, n (%): G1: 9 (24) G2: 14 (33) G3: 9 (22) G4: 0</p> <p>Constipation, n (%): G1: 6 (16) G2: 10 (24) G3: 4 (10) G4: 0</p> <p>Headache, n (%): G1: 1 (3) G2: 3 (7) G3: 6 (15) G4: 0</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Anderson et al., 1999</p> <p>Country and setting: US, multi-center</p> <p>Enrollment period: NR</p> <p>Funding: Oros Oxybutynin Study group Alza Corp</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: Oxybutynin CR vs. Oxybutynin IR 1 wk washout period prior to initiation of study</p> <p>Groups: G1: Oxybutynin 5, 10, 15, 20, 25, or 30 mg CR qd G2: Oxybutynin 5 mg IR daily to qid</p> <p>N at enrollment: G1: 53 G2: 52</p> <p>N at follow-up: G1: 46 G2: 47</p> <p>Age, yrs ± SD: G1: 59.2 ± 10.6 G2: 59.6 ± 10.0</p> <p>Race/ethnicity: NR</p> <p>Women, N (%): G1: 50 (94) G2: 47 (90)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Community dwelling • Symptoms of urge incontinence or mixed incontinence with primary urge symptoms • ≥6 episodes UUI/ wk when not taking medications • Previous favorable response with oxybutynin <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Myasthenia gravis • Narrowing of GI tract • glaucoma • Pregnant/ lactating • Positive urine drug screen • Receiving meds other than oxybutynin, hyoscamine, or propantheline which was considered effective in UUI treatment • At risk for complete urinary retention caused by anticholinergic treatment • Prostate surgery in last 9 mos • BOO • PVR>100 	<p>UUI episodes/ week, mean ± SD: G1: 27.4 ± 24.0 G2: 23.4 ± 16.3</p> <p>Total incontinence episodes, mean ± SD: G1: 29.3 ± 24.2 G2: 26.3 ± 18.5</p> <p>Voids/ week, mean ± SD: G1: 48.3 ± 18.4 G2: 51.5 ± 19.4</p> <p>Voided volume/ void (mL), mean ± SD: G1: 134.2 ± 82 G2: 161.2 ± 92</p> <p>PVR/ void (mL), mean ± SD: G1: 15.4 ± 21 G2: 18.0 ± 24</p> <p>Total bladder volume/ void (mL), mean ± SD: G1: 149.7 ± 90 G2: 179.2 ± 95</p>	<p>UUI episodes/ week, mean: G1: 4.8 G2: 3.1 <i>P</i> = 0.6</p> <p>Total incontinence episodes, mean: G1: 6 G2: 3.8 <i>P</i> = 0.7</p> <p>Percentage reduction in UUI episodes/ week (%): G1: 84 G2: 88 <i>P</i> = 0.7</p> <p>No UUI episodes/ week, %: G1: 52 G2: 51 <i>P</i> = 0.7</p> <p>Totally continent, %: G1: 41 G2: 41 <i>P</i> = 0.9</p> <p>Voided volume/ void (mL), mean ± SD: G1: 176.6 ± 138 G2: 194.5 ± 152 <i>P</i> = 0.1</p> <p>PVR/ void (mL), mean ± SD: G1: 33.5 ± 77 G2: 36.1 ± 59 <i>P</i> = 0.2</p> <p>Total bladder volume/ void (mL), mean ± SD: G1: 210.1 ± 148 G2: 230.6 ± 176 <i>P</i> = 0.9</p> <p>≥1 anticholinergic event, n (%): G1: 46 (87) G2: 49 (94)</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: -</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: ++</p> <p>Drop-out rates: +</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: -</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Anderson et al., 1999 (continued)				<p>Dry Mouth, n (%): G1: 36 (68) G2: 45 (87) <i>P</i> = 0.04</p> <p>Moderate-severe dry mouth, n (%): G1: 13 (25) G2: 24 (46) <i>P</i> = 0.03</p> <p>Somnolence, n (%): G1: 20 (38) G2: 21 (40) <i>P</i> = 0.8</p> <p>Blurred vision, n (%): G1: 15 (28) G2: 9 (17) <i>P</i> = 0.3</p> <p>Constipation, n (%): G1: 16 (30) G2: 16 (31) <i>P</i> = 1.0</p> <p>Dizziness, n (%): G1: 15 (28) G2: 20 (38) <i>P</i> = 0.3</p> <p>Impaired urination, n (%): G1: 13 (25) G2: 15 (29) <i>P</i> = 0.7</p> <p>Nervousness, n (%): G1: 13 (25) G2: 12 (23) <i>P</i> = 1.0</p> <p>Nausea, n (%): G1: 10 (19) G2: 9 (17) <i>P</i> = 1.0</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Anderson et al., 2006</p> <p>[See evidence table for Diokno et al. 2003]</p> <p>Country and setting: US, Multicenter</p> <p>Enrollment period: NR</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: 1 of 6 ALZA Corp (1)</p>	<p>Design: RCT</p> <p>Intervention: Extended release antimuscarinic medications in women with or without prior antimuscarinic treatment for OAB</p> <p>Groups: G1: Prior antimuscarinic G1a: Oxybutynin ER 10 mg po daily G1b: Tolterodine ER 4mg po daily G2: No prior anticholinergic treatment G2a: Oxybutynin ER 10 mg po daily G2b: Tolterodine ER 4mg po daily Ga: Oxybutynin ER 10 mg po daily Gb: Tolterodine ER 4mg po daily</p> <p>N at enrollment: Total: 790 G1: 373 G1a: 180 G1b: 193 G2: 417 G2a: 211 G2b: 206</p> <p>N at follow-up, 12 weeks: G1: 341 G1a: 165 G1b: 176 G2: 355 G2a: 174 G2b: 181</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Women • Age ≥ 18 • Mean of 21-60 UUI episodes per week and mean of ≥ 10 voids per 24 hr <p>Exclusion criteria: NR</p>	<p>UUI episodes/wk, mean ± SD: G1a: 36.8 ± 16.4 G1b: 37.4 ± 14.0 G2a: 37.5 ± 14.0 G2b: 36.2 ± 13.9</p> <p>Incontinence episodes/wk, mean ± SD: G1a: 41.5 ± 19.0 G1b: 43.0 ± 18.0 G2a: 45.0 ± 19.4 G2b: 41.9 ± 17.9</p> <p>UUI episodes/wk, mean ± SD: G1a: 11.4 ± 17.9 G1b: 13.3 ± 15.1 G2a: 10.2 ± 13.7 G2b: 9.3 ± 13.3 G1a/G1b: <i>P</i> = 0.3</p> <p>UUI episodes/wk, completed wk 12, mean ± SD: G1a: 9.9 ± 14.1 G1b: 12.9 ± 14.9 <i>P</i> = 0.049</p> <p>No UUI, 12 wk, %: G1a: 25.2 G1b: 16.4 G2a: 29.4 G2b: 26.4 G1a/G1b: <i>P</i> = 0.046 G2a/G2b: <i>P</i> = 0.495</p> <p>Incontinence episodes/wk, mean ± SD: G1a: 12.7 ± 18.7 G1b: 16.5 ± 19.8 G2a: 11.9 ± 15.1 G2b: 11.3 ± 16.0 G1a/G1b: <i>P</i> = 0.09 for LOCF (<i>P</i> = 0.012 if completed wk 12) G2a/G2b: <i>P</i> = 0.886</p> <p>UUI episodes/wk, mean ± SD: G1a: 68.4 ± 17.2 G1b: 72.8 ± 25.4 G2a: 64.8 ± 22.0 G2b: 69.4 ± 21.3 G1a/G1b: <i>P</i> = 0.05 for LOCF (<i>P</i> = 0.026 wk 12) G2a/G2b: <i>P</i> = 0.035 for LOCF (<i>P</i> = 0.026 wk 12)</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL</p> <p>VALIDITY: poor</p> <p>Randomization: -</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: ++</p> <p>Drop-out rates: ++</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL</p> <p>VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Anderson et al., 2006 (continued)	<p>Women, %: 100</p> <p>Age, mean ± SD: G1a: 62.6 ± 12.9 G1b: 62 ± 12.6 G2a: 57.5 ± 13.4 G2b: 58.8 ± 12.4</p> <p>Race/ethnicity, %: White: G1a: 87 G1b: 88 G2a: 82 G2b: 84 Black: G1a: 7 G1b: 9 G2a: 9 G2b: 8.7 Asian: G1a: 0.6 G1b: 0 G2a: 0 G2b: 1 Hispanic: G1a: 5.6 G1b: 3.1 G2a: 8.1 G2b: 6.3 Other G1a: 0 G1b: 0.5 G2a: 0.9 G2b: 0</p>			<p>Dry mouth, % G1a: 32 G1b: 19 G2a: 27.5 G2b: 25.2 G1a/G1b: <i>P</i> = 0.004 G2a/G2b: <i>P</i> = 0.6</p> <p>Constipation, %: G1a: 7.8 G1b: 5.2 G2a: 5.2 G2b: 10.2</p> <p>Diarrhea, %: G1a: 7.8 G1b: 5.7 G2a: 8.1 G2b: 6.8</p> <p>Headache, %: G1a: 4.4 G1b: 5.2 G2a: 6.6 G2b: 6.8</p> <p>Discontinued due to AE, n (%): G1a: 7 (3.9) G1b: 6 (3.1) G2a: 13 (6.2) G2b: 13 (6.3)</p> <p>Withdrew, %: G1: 3.5 G2: 6.2</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Appell, Abrams et al., 2001</p> <p>Country and setting: Multinational, Multicenter</p> <p>Enrollment period: NR</p> <p>Funding: Pharmacia</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Open-label extension study (after 12-wk RCT and 1 wk washout)</p> <p>Intervention: Tolterodine 2 mg b.i.d. with option for patients to self-lower their dosage to 1 mg b.i.d.</p> <p>Groups: NA</p> <p>N at enrollment: 854</p> <p>N at follow-up, n (%): 594 (70)</p> <p>Age, yrs, mean (range): 59.5 (19-89)</p> <p>Race/ethnicity: NR</p> <p>Women, %: 76</p> <p>Previous drug therapy for OAB, n (%): 472 (55)</p> <p>Previous lower urinary tract surgery, n (%): 291 (34)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • ≥18 years old • Cystometric evidence of detrusor overactivity (phasic detrusor contraction with amplitude 10+ cm H2O) • ≥ 8 voids/day • ≥ 1 urinary incontinence episode/day <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • SU1 • Hepatic or renal disease • Recurrent or symptomatic UTIs • Interstitial cystitis • Hematuria secondary to malignant disease • Contraindication to anti-muscarinic therapy • Serious AE on oxybutynin • Clinically significant voiding difficulty w/ treatment of urinary retention • Treatment in 14 days prior to baseline visit • Initiation of antimuscarinic or any drug for UI during study • Electro-stimulation or bladder training • Indwelling or intermittent catheter • Total voided volume >3L/day 	<p>Detrusor instability, %: 93.7</p> <p>UUI episodes, n (%): 724 (85)</p> <p>Symptom duration > 5 yrs, n (%): 412 (48)</p> <p>Urgency, n (%): 841 (98)</p> <p>Severe/very severe problems, n (%): 384 (45)</p> <p>UUI episodes/day, mean (range): 3.5 (0.1-24.0)</p> <p>Voids/day, mean (range): 11.4 (5.3-37.0)</p> <p>Voided volume (mL), mean (range): 159 (25-423)</p> <p>Adverse events at end of 12-wk RCT, n (%): Any: 358 (76) ANS: 203 (43) CNS: 59 (12) GI: 125 (26) Respiratory: 68 (14) Urinary: 50 (11) Dry mouth: 187 (39) UTI: 26 (5) Headache: 49 (10) Constipation: 31 (7) Abdominal pain: 36 (8) URI: 28 (6)</p>	<p>UUI episodes/day, 3 mos, mean (range): 1.3 (0.0-24.0)</p> <p>UUI episodes/day, mean change (95% CI): -2.1 (-2.4, -1.9) P = 0.0001</p> <p>UUI episodes/day, 3 mos, mean (range): 1.5 (0.0-24.0)</p> <p>UUI episodes/day, mean change (95% CI): -2.0 (-2.2, -1.7) P = 0.0001</p> <p>UUI episodes/day, median % change: -76</p> <p>Voids/day, 3 mos, mean (range): 8.8 (2.0-23.4)</p> <p>Voids/day, 3 months, mean change (95% CI): -2.6 (-2.8, -2.3) P = 0.0001</p> <p>Voids/day, 9 mos, mean (range): 8.9 (1.9-31.6)</p> <p>Voids/day, 9 months, mean change (95% CI): -2.5 (-2.9, -2.4) P = 0.0001</p> <p>Voids/day, 9 mos, median % change: -22</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: -</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: +</p> <p>Drop-out rates: +</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Appell, Abrams et al., 2001 (continued)				<p>Voided volume (mL), 3 mos, mean (range): 201 (33-444)</p> <p>Voided volume (mL), 3 mos, mean change (95% CI): 41 (36, 45) <i>P</i> = 0.0001</p> <p>Voided volume (mL), 9 mos, mean (range): 199 (34-514)</p> <p>Voided volume (mL), 9 mos, mean change (95% CI): 40 (35, 45) <i>P</i> = 0.0001</p> <p>Voided volume (mL), 9 mos, median % change: 22</p> <p>Improvement, 9 mos, %: 65</p> <p>Any adverse event, n (%): 652 (76)</p> <p>ANS, n (%): 268 (31)</p> <p>General, n (%): 219 (26)</p> <p>CNS/PNS, n (%): 82 (10)</p> <p>GI, n (%): 201 (24)</p> <p>Respiratory, n (%): 139 (16)</p> <p>Urinary, n (%): 165 (19)</p> <p>Dry mouth, n (%): 236 (28)</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Appell, Abrams et al., 2001 (continued)				<p>Mild dry mouth, %: 19</p> <p>Moderate dry mouth, %: 7</p> <p>Severe dry mouth, %: 2</p> <p>UTI, n (%): 106 (12)</p> <p>Headache, n (%): 57 (7)</p> <p>Constipation, n (%): 57 (7)</p> <p>Abdominal pain, n (%): 50 (6)</p> <p>Upper respiratory tract infection, n (%): 45 (5)</p> <p>Serious adverse events, n: 72</p> <p>Discontinued due to AE, n (%): 73 (9)</p> <p>Reduced dosage to 1 mg b.i.d., n (%): 108 (13)</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Appell, Sand et al., 2001</p> <p>Country and setting: US, 37 sites</p> <p>Enrollment period: March 2000 to October 2000</p> <p>Funding: ALZA Corp</p> <p>Author industry relationship disclosures: 12 of 12 ALZA (12) Pharmacia (9)</p>	<p>Design: RCT stratified by UUI severity (mild \leq 21/wk, or moderate $>$21/wk)</p> <p>Intervention: Oxybutynin ER vs Tolterodine</p> <p>Groups: G1: Oxybutynin ER 10 mg G2: Tolterodine 4mg (2mg b.i.d.)</p> <p>N at enrollment: G1: 185 G2: 193</p> <p>N at follow-up: G1: 160 G2: 172</p> <p>Women, %: G1: 82 G2: 85</p> <p>Age, mean \pm SD: G1: 58.6 \pm 13.4 G2: 59.6 \pm 13.2</p> <p>Race/ethnicity, %: White: G1: 87.6 G2: 86.0 Black: G1: 5.4 G2: 6.7 Hispanic: G1: 3.8 G2: 5.2 Asian: G1: 2.2 G2: 1.6 Other: G1: 1.1 G2: 0.5</p> <p>Follow-up: 12 week</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • 7-50 urge urinary incontinence episodes per week • \geq 10 voids/day • Majority of leakage urge-related <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Non-OAB cause for incontinence • Delivered baby $<$ 6 mos prior • PVR $>$150 mL • Significant medical comorbidities • Decreased hepatic function • Renal impairment • Myasthenia gravis • Gastroparesis • Hematuria • Narrow-angle glaucoma • POP to hymen • Known allergy • Pregnant or lactating 	<p>UUI episodes/week, mean \pm SD: G1: 25.5 \pm 14.6 G2: 24.6 \pm 15.1 $P = 0.36$</p> <p>Incontinence episodes/week, mean \pm SD: G1: 28.4 \pm 17.8 G2: 28.0 \pm 18.3 $P = 0.62$</p> <p>voids/week, mean \pm SD: G1: 92.9 \pm 23.0 G2: 91.8 \pm 20.0 $P = 0.77$</p> <p>Received antimuscarinic med before study, n (%): G1: 75 (40.5) G2: 73 (37.8)</p>	<p>UUI episodes/week, mean \pm SD (95% CI): G1: 6.1 \pm 9.7 (4.4, 7.3) G2: 7.8 \pm 11.1 (6.7, 9.5) $P = 0.03$</p> <p>Incontinence episodes/week, mean \pm SD (95% CI): G1: 7.1 \pm 12.0 (5.2, 8.6) G2: 9.3 \pm 13.4 (8.0, 11.3) $P = 0.02$</p> <p>voids/week, mean \pm SD (95% CI): G1: 67.1 \pm 22.1 (64.6, 70.0) G2: 71.5 \pm 20.5 (69.1, 74.2) $P = 0.02$</p> <p>Discontinuation due to AE, n (%): G1: 14 (7.6) G2: 15 (7.8) $P > 0.99$</p> <p>Dry mouth, n (%): G1: 52 (28.1) G2: 64 (33.2)</p> <p>Constipation, n (%): G1: 13 (7) G2: 12 (6.2)</p> <p>Impaired urination/retention, n (%): G1: 6 (3.2) G2: 6 (3.1) $P \geq 0.99$</p> <p>Blurred vision, n (%): G1: 4 (2.2) G2: 2 (1.0) $P = 0.44$</p> <p>Dizziness, n (%): G1: 9 (4.9) G2: 8 (4.1) $P = 0.81$</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: NA</p> <p>Masking: NA</p> <p>Pt selection criteria: +</p> <p>Loss to followup: -</p> <p>Drop-out rates: -</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: ++</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Appell, Sand et al., 2001 (continued)				<p>Somnolence, n (%): G1: 8 (4.3) G2: 3 (1.6) <i>P</i> = 0.13</p> <p>Asthenia, n (%): G1: 3 (1.6) G2: 7 (3.6) <i>P</i> = 0.34</p> <p>Insomnia, n (%): G1: 1 (0.5) G2: 3 (1.6) <i>P</i> = 0.62</p> <p>Headache, n (%): G1: 15 (8.1) G2: 17 (8.8) <i>P</i> = 0.86</p> <p>Dyspepsia, n (%): G1: 11 (5.9) G2: 10 (5.2) <i>P</i> = 0.82</p> <p>Nausea, n (%): G1: 6 (3.2) G2: 3 (1.6) <i>P</i> = 0.33</p> <p>Vomiting, n (%): G1: 3 (1.6) G2: 3 (1.6) <i>P</i> > 0.99</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Arruda et al., 2008</p> <p>Country and setting: Brazil, community</p> <p>Enrollment period: August 2001 to September 2005</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: None</p>	<p>Design: RCT</p> <p>Intervention: Oxybutynin vs. functional electrostimulation vs. pelvic floor training</p> <p>Groups: G1: Oxybutynin 5 mg b.i.d. G2: Ambulatory stimulation applied vaginally G3: Pelvic floor exercises with a therapist and at home</p> <p>N Screened: 81</p> <p>N at enrollment: G1: 22 G2: 21 G3: 21</p> <p>N at follow-up: G1: 22 G2: 21 G3: 21</p> <p>Age, range: 35-80</p> <p>Race/ethnicity: NR</p> <p>Women, %: 100</p> <p>Length of follow up: 12 weeks</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> Community-dwelling Dx of OAB and DO Capable of completing a bladder diary and performing a pelvic muscle floor contraction For those with MUI, urge was predominant <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Hx of psychiatric or neurologic illness Persistent UTI Inability to comply with regular follow-up visits Current pregnancy Postvoid residual volume > 100 ml Contraindications to anticholinergic therapy Cardiac pacemaker Type III SUI Uncontrolled metabolic conditions or indwelling catheterization Using medications including anticholinergic drugs, calcium antagonists, beta agonists, dopamine agonists, striated muscle relaxants or estrogens Any uterine prolapsed 	<p>UII episodes/week, mean ± SD: G1: 13.8 ± 11.6 G2: 13.5 ± 15.6 G3: 16.4 ± 17.2</p> <p>Nocturia episodes/week, mean ± SD: G1: 1.7 ± 1.5 G2: 1.9 ± 1.9 G3: 1.4 ± 1.2</p> <p>Pads/day, mean ± SD: G1: 2.6 ± 2.7 G2: 2.3 ± 2.4 G3: 2.7 ± 1.5</p> <p>Voids/day, mean ± SD: G1: 7.7 ± 2.6 G2: 8.6 ± 3.4 G3: 6.8 ± 2.2</p> <p>Residual volume mean mL ± SD: G1: 3.2 ± 6.3 G2: 1.0 ± 2.6 G3: 1.8 ± 3.3</p> <p>Volume, first desire to void, mean mL ± SD: G1: 117.7 ± 68.9 G2: 102.4 ± 51.1 G3: 86.7 ± 38.9</p> <p>Maximal cystometric capacity, mean mL ± SD: G1: 410.4 ± 194.1 G2: 350.0 ± 212.9 G3: 424.0 ± 149.0</p>	<p>UII episodes/week, mean ± SD: G1: 7.0 ± 10.6 G2: 7.9 ± 13.7 G3: 7.8 ± 15.3 G1/BL: P = 0.007 G2/BL: P = 0.039 G3/BL: P = 0.035</p> <p>Urgency resolved, n (%): G1: 14 (63.6) G2: 11 (52.4) G3: 12 (57.1) P = 0.754</p> <p>Satisfied, n (%): G1: 17 (77.3) G2: 11 (52.4) G3: 16 (76.2) P = 0.142</p> <p>Nocturia episodes/week, mean ± SD: G1: 0.9 ± 0.8 G2: 1.2 ± 1.3 G3: 1.0 ± 1.1 G1/BL: P = 0.003 G2/BL: P = 0.036 G3/BL: P = 0.086</p> <p>Pads/day, mean ± SD: G1: 0.9 ± 1.5 G2: 0.9 ± 1.7 G3: 0.8 ± 1.3 G1/BL: P < 0.001 G2/BL: P = 0.004 G3/BL: P < 0.001</p> <p>Voids/day, mean ± SD: G1: 6.4 ± 1.6 G2: 7.9 ± 2.3 G3: 7.1 ± 2.1 G1/BL: P = 0.014 G2/BL: P = 0.291 G3/BL: P = 0.441</p> <p>Residual volume, mean mL ± SD: G1: 4.7 ± 9.4 G2: 1.1 ± 2.5 G3: 2.1 ± 3.5 G1/BL: P = 0.425 G2/BL: P = 0.760 G3/BL: P = 0.297</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: -</p> <p>Method and blinding: -</p> <p>Pt selection criteria: +</p> <p>Loss to followup: +</p> <p>Drop-out rates: -</p> <p>Power calculation: +</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: +</p> <p>Length of followup: +</p> <p>Measurement methods: -</p> <p>Measurement reliability: -</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Arruda et al., 2008 (continued)			Involuntary detrusor contraction volume mean mL ± SD: G1: 189.5 ± 114.1 G2: 220.0 ± 127.2 G3: 239.3 ± 163.0	Volume first desire to void, mean mL ± SD: G1: 157.3 ± 63.8 G2: 123.8 ± 59.0 G3: 137.6 ± 76.7 G1/BL: <i>P</i> = 0.019 G2/BL: <i>P</i> = 0.091 G3/BL: <i>P</i> = 0.017	
			Involuntary detrusor contraction maximal pressure (mm H₂O): G1: 39.4 ± 26.1 G2: 43.7 ± 22.9 G3: 34.2 ± 19.8	Maximal cystometric capacity mean mL ± SD: G1: 517.3 ± 191.7 G2: 436.6 ± 178.7 G3: 489.0 ± 141.3 G1/BL: <i>P</i> = 0.001 G2/BL: <i>P</i> = 0.017 G3/BL: <i>P</i> = 0.113	
				Involuntary detrusor contraction volume (mL): G1: 188.6 ± 183.2 G2: 73.3 ± 112.4 G3: 114.3 ± 154.2 G1/BL: <i>P</i> = 0.986 G2/BL: <i>P</i> = 0.001 G3/BL: <i>P</i> = 0.044	
				Involuntary detrusor contraction maximal pressure, mm H₂O ± SD: G1: 19.6 ± 20.9 G2: 22.4 ± 30.1 G3: 17.2 ± 25.5 G1/BL: <i>P</i> < 0.001 G2/BL: <i>P</i> = 0.002 G3/BL: <i>P</i> = 0.027	
				Normal urodynamic evaluation, n (%): G1: 8 (36.4) G2: 12 (57.1) G3: 11 (52.4) <i>P</i> = 0.358	
				Persistent improvement at 1 year: G1: 10/17 G2: 4/11 G3: 9/16	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Arruda et al., 2008 (continued)				<p>Dry mouth, n (%): G1: 16 (72.7) G2: 0 G3: 0</p> <p>Difficulty voiding, n (%): G1: 2 (9.1) G2: 0 G3: 0</p> <p>Dizziness, n (%): G1: 1 (4.5) G2: 0 G3: 0</p> <p>Blurred vision, n (%): G1: 1 (4.5) G2: 0 G3: 0</p> <p>Constipation, n (%): G1: 1 (4.5) G2: 0 G3: 0</p> <p>Dry mouth, n (%): G1: 16 (72.7) G2: 0 G3: 0</p> <p>Difficulty voiding, n (%): G1: 2 (9.1) G2: 0 G3: 0</p> <p>Dizziness, n (%): G1: 1 (4.5) G2: 0 G3: 0</p> <p>Blurred vision, n (%): G1: 1 (4.5) G2: 0 G3: 0</p> <p>Constipation, n (%): G1: 1 (4.5) G2: 0 G3: 0</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Barkin et al., 2004</p> <p>Country and setting: Canada, Community</p> <p>Enrollment period: NR</p> <p>Funding: Purdue Pharma</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: Oxybutynin CR versus Oxybutynin IR</p> <p>3 week washout period followed by 2 weeks of dose titration plus 4 weeks stable dose period</p> <p>Groups: G1: Oxybutynin CR 15 mg daily Final titration dose, n (%): 5 mg: 3 (6) 10mg: 10 (19) ≥ 15mg: 40 (75) G2: Oxybutynin IR 5 mg t.i.d. Final titration dose, n (%): 5 mg: 6 (15) 10mg: 14 (34) ≥ 15mg: 21 (51)</p> <p>N at enrollment: G1: 65 G2: 60</p> <p>N at follow-up: G1: 53 G2: 41</p> <p>Women, n (%): G1: 48 (91) G2: 37 (90)</p> <p>Age, yrs ± SD: G1: 58.0 ± 12.4 G2: 60.6 ± 14.8</p> <p>Race/ethnicity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 18 • ≥ 7 UUI/week during washout period • ≥ 8 voids/day during washout period <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • PVR > 100mL • Unstable dosage of any drug with anticholinergic or diuretic/antidiuretic side effects • Allergy or previous life threatening side effects with anticholinergic/antispasmodic meds • SU1 • Conditions contraindicating anticholinergic therapy • Daily fluid intake > 3 liters • Hepatic/ renal disease • IC/ painful bladder syndrome • Urinary retention • Hematuria • UTI • Recurrent UTI • Indwelling catheter • Bladder training within 14 days of screening • Drug/ alcohol abuse • Untreated psychiatric disease effecting completion of bladder diary 	<p>Purdue Urgency Questionnaire, frequency, mean score ± SD: G1: 3.3 ± 1.1 G2: 3.2 ± 0.9</p> <p>Purdue Urgency Questionnaire, severity, mean score ± SD: G1: 3.8 ± 1.1 G2: 3.7 ± 1.1</p> <p>Incontinence episodes/week, mean ± SD: G1: 24.3 ± 19.0 G2: 23.0 ± 17.7</p> <p>Pads/day, mean: G1: 2.3 G2: 2.4</p> <p>Voids/day, mean ± SD: G1: 11.4 ± 2.9 G2: 11.0 ± 3.1</p> <p>Voided volume (mL), mean ± SD: G1: 177 ± 77 G2: 221 ± 137</p> <p>Fluid intake (cups/day), mean: G1: 6.8 G2: 6.6</p> <p>IIQ, mean score ± SD: G1: 2.6 ± 0.8 G2: 2.3 ± 0.8</p> <p>UDI, mean score ± SD: G1: 2.6 ± 0.6 G2: 2.5 ± 0.4</p>	<p>Purdue Urgency Questionnaire, frequency, mean ± SD: G1: 2.3 ± 1.2 G2: 1.9 ± 0.9 G1/G2: P = 0.116 G1/BL: P < 0.001 G2/BL: P < 0.001</p> <p>Purdue Urgency Questionnaire, severity, mean ± SD: G1: 2.3 ± 1.0 G2: 2.3 ± 1.3 G1/G2: P = 0.255 G1/BL: P < 0.001 G2/BL: P < 0.001</p> <p>Incontinence episodes/ week, mean ± SD: G1: 10.4 ± 18.8 G2: 6.1 ± 8.8 G1/G2: P = 0.404 G1/BL: P < 0.001 G2/BL: P < 0.001</p> <p>Pads/day, mean: G1: 1.7 G2: 1.9 G1/G2: P = NS G1/BL: P < 0.001 G2/BL: P < 0.001</p> <p>Voids/day, mean ± SD: G1: 9.6 ± 2.6 G2: 8.6 ± 2.8 G1/G2: P = 0.286 G1/BL: P < 0.001 G2/BL: P < 0.001</p> <p>Voided volume (mL), mean ± SD: G1: 202 ± 80 G2: 261 ± 119 G1/G2: P = 0.533 G1/BL: P = 0.064 G2/BL: P = 0.077</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: -</p> <p>Drop-out rates: -</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: +</p> <p>Length of followup: -</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Barkin et al., 2004 (continued)		<ul style="list-style-type: none"> • Untreated constipation • BOO • Pregnant or breast feeding • Women using reliable method of birth control 		<p>Fluid intake (cups/day), mean: G1: 7.0 G2: 7.7 G1/G2: <i>P</i> = NS G1/BL: <i>P</i> = NS G2/BL: <i>P</i> = 0.032</p> <p>IIQ score, mean ± SD: G1: 1.9 ± 1.7 G2: 1.6 ± 0.6 G1/G2: <i>P</i> = NS G1/BL: <i>P</i> < 0.001 G2/BL: <i>P</i> < 0.001</p> <p>UDI scores, mean ± SD: G1: 2.0 ± 0.6 G2: 1.8 ± 0.5 G1/G2: <i>P</i> = NS G1/BL: <i>P</i> < 0.001 G2/BL: <i>P</i> < 0.001</p> <p>Dry mouth, overall, n (%): G1: 44 (68) G2: 43 (72)</p> <p>Dry mouth, moderate/severe, n (%): G1: 25 (38) G2: 27 (45)</p> <p>Dry throat, n (%): G1: 23 (35) G2: 24 (40)</p> <p>Dry skin, n (%): G1: 11 (17) G2: 7 (12)</p> <p>Diarrhea, n (%): G1: 9 (14) G2: 3 (5)</p> <p>Headache, n (%): G1: 8 (12) G2: 13 (22)</p> <p>UTI, n (%): G1: 8 (12) G2: 11 (18)</p> <p>Dizziness, n (%): G1: 7 (11) G2: 11 (18)</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Barkin et al., 2004 (continued)				<p>Dyspepsia, n (%): G1: 7 (11) G2: 10 (17)</p> <p>Rhinitis, n (%): G1: 7 (11) G2: 9 (15)</p> <p>Abdominal pain, n (%): G1: 6 (9) G2: 6 (10)</p> <p>Asthenia, n (%): G1: 5 (18) G2: 9 (15)</p> <p>Constipation, n (%): G1: 5 (8) G2: 6 (10)</p> <p>Taste perversion, n (%): G1: 5 (8) G2: 7 (12)</p> <p>Cough increased, n (%): G1: 4 (6) G2: 8 (13)</p> <p>Dysphagia, n (%): G1: 4 (6) G2: 8 (13)</p> <p>Dry eyes, n (%): G1: 2 (3) G2: 9 (15)</p> <p>Nausea, n (%): G1: 3 (5) G2: 10 (17)</p> <p>Discontinued due to AEs, n (%): G1: 11 (17) G2: 12 (20)</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Bemelmans et al., 2000</p> <p>Country and setting: Netherlands, Primary Care</p> <p>Enrollment period: May 1995 to September 1997</p> <p>Funding: Byk Nederland (Distributor Oxybutynin)</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Prospective case series</p> <p>Intervention: Oxybutynin Run in study with all subjects enrolled in G1 with assessment at 2 wks treatment with diary followed by movement to G2 if effect was unsatisfactory or capable of improvement Tx x 4 more wks followed by voiding diary.</p> <p>Groups: G1: Responders to oxybutynin 2.5 mg t.i.d. G2: Non responders to oxybutynin 2.5 mg t.i.d. increased to xybutynin 5 mg t.i.d.</p> <p>N at enrollment: Total: 416 G1: 320 G2: 96</p> <p>N at follow-up: 350</p> <p>Women, N (%): 346 (83.2)</p> <p>Age, mean (range): 61.3 (16.4-91.5)</p> <p>Race/ethnicity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Complaint of involuntary urine loss • Score of ≥ 14 on symptom score questionnaire indicating UUI (determined by unvalidated questionnaire) <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Age ≤ 15 • Concurrent use of bladder medication • Urinary retention • Hematuria • UTI • SUI 	<p>Incontinence episodes/day, mean \pm SD: G1: 5.6 \pm 3.5 G2: 7.2 \pm 3.9</p> <p>Voids/day, mean \pm SD: G1: 9.6 \pm 3.5 G2: 11.1 \pm 3.9</p>	<p>Incontinence episodes/day, mean \pm SD: G1: 1.5 \pm 1.5 G2: 2.3 \pm 2.3 G1/BL: $P < 0.001$ G2/BL: $P < 0.001$</p> <p>Voids/day, mean \pm SD: G1: 5.8 \pm 2.3 G2: 6.8 \pm 2.9 G1/BL: $P < 0.001$ G2/BL: $P < 0.001$</p> <p>Side effects, n (%): G1: 83 (26) G2: 40 (42)</p> <p>Severity of side effects, %: Mild: 72 Severe: 21 Extremely severe: 4</p> <p>Side effects, %: Dry mouth: 23 Blurred vision: 11 Nausea/ vomiting: 6 Dizziness: 6 Headache: 6 Anorexia: 3 Constipation: 3 Drowsiness: 3 Tachycardia: 1 Urinary retention: <1 Allergic Reaction: <1</p> <p>Discontinued due to AEs, n (%): 66 (16)</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: NA</p> <p>Masking: NA</p> <p>Pt selection criteria: +</p> <p>Loss to followup: ++</p> <p>Drop-out rates: ++</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: +</p> <p>Length of followup: -</p> <p>Measurement methods: +</p> <p>Measurement reliability: -</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Boone et al., 2002</p> <p>Country and setting: US, Multicenter (86.0% of physicians in private-practice)</p> <p>Enrollment period: NR</p> <p>Funding: Alza Pharmaceuticals</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Prospective observational</p> <p>Intervention: Non-specific/non-specified pharmacologic therapy</p> <p>Groups: G1: persons who started drug therapy (any) at baseline and continued therapy for ≥ 3 mos G2: persons who started drug therapy (any) at baseline but did not continue for 3 mos G3: persons who did not start drug therapy at baseline but were receiving it at 3 mos G4: persons who did not start drug therapy at baseline and were not receiving therapy at 3 mos G5: persons who were on drugs at 3 mos and 6 mos G6: persons who were on drugs at 3 mos and not at 6 mos G7: persons who were on not drugs at 3 mos and on drug at 6 mos G8: persons on no drugs at 3 mos and 6 mos</p> <p>N at enrollment: Total: 213</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 18 • Fluent in English • US resident • Urgency and ≥ 1 incontinence episodes/day • ≥ 8 voids/day • ≥ 3 nocturia episodes/day <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Neurogenic condition causing incontinence • Persistent or recurrent UTI • History of interstitial cystitis • Bladder cancer • Previous bladder augmentation surgery • Spinal injury • Alzheimer's disease • Other dementia • Obstruction of urinary tract • Primary diagnosis of SUI • Enrolled in another study involving medication • Pharmacologic treatment for OAB during previous year 	<p>Incontinence episodes/day, mean ± SD: Total: 2.2 ± 3.4 G1-G2: 2.34 G3-G4: 2.09</p> <p>Duration of symptoms (yrs), mean ± SD: Total: 8.1 ± 10.3 G1-G2: 8.5 ± 12.0 G3-G4: 7.8 ± 9.1</p> <p>Symptom severity, n (%): Mild: G1-G2: 11 (22.0) G3-G4: 18 (25) Moderate G1-G2: 26 (52.0) G3-G4: 43 (59.7) Severe G1-G2: 13 (26.0) G3-G4: 11 (15.3)</p> <p>Symptom worsening, n (%): G1-G2: 28 (56.0) G3-G4: 37 (51.4)</p> <p>U-UDI score, mean ± SD: G1-G2: 1.99 ± 0.79 G3-G4: 1.99 ± 0.88</p> <p>Non-drug interventions, mean ± SD: G1-G2: 0.40 ± 0.49 G3-G4: 0.47 ± 0.65</p> <p>Pads/week, mean ± SD: G1-G2: 0.30 ± 1.34 G3-G4: 0.17 ± 0.56</p>	<p>Improvement in OAB sx, 3 mos, n (%): G1: 30 (76.9) G2: 7 (63.6) G3: 10 (52.6) G4: 19 (35.8) G1-G2/G3-G4: <i>P</i> < 0.001</p> <p>Moderate to great improvement in OAB sx, n (%): G1-G2: 26 (52.0) G3-G4: 21 (29.2) G1-G2/G3-G4: <i>P</i> = 0.014</p> <p>Pads/week, 3 mos, mean % change: G1-G2: -25.4 G3-G4: 135.2</p> <p>Physician visits, mean: G1: 0.15 G2: 0.64 G3: 1.16 G4: 0.26 G1-G2/G3-G4: <i>P</i> = 0.096</p> <p>Patients prescribed nondrug intervention(s), 3 mos, n (%): G1-G2: 5 (10.0) G3-G4: 16 (22.2) G1-G2/G3-G4: <i>P</i> = 0.092</p> <p>Patients with improvement in OAB sx, 4-6 mos, n (%): G5: 35 (81.4) G6: 1 (12.5) G7: 2 (28.6) G8: 12 (28.6)</p> <p>Pads/week, 4-6 mo, % change: G5: 23 G6: 25 G7: 0 G8: 16</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: NA</p> <p>Masking: NA</p> <p>Pt selection criteria: +</p> <p>Loss to followup: -</p> <p>Drop-out rates: -</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: +</p> <p>Baseline OAB status: NR</p> <p>Baseline characteristics: -</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: -</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Boone et al., 2002 (continued)	N at follow-up:			Physician visits, 4-6 mo, mean: G5: 0.16 G6: 0.38 G7: 0.43 G8: 0.05	
	G1: 39			Predictors of OAB symptom improvement, OR (95 %CI):	
	G2: 11			Rx at baseline:	
	G3: 19			4.3 (1.8-9.9)	
	G4: 53			Incontinence	
	G5: 43			episodes/day:	
	G6: 8			3.2 (1.2-8.4)	
	G7: 7				
	G8: 42				
	Women, n (%):				
	G1-G2: 40 (80)				
	G3-G4: 64 (88.9)				
	Age, mean ± SD:				
	G1-G2: 62.3 ±				
	11.9				
	G3-G4: 60.3 ±				
	16.4				
	Race/ethnicity, n				
	(%):				
	White: 94 (77.7)				
	Black: 17 (14.0)				
	Hispanic: 7 (5.8)				
	Asian: 2 (1.7)				
	Other: 1 (0.8)				

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Burgio et al., 1998</p> <p>Country and setting: US, academic health center outpatient geriatric medicine clinic</p> <p>Enrollment period: July 1989 to August 1995</p> <p>Funding: National Institutes on Aging</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT, placebo controlled</p> <p>Computer-generated random numbers using a block size of 6, w/ prior stratification by type and severity of incontinence</p> <p>Intervention: Biofeedback-assisted behavioral vs. drug treatment (oxybutynin chloride; possible range of doses 2.5 mg/d-5.0 mg t.i.d.) vs. placebo</p> <p>All patients had 4 visits over an 8-week period. Patients in G1 had biofeedback added to behavioral training in absence of 50% improvement by session 3.</p> <p>Groups: G1: Behavioral ± biofeedback G2: Pharmacologic G3: Placebo</p> <p>N at enrollment: 468 screened 271 not eligible 197 randomized G1: 65 G2: 67 G3: 65</p> <p>N at follow-up: G1: 61 G2: 55 G3: 53</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> Community-dwelling women at least age 55 Ambulatory At least 2 urge accidents per week by baseline bladder diary Urge incontinence as predominant pattern Urodynamic evidence of bladder dysfunction <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Continual leakage Postvoid residual urine volume >200mL Uterine prolapse past the introitus Narrow-angle glaucoma Unstable angina Decompensated congestive heart failure Hx of malignant arrhythmias MMSE <20 (Dementia) 	<p>Duration of symptoms, mean yrs ± SD: G1: 9.4 ± 10.8 G2: 9.8 ± 11.9 G3: 12.7 ± 15.9</p> <p>Restricted activity, (%): G1: 30.8 G2: 32.8 G3: 38.5</p> <p>UUI only, %: G1: 49.2 G2: 49.3 G3: 47.7</p> <p>Accidents per week, mean ± SD: G1: 15.8 ± 14.5 G2: 15.9 ± 14.1 G3: 15.4 ± 13.4 P = .98</p> <p>Severity classification, %: Mild (<5 accidents/wk) G1: 18.5 G2: 17.9 G3: 18.5 Moderate (5-10 accidents/wk) G1: 29.2 G2: 29.9 G3: 27.7 Severe (>10 accidents/wk) G1: 52.3 G2: 52.2 G3: 53.8</p> <p>Previous treatment with surgery, %: G1: 20.0 G2: 26.9 G3: 29.2</p>	<p>Accidents per week, mean ± SD: G1: 2.8 ± 4.7 G2: 5.7 ± 9.8 G3: 8.2 ± 11.6 P = .005</p> <p>Percent reduction, mean ± SD: G1: 80.7 ± 24.8 G2: 68.5 ± 37.2 G3: 39.4 ± 80.0 P < 0.001</p> <p>Percent reduction, range: G1: -0.9 – 100 G2: -85.7 – 100 G3: -400.0 - 100</p> <p>Patient perceptions of progress in treatment, %: Much better G1: 74.1 G2: 50.9 G3: 226.9 Better: G1: 25.9 G2: 30.9 G3: 38.5 About the same: G1: 0.0 G2: 16.4 G3: 28.8 Worse: G1: 0.0 G2: 1.8 G3: 5.8</p> <p>Estimate of % improvement, mean ± SD: G1: 81.6 ± 18.6 G2: 66.4 ± 35.4 G3: 45.1 ± 36.6</p> <p>Having fewer accidents, %: G1: 100.0 G2: 87.3 G3: 67.3</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: +</p> <p>Drop-out rates: +</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Burgio et al., 1998 (continued)	<p>Women, %: 100</p> <p>Age, yrs ± SD: G1: 67.3 ± 7.6 G2: 68.2 ± 7.5 G3: 67.6 ± 7.6</p> <p>Race/ethnicity,: NR</p> <p>Parity mean ± SD: G1: 2.8 ± 2.0 G2: 2.1 ± 1.4 G3: 2.7 ± 1.8 P < 0.05</p>		<p>Previous treatment with medication, %: G1: 27.7 G2: 35.8 G3: 30.8</p> <p>Using estrogen, %: G1: 32.3 G2: 38.8 G3: 35.4</p> <p>Using diuretics, %: G1: 20.0 G2: 14.9 G3: 12.3</p>	<p>Accidents are smaller, %: G1: 87.3 G2: 78.8 G3: 54.0</p> <p>Able to wear less protection, %: G1: 76.0 G2: 56.0 G3: 334.1</p> <p>Comfortable enough with treatment to continue indefinitely, %: G1: 96.5 G2: 54.7 G3: 43.1</p> <p>Patient satisfaction with progress, %: Completely satisfied: G1: 77.6 G2: 54.7 G3: 43.1 Somewhat satisfied: G1: 22.4 G2: 40.0 G3: 34.0 Not at all satisfied: G1: 0.0 G2: 10.9 G3: 38.0</p> <p>Wish to receive another form of treatment, %: G1: 14.0 G2: 75.5 G3: 75.5</p> <p><i>P</i> < 0.001 for all comparisons</p> <p>Adverse effects, p compared to placebo G3:</p> <p>Dry mouth, %: G1: 34.9 G2: 96.9 G3: 54.8 <i>P</i> < 0.001</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Burgio et al., 1998 (continued)				<p>Inability to void, %: G1: 6.3 G2: 21.5 G3: 3.2 <i>P</i> = 0.002</p> <p>Constipation, %: G1: 22.2 G2: 38.5 G3: 37.1 <i>P</i> = 0.10</p> <p>Blurred vision, %: G1: 9.5 G2: 15.4 G3: 9.7 <i>P</i> = 0.50</p> <p>Confusion, %: G1: 6.3 G2: 7.7 G3: 11.3 <i>P</i> = 0.59</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Burgio et al. 2000 [See Burgio et al., 1998]</p> <p>Country and setting: US, University based</p> <p>Enrollment period: two weeks after completion of Burgio et al. 1998</p> <p>Funding: National Institute on Aging</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Modified crossover of RCT</p> <p>Intervention: Participants whose treatment was not completely successful were given the opportunity to switch or use combined treatment; further reductions in incontinence were measured.</p> <p>Groups: <u>Treatment</u> <u>Changes:</u> G1: Previous oxybutynin to behavioral modification alone G2: Previous behavior alone to 2.5 mg oxybutynin t.i.d. + behavioral therapy G3: Previous oxybutynin alone to 2.5 mg oxybutynin t.i.d. + behavioral therapy G4: Placebo to behavioral G5: Placebo to oxybutynin</p> <p>N at enrollment G1: 19 G2: 8 G3: 27 G4: 34 G5: 10</p> <p>N at follow-up: G1: 18 G2: 8 G3: 26 G4: NR G5: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> Community-dwelling women at least age 55 Ambulatory At least 2 urge accidents per week by baseline bladder diary Urge incontinence as predominant pattern Urodynamic evidence of bladder dysfunction Not completely dry or satisfied with previous, 8-wk treatment <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Continual leakage Postvoid residual urine volume >200mL Uterine prolapse past the introitus Narrow-angle glaucoma Unstable angina Decompensated congestive heart failure Hx of malignant arrhythmias MMSE <20 (Dementia) 	<p>Incontinence, after previous study (at baseline), mean % reduction: G1: 59.1 G2: 57.5 G3: 72.7 G4: 22.9 G5: 44.8</p>	<p>Incontinence, final mean % reduction: G1: 77.1, $P = 0.109$ G2: 88.5, $P = 0.034$ G3: 84.3, $P = 0.001$ G4: 63.9, $P = 0.002$ G5: 76.5, $P = 0.012$</p> <p>Note: 29.2 of G3 declined to continue with drug therapy once they received behavioral modification.</p> <p>Numbers were too low to compare across groups.</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: +</p> <p>Drop-out rates: +</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Burgio et al., 2001</p> <p>[See Burgio et al., 1998]</p> <p>Country and setting: US, University based</p> <p>Enrollment period: [See Burgio et al., 1998]</p> <p>Funding: NIH</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: Psychological distress associated with pharmacologic treatment vs. behavioral vs. placebo</p> <p>Groups: G1: Behavioral training with biofeedback, 4 visits G2: Oxybutynin 2.5 mg po t.i.d., up to a max of 5 mg t.i.d. G3: Placebo</p> <p>N at enrollment: 197 women</p> <p>N treated: 169</p> <p>N at follow-up: 155 (completed both pre and post-treatment psychological assessment)</p> <p>Age yrs, mean ± SD: 67.5 ± 7.2</p> <p>Women, %: 100</p> <p>Race/ethnicity, %: White: 97 African American: 3</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • ≥ 55 yrs old • Ambulatory • UUI ≥2x/wk (2 wk bladder diary), persisting x 3 mo • Predominant UUI (vs. other forms incontinence) • Urodynamic evidence of bladder dysfunction (DI or maximal capacity ≤350 mL) <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Contraindication to oxybutynin or behavioral treatment • Continual leakage • PVR > 200 mL • Uterine prolapse beyond the introitus • Decompensated CHF • Hx of malignant arrhythmias • Impaired mental status (MMSE <20) 	<p>SCL-90-R scores ± SD:</p> <p>Somatization G1: 56.0 (10.6) G2: 51.4 (10.8) G3: 52.4 (11.1)</p> <p>Obsessive-Compulsive: G1: 56.5 (10.7) G2: 56.6 (11.4) G3: 57.7 (10.0)</p> <p>Interpersonal Sensitivity G1: 53.8 (11.0) G2: 51.4 (11.9) G3: 50.4 (12.0)</p> <p>Depression G1: 54.7 (10.0) G2: 52.5 (9.7) G3: 51.0 (11.9)</p> <p>Anxiety G1: 48.7 (13.9) G2: 46.8 (12.0) G3: 47.2 (12.8)</p> <p>Hostility G1: 49.3 (10.7) G2: 45.9 (10.1) G3: 48.3 (10.4)</p> <p>Phobia G1: 47.5 (10.2) G2: 46.7 (10.3) G3: 45.7 (8.5)</p> <p>Paranoid Ideation G1: 48.6 (12.4) G2: 49.3 (11.1) G3: 46.1 (12.8)</p> <p>Psychoticism G1: 54.1 (10.7) G2: 52.1 (10.3) G3: 50.5 (11.2)</p> <p>Global Severity G1: 54.2 (11.1) G2: 52.5 (10.3) G3: 52.4 (11.2)</p>	<p>Reduction in incontinence episodes: G1: 83.3% G2: 74.4% G3: 41.4% P<0.001</p> <p>SCL-90-R scores ± SD:</p> <p>Somatization G1: 51.8 (11.4) G2: 51.2 (9.8) G3: 49.8 (13.0)</p> <p>Obsessive-Compulsive: G1: 53.8 (13.9) G2: 53.9 (10.9) G3: 55.4 (11.0)</p> <p>Interpersonal Sensitivity G1: 49.5 (12.0) G2: 48.9 (11.2) G3: 49.2 (11.3)</p> <p>Depression G1: 51.5 (11.5) G2: 50.6 (10.7) G3: 51.4 (11.2)</p> <p>Anxiety G1: 46.1 (14.6) G2: 44.5 (12.3) G3: 45.8 (12.9)</p> <p>Hostility G1: 44.9 (10.8) G2: 44.6 (10.5) G3: 47.3 (11.2)</p> <p>Phobia G1: 47.1 (11.2) G2: 45.0 (8.3) G3: 45.1 (8.5)</p> <p>Paranoid Ideation G1: 45.8 (10.9) G2: 47.2 (11.6) G3: 47.2 (12.0)</p> <p>Psychoticism G1: 49.2 (11.7) G2: 50.4 (9.7) G3: 49.6 (10.3)</p> <p>Global Severity G1: 50.8 (12.8) G2: 50.4 (10.0) G3: 51.4 (10.9)</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Methods and blinding: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: +</p> <p>Drop-out rates: +</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Burgio et al., 2001 (continued)			<p>155 participants were compared to the 42 who did not complete intervention and psychological assessment, higher scores (greater distress) on 6 of 10 SCL-90-R scales (somatization, obsessive/compulsive, depression, hostility, paranoid ideation, global severity index), all p values <0.05</p> <p>Normal range, score 0-63 >75% in normal range (including dropouts) on 9 of 10 scales</p> <p>Highest impairment rate: 33% scored abnormal for obsessive-compulsive</p>	<p>Correlations between reduction of incontinence and changes in psychological symptoms</p> <p>Somatization G1: 0.28* G2: -0.09 G3: 0.17</p> <p>Obsessive-Compulsive G1: 0.01 G2: -0.14 G3: 0.02</p> <p>Interpersonal Sensitivity G1: -0.09 G2: 0.04 G3: 0.13</p> <p>Depression G1:-0.04 G2: 0.03 G3: 0.07</p> <p>Anxiety G1: -0.10 G2: -0.01 G3: 0.34*</p> <p>Hostility G1: -0.10 G2: 0.09 G3: 0.11</p> <p>Phobic anxiety G1:-0.21 G2: -0.17 G3: 0.02</p> <p>Paranoid Ideation G1: 0.14 G2: -0.04 G3: -0.06</p> <p>Psychoticism G1: -0.01 G2: -0.01 G3: 0.13 ***P=0.001</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Burgio et al., 2001 (continued)				Global Severity Index G1: 0.01 G2: 0.06 G3: 0.45*** *p<0.05	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Burgio et al., 2008</p> <p>Country and setting: US, Academic medical centers</p> <p>Enrollment period: July 2004 to January 2006</p> <p>Funding: NIH Pfizer</p> <p>Author industry relationship disclosures: 20 of 29 Allergan (3) Alza (1) Astellas Pharma (7) Bionovo (1) Bristol-Meyers Squibb (1) Dynogen (1) Elan (1) Ethicon (2) GSK (4) Johnson & Johnson (3) Lilly (7) Medtronic (1) Merck (1) Novartis (6) Ortho-McNeil (3) Pfizer (>10) Procter & Gamble (3) Q-Med (1) Renesa (1) Sanofi (1) Solace (1) Watson (1)</p>	<p>Design: RCT</p> <p>Intervention: Pharmacologic vs. pharmacologic plus behavioral interventions for 10 wks (Stage 1), followed by no drug therapy (Stage 2) with assessments at 10 wks and 8 mos*</p> <p>Groups: G1: Tolterodine tartrate (ER capsules), 4 mg/d plus behavioral training: pelvic floor muscle control and exercises, behavioral strategies to diminish urgency, suppress bladder contractions and prevent both stress and urge incontinence; delayed voiding; fluid management; handout with hints G2: Tolterodine tartrate (ER capsules), 4 mg/d</p> <p>N at enrollment: 4043 screened 2612 not eligible 870 declined 561 consented 254 excluded 307 randomized G1: 153 G2: 154</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Women • Community-dwelling • UUI only, or urge-predominant • ≥ 7 episodes of incontinence in a 7-day bladder diary • Persistent incontinence for at least 3 mos • No current use of antimuscarinics or other medications that could affect UI • No evidence of neurogenic etiology <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Age < 21 • Pregnant, planning a pregnancy in next 8 mos, or not using birth control • < 6 mos postpartum or termination after 20 wks gestation • Unable to contract pelvic floor muscles during evaluation • Behavioral therapy of > 2 mos in past 2 yrs • Continual leakage or always damp • Hypersensitive to study drug • Systemic disease that affects bladder function 	<p>UUI, 7-13 episodes/week, n (%): G1: 2 (1.3) G2: 2 (1.3)</p> <p>UUI, ≥ 14 episodes/week, n (%): G1: 2 (1.3) G2: 4 (2.6)</p> <p>MUI, 7-13 episodes/week, n (%): G1: 46 (29.9) G2: 46 (30.1)</p> <p>MUI, ≥ 14 episodes/week, n (%): G1: 104 (67.5) G2: 101 (66.0)</p> <p>Adjusted incontinence episodes/week, mean: G1: 23.1 G2: 23.2</p> <p>Previous non-surgical treatment for incontinence, n (%): G1: 19 (12) G2: 22 (14)</p>	<p>Success, n (%): G1: 43 (28) G2: 41 (27)</p> <p>Failure, n (%): G1: 75 (49) G2: 78 (51)</p> <p>Success rate, 8 months, lifetable analysis, % (95% CI): G1: 41 (32, 50) G2: 41 (33, 50) G1/G2: 0 (-12, 12)</p> <p>Success rate, 8 months, complete cases, % (95% CI): G1: 36 (27, 45) G2: 34 (25, 43) G1/G2: 2 (-10, 14)</p> <p>Success rate, 8 mos, assuming missings were failures, % (95% CI): G1: 28 (21, 35) G2: 27 (20, 34) G1/G2: 1 (-9, 11)</p> <p>Adjusted incontinence episodes/week, after Stage 1, mean: G1: 2.7 G2: 4.7</p> <p>Adjusted incontinence episodes/week, after Stage 1, mean change: G1: -20.4 G2: -18.5 G1/G2: -1.9 (-5.9, 2.0)</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Masking: -</p> <p>Pt selection criteria: +</p> <p>Loss to followup: -</p> <p>Drop-out rates: ++</p> <p>Power calculation: +</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: ++</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Burgio et al., 2008 (continued)	<p>N at follow-up: G1: 153 Completed treatment: 107 Outcome known at 8 mos: 119 G2: 154 Completed treatment: 101 Outcome known at 8 mos: 118</p> <p>Age, mean ± SD: G1: 55.8 ± 14.2 G2: 58.0 ± 13.5</p> <p>Women, %: 100</p> <p>Race/ethnicity, n (%): Hispanic: G1: 13 (9) G2: 17 (11) NH White: G1: 105 (69) G2: 85 (56) NH Black: G1: 22 (14) G2: 35 (23) Other: G1: 13 (9) G2: 15 (10)</p> <p>BMI, kg/m² ± SD: G1: 33.2 ± 9.5 G2: 32.3 ± 7.6</p>	<ul style="list-style-type: none"> • Currently using catheter to empty bladder • Postvoid residual volume > 150mL • Treatment for prolapse with pessary < 3 mos • Incontinence, vaginal, bladder or prolapse surgery in the past 6 mos • Urethral diverticulum • Previous augmentation cystoplasty or artificial sphincter • Neuromodulation for pelvic indications • Using anticholinergic agents, cholinergic agonists, tricyclic antidepressants, or duloxetine in ≤ 4 wks • Using diuretics with dosage change past 3 mos • Uncontrolled medical problem • History of bladder or pelvic cancer or pelvic radiation therapy • Glaucoma • Gastric retention • Non-ambulatory • Participation in other intervention trial that might influence outcome 		<p>Achieved 70% reduction in incontinence episodes, per bladder diary, 10 weeks (%): G1: 69 G2: 58 G1/G2: 11 (-0.3, 22.1)</p> <p>Totally dry, per bladder diary, 10 weeks (%): G1: 21 G2: 17</p> <p>Voids/day, mean change: G1: 0.5 G2: -0.4 G1/G2: 0.9 (0.3, 1.5)</p> <p>Symptom Distress Scores: G1/G2: <i>P</i> < 0.0001</p> <p>Symptom Bother Scores (OAB-q), Stage 1, mean change: G1: -36.7 G2: -30.4 G1/G2: <i>P</i> < 0.0001</p> <p>Symptom Bother Scores (OAB-q), Stage 2, mean change: G1: -30.9 G2: -20.4 G1/G2: <i>P</i> < 0.0001</p> <p>Patient completely satisfied, Stage 1, %: G1: 53 G2: 40 G1/G2: 13 (1, 25)</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Burgio et al., 2008 (continued)				<p>Patient completely satisfied, 8 months, %: G1: 33 G2: 20 G1/G2: 13 (2, 24)</p> <p>Patient better or much better, Stage 1, %: G1: 90 G2: 77 G1/G2: 13 (4, 22)</p> <p>Patient better or much better, 8 months, %: G1: 69 G2: 43 G1/G2: 26 (14 , 38)</p> <p>Persistence in perceived improvement, 8 mos, women with improvement at Stage 1: G1: 72 G2: 54 G1/G2: 17 (4, 30)</p> <p>Harms: G1: 3 participants 1: blurred vision, syncope, night sweats, stomach cramping and weakness 2: 2 episodes of small-bowel obstruction and an allergic reaction (pruritus and rash) 3: tachycardia during stage 2 G2: 3 participants 1: small bowel obstruction 2: peripheral edema 3: renal cell carcinoma diagnosis during stage 2</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Capo et al. 2008</p> <p>Country and setting: US, 207 centers</p> <p>Enrollment period: NR</p> <p>Funding: Astellas Pharma US, Inc., and GlaxoSmithKline</p> <p>Author industry relationship disclosures: 5 of 5 Astellas (4) Boehringer Mannheim Pharmaceuticals (1) Cephalon Inc. (1) GlaxoSmithKline (3) Reliant Pharmaceuticals (1) Schering Plough Corporation (1) Abbott (1) Akros (1) AstraZenica (1) Johnson & Johnson (1) Lilly (1) Manking (1) Merck (1) Novartis (1) NovoNordisk (1) Pfizer Inc. (1) Solvay (1) Takeda & TAP (1) Pri Med (1) Lucente (1)</p>	<p>Design: RCT</p> <p>Intervention: 12-wk treatment with 5 mg solifenacin succinate. Dosage could be increased at wk 4 and wk 8 to 10 mg, maintained or decreased in response to pt perceived efficacy and tolerability</p> <p>Groups: G1: Hispanic subgroup of G2 G2: full study population including G1</p> <p>N at enrollment: G1: 94 G2: 2205</p> <p>N at follow-up, range: G1: 47-80 G2: 1254-2005</p> <p>Age, yrs ± SD: G1: 51.8 ± 12.8 G2: 59.7 ± 14.4</p> <p>Weight in lbs, mean ± SD: G1: 171.1±36.4 G2: 182.2±47.0</p> <p>Race/ethnicity, n (%): Hispanic: G1: 94 (100) G2: 94 (4.3) White: G2: 1761 (79.9) Black: G2: 274 (12.4) Asian: G2: 59 (2.7) Other: G2: 17 (0.8)</p> <p>Women, N (%): G1: 74 (78.7) G2: 1813 (82.2)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • ≥ 18 yrs of age • OAB symptoms ≥3 mos • Toilet without difficulty • Washout of other OAB meds ≥ 7 days • Non-drug treatment of OAB if established ≥ 4 wks prior to study and continued <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • SUI • Stress predominant MUI • UTI or chronic inflammation clinically significant outflow • Obstruction due to BPH • Narrow-angle glaucoma • Urinary or gastric retention • Severe renal or hepatic impairment • Chronic severe constipation or gastrointestinal obstructive disease • Bladder cancer • Women not using a reliable contraceptive • Hypersensitivity to the study medication, any of its components, or to anticholinergic medication 	<p>Urge incontinence, n (%): G1: 63 (67.0) G2: 1586 (71.9)</p> <p>Urinary urgency, n %: G1: 88 (93.6) G2: 2007 (91.0)</p> <p>Frequency, n (%): G1: 86 (91.5) G2: 1969 (89.3)</p> <p>Nocturia, n (%): G1: 79 (84.0) G2: 1792 (81.3)</p> <p>Patient perception of bladder condition, mean score: G1: 4.0 G2: 4.0</p> <p>Most bothersome OAB symptom-frequency, n %: G1: 38 (40.4) G2: 619 (28.1)</p> <p>Most bothersome OAB symptom-urinary urgency, n %: G1: 22 (23.4) G2: 508 (23.0)</p> <p>Most bothersome OAB symptom-urge incontinence, n %: G1: 17 (18.1) G2: 602 (27.3)</p> <p>Most bothersome OAB symptom-nocturia, n %: G1: 12 (12.8) G2: 337 (15.3)</p>	<p>Increased dosage 5 mg/day, n (%)-Wk 4: G1: 46 (48.9)</p> <p>Increased dosage 10 mg/day, n (%)-Wk 4: G1: 37 (39.4)</p> <p>Discontinued, n Wk 4: G1: 1</p> <p>Increased dosage 5 mg/day, n (%)-Wk 8: G1: 31 (33.0)</p> <p>Increased dosage 10 mg/day, n (%)-Wk 8: G1: 39 (41.5)</p> <p>Discontinued-Wk 8, n: 1</p> <p>Returned to 5 mg/day-Wk 8, n: 5</p> <p>Increased dosage to 10 mg/day-Wk 8, n: 12</p> <p>Patient Perception of Bladder, mean Wk 4: G1: 3.3* G2: 3.3*</p> <p>Patient Perception of Bladder, mean Wk 8: G1: 2.8* G2: 2.8*</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: NA</p> <p>Masking: NA</p> <p>Pt selection criteria: +</p> <p>Loss to followup: -</p> <p>Drop-out rates: -</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: +</p> <p>Baseline OAB status: NR</p> <p>Baseline characteristics: -</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Capo et al., 2008 (continued)	Parity : NR		Most bothersome OAB symptom- none specified, n %: G1: 5 (5.3) G2: 139 (6.3)	Patient Perception of Bladder, mean Wk 12: G1: 3.0* G2: 2.9* * <i>P</i> <0.001 vs.baseline Symptom bother from VAS, Change from Baseline, mean scores (mm), (95% CI) p-value: Urinary urgency: G1: -39.2 (-45.9, -32.5) <i>P</i> < 0.001 G2: -39.5 (-41.0, -38.1) <i>P</i> < 0.001 Urge Incontinence: G1: -37.7 (-46.0, -29.4) <i>P</i> < 0.001 G2: -40.1 (-41.8, -38.4) <i>P</i> < 0.001 Frequency: G1: -40.0 (-46.9, -33.2) <i>P</i> < 0.001 G2: -41.8 (-43.3, -40.3) <i>P</i> < 0.0001 Nocturia: G1: -43.2 (-50.4, -36.1) <i>P</i> < 0.001 G2: -36.9 (-38.4, -35.4) <i>p</i> < 0.001 OAB-q subscales, change from baseline, mean (95% CI) p-value:	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Capo et al., 2008 (continued)				<p>Symptom bother: G1: -29.1 (-34.8, -23.4) <i>P</i> < 0.001, G2: -29.6 (-30.7, -28.6) <i>P</i> < 0.001</p> <p>Coping: G1: 24.0 (18.0, 30.1) <i>P</i> < 0.001 G2: 27.4 (26.2, 28.5) <i>P</i> < 0.001</p> <p>Concern: G1: 25.8 (19.5, 32.2) <i>P</i> < 0.001 G2: 29.6 (28.4, 30.8) <i>P</i> < 0.001</p> <p>Sleep: G1: 25.7 (19.4, 32.0) <i>P</i> < 0.001 G2: 27.3 (26.1, 28.5) <i>P</i> < 0.001</p> <p>Social Interaction G1: 15.9 (11.5, 20.3) <i>P</i> < 0.001 G2: 14.7 (13.7, 15.6) <i>P</i> < 0.001</p> <p>Overall Health-related QoL: G1: 23.2 (18.0, 28.5) <i>P</i> < 0.001 G2: 25.4 (24.4, 26.4) <i>P</i> < 0.001</p> <p>Any Adverse Event, n (%): G1: 48 (51.4) G2: 1321 (59.4)</p> <p>Dry mouth, n (%): G1: 17 (18.1) G2: 477 (21.4)</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Capo et al., 2008 (continued)				<p>Constipation, n (%): G1: 12 (12.8) G2: 295 (13.3)</p> <p>Headache, n (%): G1: 3 (3.2) G2: 76 (3.4)</p> <p>Dizziness, n (%): G1: 2 (2.1) G2: 27 (1.2)</p> <p>Blurred vision, n (%): G1: 4 (4.3) G2: 57 (2.6)</p> <p>Palpitations, n (%): G1: 2 (2.1) G2: 6 (0.3)</p> <p>Upper respiratory tract infection, n (%): G1: 3 (3.2) G2: 68 (3.1)</p> <p>UTI, n (%): G1: 3 (3.2) G2: 76 (3.4)</p> <p>Bronchitis, n (%): G1: 3 (3.2) G2: 31 (1.4)</p> <p>Nasopharyngitis, n (%): G1: 2 (2.1) G2: 50 (2.3)</p> <p>Influenza, n (%): G1: 2 (2.1) G2: 11 (0.5)</p> <p>Insomnia, n (%): G1: 2 (2.1) G2: 17 (0.8)</p> <p>Depression, n (%): G1: 2 (2.1) G2: 19 (0.9)</p> <p>Hypertension, n (%): G1: 2 (2.1) G2: 16 (0.7)</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Cardozo et al., 1993</p> <p>Country and setting: UK, Denmark, Germany, Netherlands, 4 centers</p> <p>Enrollment period: NR</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: Oral oestriol vs. placebo</p> <p>Groups: G1: 3 mg oral estradiol 3mg qd x 12 wks G2: placebo x 12 wks</p> <p>N at enrollment: G1: 34 G2: 30</p> <p>N at follow-up: G1: 31 G2: 25</p> <p>Women, %: 100</p> <p>Age, mean ± SD: G1: 58.9 ± 10.4 G2: 60.2 ± 9.8</p> <p>Race/ethnicity: NR</p> <p>Height (cm), mean ± SD: G1: 161.9 ± 7.1 G2: 162.9 ± 6.6</p> <p>Weight (kg), mean ± SD: G1: 70.9 ± 11.6 G2: 69.1 ± 12.6</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Women • Ambulant • Postmenopausal • Sensory and/or motor urge incontinence (confirmed by urodynamics) <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Symptoms present for > 3 years before menopause • Voiding difficulty • Pelvic anatomical defect requiring surgery • Neurological disease • Recent estrogen usage (< 6 mos) • Medication use that could affect bladder or urethral function • Contraindication to estrogen 	<p>Daytime incontinence episodes/day, mean ± SD: G1: 3.4 ± 4.2 G2: 3.2 ± 4.7</p> <p>Nocturnal incontinence episodes/day, mean ± SD: G1: 1.6 ± 6.0 G2: 1.0 ± 2.3</p> <p>Daytime voids/day, mean ± SD: G1: 10.2 ± 5.0 G2: 10.6 ± 5.4</p> <p>Nocturia s/day, mean ± SD: G1: 2.0 ± 1.4 G2: 2.6 ± 2.2</p> <p>Diurnal frequency, %: G1: 26 G2: 20</p> <p>Nocturia, %: G1: 25 G2: 20</p> <p>Urgency, %: G1: 29 G2: 25</p> <p>Dysuria, %: G1: 9 G2: 4</p> <p>SUI, %: G1: 11 G2: 10</p> <p>UUI, %: G1: 25 G2: 23</p> <p>Urgency severity, mean score ± SD: G1: 2.6 ± 0.5 G2: 2.5 ± 0.8</p> <p>Nocturia severity mean score ± SD: G1: 1.6 ± 0.8 G2: 1.7 ± 1.1</p>	<p>Daytime incontinence episodes/day, 1 mo, mean ± SD: G1: 3.0 ± 3.5 G2: 2.1 ± 2.6</p> <p>Daytime incontinence episodes/day, 3 mos, mean ± SD: G1: 2.7 ± 3.5 G2: 1.3 ± 2.3</p> <p>Nocturnal incontinence episodes/day, 1 mo, mean ± SD: G1: 0.6 ± 1.2 G2: 0.7 ± 2.0</p> <p>Nocturnal incontinence episodes/day, 3 mos, mean ± SD: G1: 0.3 ± 0.7 G2: 0.3 ± 0.8</p> <p>Daytime voids/day, 1 mo, mean ± SD: G1: 8.9 ± 3.0 G2: 10.6 ± 5.4</p> <p>Daytime voids/day, 3 mos, mean ± SD: G1: 8.6 ± 3.4 G2: 7.2 ± 2.2 G1/BL: <i>P</i> < 0.05 G2/BL: <i>P</i> < 0.01</p> <p>Nocturia episodes/day, 1 mo, mean ± SD: G1: 2.0 ± 1.6 G2: 1.5 ± 1.2</p> <p>Nocturia episodes/day, 3 mos, mean ± SD: G1: 1.5 ± 1.1 G2: 1.4 ± 1.3 G2/BL: <i>P</i> < 0.05</p> <p>Diurnal frequency, cure %: G1: 6 G2: 9</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: +</p> <p>Drop-out rates: ++</p> <p>Power calculation: +</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Cardozo et al., 1993 (continued)			<p>Bladder volume (mL), first desire to void, mean ± SD: G1: 161 ± 74 G2: 163 ± 86</p> <p>Detrusor pressure (cm H₂O), mean ± SD: G1: 17 ± 12 G2: 17 ± 17</p> <p>Max cystometric capacity (mL), mean ± SD: G1: 328 ± 125 G2: 322 ± 113</p> <p>Max pressure (cm H₂O), mean ± SD: G1: 31 ± 27 G2: 21 ± 79</p>	<p>Nocturia, cure %: G1: 10 G2: 8 <i>P</i> < 0.01</p> <p>Urgency, cure %: G1: 7 G2: 9 <i>P</i> < 0.01</p> <p>Dysuria, cure %: G1: 5 G2: 4</p> <p>SUI, cure %: G1: 6 G2: 6</p> <p>UUI, cure %: G1: 11 G2: 7 <i>P</i> < 0.01</p> <p>Urgency severity, 1 mo, mean score ± SD: G1: 1.8 ± 0.8 G2: 1.6 ± 1.0</p> <p>Urgency severity, 3 mos, mean score ± SD: G1: 1.5 ± 1.1 G2: 1.4 ± 1.1 G1/BL: <i>P</i> < 0.01 G2/BL: <i>P</i> < 0.01</p> <p>Nocturia severity, 1 mo, mean score ± SD: G1: 1.3 ± 0.8 G2: 1.3 ± 1.0</p> <p>Nocturia severity, 3 mos, mean score ± SD: G1: 1.0 ± 0.8 G2: 1.0 ± 1.1 G1/BL: <i>P</i> < 0.01</p> <p>Bladder volume (mL), first desire to void, 3 mos, mean ± SD: G1: 206 ± 142 G2: 202 ± 142 G1/BL: <i>P</i> < 0.05</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Cardozo et al., 1993 (continued)				<p>Detrusor pressure (cm H₂O), 3 months, mean ± SD: G1: 17 ± 15 G2: 22 ± 39</p> <p>Max cystometric capacity (mL), mean ± SD: G1: 333 ± 163 G2: 365 ± 136</p> <p>Max pressure (cm H₂O), mean ± SD: G1: 25 ± 20 G2: 32 ± 65 G1/BL: <i>P</i> < 0.05</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Cardozo et al., 2004</p> <p>[See evidence table for Kelleher et al., 2005]</p> <p>Country and setting: 84 centers in Denmark, UK, Czech Republic, Australia, Netherlands, Russia, US</p> <p>Enrollment period: NR</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: Solifenacin 5 mg vs. solifenacin 10 mg vs. placebo, 12 weeks once daily therapy</p> <p>Groups: G1: Solifenacin 5 mg qd G2: Solifenacin 10 mg qd G3: Placebo</p> <p>N at enrollment: 911</p> <p>N treated: G1: 299 G2: 307 G3: 301</p> <p>N at follow-up, first visit: G1: 286 G2: 290 G3: 281</p> <p>N at follow-up, study completion: G1: 277 G2: 283 G3: 270</p> <p>Women, n (%): G1: 237 (82.9) G2: 238 (82.1) G3: 227 (80.8)</p> <p>Age, mean ± SD (range): G1: 55.4 ± 13.8 (19-85) G2: 55.9 ± 14.2 (18-83) G3: 56.1 ± 13.3 (18-82)</p> <p>Race/ethnicity, n (%): White: G1: 280 (97.9) G2: 281 (96.9) G3: 274 (97.5)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 18 • Symptoms of OAB (including urinary frequency with urgency and/or UUI) for ≥3 months • ≥ 8 voids/day • ≥ 3 episodes urinary incontinence during 3-day diary period <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • BOO • PVR volume >200 mL • Stress predominant MUI • Neurological cause for detrusor overactivity • UTI or bladder stones • Previous pelvic irradiation • Malignant disease of pelvic organs • Contraindication to antimuscarinic medication • Narrow-angle glaucoma • Urinary or gastric retention • Non drug treatment for OAB 2 wks before study • Diabetic neuropathy • Drugs to treat incontinence • Drugs with cholinergic or anticholinergic side-effects 	<p>UUI, n (%): 405 (47)</p> <p>Urinary incontinence, n (%): 491 (57)</p> <p>Voids/day, range: 12.05-12.31</p> <p>Duration of symptoms (months), median (range): G1: 27.0, 4-383 G2: 28.0, 4-314 G3: 29.0, 5-327)</p> <p>Prior drug therapy, n (%): G1: 101 (35.3) G2: 94 (32.5) G3: 95 (33.8)</p> <p>Any other non-drug therapy, n (%): G1: 64 (22.4) G2: 77 (26.6) G3: 94 (33.5)</p>	<p>UUI episodes/day, mean change (% change): G1: -1.3 (-62.7) G2: -1.21 (-57.1) G3: -0.91 (-42.5) G1/BL: <i>P</i> = 0.014* G2/BL: <i>P</i> = 0.042*</p> <p>Urgency episodes/day, mean change (95% CI for difference): G1: -2.84 (-1.44, -0.28) G2: -2.90 (-1.49, -0.35) G3: -1.98 G1/G3: <i>P</i> = 0.003 G2/G3: <i>P</i> = 0.002</p> <p>Incontinence episodes/day, mean change (% change): G1: -1.63 (-60.7) G2: -1.57 (-51.9) G3: -1.25 (-27.9) G1/BL: <i>P</i> = 0.002* G2/BL: <i>P</i> = 0.016*</p> <p>Incontinent patients with no incontinence at end of study, %: G1: 50.3 G2: 49.7</p> <p>Voids/day, mean change (95% CI for difference): G1: -2.37 (-1.27, -0.29) G2: -2.81 (-1.71, -0.72) G3: -1.59 G1/G3: <i>P</i> = 0.0018 G2/G3: <i>P</i> = 0.0001</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: fair</p> <p>Randomization: -</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: ++</p> <p>Drop-out rates: ++</p> <p>Power calculation: +</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Cardozo et al., 2004 (continued)	<p>Black: G1: 2 (0.7) G2: 2 (0.7) G3: 3 (1.1) Asian: G1: 4 (1.4) G2: 6 (2.1) G3: 2 (0.7) Other: G1: 0 G2: 1(0.3) G3: 2 (0.7)</p> <p>Weight (kg), mean ± SD: G1: 74.1 ± 15.0 G2: 74.6 ± 15.4 G3: 74.1 ± 14.4</p>	<ul style="list-style-type: none"> • Participation in a clinical trial within 30 days • Childbearing potential, pregnant or nursing or not using reliable contraceptive methods 		<p>Nocturia episodes/day, mean change (95% CI for difference): G1: -0.58 G2: -0.71 (-0.38, -0.01) <i>P</i> = 0.005 G3: -0.52 G1/G3: <i>P</i> = 0.48 G2/G3: <i>P</i> = 0.036</p> <p>KHQ, general health perception, mean change: G1: -5.8 G2: -4.5 G3: -2.3</p> <p>KHQ, incontinence impact, mean change: G1: -27.0 G2: -30.3 G3: -20.1 G1/BL: <i>P</i> < 0.01 G2/BL: <i>P</i> < 0.001</p> <p>KHQ, role limitations, mean change: G1: -21.9 G2: -24.4 G3: -15.9 G1/BL: <i>P</i> < 0.01 G2/BL: <i>P</i> < 0.001</p> <p>KHQ, physical limitations, mean change: G1: -19.2 G2: -22.3 G3: -15.4 G2/BL: <i>P</i> < 0.001</p> <p>KHQ, social limitations, mean change: G1: -12.1 G2: -11.7 G3: -9.0</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Cardozo et al., 2004 (continued)				KHQ, personal relationships, mean change: G1: -8.5 G2: -8.3 G3: -9.4	
				KHQ, emotions, mean change: G1: -16.4 G2: -16.6 G3: -12.2 G1/BL: $P < 0.05$ G2/BL: $P < 0.05$	
				KHQ, sleep/energy, mean change: G1: -15.4 G2: -15.9 G3: -11.4 G1/BL: $P < 0.05$ G2/BL: $P < 0.05$	
				KHQ, severity measures, mean change: G1: -9.6 G2: -11.8 G3: -7.2 G2/BL: $P < 0.01$	
				KHQ, symptom severity, mean change: G1: -3.4 G2: -3.4 G3: -2.4 G1/BL: $P < 0.01$ G2/BL: $P < 0.01$	
				Voided volume (mL), mean change (95% CI for difference): G1: 30.8 (12.4-27.2) G2: 36.0 (17.7-33.0) G3: 10.67 G1/G3: $P = 0.0001$ G2/G3: $P = 0.0001$	
				Dry mouth, n (%): G1: 23 (7.7) G2: 71 (23.1) G3: 7 (2.3)	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Cardozo et al., 2004 (continued)				<p>Dry mouth, mild, n (%): G1: 18 (6.0) G2: 55 (17.9) G3: 5 (1.7)</p> <p>Dry mouth, moderate, n (%): G1: 5 (1.7) G2: 15 (5.2) G3: 1 (0.3)</p> <p>Dry mouth, severe, n (%): G1: 0 G2: 0 G3: 1 (0.3)</p> <p>Constipation, n (%): G1: 11 (3.7) G2: 28 (9.1) G3: 6 (2.0)</p> <p>Constipation, mild, n (%): G1: 9 (3.0) G2: 16 (5.2) G3: 5 (1.7)</p> <p>Constipation, moderate, n (%): G1: 5 (1.7) G2: 15 (5.2) G3: 1 (0.3)</p> <p>Constipation, severe, n (%): G1: 0 G2: 2 (0.7) G3: 0</p> <p>Blurred vision, n (%): G1: 12 (4.0) G2: 18 (5.9) G3: 7 (2.3)</p> <p>Blurred vision, mild, n (%): G1: 7(2.3) G2: 14 (4.6) G3: 6 (2.0)</p> <p>Blurred vision, moderate, n (%): G1: 4 (1.3) G2: 4 (1.3) G3: 1 (0.3)</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Cardozo et al., 2004 (continued)				Blurred vision, severe, n (%): G1: 1 (0.3) G2: 0 G3: 0 Discontinued due to AEs, n (%): G1: 7 (2.3) G2: 12 (3.9) G3: 10 (3.3)	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Cardozo et al., 2008</p> <p>Country and setting: Europe, Community</p> <p>Enrollment period: April 2004 to October 2005</p> <p>Funding: Astellas Pharma Europe Ltd</p> <p>Author industry relationship disclosures: 9 of 9 Apogepha (1) Astellas (9) BARD (1) Bayer-Schering (1) Boston Scientific (1) Cook (1) Gynecare (1) J&J (1) Lilly (3) Merckle Recordati (1) Organon (1) Pfizer (3) Plethora (1) Recordati (1) Sanofi-Aventis (1) UCB (1)</p>	<p>Design: Multicenter randomized double blind placebo-controlled 2 week run in</p> <p>Intervention: Solifenacin 5 or 10 mg daily (pts requesting dose increase to 10 mg at 8 weeks were randomized in a 1:1 ratio to 5 mg or 10 mg)</p> <p>Groups: G1: Solifenacin 5 or 10 mg daily G2: Placebo</p> <p>N at enrollment: G1: 640 G2: 223</p> <p>N at follow-up: G1: 505 G2: 223</p> <p>Women, %: G1: 89.1 G2: 85.7</p> <p>Age, mean: G1: 57.7 G2: 57.9</p> <p>Race/ethnicity, %: White: G1: 98.4 G2: 99.1 Black: G1: 0.4 G2: 0.4 Asian: G1: 0.8 G2: 0 Other: G1: 0.4 G2: 0.4</p> <p>Follow-up: 16 weeks</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 18 • Male or female • OAB symptoms of urgency, frequency, urge incontinence ≥ 3 months • ≥ 3 episodes of urgency with or without incontinence in last 3 days • Mean of ≥ 8 voids/day <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Clinically significant BOO • PVR > 200 mL • UTI • Bladder stones • IC • Previous irradiation • Malignant pelvic disease • Sexually active women practicing an unreliable method of birth control • Pregnancy/lactation • Conditions contraindicating anticholinergic medication 	<p>UUI episodes/day, mean: G1: 1.63 G2: 2.01</p> <p>Severe urgency episodes (PPIUS 3+4)/day, mean: G1: 5.11 G2: 5.54</p> <p>Urgency episodes (PPIUS 1-4)/day, mean: G1: 10.72 G2: 10.81</p> <p>Maximum urgency intensity (PPIUS)/day, mean: G1: 3.53 G2: 3.59</p> <p>Incontinence episodes/day, mean: G1: 1.58 G2: 1.96</p> <p>Voids/day, mean: G1: 11.10 G2: 11.14</p> <p>PPBC score, mean: G1: 5 G2: 5</p> <p>Urgency bother score, VAS (mm), mean: G1: 68 G2: 69</p> <p>Treatment satisfaction score, VAS (mm), mean: G1: 29 G2: 29</p>	<p>UUI episodes/day, mean change ± SD: G1: -1.7 ± 2.2 (N=326) G2: -1.3 ± 2.0 (N=158) P = 0.0002</p> <p>Severe urgency episodes (PPIUS 3+4)/day, mean ± SD: G1: -2.6 ± NR (N=503) G2: -1.8 ± NR (N= 216) P < 0.0001</p> <p>Urgency episodes (PPIUS 1-4)/day, mean ± SD: G1: -2.3 ± 3.0 (N=503) G2: -1.6 ± 3.3 (N=216) P = 0.0006</p> <p>Maximum urgency intensity (PPIUS)/day, mean change ± SD: G1: -0.8 ± 0.9 (N=503) G2: -0.6 ± 0.9 (N=216) P = 0.0006</p> <p>Incontinence episodes/day, mean change ± SD: G1: -1.7 ± 2.2 (N=329) G2: -1.4 ± 2.0 (N=158) P = 0.0003</p> <p>Voids/day, mean change ± SD: G1: -2.1 ± 2.6 (N= 502) G2: -1.3 ± 2.7 (N= 216) P < 0.0001</p>	<p>Quality: Overall quality score: good</p> <p>INTERNAL VALIDITY: good</p> <p>Randomization: +</p> <p>Method and blinding: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: ++</p> <p>Drop-out rates: ++</p> <p>Power calculation: +</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: ++</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Cardozo et al., 2008				<p>PPBC score, median % change: G1: -33 G2: -20 <i>P</i> < 0.0001</p> <p>Urgency bother score, VAS (mm), median % change: G1: -59 G2: -35 <i>P</i> < 0.0001</p> <p>Treatment satisfaction score, VAS (mm), median % change: G1: 66 G2: 34 <i>P</i> < 0.0001</p> <p>Withdrawn due to AEs, n: G1: 17 G2: 6</p> <p>Dry mouth, %: G1: 15.8 G2: 2.7 <i>P</i> < 0.001</p> <p>Constipation, %: G1: 6.9 G2: 2.2 <i>P</i> = NS</p> <p>Blurred Vision, %: G1: 0.8 G2: 0.9 <i>P</i> = NS</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Chancellor et al., 2000</p> <p>Country and setting: 167 centers in Europe, North America, Australia</p> <p>Enrollment period: NR</p> <p>Funding: Pharmacia & Upjohn</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: randomized, double-blind, placebo-controlled, parallel-group</p> <p>Intervention: tolterodine vs. placebo</p> <p>Groups: G1: tolterodine 2 mg b.i.d. x 12 wks G2: placebo x 12 wks</p> <p>N at enrollment: G1: 514 G2: 508</p> <p>N at follow-up: G1: 514 G2: 508</p> <p>Age, mean yrs (range): G1: 60 (22-92) G2: 61 (21-93)</p> <p>Women, N (%): 818 (80%)</p> <p>Race/ethnicity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • ≥18 years • ≥5 incontinence episodes/wk • Average ≥8 voids/24 hr • OAB symptoms ≥6 mos <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Stress incontinence • Total daily urine volume >3L • Contra-indications to antimuscarinic therapy • Hepatic/renal disorders • Symptomatic or history of recurrent UTI • BOO • Haematuria • Interstitial cystitis • Receiving bladder training, electro-stimulation therapy • Indwelling or intermittent catheter • Pregnant or nursing • Women not using reliable contraception • Concomitant treatment for OAB (other than ERT for at least 2 mos) • Use of agents or drugs with potential to inhibit cytochrome P4503A4 isoform 	<p>Previous drug therapy for OAB, %: G1: 54 G2: 52</p> <p>Good efficacy response, %: G1: 62 G2: 59</p> <p>Incontinence episodes/week, mean (range): G1: 23.2 (0-168) G2: 23.3 (0-168)</p> <p>≥5 incontinence episodes/week, n (%): G1: 494 (97) G2: 498 (97)</p> <p>voids/day, mean (range): G1: 11.1 (2-48.6) G2: 11.3 (2-37.4)</p> <p>≥8 voids/day, n (%): G1: 469 (91) G2: 467 (92)</p> <p>Voided volume (mL), mean (range): G1: 137 (38-283) G2: 136 (21-374)</p> <p>Pads/day, mean (range): G1: 1.4 (0-25) G2: 1.5 (0-22)</p>	<p>Incontinence episodes/week, mean change ± SD: G1: -10.6 ± 16.9 G2: -6.9 ± 15.4 P = 0.0005*</p> <p>voids/day, mean change ± SD: G1: -1.7 ± 3.3 G2: -1.2 ± 2.9 P = 0.0079*</p> <p>Voided volume (mL), mean change ± SD: G1: +29 ± 47 G2: +14 ± 41 P = 0.0001*</p> <p>Pads/day, mean change ± SD: G1: -0.5 ± 1.8 G2: -0.2 ± 1.4 P = 0.0035*</p> <p>Improvement, n (%): G1: 313 (61) G2: 218 (43)</p> <p>No change, n (%): G1: 158 (31) G2: 225 (44)</p> <p>Deterioration, n (%): G1: 43 (8) G2: 65 (13)</p> <p>Mild dry mouth, n (%): G1: 93 (18) G2: 31 (6)</p> <p>Moderate dry mouth, n (%): G1: 50 (10) G2: 8 (2)</p> <p>Severe dry mouth, n (%): G1: 12 (2) G2: 0</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: +</p> <p>Drop-out rates: +</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Chancellor et al., 2000 (continued)				Constipation, n (%): G1: 35 (7) G2: 22 (4) Headache, n (%): G1: 19 (4) G2: 23 (5)	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Chancellor, Kianifard, et al., 2008</p> <p>Country and setting: US</p> <p>Enrollment period: May 2005 to February 2006</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: 6 of 7 Genaera (1) Novartis (6)</p>	<p>Design: Randomized open-label</p> <p>Intervention: Behavioral modification in addition to darifenacin (allowed to increase dose from darifenacin 7.5 mg to 15 mg daily after 2 weeks)</p> <p>Groups*: G1: Darifenacin G2: Darifenacin and behavioral modification</p> <p>N at enrollment: G1: 190 G2: 205</p> <p>N at follow-up, 12 weeks: G1: 173 G2: 175</p> <p>Age, mean ± SD: G1: 57.4 ± 13.1 G2: 58.4 ± 14.6</p> <p>Race/ethnicity: White: G1: 90 G2: 88.3 Other (non-white): G1: 10 G2: 11.7</p> <p>Women, %: G1: 90 G2: 88.3</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • ≥ 8 voids/ day • ≥ 2 UUI episodes/ day • ≥ 2 episodes of urgency/day <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Any drug with bladder effects for 2 weeks prior to study participation • Participation in any formal bladder-training program within 30 days of screening • Predominant SUI • Bladder or neurologic condition that could affect bladder function or in which use of anticholinergics was contra-indicated 	<p>UUI episodes/day, mean ± SD: G1: 2.78 ± 2.57 G2: 3.00 ± 2.56</p> <p>UUI episodes/day, median: G1: 2.33 G2: 2.58</p> <p>Mean UUI episodes/day, total population, n (%): 0 episodes: G1: 36 (19.1) G2: 29 (14.2) 1-6 episodes: G1: 138 (73.0) G2: 155 (76.0) 7-13 episodes: G1: 15 (7.9) G2: 20 (9.8) ≥14 episodes: G1: 0 (0.0) G2: 0 (0.0)</p> <p>Mean UUI episodes/day, age ≥ 65, n (%): 0 episodes: G1: 9 (17.0) G2: 8 (9.9) 1-6 episodes: G1: 40 (75.5) G2: 63 (77.8) 7-13 episodes: G1: 4 (7.5) G2: 10 (12.3)</p> <p>Urgency episodes/day, mean ± SD: G1: 10.88 ± 3.80 G2: 10.58 ± 4.00</p> <p>Urgency episodes/day, median: G1: 10.67 G2: 10.33</p> <p>Voids/day, mean ± SD: G1: 11.92 ± 3.03 G2: 11.75 ± 3.37</p>	<p>UUI episodes/day mean change ± SD: G1: -1.89 ± 2.29 G2: -2.10 ± 2.32 G1/G2: <i>P</i> = 0.268</p> <p>UUI episodes/day median change (95% CI): G1: -1.33 (-2.00, -1.00) G2: -2.00 (-2.00, -1.33)</p> <p>Urgency episodes/day mean change ± SD: G1: -2.87 ± 3.59 G2: -2.68 ± 3.54 G1/G2: <i>P</i> = 0.882</p> <p>Urgency episodes/day median change (95% CI): G1: -2.33 (-3.00, -1.67) G2: -2.67 (-3.00, -2.00)</p> <p>Voids/day, mean change ± SD: G1: -2.96 ± 2.91 G2: -2.82 ± 2.87 G1/G2: <i>P</i> = 0.681</p> <p>Voids/day, median change (95% CI): G1: -2.67 (-3.33, -2.00) G2: -2.67 (-3.00, -2.33)</p> <p>Nocturia episodes/day, mean change ± SD: G1: -0.65 ± 1.26 G2: -0.67 ± 1.21 G1/G2: <i>P</i> = 0.315</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Masking: -</p> <p>Pt selection criteria: +</p> <p>Loss to followup: +</p> <p>Drop-out rates: -</p> <p>Power calculation: +</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Chancellor, Kianifard, et al., 2008 (continued)			<p>Voids/day, median: G1: 11.33 G2: 11.33</p> <p>Nocturia episodes/day, mean ± SD: G1: 1.77 ± 1.43 G2: 1.87 ± 1.35</p> <p>Nocturia episodes/day, median: G1: 1.67 G2: 1.67</p> <p>Pads used/day, mean ± SD: G1: 1.12 ± 1.93 G2: 0.99 ± 1.67</p> <p>Pads used/day, median: G1: 0 G2: 0</p>	<p>Nocturia episodes/day, median change (95% CI): G1: -0.67 (-0.67, -0.33) G2: -0.67 (-0.67, -0.33)</p> <p>Pads used/day mean change ± SD: G1: -0.72 ± 1.54 G2: -0.61 ± 1.28 G1/G2: P = 0.978</p> <p>Pads used/day median change (95% CI): G1: 0 (0,0) G2: 0 (0,0)</p> <p>Side effects, %: Constipation: 18.5 Dry mouth 25 UTI: 4.8 Headache: 3.8</p> <p>Discontinued due to adverse event(s), n (%) G1: 6 (3.2) G2: 21 (10.25)</p> <p>Discontinued due to constipation, %: 2</p> <p>Discontinued due to dry mouth, %: 1.8</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Chancellor, Zinner et al., 2008</p> <p>Country and setting: USA, multicenter</p> <p>Enrollment period: NR</p> <p>Funding: Astellas Pharma US, Glaxo-SmithKline</p> <p>Author industry relationship disclosures: 12 of 12 Abbot (1) AEterna Zentaris (1) Akros (1) Allergan (3) American Medical Systems (2) Amgen (1) Astellas (10) AstraZeneca (1) Bard (1) Bayer (1) Boehringer Ingelheim (3) Cephalon (1) Coloplast (1) Cook (1) CooperSurgical (1) Eli Lilly (2) Ferring GSK (6) Gynecare (1) Indevus (2) J&J (1) Mankind (1) Medtronic (1) Merck (1) Novartis (3) Novo Nordisk (1) Ortho-McNeil (1)</p>	<p>Design: Prospective cohort, open label, flexible-dose, 14 day washout</p> <p>Intervention: Solifenacin 5-10 mg daily</p> <p>Groups: G1: Solifenacin 5-10 mg daily</p> <p>N at enrollment: G1: 440</p> <p>N at follow-up (%): G1: 390 (88.4)</p> <p>Age, mean ± SD: G1: 61.4 ± 13.8</p> <p>Race/ethnicity, n (%): White: G1: 392 (88.9)</p> <p>Women, N (%): G1: 389 (88.2)</p> <p>BMI, kg/m² ± SD: G1: 30.1 ± 7.1</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 18 • OAB symptoms > 3 months • Treatment with tolterodine ER 4 mg, solifenacin, darifenacin, or trospium x ≥ 4 weeks • Desired change in therapy • ≥ 3 mean UUI episodes/day <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Treatment < 4 weeks with tolterodine, solifenacin, darifenacin, or trospium • SUI • MUI with primary stress • UTI • Chronic bladder inflammation • Bladder cancer • Severe constipation • Elevated PVR • Neurological deficit • Renal/ hepatic disease • Narrow angle glaucoma • Urinary retention • Gastric retention • Hypersensitivity to drugs • BOO • Women of childbearing age/ lactation 	<p>Urgency episodes/day, mean change ± SD: G1: 6.9 ± 4.4</p> <p>Incontinence episodes/day, mean ± SD: G1: 3.8 ± 3.6</p> <p>Voids/day, mean ± SD: G1: 11.3 ± 3.8</p> <p>Nocturia episodes/day, mean ± SD: G1: 1.9 ± 1.2</p> <p>PPBC score, mean: G1: 4.2</p> <p>OAB-q score, mean: Symptom bother: 57.3 Coping: 52.2 Concern: 51.3 Sleep: 50.6 Social Interaction: 77.7 Total score: 56.7</p>	<p>Urgency episodes/day, mean change ± SD (95% CI): G1: -4.2 ± 4.2 (-4.6, -3.8) G1/BL: P < 0.001</p> <p>Incontinence episodes/day, mean change ± SD (95% CI): G1: -2.6 ± 3.2 (-3.0, -2.3) G1/BL: P < 0.001</p> <p>Voids/day, mean change ± SD (95% CI): G1: -2.3 ± 3.2 (-2.6, -2.0) G1/BL: P < 0.001</p> <p>Nocturia episodes/day, mean change ± SD (95% CI): G1: -0.8 ± 1.0 (-0.9, -0.6) G1/BL: P < 0.001</p> <p>PPBC score, mean change (95% CI): G1: -1.2 (-1.3, 1.0) G1/BL: P < 0.001</p> <p>OAB-q score, mean: Symptom bother: 27.8 Coping: 80.1 Concern: 81.1 Sleep: 75.1 Social Interaction: 92.7 Total score: 81.9 G1/BL: P < 0.001 all domains and total score</p> <p>Dry Mouth, n (%): G1: 77 (17.5)</p> <p>Constipation, n (%): G1: 51 (11.6)</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: fair</p> <p>Randomization: -</p> <p>Method and blinding: NA</p> <p>Pt selection criteria: +</p> <p>Loss to followup: +</p> <p>Drop-out rates: +</p> <p>Power calculation: +</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Chancellor, Zinner, et al., 2008 (continued)				Blurred vision, n (%): G1: 10 (2.3)	
Author industry relationship disclosures:				UTI, n (%): G1: 19 (4.3)	
Pfizer (7)				Headache, n (%): G1: 13 (2.9)	
Pharmacia (1)				URI, n (%): G1: 11 (2.5)	
Pri Med (1)					
Reliant (1)					
Solvay (1)					
Sanofi-Aventis (1)					
Schering-Plough (1)					
Takeda (1)					
TAP (1)					
Watson (1)					

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Chapple et al., 2004</p> <p>Country and setting: 98 centers – United States, United Kingdom, Poland, Russia, New Zealand, Belgium, Netherlands</p> <p>Enrollment period: NR</p> <p>Funding: Yamanouchi Pharmaceutical Co., Ltd, Tokyo, Japan</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT, double-blind placebo controlled, with 2-wk placebo run-in period</p> <p>Intervention: Tolterodine vs. solifenacin vs. placebo</p> <p>Groups: G1: Tolterodine 2 mg b.i.d. G2: Solifenacin 5 mg qd G3: Solifenacin 10 mg qd G4: Placebo qd</p> <p>N at enrollment: G1: 279 G2: 269 G3: 266 G4: 267</p> <p>N at follow-up: G1: 250 G2: 266 G3: 264 G4: 253</p> <p>Female, n (%) G1: 200 (80.0) G2: 188 (71.2) G3: 194 (72.9) G4: 193 (76.3)</p> <p>Age, yrs ± SD: G1: 56.9 +/-12.8 G2: 58.1± 13.4 G3: 57.2 ±13.4 G4: 57.8 ± 13.7</p> <p><65 years of age, n (%) G1: 169 (63.5) G2: 172 (65.2) G3: 172 (68.8) G4: 168 (66.4)</p> <p>≥65 years of age, n (%) G1: 97 (36.5) G2: 92 (34.8) G3: 78 (31.2) G4: 85 (33.6)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 18 • Symptoms of OAB (including urinary frequency with urgency and/or UI) for ≥ 3 months • ≥ 8 voids per 24 hr • ≥ 3 episodes UII during 3-day diary period <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Clinically significant BOO • Postvoid residual volume >200 mL • Stress predominant factor • Neurological cause for detrusor overactivity • UTI or bladder stones • Previous pelvic irradiation • Malignant disease of pelvic organs • Contraindication to antimuscarinic medication (including narrow-angle glaucoma, urinary or gastric retention) • Non pharmacological treatment for OAB 2 wks before study • Diabetic neuropathy • Use of drugs intended to treat incontinence 	<p>UII only, n (%): G1: 142 (56.8) G2: 162 (61.4) G3: 172 (64.7) G4: 177(70.0)</p> <p>MUI, n (%): G1: 90 (36.0) G2: 81 (30.7) G3: 79 (29.7) G4: 59(23.3)</p> <p>No incontinence, n (%): G1: 18 (7.2) G2: 20 (7.6) G3: 15 (5.6) G4: 17 (6.7)</p> <p>UII episodes/day, mean ± SD: G1: 2.33 ± 2.50 G2: 2.14 ± 2.44 G3: 1.86 ± 1.54 G4: 2.02 ± 2.50</p> <p>Urgency episodes/day, mean ± SD: G1: 5.77 ± 4.89 G2: 5.82 ± 4.45 G3: 5.45 ± 3.87 G4: 5.30 ± 3.92</p> <p>Incontinence episodes/day, mean ± SD: G1: 2.64 ± 2.55 G2: 2.59 ± 2.88 G3: 2.32 ± 1.94 G4: 2.71 ± 2.83</p> <p>Voids/day, mean ± SD: G1: 12.08 ± 3.86 G2: 12.32 ± 3.95 G3: 12.08 ± 3.43 G4: 12.20 ± 4.11</p> <p>Voided volume (mL), mean ± SD: G1: 149.6 ± 54.6 G2: 147.2 ± 51.2 G3: 147.0 ± 50.3 G4: 143.8 ± 53.6</p>	<p>UII episodes/day, mean change ± SD: G1: -0.91 (2.01) G2: -1.41 (1.74) G3: -1.36 (2.13) G4: -0.62 (1.96) G1/G4: P = 0.239 G2/G4: P = 0.002 G3/G4: P = 0.0028</p> <p>UII episodes/day, mean % change: G1: -58 G2: -65 G3: -63 G4: -40</p> <p>UII episodes/day, estimated difference vs. tolterodine (95% CI): G2: -0.487 (-0.988, 0.014) G3: -0.436 (-0.921, 0.048)</p> <p>Urgency episodes/day, mean change ± SD: G1: -2.05 ± 3.58 G2: -2.85 ± 3.74 G3: -3.07 ± 3.90 G4: -1.41 ± 3.67 G1/G4: P = 0.0511 G2/G4: P < 0.001 G3/G4: P < 0.001</p> <p>Urgency episodes/day, mean % change: G1: -38 G2: -55 G3: -52 G4: -33</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: -</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: +</p> <p>Drop-out rates: ++</p> <p>Power calculation: +</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Chapple et al., 2004 (continued)	<p><75 years of age, n (%) G1: 236 (88.7) G2: 241 (91.3) G3: 233 (93.2) G4: 225 (88.9)</p> <p>≥75 years of age, n (%) G1: 30 (11.3) G2: 23 (8.7) G3: 17 (16.8) G4: 28 (11.1)</p> <p>Weight, mean kg ± SD G1: 74.8 +/-14.8 G2: 74.6±14.3 G3: 75.5±14.2 G4: 72.6±14.4</p> <p>Race/ethnicity, n (%): White G1: 261(98.1) G2: 260 (98.5) G3: 247(98.8) G4: 248 (98.0) Black G1: 2 (0.8) G2: 0 G3: 1 (0.4) G4: 1 (0.4) Asian G1: 2 (0.8) G2: 0 G3: 2(0.8) G4: 1 (0.4) Other G1: 1 (0.4) G2: 4 (1.5) G3: 0 G4: 3 (1.2)</p>	<ul style="list-style-type: none"> • Use of drugs with cholinergic or anticholinergic side-effects • Participation in a clinical trial within 30 days • Childbearing potential, pregnant or nursing or not using reliable contraceptive methods 	<p>Time from start of symptoms (months), mean ± SD: G1: 57.4 ± 60.5 G2: 72.6 ± 105.4 G3: 62.9 ± 82.5 G4: 61.0 ± 83.9</p> <p>Prior drug therapy, n (%): G1: 93 (34.9) G2: 106 (40.1) G3: 77 (30.8) G4: 83 (32.8)</p> <p>Any non drug therapy, n (%): G1: 92 (34.6) G2: 92 (34.8) G3: 88 (35.2) G4: 76 (30.0)</p>	<p>Urgency episodes/day, estimated difference vs. tolterodine (95% CI): G2: -0.791 (-14.34, -0.148) G3: -1.015 (-1.659, -0.370)</p> <p>Incontinence episodes/day, mean change ± SD: G1: -1.42 ± 1.82 G2: -1.45 ± 2.24 G3: -1.14 ± 2.15 G4: -0.76 ± 2.26 G1/G4: P = 0.1122 G2/G4: P = 0.008 G3/G4: P = 0.0038</p> <p>Incontinence episodes/24 h Δ from baseline, %: G1: -59 G2: -47 G3: -59 G4: -29</p> <p>Incontinence episodes/day, estimated difference vs. tolterodine (95% CI): G2: -0.276 (-0.761, 0.208) G3: -0.316 (-0.786, 0.154)</p> <p>Voids/day, mean change ± SD: G1: -1.88 ± 3.00 G2: -2.19 ± 2.87 G3: -2.61 ± 3.24 G4: -1.20 ± 3.26 G1/G4: P = 0.0145 G2/G4: P = 0.003 G3/G4: P < 0.001</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Chapple et al., 2004 (continued)				<p>Voids/day, mean % change: G1: -15 G2: -20 G3: -17 G4: -18</p> <p>Voids/day, estimated difference vs. tolterodine (95% CI): G2: -0.312 (-0.844, 0.219) G3: -0.737 (-1.269, -0.204)</p> <p>Voided volume (mL), mean change ± SD: G1: 24.4 ± 49.2 G2: 32.9 ± 47.7 G3: 39.2 ± 50.4 G4: 07.4 ± 36.3 G1/G4: <i>P</i> < 0.001 G2/G4: <i>P</i> < 0.001 G3/G4: <i>P</i> < 0.001</p> <p>Voided volume (mL), mean % change: G1: 20 G2: 25 G3: 29 G4: 9</p> <p>Voided volume (mL), estimated difference vs. tolterodine (95% CI): G2: 8.4 (0.496, 16.34) G3: 14.8 (6.855, 22.72)</p> <p>Discontinued due to AEs, n (%): G1: 5 (1.9) G2: 9 (3.2) G3: 7 (2.6) G4: 10 (3.7)</p> <p>Dry mouth, n (%): G1: 49 (18.6) G2: 39 (14.0) G3: 57 (21.3) G4: 13 (4.9)</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Chapple et al., 2004 (continued)				Constipation, n (%): G1: 7 (2.6) G2: 20 (7.2) G2: 21 (7.8) G4: 5 (1.9)	
				Blurred Vision, n (%): G1: 4 (1.5) G2: 10 (3.6) G3: 15 (5.6) G4: 7 (2.6)	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Chapple et al., 2005*</p> <p>Chapple, Fianu-Jonsson et al., 2007†</p> <p>Country and setting: European, Multicenter</p> <p>Enrollment period: May 2003 to October 2004</p> <p>Funding: Yamanouchi (now Astellas)</p> <p>Author industry relationship disclosures: NR* 1 of 9† Astellas (1) Novartis (1) Pfizer (1) Schwartz (1) UCB (1)</p>	<p>Design: RCT</p> <p>Intervention: Solifenacin 5 mg vs. tolterodine ER 4 mg</p> <p>Groups: G1: Solifenacin 5mg G2: Tolterodine ER 4mg</p> <p>G1a: G1 who elected not to increase dose at 4 weeks G2a: G2 who elected not to increase dose at 4 weeks</p> <p>N at enrollment: G1: 578 G2: 599</p> <p>N at follow-up: G1a: 297 G2a: 267</p> <p>Age, mean: G1: 56.5 G2: 56.4 G1a: 56.5 G2a: 56.9</p> <p>Race/ethnicity, %: White: G1: 99.3 G2: 99.5 G1a: 99.3 G2a: 99.3</p> <p>Women, %: G1: 85.3 G2: 88.3 G1a: 87.5 G2a: 87.1</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 18 • OAB symptoms for ≥ 3 months • Outpatient treatment • ≥ 8 voids/day • ≥ 1 incontinence episode/day or ≥ 1 urgency episode/day <p>Exclusion criteria: NR</p>	<p>UUI episodes/day, mean ± SD:* G1: 2.31 ± 2.35 G2: 2.12 ± 2.14</p> <p>UUI episodes/day, mean:† G1a: 1.97 G2a: 1.66</p> <p>Urgency episodes/day, mean ± SD:* G1: 6.01 ± 4.66 G2: 5.84 ± 4.12</p> <p>Urgency episodes/day, mean:† G1a: 5.67 G2a: 5.30</p> <p>Incontinence episodes/day, mean ± SD:* G1: 2.77 ± 2.65 G2: 2.55 ± 2.37</p> <p>Incontinence episodes/day, mean:† G1a: 2.32 G2a: 2.01</p> <p>Voids/day, mean ± SD:* G1: 11.78 ± 3.58 G2: 11.66 ± 3.25</p> <p>Voids/day, mean:† G1: 11.10 G2: 11.36</p> <p>Nocturia, episodes/day, mean ± SD:* G1: 2.02 ± 1.33 G2: 1.92 ± 1.22</p> <p>Nocturia, episodes/day, mean:† G1a: 1.83 G2a: 1.84</p>	<p>UUI episodes/day, 4 wks, mean change:† G1: -1.22 G2: -0.91 <i>P</i> = NS</p> <p>UUI episodes/day, 12 wks, mean change:* G1: -1.42 G2: -0.83 <i>P</i> = 0.001</p> <p>UUI episodes/day, 12 wks, mean change:† G1a: -1.46 G2a: -1.03</p> <p>Urgency episodes/day, 4 wks, mean change:† G1: -1.98 G2: -1.67 <i>P</i> = NS</p> <p>Urgency episodes/day, 12 wks, mean change:* G1: -2.85 G2: -2.42 <i>P</i> = 0.035</p> <p>Urgency episodes/day, 12 wks, mean change:† G1a: -3.08 G2a: -2.62</p> <p>Incontinence episodes/day, 4 wks, mean change:† G1: -1.30 G2: -0.90 <i>P</i> = 0.0181</p>	<p>Quality: Overall quality score: good</p> <p>INTERNAL VALIDITY: good</p> <p>Randomization: +</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: ++</p> <p>Drop-out rates: ++</p> <p>Power calculation: +</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Chapple et al., 2005* Chapple et al., 2007† (continued)				<p>Incontinence episodes/day, 12 wks, mean change:* G1: -1.60 G2: -1.11 <i>P</i> = 0.006</p>	
				<p>Incontinence episodes/day, 12 wks, mean change:† G1a: -1.56 G2a: -1.23</p>	
				<p>Voids/day, 4 wks, mean change:† G1: -1.71 G2: -1.47 <i>P</i> = NS</p>	
				<p>Voids/day, 12 wks, mean change:* G1: -2.45 G2: -2.24 <i>P</i> = 0.004 for non-inferiority</p>	
				<p>Voids/day, 12 wks, mean change:† G1a: -2.47 G2a: -2.49</p>	
				<p>Nocturia, episodes/day, 4 wks, mean change:† G1: -0.51 G2: -0.44 <i>P</i> = NS</p>	
				<p>Nocturia, episodes/day, 12 wks, mean change:* G1: -0.71 G2: -0.63 <i>P</i> = NS</p>	
				<p>Nocturia episodes/day, 12 wks, mean change:† G1a: -0.72 G2a: -0.69</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Chapple et al., 2005* Chapple et al., 2007† (continued)				<p>Perception of bladder condition, 4 wks, mean change:† G1: -0.96 G2: -0.88 <i>P</i> = NS</p> <p>Perception of bladder condition, 12 wks, mean change:* G1: -1.51 G2: -1.33 <i>P</i> < 0.0061</p> <p>Perception of bladder condition, 12 wks, mean change:† G1a: -1.72 G2a: -1.62</p> <p>Pad use, 4 wks, mean change:† G1: -1.21 G2: -0.80 <i>P</i> = 0.0089</p> <p>Pad use, 12 wks, mean change:* G1: -1.72 G2: -1.19 <i>P</i> < 0.0023</p> <p>Pad use, 12 wks, mean change:† G1a: -1.55 G2a: -1.40</p> <p>Dry rate, 4 wks, mean % change:† G1: 39 G2: 34 <i>P</i> = NS</p> <p>Dry rate, 12 wks, mean % change:† G1a: 65.4 G2a: 58.3 <i>P</i> = NS</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Chapple et al., 2005* Chapple et al., 2007† (continued)				<p>Voided volume (mL), 12 wks, mean change:* G1: 38 G2: 31 <i>P</i> = 0.010</p> <p>Voided volume (mL), 12 wks, mean change:† G1a: 39.95 G2a: 37.84</p> <p>Dry mouth, mild, %: *† G1: 17.5 G2: 14.8 G1a: 6.5 G2a: 5.0</p> <p>Dry mouth, moderate, %: *† G1: 10.8 G2: 7.7 G1a: 10.4 G2a: 7.0</p> <p>Dry mouth, severe, %: *† G1: 1.7 G2: 1.5 G1a: 0.7 G2a: 2.1</p> <p>Constipation, mild, %: *† G1: 3.2 G2: 1.3 G1a: 2.0 G2a: 1.0</p> <p>Constipation, moderate, %: *† G1: 2.7 G2: 1.0 G1a: 1.7 G2a: 1.4</p> <p>Constipation, severe, %: *† G1: 0.5 G2: 0.2 G1a: 0.3 G2a: 0.0</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Chapple et al., 2005*				Blurred vision, mild, %:*†	
Chapple et al., 2007† (continued)				G1: 0.7 G2: 0.7 G1a: 0.3 G2a: 0.7	
				Blurred vision, moderate, %:*†	
				G1: 0.0 G2: 1.0 G1a: 0.0 G2a: 1.7	
				Blurred vision, severe, %:*†	
				G1: 0.0 G2: 0.0 G1a: 0.0 G2a: 0.0	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Chapple, Van Kerrebroeck et al., 2007</p> <p>Country and setting: Multicenter study in Europe, Africa, and Oceania (academic and private), 150 sites in 19 countries</p> <p>Enrollment period: NR</p> <p>Funding: Schwarz BioSciences GmbH, Pfizer Inc.</p> <p>Author conflicts of interest: 9 of 9: Astellas (Yamanouchi) (2) Eli-Lilly (1) Ferring (1) GlaxoSmithKline (1) Novartis (3) Pfizer (5) Schwarz (4) UCB (1)</p>	<p>Design: RCT</p> <p>Intervention: Fesoterodine (8 mg or 4 mg) vs. tolterodine ER 4 mg</p> <p>Groups: G1: Tolterodine ER 4mg G2: Fesoterodine 8mg G3: Fesoterodine 4mg G4: Placebo</p> <p>N at enrollment: G1: 290 G2: 288 G3: 272 G4: 285</p> <p>N at follow-up: G1: 253 G2: 252 G3: 231 G4: 252</p> <p>Age, mean ± SD: G1: 57.7 ± 14.6 G2: 55.6 ± 14.1 G3: 57.1 ± 13.2 G4: 56.0 ± 13.7</p> <p>Women, %: G1: 78 G2: 82 G3: 81 G4: 81</p> <p>Race/ethnicity, %: White: G1: 98 G2: 98 G3: 96 G4: 98 Black: G1: 0 G2: 1 G3: 0 G4: <1 Asian: G1: 2 G2: 1 G3: 2 G4: 2</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • OAB symptoms with urinary urgency for ≥ 6 mos • Age ≥ 18 • ≥ 8 voids per day • ≥ 6 urgency episodes or ≥ 3 UUI episodes/day • At least moderate problems recorded via a Likert scale • Negative pregnancy test • Adequate contraception <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • SUI • Bladder stones • Interstitial cystitis • Urothelial tumors • Grade III or higher pelvic prolapse • BOO • Polyuria (>3L/d) • UTIs • PVR urine volume >100mL • Use of antimuscarinic ≤ 2 weeks • Treatment with electrostimulation n ≤ 4 weeks • Neurological disease • Clinically relevant cardiac arrhythmia • Unstable angina • QTcB interval > 500ms 	<p>UUI episodes/day, mean ± SD: G1: 3.8 ± 3.1 G2: 3.7 ± 3.0 G3: 3.8 ± 3.1 G4: 3.7 ± 3.1</p> <p>Urgency episodes/day mean ± SD: G1: 11.0 ± 3.4 G2: 11.5 ± 4.2 G3: 11.0 ± 3.4 G4: 11.4 ± 4.0</p> <p>Incontinence, %: G1: 79 G2: 81 G3: 75 G4: 76</p> <p>Continent days/wk, mean ± SD: G1: 0.6 ± 1.3 G2: 0.6 ± 1.3 G3: 0.8 ± 1.6 G4: 0.8 ± 1.5</p> <p>Voids/day, mean ± SD: G1: 11.5 ± 2.9 G2: 11.9 ± 3.8 G3: 11.6 ± 3.2 G4: 12.0 ± 3.7</p> <p>Daytime voids/day, mean ± SD: G1: 2.0 ± 1.2 G2: 2.0 ± 1.6 G3: 1.9 ± 1.3 G4: 1.8 ± 1.2</p> <p>Duration of OAB symptoms (yrs), mean ± SD: G1: 8.7 ± 10.1 G2: 7.6 ± 8.4 G3: 9.0 ± 11.2 G4: 7.9 ± 9.6</p> <p>Previous drug treatment, n (%): G1: 135 (47) G2: 118 (41) G3: 102 (38) G4: 112 (40)</p>	<p>UUI episodes/day, LS mean change (SE): G1: -1.74 (0.16) G2: -2.22 (0.16) G3: -1.95 (0.17) G4: -1.14 (0.16) G1/G4: P = 0.008 G2/G4: P < 0.001 G3/G4: P = 0.001</p> <p>UUI episodes/day, median % change: G1: -70.0 G2: -87.5 G3: -80.0 G4: -50.0 G1/G4: P = 0.105 G2/G4: P < 0.001 G3/G4: P = 0.001</p> <p>Urgency episodes/day mean change (SE): G1: -2.03 (0.19) G2: -2.36 (0.20) G3: -1.88 (0.20) G4: -1.07 (0.19) G1/G4: P < 0.001 G2/G4: P < 0.001 G3/G4: P = 0.003</p> <p>Urgency episodes/day median change, %: G1: -16.0 G2: -19.1 G3: -17.6 G4: -11.1 G1/G4: P = 0.004 G2/G4: P < 0.001 G3/G4: P = 0.002</p> <p>Continent days/wk mean change (SE): G1: 2.48 (0.20) G2: 3.32 (0.19) G3: 2.84 (0.21) G4: 2.07 (0.20) G1/G4: P = 0.139 G2/G4: P < 0.001 G3/G4: P = 0.007</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: +</p> <p>Drop-out rates: -</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Chapple, Van Kerrebroeck et al., 2007 (continued)	<p>Other: G1: <1 G2: 0 G3: 3 G4: <1</p> <p>BMI, kg/m² ± SD: G1: 27.5 ± 5.2 G2: 27.1 ± 5.2 G3: 27.5 ± 5.5 G4: 27.2 ± 5.2</p>		<p>Voided volume (mL), mean ± SD: G1: 154.3 ± 52.9 G2: 153.9 ± 56.9 G3: 160.0 ± 59.5 G4: 150.2 ± 52.0</p>	<p>Voids/day, LS mean change (SE): G1: -1.73 (0.16) G2: -1.88 (0.16) G3: -1.76 (0.17) G4: -0.95 (0.16) G1/G4: <i>P</i> = 0.001 G2/G4: <i>P</i> < 0.001 G3/G4: <i>P</i> < 0.001</p> <p>Voids/day, median % change: G1: -13.8 G2: -18.6 G3: -16.7 G4: -11.1 G1/G4: <i>P</i> = 0.005 G2/G4: <i>P</i> < 0.001 G3/G4: <i>P</i> < 0.001</p> <p>Daytime voids/day, LS mean change (SE): G1: -0.40 (0.06) G2: -0.39 (0.06) G3: -0.39 (0.06) G4: -0.32 (0.06)</p> <p>Daytime voids/day, median % change: G1: -25.0 G2: -23.1 G3: -28.6 G4: -26.8</p> <p>Treatment Response, % yes: G1: 72 G2: 79 G3: 75 G4: 53 G1/G4: <i>P</i> < 0.001 G2/G4: <i>P</i> < 0.001 G3/G4: <i>P</i> < 0.001</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Chapple, Van Kerrebroeck et al., 2007 (continued)				<p>Maximum voided volume, mean change (SE): G1: 23.64 (3.31) G2: 33.62 (3.35) G3: 27.72 (3.41) G4: 9.37 (3.33) G1/G4: (<i>P</i> = 0.003) G2/G4: (<i>P</i> <0.001) G3/G4: (<i>P</i> = 0.001)</p> <p>Adverse events, n (%): G1: 144 (50) G2: 167 (58) G3: 135 (50) G4: 107 (38)</p> <p>Dry mouth, n (%): G1: 49 (16.9) G2: 97 (33.8) G3: 59 (21.7) G4: 20 (7.1)</p> <p>Constipation, n (%): G1: 8 (2.8) G2: 13 (4.5) G3: 9 (3.3) G4: 4 (1.4)</p> <p>Headache, n (%): G1: 14 (4.8) G2: 7 (2.4) G3: 12 (4.4) G4: 14 (4.9)</p> <p>Dry eye, n (%): G1: 1 (<1) G2: 12 (4.2) G3: 6 (2.2) G4: 0</p> <p>Nasopharyngitis, n (%): G1: 10 (3.4) G2: 5 (1.7) G3: 8 (2.9) G4: 7 (2.5)</p> <p>Fatigue, n (%): G1: 10 (3.4) G2: 1 (<1) G3: 1 (<1) G4: 1 (<1)</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Chapple, Van Kerrebroeck et al., 2007 (continued)				<p>Influenza, n (%): G1: 2 (<1) G2: 2 (<1) G3: 9 (3.3) G4: 6 (2.1)</p> <p>Dry throat, n (%): G1: 3 (1) G2: 8 (2.8) G3: 1 (<1) G4: 0</p> <p>Dizziness, n (%): G1: 4 (1.4) G2: 3 (1.0) G3: 4 (1.5) G4: 7 (2.5)</p> <p>Alanine aminotransferase, n (%): G1: 0 G2: 6 (2.1) G3: 2 (<1) G4: 1 (<1)</p> <p>Nausea, n (%): G1: 6 (2.1) G2: 4 (1.4) G3: 1 (<1) G4: 1 (<1)</p> <p>Mean change in heart rate (bpm): G1: 2.89 G2: 3.9 G3: 3.3 G4: 0.8</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Chapple, DuBeau et al., 2007</p> <p>Country and setting: US, Poland, South Africa, Hungary, Sweden, United Kingdom, Germany; 73 centers</p> <p>Enrollment period: NR</p> <p>Funding: Novartis</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: Darifenacin 7.5 mg qd versus placebo x 12 weeks, with optional up-titration offered (to 15 mg qd) after 2 weeks of treatment</p> <p>Groups: G1: Darifenacin G2: Placebo</p> <p>N at enrollment: G1: 266 G2: 133</p> <p>N at follow-up (%): G1: 244 (91.7) G2: 117 (87.3)</p> <p>Age, mean ± SD (range): G1: 72 ± 5 (64-89) G2: 73 ± 5 (64-87)</p> <p>Women, n (%): G1: 206 (77.4) G2: 100 (75.2)</p> <p>Race/ethnicity: NR</p> <p>Follow-up: 12 weeks</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 65 • OAB sx for ≥ 6 months • Capable of independent toileting • Able to complete the diary independently • Completed at least 5 days of the 7-day diary during the baseline period • ≥ 1 UUI episodes/day • ≥ 10 micturitions/day <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Treatment w/ drugs known to effect urinary bladder function or the external urethral sphincter • Total urinary volume ≥ 3L • Mean voided volume > 300 ml • Clinically significant SUI or BOO • PVR > 100mL by ultrasound • Women with marked cystocele or other clinically significant Stage 3 or Stage 4 pelvic prolapsed • Participated in a bladder-training program or received electrical stimulation therapy w/in 3 mos of screening • Serious or intermittent UTI 	<p>UUI episodes/week, median (range): G1: 19.8 (4.0-142.0) G2: 21.0 (7.0-155.4)</p> <p>Urgency episodes/day, median (range): G1: 7.6 (1.0-24.4) G2: 7.4 (1.3-22.2)</p> <p>Voids/day, median (range): G1: 11.8 (7.1-25.1) G2: 12.0 (7.6-27.6)</p>	<p>UUI episodes/week, median change (% change): G1: -14.0 (-88.6) G2: -13.0 (-77.9)</p> <p>UUI episodes/week, median change (95% CI): G1/G2: -1.0 (-2.83, 1) <i>P</i> = 0.328</p> <p>Urgency episodes/day, median change (% change): G1: -3.3 (-69.6) G2: -3.1 (-55.5)</p> <p>Urgency episodes/day, median change (% change): G1/G2: -0.5 (-1.2, 0.2) <i>P</i> = 0.174</p> <p>Voids/day, median change (% change): G1: -3.0 (-25.3) G2: -2.2 (-18.5)</p> <p>Voids/day, median change (95% CI): G1/G2: -0.7 (-1.1, -0.2) <i>P</i> = 0.006</p> <p>OAB-q, total score, mean change: G1: 22.9 G2: 16.8 <i>P</i> < 0.001</p> <p>OAB-q, symptom severity score, mean change: G1: -27.9 G2: -20.7 <i>P</i> < 0.001</p>	<p>Quality: Overall quality score: good</p> <p>INTERNAL VALIDITY: good</p> <p>Randomization: +</p> <p>Method and blinding: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: ++</p> <p>Drop-out rates: +</p> <p>Power calculation: +</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Chapple, DuBeau et al., 2007 (continued)		<ul style="list-style-type: none"> • Clinically significant congenital or acquired disorder of the urinary tract • Any urinary bladder dysfunction other than OAB • Hx of chronic pain syndrome of the lower urinary tract <p>Other significant medical conditions that made participant unfit (cognitive impairment, uncontrolled severe HTN, uncontrolled severe heart failure, recent MI, uncontrolled thyroid disease</p>		<p>OAB-q, coping score, mean change: G1: 27.4 G2: 19.2 <i>P</i> < 0.001</p> <p>OAB-q, concern score, mean change: G1: 26.5 G2: 18.6 <i>P</i> < 0.001</p> <p>OAB-q, sleep score, mean change: G1: 21.1 G2: 16.8 <i>P</i> = 0.015</p> <p>OAB-q, social score, mean change: G1: 12.5 G2: 10.4 <i>P</i> = 0.047</p> <p>PVR (mL), wk 12, mean change (95% CI) : G1: 11.9 (1.7, 22.1) G2: 17.3 (-18.1, 52.8) <i>P</i> = NS</p> <p>Adverse Event, n (%) G1: 149 (56.0) G2: 60 (45.1)</p> <p>AEs attributed to treatment, n (%) G1: 99 (37.2) G2: 24 (18.0)</p> <p>Dry mouth, n (%) G1: 59 (22.2) G2: 5 (3.8)</p> <p>Constipation, n (%) G1: 41 (15.4) G2: 11 (8.3)</p> <p>Discontinued due to AEs, n (%) G1: 4 (1.6) G2: 1 (0.8)</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Chapple et al., 2008</p> <p>Country and setting: European, multicenter (academic and private)</p> <p>Enrollment period: NR</p> <p>Funding: Pfizer, Schwarz BioSciences</p> <p>Author industry relationship disclosures: 4 of 5 Pfizer (4)</p>	<p>Design: Multicenter randomized placebo-controlled</p> <p>Intervention: Tolterodine 4 mg ER vs. fesoterodine 8 mg</p> <p>Groups: G1: Tolterodine ER 4mg G2: Fesoterodine 8mg G3: Placebo</p> <p>N at enrollment: G1: 290 G2: 287 G3: 283</p> <p>Age, mean ± SD: Total: 57 ± 14</p> <p>Race/ethnicity, %: White: Total: >95</p> <p>Women, %: Total: 80</p> <p>Follow-up: 12 weeks</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 18 • OAB sx w/ urinary urgency for ≥6 mos • ≥ 8 voids/day • ≥ 6 urgency episodes or ≥ 3 UUI episodes/day • At least moderate problems recorded via a Likert scale • Negative pregnancy test • Adequate contraception throughout trial <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • LUT pathology: SI, bladder stones, interstitial cystitis, urothelial tumors • Grade III or higher pelvic prolapsed • Bladder-outlet obstruction • Polyuria (>3L/d) • Symptomatic or recurrent UTIs • PVR urine volume >100 mL • Antimuscarinic agent w/in 2 wks • Electrostimulation for bladder training in prior 4 wks • Active UTI • Underlying neurological dz • Clinically relevant cardiac arrhythmia • Unstable angina • QTcB interval > 500ms 	<p>Time since first diagnosis or onset of OAB (years), mean: Total: 8-9</p> <p>Incontinent at BL, n: G1: 213 G2: 217 G3: 203</p>	<p>KHQ severity score, mean change: G1: -12.6 G2: -14.0 G3: -9.0 G1/G3: <i>P</i> < 0.05 G2/G3: <i>P</i> < 0.05</p> <p>KHQ severity score, patients incontinent at BL, mean change: G1: -14.9 G2: -15.8 G3: -10.8 G1/G3: <i>P</i> < 0.05 G2/G3: <i>P</i> < 0.05</p> <p>KHQ emotions score, mean change: G1: -16.3 G2: -17.4 G3: -10.1 G1/G3: <i>P</i> < 0.05 G2/G3: <i>P</i> < 0.05</p> <p>KHQ emotions score, patients incontinent at BL, mean change: G1: -17.3 G2: -18.6 G3: -11.3 G1/G3: <i>P</i> < 0.05 G2/G3: <i>P</i> < 0.05</p> <p>KHQ role limitations score, mean change: G1: -22.1 G2: -21.7 G3: -11.8 G1/G3: <i>P</i> < 0.05 G2/G3: <i>P</i> < 0.05</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Method and blinding: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: NR</p> <p>Drop-out rates: NR</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Chapple et al., 2008 (continued)				<p>KHQ role limitations score, patients incontinent at BL, mean change: G1: -23.2 G2: -23.7 G3: -12.4 G1/G3: $P < 0.05$ G2/G3: $P < 0.05$</p> <p>KHQ physical limitations score, mean change: G1: -19.7 G2: -21.7 G3: -11.4 G1/G3: $P < 0.05$ G2/G3: $P < 0.05$</p> <p>KHQ Physical limitations score, patients incontinent at BL, mean change: G1: -20.5 G2: -23.3 G3: -11.1 G1/G3: $P < 0.05$ G2/G3: $P < 0.05$</p> <p>KHQ social limitations score, mean change: G1: -14.1 G2: -15.4 G3: -8.7 G1/G3: $P < 0.05$ G2/G3: $P < 0.05$</p> <p>KHQ social limitations score, patients incontinent at BL, mean change: G1: -15.7 G2: -16.2 G3: -9.5 G1/G3: $P < 0.05$ G2/G3: $P < 0.05$</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Chapple et al., 2008 (continued)				<p>KHQ sleep/energy score, mean change: G1: -11.7 G2: -13.6 G3: -9.6 G1/G3: P = NS G2/G3: P < 0.05</p> <p>KHQ sleep/energy score, patients incontinent at BL, mean change: G1: -12.5 G2: -15.3 G3: -10.4 G1/G3: P = NS G2/G3: P < 0.05</p> <p>KHQ personal relationship score, mean change: G1: -10.4 G2: -11.9 G3: -6.2 G1/G3: P = NS G2/G3: P < 0.05</p> <p>KHQ personal relationship score, patients incontinent at BL, mean change: G1: -12.7 G2: -12.3 G3: -6.8 G1/G3: P < 0.05 G2/G3: P = NS</p> <p>KHQ incontinence impact score, mean change: G1: -23.3 G2: -24.6 G3: -16.1 G1/G3: P < 0.05 G2/G3: P < 0.05</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Chapple et al., 2008 (continued)				<p>KHQ incontinence impact score, patients incontinent at BL, mean change: G1: -23.8 G2: -26.5 G3: -17.7 G1/G3: $P < 0.05$ G2/G3: $P < 0.05$</p> <p>KHQ general health score, mean change: G1: -4.3 G2: -4.0 G3: -3.8 G1/G3: $P = NS$ G2/G3: $P = NS$</p> <p>KHQ general health score, patients incontinent at BL, mean change: G1: -4.3 G2: -4.5 G3: -5.5 G1/G3: $P = NS$ G2/G3: $P = NS$</p> <p>ICIQ-SF total score, mean: G1: -3.95 G2: -4.41 G3: -2.55 G1/G3: $P < 0.05$ G2/G3: $P < 0.05$</p> <p>ICIQ-SF total score, patients incontinent at BL, mean: G1: -4.56 G2: -5.29 G3: -3.12 G1/G3: $P < 0.05$ G2/G3: $P < 0.05$</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Chapple et al., 2008 (continued)				<p>Major improvement in severity of bladder-related problems, %: G1: 34 G2: 39 G3: 25 G1/G3: <i>P</i> = 0.01 G2/G3: <i>P</i> = 0.01</p> <p>Dry mouth, n (%): G1: 49 (17) G2: 97 (34) G3: 20 (7)</p> <p>Constipation, n (%): G1: 8 (3) G2: 13 (5) G3: 4 (1)</p> <p>Nasopharyngitis, n (%): G1: 10 (3) G2: 5 (2) G3: 7 (3)</p> <p>Dry eye, n (%): G1: 1 (1) G2: 12 (4) G3: 0 (0)</p> <p>Nausea, n (%): G1: 6 (2) G2: 4 (1) G3: 1 (1)</p> <p>Fatigue, n (%): G1: 10 (3) G2: 1 (1) G3: 1 (1)</p> <p>Dry throat, n (%): G1: 3 (1) G2: 8 (3) G3: 0 (0)</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Choo et al., 2008</p> <p>Country and setting: Korea, Academic hospital</p> <p>Enrollment period: July 2005 to March 2006</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: None</p>	<p>Design: Prospective cohort</p> <p>Intervention: Tolterodine ER 4 mg qd x 12 wks</p> <p>Groups: NA</p> <p>N at enrollment: 60</p> <p>N at follow-up: 56</p> <p>Women, %: 100</p> <p>Age, mean ± SD: 54.8 ± 11.5</p> <p>Race/ethnicity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 18 • OAB for ≥ 3 mos • ≥ 8 frequency episodes/d and ≥ 2 urgency episodes/d <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Total 24-h micturition volume > 3L • PVR > 30% of voided volume or ≥ 150 mL • Stress UI • Hx of urinary retention • Active UTI • Indwelling catheter • Intermittent self-catheterisation • Neurological dz • Any condition for which antimuscarinic agents are contraindicated 	<p>UUI episodes/day, mean ± SD: 1.4 ± 2.7</p> <p>UUI episodes/day, mean ± SD: PGA < 80%: 2.4 ± 3.4 PGA ≥ 80%: 2.9 ± 4.1</p> <p>Urgency episodes/day, mean ± SD: 9.6 ± 6.8</p> <p>Urgency episodes/day, mean ± SD: PGA < 60%: 12.6 ± 7.7 PGA ≥ 60%: 7.1 ± 4.9 <i>P</i> < 0.01</p> <p>Total of all urgency scales/day, mean ± SD: 94.4 ± 43.0</p> <p>Voids/day, mean ± SD: 12.6 ± 5.3</p> <p>Voids/day, mean ± SD: PGA < 60%: 14.9 ± 6.2 PGA ≥ 60%: 10.5 ± 2.9 <i>P</i> < 0.01</p> <p>Nocturia episodes/day, mean ± SD: 1.6 ± 0.9</p> <p>Nocturia episodes/day, mean ± SD: PGA < 50%: 1.9 ± 0.8 PGA ≥ 50%: 1.6 ± 0.7</p>	<p>UUI episodes/day, 12 wks, mean ± SD: 0.4 ± 0.9 <i>P</i> < 0.01</p> <p>UUI severity, 12 wks, median % change (95% CI): -71 (-76.8, -39.2)</p> <p>UUI, PGA median rate, % (95% CI): 80 (34.2, 80.0)</p> <p>UUI episodes/day, 12 wks, mean change ± SD: PGA < 80%: -0.9 ± 1.6 PGA ≥ 80%: -2.8 ± 4.3</p> <p>UUI, PPBC, mean change ± SD: PGA < 80%: -1.6 ± 1.1 PGA ≥ 80%: -2.2 ± 1.5</p> <p>UUI, GII, %: PGA < 80%: No: 1 (11.1) Little: 5 (55.6) Much: 3 (33.3) PGA ≥ 80%: No: 0 (0) Little: 4 (40.0) Much: 6 (60.0)</p> <p>UUI, continue treatment, n (%): PGA < 80%: 5 (55.6) PGA ≥ 80%: 8 (80.0)</p> <p>Urgency episodes/day, 12 wks, mean ± SD: 5.4 ± 4.5 <i>P</i> < 0.01</p> <p>Urgency severity, 12 wks, median % change (95% CI): -55 (-60.4, -40.1)</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: -</p> <p>Methods and blinding: -</p> <p>Pt selection criteria: +</p> <p>Loss to followup: ++</p> <p>Drop-out rates: ++</p> <p>Power calculation: +</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: +</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Choo et al., 2008 (continued)			<p>Most troublesome symptoms, n (%): UUI: 9 (16.1) Urgency: 6 (10.7) Frequency: 28 (50) Nocturia: 10 (17.9) Tenesmus: 3 (5.4)</p> <p>Voided volume, (mL), mean ± SD: 157.0 ± 70.3</p>	<p>Total of all urgency scales/ day, 12 wks, mean ± SD: 66.3 ± 31.2 <i>P</i> < 0.01</p> <p>Urgency, PGA median rate, % (95% CI): 60 (46.2, 64.9)</p> <p>Urgency episodes /day, 12 wks, mean change ± SD: PGA < 60%: -4.2 ± 6.8 PGA ≥ 60%: -4.3 ± 4.7 <i>P</i> < 0.01</p> <p>Urgency, PPBC, mean change ± SD: PGA < 60%: -1.5 ± 1.0 PGA ≥ 60%: -2.3 ± 1.1 <i>P</i> < 0.01</p> <p>Urgency, GII, %: PGA < 60%: No: 4 (16.0) Little: 15 (60.0) Much: 6 (24.0) <i>P</i> < 0.01 PGA ≥ 60%: No: 0 (0) Little: 7 (22.6) Much: 24 (77.4) <i>P</i> < 0.01</p> <p>Urgency, pts want to continue treatment, n (%): PGA < 60%: 16 (64.0) PGA ≥ 60%: 25 (80.6)</p> <p>Voids/day, 12 wks, mean ± SD: 9.9 ± 4.2 <i>P</i> < 0.01</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Choo et al., 2008 (continued)				<p>Frequency severity, 12 wks, median % change (95% CI): -45 (-54.4, -36.2)</p> <p>Frequency, PGA median rate, % (95% CI): 60 (46.9, 63.6)</p> <p>Urinary frequency, 12 wks, mean change ± SD: PGA < 60%: -2.6 ± 2.9 PGA ≥ 60%: -2.7 ± 2.2 <i>P</i> < 0.01</p> <p>Frequency, PPBC, mean change ± SD: PGA < 60%: -1.4 ± 1.1 PGA ≥ 60%: -2.4 ± 1.0 <i>P</i> < 0.01</p> <p>Frequency, GII, %: PGA < 60%: No: 4 (14.8) Little: 16 (59.3) Much: 7 (25.9) <i>P</i> < 0.01 PGA ≥ 60%: No: 0 (0) Little: 6 (20.7) Much: 23 (79.3) <i>P</i> < 0.01</p> <p>Frequency, continue treatment, n (%): PGA < 60%: 18 (66.7) PGA ≥ 60%: 23 (79.3)</p> <p>Nocturia episodes /day, 12 wks, mean ± SD: 0.9 ± 0.8 <i>P</i> < 0.01</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Choo et al., 2008 (continued)				<p>Nocturia severity, 12 wks, median % change (95% CI): -52 (-59.7, -40.2)</p> <p>Nocturia, PGA median rate, % (95% CI): 50 (39.4, 57.6)</p> <p>Nocturia episodes /day, 12 wks, mean change ± SD: PGA < 50%: -0.6 ± 1.1 PGA ≥ 50%: -1.4 ± 0.9</p> <p>Nocturia, PPBC, mean change ± SD: PGA < 50%: -1.3 ± 0.9 PGA ≥ 50%: -2.3 ± 1.1 <i>P</i> < 0.01</p> <p>Nocturia, GII, %: PGA < 50%: No: 4 (20.0) Little: 13 (65.0) Much: 3 (15.0) <i>P</i> < 0.01 PGA ≥ 50%: No: 0 (0) Little: 6 (20.7) Much: 23 (79.3) <i>P</i> < 0.01</p> <p>Nocturia, pts want to continue treatment, n (%): PGA < 50%: 9 (45.0) PGA ≥ 50%: 26 (89.7) <i>P</i> < 0.01</p> <p>Nocturia severity, 12 wks, median % change (95% CI): -52 (-59.7, -40.2)</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Choo et al., 2008 (continued)				<p>Tenesmus severity, 12 wks, median % change (95% CI): -26 (-50.4, -16.9)</p> <p>Tenesmus, PGA median rate, % (95% CI): 30 (25.4, 52.2)</p> <p>Tenesmus, PPBC, mean change ± SD: PGA < 30%: -1.4 ± 0.9 PGA ≥ 30%: -2.1 ± 1.1</p> <p>Tenesmus, GII, %: PGA < 30%: No: 2 (15.4) Little: 9 (69.2) Much: 2 (15.4) <i>P</i> < 0.01 PGA ≥ 30%: No: 0 (0) Little: 5 (31.3) Much: 11 (68.8) <i>P</i> < 0.01</p> <p>Tenesmus, pts want to continue treatment, n (%): PGA < 30%: 8 (61.5) PGA ≥ 30%: 14 (87.5)</p> <p>Most troublesome symptom severity, median % change: UUI: -64.5 Urgency: -32 Frequency: -42.5 Nocturia: -56.5</p> <p>Voided volume (mL), 12 wks, mean ± SD: 189.3 ± 111.5 <i>P</i> = 0.05</p> <p>Dry mouth, n (%): 13 (21.7)</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Choo et al., 2008 (continued)				Constipation/ indigestion, n (%): 6 (10) Headache, n (%): 3 (5) UTI, n (%): 2 (3.3) Peripheral oedema, n (%): 1 (1.7)	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Colombo et al., 1995</p> <p>Country and setting: Italy; Setting</p> <p>Enrollment period: May 1990 to March 1993</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT, Computer generated random assignment</p> <p>Intervention: Oxybutynin vs. Bladder training for 6 weeks</p> <p>Groups: G1: Oxybutynin, 3 daily doses of 5 mg each for 6 weeks (dose reduced to half if substantial AEs) G2: Bladder training</p> <p>N at enrollment: G1: 42 G2: 39</p> <p>N at 6 wk follow-up: G1: 38 G2: 37</p> <p>N at 6 mo follow-up: G1: 28 G2: 27</p> <p>Age, yrs ± SD: G1: 48 (31 – 65) G2: 49 (24 – 65)</p> <p>Race/ethnicity: NR</p> <p>Women, N (%): G1: 42 (100) G2: 39 (100)</p> <p>Postmenopausal n (%): G1: 16 (38) G2: 20 (51)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Socially embarrassing (severe) urinary urge incontinence • On cystometry: detrusor instability, or low-compliance bladder (LCB), or sensory bladder <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Stable bladder at cystometry • Neurologic disease • Detrusor hyperreflexia • Age greater than 65 y • Coexisting genuine SUI • Genital prolapse • Postvoid residual volume >50mL • Previous gynecologic or urogynecologic surgery • Prior use of any drug to treat UUI • Urethral diverticula • Fistulas • Urinary tract neoplasia • Cystitis • Bladder stones • Previous pelvic radiotherapy 	<p>Detrusor instability, n (%): G1: 14 (37) G2: 13 (35)</p> <p>Low compliance bladder, n (%): G1: 9 (24) G2: 8 (22)</p> <p>Sensory bladder, n (%): G1: 15 (39) G2: 16 (43)</p> <p>Daily UUI episodes, range 9 – 17</p> <p>Diurnal frequency, n (%): G1: 32 (84) G2: 29 (78)</p> <p>Nocturia, n (%): G1: 11 (29) G2: 18 (49)</p> <p>Volume at first desire (mL): G1: 120 ± 59 G2: 134 ± 61</p> <p>Volume at very strong desire (mL): G1: 317 ± 92 G2: 332 ± 87</p>	<p>Clinically cured overall, n (%): G1: 28 (74) G2: 27 (73)</p> <p>Cured among DI, (%): G1: 13 (93) G2: 8 (62) <i>P</i> = 0.07</p> <p>Cured among LCB: G1: 6 (67) G2: 6 (75) <i>P</i> = 0.56</p> <p>Cured among sensory bladder: G1: 9 (60) G2: 13 (81) <i>P</i> = 0.18</p> <p>Cured among 18 G1 patients requiring dosage halving: 12 (67%)</p> <p>Diurnal frequency resolved, n (%): G1: 18 (56) G2: 20 (69)</p> <p>Nocturia resolved n (%): G1: 3 (27) G2: 11 (61)</p> <p>Volume at first desire (mL), mean ± SD: G1: 179 ± 32 <i>P</i> = .0009 G2: 178 ± 49 <i>P</i> = .001</p> <p>Volume at very strong desire (mL) mean ± SD: G1: 408 ± 76 <i>P</i> = .00001 G2: 403 ± 69 <i>P</i> = .0002</p> <p>Patients still cured at 6 mos,n: G1: 16 G2: 26</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Masking: -</p> <p>Pt selection criteria: +</p> <p>Loss to followup: ++</p> <p>Drop-out rates: +</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: -</p> <p>Measurement methods: +</p> <p>Measurement reliability: -</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Colombo et al., 1995 (continued)				<p>Patients still cured at 6 mos among DI, n: G1: 8 G2: 8</p> <p>Patients still cured at 6 mos among LCB, n: G1: 4 G2: 6</p> <p>Patients still cured at 6 mos among sensory bladder, n: G1: 4 G2: 12</p> <p>Treatment discontinued in 6 cases: G1: 4 (3 cases of severe dry mouth, 1 case of previously unknown glaucoma) G2: 2 (treatment was time consuming)</p> <p>Other adverse effects G1: 18 (47%) with AE requiring halving of dosage: -dry mouth (n=15) -constipation (n=6) -nausea (n=5) -dizziness (n=2) - decrease in visual acuity (n=1) - tachycardia (n=1)</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Davila, et al., 2001</p> <p>Country and setting: US, Specialty treatment centers</p> <p>Enrollment period: 6 weeks</p> <p>Funding: Watson Laboratories, Inc</p> <p>Author industry relationship disclosures: 2 of 3 Pharmadigm (1) Watson Laboratories (2)</p>	<p>Design: RCT</p> <p>Intervention: Transdermal Oxybutynin vs. Oral Oxybutynin 2 week washout</p> <p>Groups: G1: Transdermal OXY 1.3mg/day G2: OXY 2.5mg b.i.d./t.i.d.</p> <p>N at enrollment: G1: 38 G2: 38</p> <p>N at follow-up: G1: 38 G2: 36</p> <p>Women, n (%): G1: 33 (87) G2: 37 (97)</p> <p>Age, mean ± SD: G1: 64 ± 15 G2: 63 ± 13</p> <p>Race/ethnicity, n (%): White: 72 (95) Black: 4 (5)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 18 • History of UUI or MUI with predominant urge symptoms • ≥ 3 episodes UI/day with ≥ 30% increase in UUI episodes/ day during 2 week washout • Diagnosis of motor urge incontinence • Improvement during ≥ 6 wks with oral oxybutynin • UDS confirmed motor urge incontinence <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Allergy to oxybutynin • Intolerability of transdermal system • Pregnancy or breastfeeding • Overflow incontinence secondary to underactive or non-contractile detrusor • BOO • Impaired bladder compliance including tonic increase in pressure > 15 cmH₂O during filling cystometry • Medical conditions or drug therapies that could cause incontinence 	<p>Incontinence episodes/day, mean ± SD: G1: 7.2 ± 4.5 G2: 7.2 ± 4.1</p> <p>Bladder volume (mL), first contraction, mean ± SD: G1: 165 ± 158 G2: 267 ± 187</p> <p>Max cystometric capacity (mL), mean ± SD: G1: 244 ± 168 G2: 342 ± 167</p>	<p>Incontinence episodes/day, mean ± SD: G1: 2.4 ± 2.4 G2: 2.6 ± 3.3 G1/BL: <i>P</i> < 0.001 G2/BL: <i>P</i> < 0.001 G1/G2: <i>P</i> = NS</p> <p>Patient assessment of efficacy, VAS score, mean change ± SD: G1: 5.8 ± 4.2 G2: 6.0 ± 3.3 G1/BL: <i>P</i> < 0.001 G2/BL: <i>P</i> < 0.001 G1/G2: <i>P</i> = 0.9</p> <p>Bladder volume (mL), first contraction, mean ± SD: G1: 229 ± 189 G2: 302 ± 198 G1/BL: <i>P</i> = 0.0055 G2/BL: <i>P</i> = 0.1428 G1/G2: <i>P</i> = 0.57</p> <p>Max cystometric capacity (mL), mean ± SD: G1: 297 ± 176 G2: 387 ± 162 G1/BL: <i>P</i> = 0.0011 G2/BL: <i>P</i> = 0.0538 G1/G2: <i>P</i> = NR</p> <p>Dry mouth, n (%): G1: 15 (39) G2: 31 (82)</p> <p>Constipation, n (%): G1: 8 (21) G2: 19 (50)</p> <p>Somnolence, n (%): G1: 7 (18) G2: 14 (37)</p> <p>Nausea, n (%): G1: 3 (8) G2: 10 (26)</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: -</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: +</p> <p>Drop-out rates: -</p> <p>Power calculation: +</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: -</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Davila et al. 2001 (continued)		<ul style="list-style-type: none"> Medical conditions that could be worsened by oxybutynin 		<p>Dizziness, n (%): G1: 6 (16) G2: 10 (26)</p> <p>Blurred vision, n (%): G1: 7 (18) G2: 9 (24)</p> <p>Urinary retention, n (%): G1: 9 (24) G2: 13 (34)</p> <p>Impaired vision, n (%): G1: 9 (24) G2: 9 (24)</p> <p>Palpitation, n (%): G1: 3 (8) G2: 5 (13)</p> <p>Dry mouth by unvalidated questionnaire, %: G1: 38 G2: 94 <i>P</i> < 0.001</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Diokno et al., 1995</p> <p>Country and setting: US, private clinic</p> <p>Enrollment period: January 1992 to December 1992</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Cohorts with comparison; (Series of patients self selected into two groups)</p> <p>Intervention: Bladder training vs. anticholinergic or antispasmodic</p> <p>Groups: G1: Bladder training G2: oxybutynin 2.5-5.0 mg b.i.d. or t.i.d.</p> <p>N at enrollment: G1: 39 G2: 33</p> <p>N at follow-up: G1: 26 G2: 28</p> <p>Age, mean (range): 64 (20-93)</p> <p>Race/ethnicity: NR</p> <p>Women, N (%): 72 (100)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> Incontinent by AHCPR guidelines* Incontinence persisted after treatment for transient incontinence <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Post-void residual of more than 150 mL 	<p>UII, n (%): G1: 27 (49) G2: 28 (51)</p> <p>MUI with predominant urge, n (%): G1: 12 (71) G2: 5 (29)</p>	<p>Continent, n (%): G1: 1 (4) G2: 1 (4)</p> <p>Improved, n (%): G1: 22 (85) G2: 19 (68)</p> <p>No Change, n (%): G1: 3 (11) G2: 8 (28)</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: NA</p> <p>Masking: NA</p> <p>Pt selection criteria: -</p> <p>Loss to followup: -</p> <p>Drop-out rates: NR</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: poor</p> <p>Age: -, NR</p> <p>Baseline OAB status: NR</p> <p>Baseline characteristics: -</p> <p>Length of followup: ++</p> <p>Measurement methods: -</p> <p>Measurement reliability: -</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Diokno et al., 2002</p> <p>Country and setting: US, Community-dwelling</p> <p>Enrollment period: NR</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: 1 of 7 ALZA Corp (1)</p>	<p>Design: Multicenter, open-label, prospective cohort with 12 mos follow-up</p> <p>Intervention: Oxybutynin chloride ER</p> <p>Groups: NA</p> <p>N at enrollment: 1,067</p> <p>N at follow-up: 493</p> <p>Women, n (%): 904 (84.7)</p> <p>Age, mean ± SD: 64 ± 14.4</p> <p>Age, n (%): < 40: 65 (6.1) 40-65: 415 (38.9) >65-74: 313 (29.3) ≥ 75: 274 (25.7)</p> <p>Race/ethnicity, n (%): White: 950 (89.0) Black: 82 (7.7) Hispanic: 28(2.6) Asian: 2 (0.2) Native American: 5 (0.5)</p> <p>Follow-up: 12 months</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Community-dwelling • Urge or mixed incontinence • Age ≥ 18 • Generally in good health <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • UTI • Prostatitis • Urinary tract obstruction • Bladder tumor • Bladder stone • Prostate cancer • Contraindication to study drug • Current drug or alcohol abuse • Pregnant or breastfeeding • Pre-existing severe GI narrowing • Taken an investigational drug ≤ 1 mo • Known allergy or hypersensitivity to oxybutynin chloride • Not able to swallow medication whole 	<p>Incontinence impact, mean score ± SD: 1.4 ± 0.72</p> <p>Sleep impact, mean score ± SD: 3.3 ± 1.67</p> <p>Bother score, estimated from figure, mean: 72</p>	<p>Incontinence impact, end of dose adjustment, mean change ± SD: -0.5 ± 0.70 <i>P</i> < 0.001</p> <p>Incontinence impact, 90 days, mean change ± SD: -0.6 ± 0.76 <i>P</i> < 0.001</p> <p>Incontinence impact, final measure, mean change ± SD: -0.5 ± 0.77 <i>P</i> < 0.001</p> <p>Sleep impact, end of dose adjustment, mean change ± SD: -1.0 ± 1.73 <i>P</i> < 0.001</p> <p>Sleep impact, 90 days, mean change ± SD: -1.2 ± 1.82 <i>P</i> < 0.001</p> <p>Sleep impact, final measure, mean change ± SD: -1.1 ± 1.81 <i>P</i> < 0.001</p> <p>Bother score, end of dose adjustment, estimated from figure, mean: 44 <i>P</i> < 0.001</p> <p>Bother score, 90 days, estimated from figure, mean: 42 <i>P</i> < 0.001</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor Randomization: NA Masking: NA Pt selection criteria: + Loss to followup: - Drop-out rates: - Power calculation: - Statistical issues: -</p> <p>EXTERNAL VALIDITY: fair Age: + Baseline OAB status: NR Baseline characteristics: - Length of followup: ++ Measurement methods: + Measurement reliability: + Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Diokno et al., 2002 (continued)				<p>Bother score, 12 mos, estimated from figure, mean: 39 $P < 0.001$</p> <p>Headache, %: 1-90 days: 6.2 91-180 days: 0.6 181-270 days: 0.4 271-364 days: 0.4 Total: 7.6</p> <p>Dizziness, %: 1-90 days: 5.0 91-180 days: 0.4 181-270 days: 0.2 271-364 days: 0 Total: 5.6</p> <p>Blurred vision, %: 1-90 days: 2.8 91-180 days: 0.4 181-270 days: 0.1 271-364 days: 0 Total: 3.3</p> <p>Somnolence, %: 1-90 days: 2.8 91-180 days: 0 181-270 days: 0.2 271-364 days: 0 Total: 3.0</p> <p>Confusion, %: 1-90 days: 1.1 91-180 days: 0.1 181-270 days: 0.2 271-364 days: 0 Total: 1.4</p> <p>Discontinued due to AEs, n (%): 256 (24)</p> <p>Discontinued due to lack of efficacy, n (%): 108 (10.1)</p> <p>Discontinued due to administrative reasons, n (%): 90 (8.4)</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Diokno et al., 2003*</p> <p>Chu et al., 2005†</p> <p>Country and setting: US, Academic medical center</p> <p>Enrollment period: November 2000 to October 2001</p> <p>Funding: ALZA Corporation, Ortho-McNeil Pharmaceuticals</p> <p>Author industry relationship disclosures: 5 of 7* ALZA (1) Indevus (1) Ortho-McNeil (4) Pharmacia (1) Watson (1) NR†</p>	<p>Design: RCT</p> <p>Intervention: Oxybutynin ER vs Tolterodine ER 12 week treatment period</p> <p>Groups: G1: Oxybutynin ER 10 mg daily G2: Tolterodine ER 4mg daily</p> <p>N at enrollment: G1: 391 G2: 399</p> <p>N at follow-up: G1: 339 G2: 357</p> <p>Age, mean (range) : G1: 60 (23-92) G2: 60 (18-85)</p> <p>Race/ethnicity, N (%): White: G1: 329 (84) G2: 342 (86) Black: G1: 32 (8) G2: 35 (9) Hispanic: G1: 27 (7) G2: 19 (5) Other: G1: 3 (1) G2: 3 (1)</p> <p>Women, N (%): G1: 391 (100) G2: 399 (100)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 18 • ≥ 21 episodes UUI/ week • ≥ 10 voids/ day • Urge > non-urge incontinence <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Treatable GU conditions causing incontinence • PVR > 150 mL by ultrasound x 2 • Risk of developing complete urinary retention • Medical problems worsened by anticholinergic effects • Hematuria • Uncontrolled narrow angle glaucoma • Obstructive uropathy • Reduced GI motility • Hypersensitivity to medications 	<p>UUI episodes/wk, mean ± SD: G1: 37.2 ± 15.2 G2: 36.9 ± 14.1</p> <p>Total incontinence episodes/wk, mean ± SD: G1: 43.3 ± 19.3 G2: 42.6 ± 18.2</p> <p>Voids/wk, mean ± SD: G1: 94.8 ± 25.4 G2: 96.2 ± 24.5</p>	<p>UUI episodes/wk, mean: G1: 10.8 G2: 11.2 <i>P</i> = 0.28</p> <p>Incontinence episodes/wk, mean: G1: 12.3 G2: 13.8 <i>P</i> = 0.08</p> <p>Voids/wk, mean: G1: 66.4 G2: 71.1 <i>P</i> = 0.03</p> <p>UUI episodes/wk, % reduction ± SD: G1: 72.0 ± 34.0 G2: 70.2 ± 33.2 <i>P</i> = 0.13</p> <p>Incontinent episodes/wk, % reduction ± SD: G1: 72.8 ± 31.0 G2: 69.1 ± 34.9 <i>P</i> = 0.08</p> <p>Voids/wk, % reduction ± SD: G1: 27.7 ± 19.7 G2: 24.9 ± 18.9 <i>P</i> = 0.05</p> <p>Total dryness, %: G1: 23 G2: 16.8 <i>P</i> = 0.03</p> <p>No UUI episodes, %: G1: 26.7 G2: 20.9 <i>P</i> = 0.06</p> <p>Dry mouth, any degree, n (%): G1: 116 (29.7) G2: 89 (22.3) <i>P</i> = 0.02</p> <p>Dry Mouth, mild, n (%): G1: 87 (22.3) G2: 69 (17.3)</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: fair</p> <p>Randomization: -</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: ++</p> <p>Drop-out rates: ++</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Diokno et al., 2003* Chu et al., 2005† (continued)				<p>Dry mouth, moderate-severe, n (%): G1: 29 (7.4) G2: 20 (5.0)</p> <p>Constipation, n (%): G1: 25 (6.4) G2: 31 (7.8)</p> <p>Diarrhea, n (%): G1: 31 (7.9) G2: 25 (6.3)</p> <p>Headache, n (%): G1: 22 (5.6) G2: 24 (6.0)</p> <p>UTI, n (%): G1: 20 (5.1) G2: 13 (3.3)</p> <p>CNS AE, n (%):† G1: 35 (9.0) G2: 33 (8.3)</p> <p>Dizziness, n (%):† G1: 15 (3.8) G2: 10 (2.5)</p> <p>Dizziness, mild,** %:† G1: 1.8 G2: 1.5</p> <p>Dizziness, moderate,** %:† G1: 0.8 G2: 0.3</p> <p>Somnolence, n (%):† G1: 4 (1.0) G2: 9 (2.3)</p> <p>Somnolence, mild,** %:† G1: 0.5 G2: 1.5</p> <p>Somnolence, moderate,** %:† G1: 0.3 G2: 0.5</p> <p>Insomnia, n (%):† G1: 7 (1.8) G2: 3 (0.8)</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Diokno et al., 2003*				Insomnia, mild** %:† G1: 0.8 G2: 0	
Chu et al., 2005† (continued)				Insomnia, moderate,** %:† G1: 0.5 G2: 0	
				Depression, n (%) :† G1: 5 (1.3) G2: 3 (0.8)	
				Hypertonia, n (%) :† G1: 2 (0.5) G2: 4 (1.0)	
				Hypertonia, mild,** %:† G1: 0 G2: 0	
				Hypertonia, moderate,** %:† G1: 0.3 G2: 0	
				Anxiety, mild,** %:† G1: 0.5 G2: 0	
				Anxiety, moderate,** %:† G1: 0.3 G2: 0	
				Nervousness, mild,** %:† G1: 0 G2: 0	
				Nervousness, moderate,** %:† G1: 0.3 G2: 0	
				Tremor, mild,** %:† G1: 0.3 G2: 0.3	
				Tremor, moderate,** %:† G1: 0 G2: 0	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Diokno et al., 2003*				Confusion, mild,** %:† G1: 0.3 G2: 0.3	
Chu et al., 2005† (continued)				Confusion, moderate,** %:† G1: 0 G2: 0	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Dmochowski et al., 2002</p> <p>Country and setting: US, Community</p> <p>Enrollment period: NR</p> <p>2 week washout 12 week treatment period</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: 7 of 7 Abbot (1) Alza (3) Amgen (1) AstraZeneca (1) Bayer (1) Bioform (1) Genyx (1) Glaxo (1) Interneuron (2) Lilly (2) Merck (1) Otsulta (1) Pharmacia (3) Pfizer (1) Praecis (2) Roche (1) Seprecor (1) Surx (1) Synthelabo (1) Vivus (1) Watson (7) Yamanouchi (1)</p>	<p>Design: RCT Followed by: 12 week open label dose titration</p> <p>Intervention: Oxybutynin TDS vs. placebo</p> <p>Groups: G1: OXY TDS 1.3 mg G2: OXY TDS 2.6 mg G3: OXY TDS 3.9 mg G4: placebo</p> <p>N at enrollment: G1: 130 G2: 133 G3: 125 G4: 132</p> <p>N at follow-up: G1: 128 G2: 131 G3: 123 G4: 130</p> <p>Women, N (%): G1: 120 (92.3) G2: 123 (92.5) G3: 114 (91.2) G4: 121 (91.7)</p> <p>Age, mean ± SD: G1: 61.5 ± 11.8 G2: 61.9 ± 13.5 G3: 59.4 ± 14.5 G4: 62.7 ± 13.1</p> <p>Race/ethnicity, n (%): White: G1: 119 (91.5) G2: 118 (88.7) G3: 118 (94.4) G4: 118 (89.4) Black: G1: 7 (5.4) G2: 10 (7.5) G3: 3 (2.4) G4: 11 (8.3)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 18 • History of OAB • ≥ 10 episodes UUI either pure urge or predominant urge on 7 day voiding diary • ≥ 56 voids/ 7 day diary • Average recorded urinary volume of ≤ 350 mL <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Incontinence related to chronic illness • Concomitant medications • History of lower urinary tract surgery in ≤ 6 mos • IC • Urethral syndrome • Painful bladder syndrome • Overflow urinary incontinence • Alcohol/drug abuse • Known hypersensitivity to oxybutynin, similar compounds or transdermal systems • Active skin disorder • Narrow angle glaucoma • Excessive caffeine intake > 5 cups/day 	<p>Incontinence episodes/week, median: G1: 31.0 G2: 30.0 G3: 31.0 G4: 30.0</p> <p>voids/day, mean ± SD: G1: 12.4 ± 2.9 G2: 12.1 ± 3.3 G3: 12.3 ± 3.3</p> <p>Voided volume (mL), mean: G1: 175 G2: 165 G3: 156 G4: 170</p> <p>Open Label: Incontinence episodes/week, open label, median: G1: 30.0 G2: 29.0 G3: 37.0 G4: NA</p> <p>Prior years incontinent, years, mean ± SD: G1: 9.1 ± 10.3 G2: 8.9 ± 8.8 G3: 9.9 ± 9.8 G4: 9.1 ± 9.1</p> <p>IIQ (QoL), total score, mean: G1: 167 G2: 161 G3: 144 G4: 160</p> <p>Prior anticholinergic treatment, n, (%): G1: 30 (23.1) G2: 28 (21.1) G3: 28 (22.4) G4: 27 (20.5)</p>	<p>Incontinence episodes/ week, median change: G1: -16.0 G2: -14.0 G3: -19.0 G4: -14.5 G3/G4: P < 0.0165</p> <p>voids/day, mean change ± SD: G1: -1.8 ± 2.6 G2: -1.8 ± 2.4 G3: -2.3 ± 2.5 G4: -1.7 ± 3.0 G3/G4: P = 0.0457</p> <p>Voided volume, mL, mean increase ± SD: G1: -2 G2: 19 G3: 24 G4: 6 G3/G4: P = 0.0063 G2/G4: P = 0.0157</p> <p>Open Label: Incontinence episodes/week, median change: G1: -18.0 G2: -17.0 G3: -19.0 G4: NA G3/G4: P = 0.0327</p> <p>IIQ (QoL), total score, mean ± SD: G1: NR G2: NR G3: 78.8 ± 51.9 G4: 94.7 ± 50.0 G3/G4: P = 0.0266</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: - Masking: + Pt selection criteria: + Loss to followup: + Drop-out rates: - Power calculation: + Statistical issues: + EXTERNAL VALIDITY: good</p> <p>Age: + Baseline OAB status: + Baseline characteristics: ++ Length of followup: + Measurement methods: + Measurement reliability: + Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Dmochowski et al., 2002 (continued)	Other: G1: 4 (3.1) G2: 5 (3.8) G3: 4 (3.2) G4: 3 (2.3)			Dry mouth, n (%): G1: 6 (4.6) G2: 9 (6.8) G3: 12 (9.6) G4: 11 (8.3) Dizziness, n (%): G1: 2 (1.5) G2: 4 (3.0) G3: 5 (4.0) G4: 5 (3.8) Dysuria, n (%): G1: 1 (0.8) G2: 3 (2.3) G3: 3 (2.4) G4: 0 (0) Somnolence, n (%): G1: 1 (0.8) G2: 0 (0) G3: 2 (1.6) G4: 1 (0.8) Nausea, n (%): G1: 6 (4.6) G2: 5 (3.8) G3: 2 (1.6) G4: 7 (5.3) Constipation, n (%): G1: 7 (5.4) G2: 3 (2.3) G3: 1 (0.8) G4: 4 (3.0) Palpitations, n (%): G1: 1 (0.8) G2: 0 (0) G3: 1 (0.8) G4: 0 (0) Vision abnormal, n (%): G1: 3 (2.3) G2: 2 (1.5) G3: 0 (0) G4: 2 (1.5) Localized application site reactions, n (%): G1: 32 (26.4) G2: 7 (5.7) G3: 8 (6.9)	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Dmochowski et al., 2002 (continued)				<p>Application site erythema, n (%): G1: 23 (19) G2: 29 (23.6) G3: 14 (12.0)</p> <p>Application site erythema, mild, double blind period, n (%): G1: 79 (31.5) G2: 29 (36.2) G3: NA</p> <p>Application site erythema, moderate, double blind period, n (%): G1: 46 (18.3) G2: 46 (18.1) G3: NA</p> <p>Application site erythema, severe, double blind period, n (%): G1: 6 (2.4) G2: 8 (3.1) G3: NA</p> <p>Application site erythema, mild, open label, n (%): G1: 18 (34.6) G2: 58 (38.4) G3: 87 (43.9)</p> <p>Application site erythema, moderate, open label, n (%): G1: 7(13.5) G2: 23 (15.2) G3: 24 (12.1)</p> <p>Application site erythema, severe, open label, n (%): G1: 0 (0) G2: 2 (1.3) G3: 0 (0)</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Dmochowski et al., 2003</p> <p>Country and setting: US, Community</p> <p>Enrollment period: NR</p> <p>2 week washout 12 week treatment period</p> <p>Funding: Watson Pharma</p> <p>Author industry relationship disclosures: 6 of 6 Watson Pharma (6)</p>	<p>Design: RCT</p> <p>Intervention: Oxybutynin TDS vs. Tolterodine ER vs. placebo, 2 week washout plus 12 weeks treatment period; drug/placebo applied transdermally twice weekly to the abdomen and oral capsule ingested once daily</p> <p>Groups: G1: OXY TDS 3.9 mg/day G2: Tolterodine ER 4 mg daily G3: placebo</p> <p>N at enrollment: G1: 121 G2: 123 G3: 117 Total: 361</p> <p>N at follow-up: G1: NR G2: NR G3: NR Total: 320 (89%)</p> <p>Age, mean yrs ± SD: G1: 63.1 ± 12.0 G2: 62.9 ± 13.5 G3: 64.5 ± 12.3</p> <p>Race/ethnicity, n (%): White: G1: 111 (91.7) G2: 120 (97.6) G3: 110 (94.0) Black: G1: 8 (6.6) G2: 1(0.8) G3: 4 (3.4)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • ≥ 18yo men and women • Current pharmacological treatment for OAB with beneficial response • ≥4 episodes UUI episodes either pure urge or predominant urge on 3 day voiding diary • ≥ 24 voids/ 3 day diary • Average recorded urinary volume of ≤350 mL <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • History of lower urinary tract surgery in previous 6 months • IC • Urethral syndrome • Painful bladder syndrome • Overflow urinary incontinence 	<p>Incontinence episodes/ day, mean ± SD: G1: 4.7 ± 2.9 G2: 5.0 ± 2.9 G3: 5.0 ± 3.2</p> <p>Incontinence episodes/day, median: G1: 4 G2: 4 G3: 4</p> <p>Voids/day, mean ± SD: G1: 12.4 ± 2.9 G2: 12.1 ± 3.3 G3: 12.3 ± 3.3</p> <p>Voids/day, median: G1: 12 G2: 12 G3: 12</p> <p>Voided volume (mL), mean ± SD: G1: 165 ± 62 G2: 165 ± 61 G3: 175 ± 68</p> <p>Voided volume, mL, median: G1: 160 G2: 150 G3: 171</p> <p>Global assessment of disease, QOL, mean ± SD: G1: 65 ± 19 G2: 63 ± 20 G3: 65 ± 20</p> <p>IIQ (QOL), travel domain, mean ± SD: G1: 49 ± 28 G2: 47 ± 29 G3: 43 ± 29</p>	<p>Incontinence episodes/ day, mean ± SD: G1: 1.9 ± 2.7 G2: 1.9 ± 3.0 G3: 2.9 ± 3.8 G1/G3: P = 0.0137 G2/G3: P = 0.0011 G1/G2: P = 0.5878</p> <p>Incontinence episodes/day, mean change ± SD: G1: 2.9 ± 3.0 G2: 3.2 ± 2.8 G3: 2.1 ± 3.0</p> <p>Incontinence episodes/day, median G1: 1 G2: 1 G3: 2</p> <p>Incontinence episodes/day, median change G1: 3 G2: 3 G3: 2</p> <p>Voids/day, mean ± SD: G1: 10.4 ± 3.2 G2: 9.9 ± 3.1 G3: 10.9 ± 3.8 G1/G3: P = 0.101 G2/G3: P = 0.0025 G1/G2: P = 0.276</p> <p>Voids/day, mean change ± SD: G1: -1.9 ± 2.7 G2: -2.2 ± 2.6 G3: -1 ± 1.4</p> <p>Voids/day, median: G1: 10 G2: 10 G3: 10</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: - Masking: + Pt selection criteria: + Loss to followup: + Drop-out rates: + Power calculation: + Statistical issues: + EXTERNAL VALIDITY: good</p> <p>Age: + Baseline OAB status: + Baseline characteristics: ++ Length of followup: + Measurement methods: + Measurement reliability: + Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Dmochowski et al., 2003 (continued)	<p>Other: G1: 2 (1.6) G2: 2 (1.6) G3: 3 (2.6)</p> <p>Women, N (%): G1: 109 (90.1) G2: 117 (95.1) G3: 109 (93.2)</p> <p>Prior antimuscarinic treatment, Tolterodine, n, (%): G1: 57 (47) G2: 60 (49) G3: 54 (46)</p> <p>Prior antimuscarinic treatment, Oxybutynin, n, (%): G1: 61 (51) G2: 59 (48) G3: 59 (50)</p> <p>Prior antimuscarinic treatment, Other, n, (%): G1: 7 (6) G2: 6 (5) G3: 6 (5)</p>		<p>UDI (QOL), irritative symptoms, mean \pm SD: G1: 62 \pm 20 G2: 66 \pm 18 G3: 63 \pm 20</p>	<p>Voids/day, median change: G1: 2 G2: 2 G3: 1</p> <p>Voided volume, (mL), mean \pm SD: G1: 198 \pm 84 G2: 193 \pm 75 G3: 182 \pm 84 G1/G3: $P = 0.0010$ G2/G3: $P = 0.0017$ G1/G2: $P = 0.7690$</p> <p>Voided volume (mL), mean change \pm SD: G1: 32 \pm 55 G2: 29 \pm 57 G3: 9 \pm 63</p> <p>Voided volume (mL), median: G1: 188 G2: 189 G3: 165</p> <p>Voided volume (mL), median change: G1: 24 G2: 29 G3: 5.5</p> <p>Global assessment of disease, QOL, change \pm SD: G1: 30 \pm 30 G2: 33 \pm 28 G3: 21 \pm 31 G1/G3: $P = 0.0106$ G2/G3: $P = 0.001$ G1/G2: $P = 0.1861$</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Dmochowski et al., 2003 (continued)				<p>IIQ (QOL) travel domain, change \pm SD: G1: 23 \pm 25 G2: 22 \pm 29 G3: 11 \pm 30 G1/G3: <i>P</i> = 0.0018 G2/G3: <i>P</i> = 0.0045</p> <p>UDI (QOL), irritative symptoms, change \pm SD: G1: 25 \pm 26 G2: 28 \pm 26 G3: 18 \pm 24 G1/G3: <i>P</i> = 0.0156 G2/G3: <i>P</i> = 0.0010</p> <p>Treatment compliance with assigned dosage regimen: 92%:</p> <p>Frequency decreased to a greater extent for patients with > 14 micturations per day at baseline G1: -2.9/day, <i>P</i> = 0.0036 (data for other groups not reported)</p> <p>Adverse effects:</p> <p>Dry mouth, %: G1: 4.1 G2: 7.3 G3: 1.7 G1/G3: <i>P</i> = 0.2678 G2/G3: <i>P</i> = 0.0379</p> <p>Constipation, (%): G1: 3.3 G2: 5.7 G3: NR</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Dmochowski et al., 2003 (continued)				<p>Mild systemic adverse effects, n (%): G1: 15 (12.4) G2: 13 (10.6) G3: 6 (5.1)</p> <p>Moderate systemic adverse effects, n (%): G1: 7 (5.8) G2: 13 (10.6) G3: 7 (6.0)</p> <p>Severe systemic adverse effects, n (%): G1: 1 (0.8) G2: 3 (2.4) G3: 1 (0.9)</p> <p>Mild localized application site reactions, n (%): G1: 9 (7.4) G2: 2 (1.6) G3: 5 (4.3)</p> <p>Moderate localized application site reactions, n (%): G1: 17 (14.0) G2: 4 (3.3) G3: 2 (1.7)</p> <p>Severe localized application site reactions, n (%): G1: 6 (5.0) G2: 1 (0.8) G3: 1 (0.9)</p> <p>Treatment discontinuation due to adverse effects, n (%): G1: 13 (10.7) (12 due to application site reactions, 1 due to hot flushes) G2: 2 (1.6) (1 due to fatigue and 1 due to dizziness) G3: NR</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Dmochowski et al., 2003 (continued)				<p>Postvoid residual >150 mL at end of treatment: G1: 4 G2: 4 G3: 3 No reports of symptomatic urinary retention</p> <p>Withdraw due to AEs: 23</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Dmochowski et al., 2007</p> <p>Country and setting: International, multicenter</p> <p>Enrollment period: NR</p> <p>Funding: Pfizer Inc.</p> <p>Author industry relationship disclosures: 5 of 5 Astellas (2) Esprit (1) Lilly (1) Merck (1) Novartis (2) Ortho-McNeil (1) Pfizer (5) Watson (1)</p>	<p>Design: RCT (subanalysis)</p> <p>Intervention: Tolterodine ER vs. placebo</p> <p>Groups: G1: Tolterodine ER 4mg po qAM G2: Placebo qAM</p> <p>N at enrollment: Total: 1015 G1: 507 G2: 508</p> <p>N at follow-up: G1: 507 G2: 507</p> <p>Age, yrs (range): G1: 60 (20-89) G2: 61 (22-93)</p> <p>Race/ethnicity, %: White: G1: 95 G2: 94 Black: G1: 3 G2: 4 Other: G1: 1 G2: 2</p> <p>Women, %: 81</p> <p>Follow-up: 12 wks</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥18 • ≥ 8 voids/day and ≥ 5 episodes/wk UUI x 6 mos <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Hepatic, renal disease • UTI, recurrent UTIs • SUI • BOO • Indwelling catheter • Self catheterization • Taking anticholinergic medication • Mean voided volume > 200 mL or total daily voided volume > 3000 mL 	<p>UUI episodes/wk, mean ± SD: G1: 22.1 ± 22.3 G2: 23.3 ± 20.7</p> <p>UUI episodes/wk, 24:00-06:00, mean ± SD: G1: 3.5 ± 5.0 G2: 3.9 ± 5.0</p> <p>UUI episodes/wk, 06:00-12:00, mean ± SD: G1: 6.9 ± 8.1 G2: 6.7 ± 6.7</p> <p>UUI episodes/wk, 12:00-18:00, mean ± SD: G1: 6.9 ± 8.2 G2: 7.2 ± 6.9</p> <p>UUI episodes/wk, 18:00-24:00, mean ± SD: G1: 5.3 ± 8.1 G2: 5.5 ± 5.8</p> <p>Voids/day, mean ± SD: G1: 10.9 ± 4.2 G2: 11.3 ± 3.8</p> <p>Voids/day, 24:00-06:00, mean ± SD: G1: 1.7 ± 1.0 G2: 1.8 ± 1.1</p> <p>Voids/day, 06:00-12:00, mean ± SD: G1: 3.3 ± 1.4 G2: 3.4 ± 1.3</p> <p>Voids/day, 12:00-18:00, mean ± SD: G1: 3.1 ± 1.4 G2: 3.2 ± 1.3</p> <p>Voids/day, 18:00-24:00, mean ± SD: G1: 2.8 ± 1.3 G2: 2.9 ± 1.1</p>	<p>UUI episodes/wk, mean change ± SD: G1: -11.8 ± 17.8 G2: -6.9 ± 15.4 P ≤ 0.001 Benefit ratio 1.71</p> <p>UUI episodes/wk, 24:00-06:00, mean change ± SD: G1: -1.6 ± 4.0 G2: -1.0 ± 3.5 P ≤ 0.001 Benefit ratio 1.53</p> <p>UUI episodes/wk, 06:00-12:00, mean change ± SD: G1: -3.7 ± 7.1 G2: -2.0 ± 5.4 P ≤ 0.001 Benefit ratio 1.82</p> <p>UUI episodes/wk, 12:00-18:00, mean change ± SD: G1: -3.9 ± 7.1 G2: -2.2 ± 5.7 P ≤ 0.001 Benefit ratio 1.82</p> <p>UUI episodes/wk, 18:00-24:00, mean change ± SD: G1: -3.0 ± 6.9 G2: -1.6 ± 4.5 P ≤ 0.001 Benefit ratio 1.89</p> <p>Voids/day, mean change ± SD: G1: -1.8 ± 3.4 G2: -1.2 ± 2.9 P ≤ 0.001 Benefit ratio 1.46</p> <p>Voids/day, 24:00-06:00, mean change ± SD: G1: -0.22 ± 0.7 G2: -0.17 ± 0.8 P ≤ 0.05 Benefit ratio 1.29</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: +</p> <p>Drop-out rates: -</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Dmochowski et al., 2007 (continued)			<p>Voided volume (mL), mean ± SD: G1: 141 ± 43 G2: 136 ± 42</p> <p>Voided volume (mL), 24:00-06:00, mean ± SD: G1: 201 ± 99 G2: 190 ± 96</p> <p>Voided volume (mL), 06:00-12:00, mean ± SD: G1: 141 ± 52 G2: 139 ± 55</p> <p>Voided volume (mL), 12:00-18:00, mean ± SD: G1: 126 ± 49 G2: 119 ± 45</p> <p>Voided volume (mL), 18:00-24:00, mean ± SD: G1: 129 ± 50 G2: 123 ± 47</p>	<p>Voids/day, 06:00-12:00, mean change ± SD: G1: -0.57 ± 1.2 G2: -0.35 ± 1.0 <i>P</i> ≤ 0.001 Benefit ratio 1.60</p> <p>Voids/day, 12:00-18:00, mean change ± SD: G1: -0.55 ± 1.2 G2: -0.39 ± 1.0 <i>P</i> ≤ 0.001 Benefit ratio 1.42</p> <p>Voids/day, 18:00-24:00, mean change ± SD: G1: -0.43 ± 1.1 G2: -0.30 ± 0.8 <i>P</i> ≤ 0.002 Benefit ratio 1.43</p> <p>Voided volume (mL), mean change ± SD: G1: 34.1 ± 50.5 G2: 14.0 ± 41.3 <i>P</i> ≤ 0.001 Benefit ratio 2.44</p> <p>Voided volume (mL), 24:00-06:00, mean change ± SD: G1: 33.8 ± 105.5 G2: 16.7 ± 87.7 <i>P</i> ≤ 0.002 Benefit ratio 2.03</p> <p>Voided volume (mL), 06:00-12:00, mean change ± SD: G1: 38.2 ± 63.9 G2: 15.5 ± 58.6 <i>P</i> ≤ 0.001 Benefit ratio 2.47</p> <p>Voided volume (mL), 12:00-18:00, mean change ± SD: G1: 31.5 ± 59.4 G2: 12.4 ± 42.4 <i>P</i> ≤ 0.001 Benefit ratio 2.55</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Dmochowski et al., 2007 (continued)				Voided volume (mL), 18:00-24:00, mean change ± SD: G1: 29.2 ± 55.0 G2: 12.0 ± 49.6 <i>P</i> ≤ 0.001 Benefit ratio 2.44	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Dmochowski et al., 2008</p> <p>Country and setting: US, 62 sites</p> <p>Enrollment period: September 2005 to June 2006</p> <p>Funding: Esprit Pharma and Indevus Pharmaceuticals Inc.</p> <p>Author industry relationship disclosures: 4 of 4 Allergan (4) Astellas (3) Eli Lilly (1) Esprit Pharma (4) GSK (2) Indevus (4) Medtronic (1) Novartis (3) Ortho (2) Pfizer (2) Schwarz Pharma (1) Watson (4)</p>	<p>Design: RCT</p> <p>Intervention: Trospium vs. placebo</p> <p>Groups: G1: Trospium 60 mg qd G2: Placebo</p> <p>N at enrollment: G1: 280 G2: 284</p> <p>N at follow-up: G1: 267 G2: 276</p> <p>Women, n (%): G1: 230 (82.1) G2: 249 (87.7)</p> <p>Age, mean (SE): G1: 61.2 (0.7) G2: 58.4 (0.7)</p> <p>Race/ethnicity, n (%): White: G1: 245 (87.5) G2: 234 (82.4) Black: G1: 18 (16.4) G2: 28 (9.9) Hispanic: G1: 12 (4.3) G2: 14 (4.9) Asian: G1: 2 (0.7) G2: 3 (1.1) Other: G1: 3 (1.1) G2: 5 (1.8)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 18 • Symptoms of OAB ≥ 6 mos duration • Urinary frequency (mean of ≥ 10 toilet voids per day) • Urgency (1 or more episodes of severe urgency associated with a toilet void) • UUI (a mean of 1 or more UUI episodes per day) <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Total voided volumes > 3000 mL/day • Mean voided volume > 250 mL • Predominantly SUI • Insensate, or overflow incontinence • Neurogenic bladder • Indwelling or intermittent catheterization • Renal disease (serum creatinine >1.5 mg/dL) • Uninvestigated hematuria • UTI • History of > 3 UTIs in the previous 12 mos • Urinary retention (PVR > 100 mL) • Cancer • Interstitial cystitis 	<p>UUI episodes/day, mean (SE): G1: 4.0 (0.2) G2: 4.0 (0.2)</p> <p>Urgency severity, mean (SE): G1: 1.82 (0.03) G2: 1.83 (0.03)</p> <p>voids/day, mean (SE): G1: 12.8 (0.2) G2: 12.9 (0.2)</p> <p>Prior anticholinergic use for OAB, mean (SE): G1: 142 (50.7) G2: 156 (54.9)</p> <p>Voided volume (mL), mean (SE): G1: 149.1 (2.9) G2: 151.8 (2.8)</p>	<p>UUI episodes/day, 12 wks, mean change (SE): G1: -2.4 (0.2) G2: -1.6 (0.2) G1/BL: $P \leq 0.001$</p> <p>Urgency severity, 12 wks, mean change (SE): G1: -0.28 (0.03) G2: -0.13 (0.03) G1/BL: $P \leq 0.001$</p> <p>voids/day, 12 wks, mean change (SE): G1: -2.5 (0.2) G2: -1.8 (0.2) G1/BL: $P \leq 0.001$</p> <p>Voided volume (mL), 12 wks, mean change (SE): G1: 31.5 (3.4) G2: 17.8 (3.3) G1/BL: $P \leq 0.01$</p> <p>Treatment emergent AE, n (%): G1: 154 (55) G2: 130 (45.8)</p> <p>Mild to Moderate AEs, %: G1: 87% G2: 91.5%</p> <p>Dry mouth, n (%): G1: 36 (12.9) G2: 12 (4.6)</p> <p>Constipation, n (%): G1: 21 (7.5) G2: 5 (1.8)</p> <p>Headache, n (%): G1: 5 (1.8) G2: 6 (2.1)</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: - Masking: + Pt selection criteria: + Loss to followup: ++ Drop-out rates: + Power calculation: - Statistical issues: + EXTERNAL VALIDITY: good</p> <p>Age: + Baseline OAB status: + Baseline characteristics: ++ Length of followup: + Measurement methods: + Measurement reliability: + Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Dmochowski et al., 2008 (continued)		<ul style="list-style-type: none"> • PSA > greater than 4 ng/mL • Prostate cancer, or chronic prostatitis 			

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Drutz et al., 1999</p> <p>Country and setting: 25 centers United States and Canada</p> <p>Enrollment period: NR</p> <p>Funding: Pharmacia & Upjohn AB, Uppsala, Sweden</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: Tolterodine vs. oxybutynin vs. placebo</p> <p>Groups: 2 wk washout/run-in period G1: Tolterodine 2 mg b.i.d. x 12 wks G2: Oxybutynin 5 mg t.i.d. x 12 wks G3: placebo</p> <p>N at enrollment: G1: 109 G2: 112 G3: 56</p> <p>N at follow-up: Protocol correct G1: 70 G2: 41 G3: 36</p> <p>Age, mean range: G1: 63.0 (31-88) G2: 66.3 (23-91) G3: 62.1 (26-87)</p> <p>Women, %: G1: 81 G2: 91 G3: 80</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 18 yrs • Provided written informed consent • Post-menopausal • Surgically sterile • Adequate contraception • Detrusor overactivity on subtracted cystometry • Urinary frequency (58 voids on average per 24 hrs) and either urge incontinence (51 incontinence episode on average per 24 hrs <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • SU1 • Hepatic or renal disease • Recurrent UTIs • Interstitial cystitis • Hematuria or hematuria secondary to malignant disease • Indwelling catheter or intermittent catheterization • Treatment with investigational drug in ≤ 2 mos • Treatment with tolterodine or electro-stimulation therapy or BT ≤ 14 days 	<p>Duration of symptoms >5 years (%): G1: 47 (43) G2: 41 (37) G3: 25 (45)</p> <p>Instability hyperreflexia (%): G1: 101:8 (93:7) G2: 100:8 (89:7) G3: 53:3 (95:5)</p> <p>Urgency (%): G1: 104 (95) G2: 111 (99) G3: 55 (98)</p> <p>Previous drug therapy for urge incontinence, (%): G1: 49 (45) G2: 50 (45) G3: 31 (55)</p> <p>Good response to previous drug therapy, n (%): G1: 20 (41) G2: 20 (40) G3: 12 (39)</p> <p>Previous surgery affecting the lower urinary tract (%): G1: 29 (27) G2: 50 (45)* G3: 19 (34)</p> <p>Patients with incontinence episodes, n (%): G1: 90 (83) G2: 103 (92) G3: 50 (89)</p> <p>Incontinence episodes/day, mean (range): G1: 3.7 (0.1-19.9) G2: 3.4 (0.1-13.4) G3: 3.5 (0.1-14.9)</p>	<p>Incontinence episodes/day, mean change ± SD: G1: -1.7 ± 2.0 G2: -1.7 ± 1.7 G3: -1.0 ± 2.2 G1/G3: <i>P</i> = 0.063</p> <p>Mean difference (SEM): G1/G2: 0.0 (0.4) (95% CI) G1/G2: -0.7 to 0.7</p> <p>Voided volume (mL), mean change ± SD: G1: -2.0 ± (2.5) G2: -2.0 ± (2.3) G3: -1.1 ± (2.9) G1/G3: <i>P</i> = 0.036</p> <p>Mean difference (SEM): G1/G2: 0.0 (0.4) (95% CI) G1/G2: -0.8 to 0.8</p> <p>Voided volume (mL), mean change ± SD: G1: 34 ± 41 G2: 50 ± 43 G3: 12 ± 41 G1/G3: <i>P</i> = 0.0075</p> <p>Mean difference (SEM): G1/G2: NA (95% CI) G1/G2: NA</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: - Masking: + Pt selection criteria: + Loss to followup: - Drop-out rates: - Power calculation: + Statistical issues: + EXTERNAL VALIDITY: good</p> <p>Age: + Baseline OAB status: + Baseline characteristics: ++ Length of followup: + Measurement methods: + Measurement reliability: + Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Drutz et al., 1999 (continued)	<p>Race/ethnicity, %: Caucasian: G1: 87 G2: 94 G3: 93</p> <p>BMI, mean kg/m² (range): G1: 29.0 (18.3-56.2) G2: 28.0 (16.0-51.4) G3: 29.1 (17.2-52.8)</p>	<ul style="list-style-type: none"> • Treatment with anticholinergic drug, or any drug for urinary urge • Incontinence ≤ 14 days • Unstable dosage of any anticholinergic • Serious adverse effects on oxybutynin • Average total voided volume/24 hours of >3000 mL • Clinically significant voiding difficulty with risk of urinary retention (such as residual volume >200 mL or urine flow rate <10 mL/s) 	<p>Patients with ≥ 8 voids/day, n (%): G1: 108 (99) G2: 110 (98) G3: 55 (98)</p> <p>Voids/day, mean (range): G1: 11.6 (7.7-22.0) G2: 11.5 (7.1-31.4) G3: 11.6 (6.6-21.9)</p> <p>Volume voided/void, mean mL (range): G1: 155 (48–290) G2: 149 (42–315) G3: 160 (33–371)</p>	<p>*<i>P</i> <0.05 vs. placebo and tolterodine treatment groups.</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Dwyer et al., 2008</p> <p>Country and setting: Multinational, multicenter (academic and private)</p> <p>Enrollment period: NR</p> <p>Funding: Novartis Pharma AG</p> <p>Author industry relationship disclosures: 6 of 7 Allergan (1) Amgen (1) AMS (1) Astellas (1) Bayer (1) GSK (1) Hospira Australia (1) Novartis (5) Ocean Spray (1) Ono (1) Pfizer (2) Spectrum (1) Sunumoto (1)</p>	<p>Design: Multicenter open label extension</p> <p>Intervention: Darifenacin 7.5 mg or 15 mg (all pts received 7.5 mg dose for 2 weeks followed by individual dose adjustments)</p> <p>Groups: G1: Darifenacin continuation group 7.5 or 15 mg daily</p> <p>N at enrollment: G1: 303</p> <p>N at follow-up: G1: 300</p> <p>Women, N (%): G1: 262 (86.5)</p> <p>Age, mean (range): G1: 57.1 (22-89)</p> <p>Race/ethnicity: NR</p> <p>Follow-up: 12 weeks followed by 2 year extension</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Successful completion of one of 2 previous, 12-wk feeder studies with no major protocol violation • “wet OAB” (urgency, frequency, and incontinence) x 6 months • Adequate contraception if of childbearing potential • Able to complete patient diaries independently • Capable of independent toileting <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Certain concomitant medications (see previous studies) • Max darifenacin dose of 7.5 mg qd for patients taking potent inhibitors of CYP450 3A4 and pts w/ moderate hepatic impairment (Child Pugh B) • Other active management strategies of OAB (bladder training) 	<p>Incontinence episodes/week, mean ± SD: G1: 20.8 ± 13.6</p> <p>Incontinence episodes/week, median: G1: 17.0</p> <p>KHQ incontinence impact score, mean: G1: 78.56</p> <p>KHQ severity score, mean: G1: 52.56</p> <p>KHQ role limitations score, mean: G1: 62.72</p> <p>KHQ physical limitations score, mean: G1: 62.67</p> <p>KHQ social limitations score, mean: G1: 34.28</p> <p>KHQ emotions score, mean: G1: 43.85</p> <p>KHQ sleep/energy score, mean: G1: 52.50</p> <p>KHQ personal relationship score, mean: G1: 26.94</p> <p>KHQ general health score, mean: G1: 27.25</p>	<p>Incontinence episodes/week, median % change: G1: -83.3 <i>P</i> < 0.001</p> <p>KHQ incontinence impact score, mean change: G1: -25.25 <i>P</i> < 0.001</p> <p>KHQ severity score, mean change: G1: -13.53 <i>P</i> < 0.001</p> <p>KHQ role limitations score, mean change: G1: -26.77 <i>P</i> < 0.001</p> <p>KHQ physical limitations score, mean change: G1: -26.35 <i>P</i> < 0.001</p> <p>KHQ social limitations score, mean change: G1: -15.51 <i>P</i> < 0.001</p> <p>KHQ emotions score, mean change: G1: -12.79 <i>P</i> < 0.001</p> <p>KHQ sleep/energy score, mean change: G1: -11.59 <i>P</i> < 0.002</p> <p>KHQ personal relationship score, mean change: G1: -8.68 <i>P</i> < 0.001</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: -</p> <p>Method and blinding: -</p> <p>Pt selection criteria: +</p> <p>Loss to followup: -</p> <p>Drop-out rates: -</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: ++</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Dwyer et al., 2008 (continued)				KHQ general health score, mean change: G1: 2.05 <i>P</i> = NS	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Elinoff et al., 2006[^]</p> <p>Roberts et al., 2006</p> <p>Country and setting: US, 82 Primary care & Ob-Gyn offices</p> <p>Enrollment period: NR</p> <p>Funding: Pfizer</p> <p>Author industry relationship disclosures: 6 of 6 Pfizer (6) Astellas (1) Novartis (1)</p>	<p>Design: Open-label, single-arm cohort</p> <p>Intervention: Tolterodine ER 4 mg qd for 12 wks</p> <p>Groups: NA</p> <p>N at enrollment: 896</p> <p>N at follow-up: 758</p> <p>Age, yrs ± SD: 56 ± 15</p> <p>Race/ethnicity, n (%): White: 691 (80) Black: 94 (11) Asian: 22 (3) Other: 53 (6)</p> <p>Women, N (%): 708 (82)</p> <p>Parity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 18 • OAB symptoms for ≥3 mos • ≥ 8 voids/day • 2+ episodes of urgency or UUI in a 3-d period • 3+ on the OAB Bother Rating Scale <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Stress, functional, or overflow incontinence • Acute UTI • Clinically significant LUT pathology • Indwelling catheter • Intermittent self-catherization • Use of anticholinergics, antimuscarinics for OAB, drugs to treat UUI, potent cytochrome P450 3A4 inhibitors, or investigational drugs 	<p>Duration of OAB (yrs), mean ± SD: 6 ± 8</p> <p>Duration on study drug (days), mean ± SD: 80 ± 19</p> <p>Daytime voids/day, mean ± SD: 9.3 ± 2.8</p> <p>Nocturia episodes/day, mean ± SD: 3.0 ± 1.7</p> <p>UUI episodes/day, mean ± SD: 2.7 ± 3.2</p> <p>Urgency episodes/day, mean ± SD: 5.0 ± 3.8</p> <p>OAB-q scores: Symptom bother: 60.0 Coping: 50.0 Concern: 51.4 Sleep: 44.4 Social Interaction: 84.0 Total HRQL: 55.2</p> <p>AUA-SI: Total: 18 Irritative: 10 Obstructive: 8</p> <p>Most bothersome symptom, n (%) Daytime frequency: 256 (30) Nocturnal frequency: 241 (28) UUI: 206 (24) Urgency: 154 (18)</p>	<p>UUI episodes/day, week 12, mean % change (95% CI):[^] -86.1 (-91.7, -80.0)*</p> <p>UUI episodes/day, week 12, UUI most bothersome, mean % change (95% CI):[^] -80.0 (-85.7, -69.2)*</p> <p>Urgency episodes/day, week 12, mean % change (95% CI):[^] -75.0 (-80.0, -71.4)*</p> <p>Urgency episodes/day, week 12, urgency most bothersome, mean % change (95% CI):[^] -78.4 (-83.3, -72.2)*</p> <p>Nocturia episodes/day, week 12, mean % change (95% CI):[^] -37.5 (-40.0, -33.3)*</p> <p>Nocturia episodes/day, week 12, nocturia most bothersome, mean % change (95% CI):[^] -40.0 (-42.9, -33.3)*</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: NA</p> <p>Masking: NA</p> <p>Pt selection criteria: +</p> <p>Loss to followup: +</p> <p>Drop-out rates: +</p> <p>Power calculation: +</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Elinoff et al., 2006 Roberts et al., 2006 (continued)				<p>Daytime voids/ day, week 12, mean % change (95% CI):[^] -29.0 (-31.0, -27.3)*</p> <p>Daytime voids/ day, week 12, daytime voids most bothersome, mean % change (95% CI):[^] -30.4 (-33.3, -27.3)*</p> <p>OAB-q scores, 12 weeks, median change (95% CI):[*] Symptom bother: -37.5 (-37.5, -35.0) Coping: 32.5 (30.0, 35.0) Concern: 34.3 (31.4, 37.1) Sleep: 28.0 (28.0, 32.0) Social Interaction: 12.0 (12.0, 16.0) Total HRQL: 28.9 (27.2, 31.2)</p> <p>AUA-SI, Median change from baseline to 12 wks (95% CI):[*] Total: -9 (-9, -8) Irritative: -5 (-5, -5) Obstructive: -4 (-4, -3) *<i>P</i> < 0.0001 vs. baseline</p> <p>All-cause AE, %: 51</p> <p>Discontinued treatment, n %: 15 (7)</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Elinoff et al., 2006				Trt-related AE, %: 23	
Roberts et al., 2006 (continued)				Dry mouth, %: 10.0 Constipation, %: 3.7 Headache, %: 3.0 UTI, %: 2.7	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Fantl et al., 1981</p> <p>Country and setting: US, University</p> <p>Enrollment period: N/A</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Retrospective chart review</p> <p>Intervention: Bladder retraining drill (BRD) and anticholinergics when BRD not satisfactory</p> <p>Groups: G1: Detrusor muscle contraction approximately 5 seconds after a cough G2: Detrusor muscle contracts spontaneously without prior provocation G3: Both cough contraction and spontaneous contraction</p> <p>N at enrollment: G1: 39 G2: 4 G3: 49</p> <p>N at follow-up: NA</p> <p>Women, %: 100</p> <p>Age, mean ± SD: 42.7 ± 10:3</p> <p>Race/ethnicity: NR</p> <p>Parity, mean: 3.5</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> Cystometric tracings showed a rise of 15 cm water or more in intravesical pressure when recorded independently and calculated by subtraction from intra-abdominal transrectal pressures <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Neuropathic conditions UTI 	<p>Urgency and frequency, n (%): 72 (78.2)</p> <p>Urinary incontinence, n (%): 65 (70.6)</p> <p>Sensation of incomplete voiding, n (%): 59 (64.1)</p> <p>Nocturia, n (%): 66 (71.7)</p> <p>Dysuria, n (%): 3 (3.2)</p> <p>MUI, n (%): 6 (6.5)</p> <p>Enuresis in childhood, n (%): 5 (5.4)</p> <p>Previously operated on for similar urologic symptoms, n (%): 34 (36.9)</p>	<p>Cured, n (%): G1: 32 (82.1) G2: 1 (25) G3: 41 (83.7)</p> <p>Cured by BRD alone, n (%): 44 (78.6)</p> <p>Cured by BRD + anticholinergics, n (%):* 30 (83.5)</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: NA</p> <p>Masking: NA</p> <p>Pt selection criteria: +</p> <p>Loss to followup: NR</p> <p>Drop-out rates: NR</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: NA</p> <p>Measurement methods: +</p> <p>Measurement reliability: -</p> <p>Intervention description: -</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Garely et al., 2006</p> <p>Country and setting: US, 207 centers</p> <p>Enrollment period: June 2004 to April 2005</p> <p>Funding: Astellas Pharma US, Inc. and GlaxoSmithKline</p> <p>Author industry relationship disclosures: 2 of 5 Astellas (2)</p>	<p>Design: Phase IIIb trial</p> <p>Intervention: Treatment with 5 mg solifenacin succinate. Dosage could be increased to 10 mg, maintained or decreased in response to pt perceived efficacy and tolerability</p> <p>N at enrollment: 2,205</p> <p>N at follow-up: 1,743</p> <p>Women, N (%): 1813 (82.2)</p> <p>Age, mean ± SD: 59.7 ± 14.4</p> <p>Race/ethnicity, n (%): White: 1,761 (79.9) Black: 274 (12.4) Other: 170 (7.7)</p> <p>Weight, mean ± SD: 182.2 ± 47.0</p> <p>Follow-up: 12 weeks</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • ≥ 18 yrs • OAB symptoms ≥ 3 months • Toilet without difficulty • Other OAB meds with washout period ≥ 7 days • Non-drug treatment of OAB if established ≥ 4 wks prior to study and continued <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • SUI • Stress predominant MUI • UTI or chronic inflammation • Outflow obstruction due to benign prostatic hyperplasia • Uncontrolled narrow-angle glaucoma • Urinary or gastric retention • Severe renal or hepatic impairment • Chronic severe constipation or diagnosed gastrointestinal obstructive disease • Bladder cancer • Pregnant or not using reliable method of birth control 	<p>UUI, physician assessment, n (%): 1,586 (71.9)</p> <p>Urgency, physician assessment, n (%): 2,007 (91.0)</p> <p>Frequency, physician assessment, n (%): 1,969 (89.3)</p> <p>Nocturia, physician assessment, n (%): 1,792 (81.3)</p> <p>Perception of bladder condition, mean: 4.4</p> <p>Most bothersome OAB symptom, n (%): Frequency: 619 (28.1) UUI: 602 (27.3) Urgency: 508 (23.0) Nocturia: 337 (15.3) None: 139 (6.3)</p> <p>Duration of OAB, n (%): 3 mo to 1 year: 349 (15.8) > 1 to 5 years: 1,124 (51.0) > 5 years: 732 (33.2)</p>	<p>Flexible drug admin dose, wk 4, n (%): Remained at 5 mg/d: 941 (45.3) 5 mg/d to 10 mg/d: 1,076 (51.8) Discontinued: 60 (2.9)</p> <p>Flexible drug admin dose, wk 8, n (%): Remained at 5 mg/d: 782 (43.3) 5 mg/d to 10 mg/d: 166 (17.6) Remained at 10 mg/d: 1,018 (56.4) 10 mg/d to 5 mg/d: 491 (8.5) Discontinued: 4 (0.2)</p> <p>Perception of bladder condition, wk 4, mean: 3.3 <i>P</i> < 0.001</p> <p>Perception of bladder condition, wk 12 early termination, mean: 2.9 <i>P</i> < 0.001</p> <p>Perception of bladder condition, endpoint, mean: 2.9 <i>P</i> < 0.001</p> <p>Urgency bother, VAS score (mm), mean (n): 29.1 (1,781)</p> <p>UUI bother, VAS score (mm), mean (n): 24.0 (1,504)</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: NA</p> <p>Masking: NA</p> <p>Pt selection criteria: +</p> <p>Loss to followup: -</p> <p>Drop-out rates: -</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Garely et al., 2006 (continued)		<ul style="list-style-type: none"> Known hypersensitivity to study medication, its components, or to anticholinergic medication 		<p>Frequency bother, VAS score (mm), mean (n): 28.3 (1,751)</p> <p>Nocturia bother, VAS score mean mm (n): 28.3 (1,659)</p> <p>OAB-q symptom severity subscale, mean change (95% CI): -29.6 (-30.7, -28.6) <i>P</i> < 0.001</p> <p>OAB-q coping subscale, mean change (95% CI): 27.4 (26.2, 28.5) <i>P</i> < 0.001</p> <p>OAB-q concern subscale, mean change (95% CI): 29.6 (28.4, 30.8) <i>P</i> < 0.001</p> <p>OAB-q sleep subscale, mean change (95% CI): 27.3 (26.1, 28.5) <i>P</i> < 0.001</p> <p>OAB-q social interaction subscale, mean change (95% CI): 14.7 (13.7, 15.6) <i>P</i> < 0.001</p> <p>OAB-q overall health-related QoL subscale, mean change (95% CI): 25.4 (24.4, 26.4) <i>P</i> < 0.001</p> <p>Adverse events, n (%): 1,321 (59.4)</p> <p>Any treatment-related AE, n (%): 928 (41.7)</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Garely et al., 2006 (continued)				<p>Dry mouth, n (%): Mild: 352 (15.8) Moderate: 100 (4.5) Severe: 25 (1.1)</p> <p>Constipation, n (%): Mild: 190 (8.5) Moderate: 85 (3.8) Severe: 20 (0.9)</p> <p>Headache, n (%): 76 (3.4)</p> <p>Blurred vision, n (%): Mild: 41 (1.8) Moderate: 15 (0.7) Severe: 1 (0)</p> <p>Dry eye, n (%): 29 (1.3)</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Garely et al., 2007*</p> <p>Mallett et al., 2007^</p> <p>Country and setting: US, multicenter (207 sites)</p> <p>Enrollment period: NR</p> <p>Funding: Astellas Pharma US Glaxo-SmithKline</p> <p>Author industry relationship disclosures: 4 of 4* Astellas (4) Pfizer (1) Novartis (1)</p> <p>4 of 4^ Astellas (3) GlaxoSmithKline (2) Novartis (1) Watson (1) Yamanouchi (1)</p>	<p>Design: Prospective case series</p> <p>Intervention: Flexibly dosed, once daily solifenacin (started on 5 mg/d; option of increasing to 10 mg/d at week 4; option to maintain 10 mg dose or decrease to 5 mg at wk 8); for 12 weeks of treatment</p> <p>10 mg dose, %: 4 wks: G1: 55 8 wks: G1: 59 (8% went back to 5mg; 19% increased to 10 mg)</p> <p>Groups: G1: Urge incontinence as most bothersome symptom G2: Black participants G3: Full study population</p> <p>N at enrollment: G1:582 G2: 274 G3: 2205</p> <p>Age, mean yrs ± SD: G1: 60.9 ± 13.1 G2: 54.7 ± 13.6 G3: 59.7 ± 14.4</p> <p>Race/ethnicity, n (%): White: G1: 493 (84.7) G3: 1761 (79.9) African American: G1: 62 (10.7) G3: 274 (12.4)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 18 • Symptoms of OAB for ≥ 3 mos • Ambulatory • Able to use the toilet without difficulty • Urgency, UUI, frequency or nocturia ≥ 3 mos <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Previous use of solifenacin 	<p>UUI, n (%): G1: 582 (100%) G2: NR G3: 1596 (71.9%)</p> <p>Urgency, n (%): G1: 534 (91.8) G2: NR G3: 2007 (91.0)</p> <p>Frequency, n (%): G1: 488 (83.8) G2: NR G3: 1969 (89.3)</p> <p>Nocturia, n (%): G1: 438 (75.3) G2: NR G3: 1792 (81.3)</p> <p>Most bothersome OAB symptom</p> <p>Urgency, n (%): G2: 48 (17.5) G3: 508 (23.0)</p> <p>UUI, n (%): G1: 582 (100) G2: 63 (23.0) G3: 602 (27.3)</p> <p>Frequency, n (%): G2: 106 (38.7) G3: 619 (28.1)</p> <p>Nocturia, n (%): G2: 34 (12.4) G3: 337 (15.3)</p> <p>None specified, n (%): G2: 23 (8.4) G3: 139 (6.3)</p> <p>OAB for 3 mos -1 yr, n (%): G1: 57 (9.8) G3: 349 (15.8%)</p> <p>OAB for 1-5 yrs, n (%): G1: 294 (50.5) G3: 1124 (51.0)</p> <p>OAB for >5 yrs: G1: 231 (39.7) G3: 732 (33.2)</p>	<p>Patient perception of bladder condition scale, 4 wks, mean: G1: 3.3* G2: 3.1* G3: 3.3*</p> <p>Patient Perception of Bladder Condition Scale, 8 wks, mean: G1: 3.0* G2: 2.7* G3: 2.9*</p> <p>Patient Perception of Bladder Condition Scale, 12 wks or early termination, mean: G1: 2.9* G2: 2.6* G3: 2.9*</p> <p>Symptom severity, mean VAS, wk 4: Urinary urgency: G1: 40.9 G2: 39.7 G3: 40.6</p> <p>UUI: G1: 36.9 G2: 29.9 G3: 32.8</p> <p>Frequency: G1: 34.5 G2: 39.3 G3: 39.9</p> <p>Nocturia: G1: 30.4 G2: 35.4 G3: 38.2</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: NA</p> <p>Masking: NA</p> <p>Pt selection criteria: -</p> <p>Loss to followup: NR</p> <p>Drop-out rates: +</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Garely et al., 2007*	Other: G1: 27 (4.6) G3: 170 (7.7)		Perception of bladder condition scale (mean): G1: 4.6 G2: 4.2 G3: 4.4	Symptom severity, mean VAS, wk 8: Urinary urgency: G1: 31.1 G2: 29.4 G3: 30.0	
Mallett et al., 2007^ (continued)	Women, n (%): G1: 536 (92.1) G2: 213 (77.7) G3: 1813 (82.2) Weight (lbs), mean ± SD (range): G2: 198 ± 51.6 (100-400) G3: 182.2 ± 47.0 (80-413)		Symptom severity, mean VAS: Urinary urgency: G1: 72.3 G2: 70.8 G3: 68.7 UUI: G1: 78.5 G2: 70.8 G3: 64.1 Frequency: G1: 65.7 G2: 70.8 G3: 70.6 Nocturia: G1: 57.9 G2: 66.5 G3: 65.2 OAB-q, symptom severity, mean (SE): G1: 63.1 (0.84) G2: 59.4 (NR) G3: 56.9 (NR) OAB-q, coping, mean (SE): G1: 48.9 (1.18) G2: 46.5 (NR) G3: 53.1 (NR) OAB-q, concern, mean (SE) : G1: 43.1 (1.12) G2: 46.6 (NR) G3: 50.8 (NR) OAB-q, sleep, mean (SE): G1: 55.2 (1.22) G2: 43.8 (NR) G3: 49.2 (NR) OAB-q, social, mean (SE): G1: 73.6 (1.14) G2: 68.9 (NR) G3: 76.0 (NR)	Urinary urgency: G1: 28.9 G2: 22.1 G3: 24.5 Frequency: G1: 24.9 G2: 27.0 G3: 28.6 Nocturia: G1: 25.4 G2: 36.4 G3: 29.0 Symptom severity, mean VAS, wk 12 (or early termination): Urinary urgency: G1: 27.5 G2: 25.3 G3: 28.0 UUI: G1: 24.3 G2: 20.0 G3: 22.7 Frequency: G1: 21.6 G2: 24.5 G3: 27.4 Nocturia: G1: 22.0 G2: 21.4 G3: 27.2	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Garely et al., 2007* Mallett et al., 2007^ (continued)			OAB-q, HRQoL, mean (SE): G1: 53.4 (1.01) G2: 50.5 (NR) G3: 56.3 (NR)	Symptom severity, VAS, mean change from baseline (95% CI) p-value: Urinary urgency*: G1: -43.1 (-45.8, -40.4) p < 0.001 G2: -43.0 (-47.6, -38.5) G3: -39.5 (-41.0, -38.1) p < 0.001 UUI*: G1: -51.7 (-54.5, -49.0) p < 0.001 G2: -42.3 (-47.8, -36.8) G3: -40.1 (-41.8, -38.4) p < 0.001 Frequency*: G1: -42.0 (-45.0, -39.0) p < 0.001 G2: -44.2 (-48.8, -39.6) G3: -41.8 (-43.3, -40.3) p < 0.001 Nocturia*: G1: -34.4 (-37.3, -31.5) p < 0.001 G2: -42.6 (-47.5, -37.8) G3: -36.9 (-38.4, -35.4) p < 0.001 OAB-q, symptom severity, mean change (\pm SE or 95% CI)*: G1: -35.9 \pm 1.08 G2: -33.6 (-37.0, -30.3) G3: -29.6 (-30.7, -28.6) OAB-q, coping, mean change (\pm SE or 95% CI)*: G1: 32.5 \pm 1.16 G2: 32.9 (29.3, 36.6) G3: 27.4 (26.2, 28.5)	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Garely et al., 2007*				OAB-q, concern, mean change (\pm SE or 95% CI)*: G1: 37.0 \pm 1.21 G2: 34.0 (30.4, 37.6) G3: 29.6 (28.4, 30.8)	
Mallett et al., 2007^ (continued)				OAB-q, sleep, mean change (\pm SE or 95% CI)*: G1: 26.0 \pm 1.19 G2: 33.8 (29.7, 37.9) G3: 27.3 (26.1, 28.5) UUI*: G1: -51.7 (-54.5, -49.0) p < 0.001 G2: -42.3 (-47.8, -36.8) G3: -40.1 (-41.8, -38.4) p < 0.001 Frequency*: G1: -42.0 (-45.0, -39.0) p < 0.001 G2: -44.2 (-48.8, -39.6) G3: -41.8 (-43.3, -40.3) p < 0.001 Nocturia*: G1: -34.4 (-37.3, -31.5) p < 0.001 G2: -42.6 (-47.5, -37.8) G3: -36.9 (-38.4, -35.4) p < 0.001	
				OAB-q, symptom severity, mean change (\pm SE or 95% CI)*: G1: -35.9 \pm 1.08 G2: -33.6 (-37.0, -30.3) G3: -29.6 (-30.7, -28.6)	
				OAB-q, coping, mean change (\pm SE or 95% CI)*: G1: 32.5 \pm 1.16 G2: 32.9 (29.3, 36.6) G3: 27.4 (26.2, 28.5)	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Garely et al., 2007* Mallett et al., 2007^ (continued)				<p>OAB-q, concern, mean change (\pm SE or 95% CI)*: G1: 37.0 \pm 1.21 G2: 34.0 (30.4, 37.6) G3: 29.6 (28.4, 30.8)</p> <p>OAB-q, sleep, mean change (\pm SE or 95% CI)*: G1: 26.0 \pm 1.19 G2: 33.8 (29.7, 37.9) G3: 27.3 (26.1, 28.5)</p> <p>OAB-q, social, mean change (\pm SE or 95% CI)*: G1: 17.7 \pm 1.03 G2: 20.1 (16.7, 23.5) G3: 14.7 (13.7, 15.6)</p> <p>OAB-q, HRQoL, mean change (\pm SE or 95% CI)*: G1: 29.6 \pm 1.02 G2: 30.8 (27.5, 34.1) G3: 25.4 (24.4, 26.4)</p> <p>Side effects, n (%): G1: 357 (61.3) G2: 128 (46.4) G3: 1321 (59.4)</p> <p>Dry mouth, n (%): G1: 104 (17.9) G2: 36 (13.0) G3: 477 (21.4)</p> <p>Constipation, n (%): G1: 85 (14.6) G2: 19 (6.9) G3: 295 (13.3)</p> <p>Nausea, n (%): G1: NR G2: 7 (2.5) G3: 39 (1.8)</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Garely et al., 2007* Mallett et al., 2007^ (continued)				<p>Headache, n (%): G1: 21 (3.6) G2: 9 (3.3) G3: 76 (3.4)</p> <p>Blurred vision, n (%): G1: 20 (3.4) G2: 7 (2.5) G3: 57 (2.6)</p> <p>Upper respiratory tract infection, n (%): G1: 27 (4.6) G2: 7 (2.5) G3: 69 (3.10)</p> <p>UTI, n (%) G1: 21 (3.6%) G2: NR G3: NR</p> <p>Rash, n (%): G1: NR G2: 6 (2.2) G3: 22 (0.99)</p> <p>Nasopharyngitis, n (%) G1: 13 (2.2%) G2: NR G3: NR</p> <p>Cough, n (%) G1: 12 (2.1%) G2: NR G3: NR</p> <p>Withdrew, n (%): G1: NR G2: 21 (7.6) G3: 216 (9.7)</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Ghei et al., 2006</p> <p>Country and setting: UK, Primary care</p> <p>Enrollment period: NR Duration 16 weeks</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Observational cohort</p> <p>Intervention: Bladder retraining vs. Bladder retraining + antimuscarinic (oxybutynin IR/ER, tolterodine IR/ER, or imipramine combined with either oxybutynin or tolterodine as combination therapy)</p> <p>Groups: G1: Bladder retraining alone G2: Antimuscarinic therapy+ bladder retraining</p> <p>N at enrollment: G1: 52 G2: 656</p> <p>N at Follow-up: G1: 46 G2: 501</p> <p>Women, n (%): G1: 45 (86) G2: 618 (94)</p> <p>Age, mean ± SD: G1: 52 ± 14 G2: 54 ± 23</p> <p>Race/ethnicity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Frequency • Urgency with or without UUI <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • SUI • Symptoms of BOO 	<p>Incontinence episodes/day, mean ± SD: G1: 0.47 ± 1.2 G2: 1.12 ± 1.7 <i>P</i> < 0.001</p> <p>Voids/day, mean ± SD: G1: 14 ± 6 G2: 11 ± 6 <i>P</i> = 0.001</p>	<p>Incontinence episodes/day, mean change difference (95% CI): G2/G1: -0.60 (-0.93, -0.27) <i>P</i> = 0.024</p> <p>Voids/day, mean change difference (95% CI): G2/G1: 2.35 (1.4, 3.3) <i>P</i> < 0.001</p> <p>Nocturia episodes/day, mean change difference (95% CI): G2/G1: 0.57 (0.15, 0.99)</p> <p>Attendance visits, n (%) Failed follow-up: G1: 6 (12) G2: 155 (23) 1 follow-up visit: G1: 10 (19) G2: 15 (2) 2 follow-up visits: G1: 18 (35) G2: 86 (13) 3 follow-up visits: G1: 9 (17) G2: 225 (35) 4 follow-up visits: G1: 9 (17) G2: 175 (27)</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: NA</p> <p>Masking: NA</p> <p>Pt selection criteria: +</p> <p>Loss to followup: -</p> <p>Drop-out rates: NR</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: +</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: -</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Giannitsas et al., 2004</p> <p>Country and setting: Greece, Specialty treatment center</p> <p>Enrollment period: NR</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Randomized for which drug to receive first two-way crossover, table of random numbers</p> <p>Intervention: Oxybutynin 15 mg t.i.d. vs. Tolterodine 4mg b.i.d.; 6 weeks treatment with 3-4 weeks washout</p> <p>Groups: G1: Oxybutynin 15 mg t.i.d. G2: Tolterodine 4mg b.i.d.</p> <p>Stratified by UDS findings: a: high volume (> 250mL); low pressure (< 25cmH₂O) b: high volume (≥ 250mL); high pressure (> 25cmH₂O) c: low volume (< 250mL); low pressure (< 25cmH₂O) d: low volume (< 250mL); high pressure (> 25cmH₂O)</p> <p>N at enrollment: Total: 128</p> <p>N at follow-up: Total: 107 Ga: 6 Gb: 25 Gc: 36 Gd: 40</p> <p>Women, %: 100</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 18 • DO on urodynamics <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Symptomatic or recurrent UTI • BOO • Neurologic disease • History of previous pelvic surgery • Narrow angle glaucoma • SUI • History of anticholinergic side effects • Interstitial cystitis • Child-bearing age without BC 	<p>Urinary voids/day, mean ± SD: Total: 8.5 ± 2.63 Ga: 7.2 ± NR Gb: 8.0 ± 2.40 Gc: 8.3 ± 2.31 Gd: 9.3 ± 2.91</p> <p>Volume (mL)/day, mean ± SD: Total: 1568.5 ± 398.64 Ga: 1756 ± NR Gb: 1594.6 ± 326.12 Gc: 1678.8 ± 402.15 Gd: 1420.5 ± 384.96</p> <p>Voided volume (mL), mean ± SD: Total: 196.1 ± 60.19 Ga: 253 ± NR Gb: 213.5 ± 53.57 Gc: 209.8 ± 57.43 Gd: 163.4 ± 51.97</p> <p>Bladder volume (mL), first void, mean ± SD: Total: 105.6 ± 39.38 Ga: 109 ± NR Gb: 128.0 ± 41.78 Gc: 94.3 ± 35.40 Gd: 101.1 ± 38.14</p> <p>Bladder volume (mL), first contraction, mean ± SD: Total: 172 ± 98.4 Ga: 258 ± NR Gb: 303.3 ± 60.51 Gc: 111.9 ± 48.11 Gd: 124.4 ± 56.66</p>	<p>Urinary voids/day, mean ± SD: G1: 7.7 ± 1.87 G2: 7.6 ± 2.18 G1a: 6.1 ± NR G2a: 6.3 ± NR G1b: 7.3 ± 1.69 G2b: 7.0 ± 1.87 G1c: 7.5 ± 1.41 G2c: 7.2 ± 1.58 G1d: 8.3 ± 2.23 G2d: 8.4 ± 2.53</p> <p>Volume (mL)/day, mean ± SD: G1: 1764.4 ± 333.03 G2: 1670.7 ± 338.6 G1a: 1862 ± NR G2a: 1720 ± NR G1b: 1715.6 ± 292.54 G2b: 1665.8 ± 251.19 G1c: 1847.9 ± 333.81 G2c: 1808.1 ± 317.59 G1d: 1694.7 ± 331.33 G2d: 1550.3 ± 373.65</p> <p>Voided volume (mL), mean ± SD: G1: 239.9 ± 64.98 G2: 236.7 ± 63.03 G1a: 321 ± NR G2a: 286 ± NR G1b: 243.3 ± 59.56 G2b: 248.3 ± 53.91 G1c: 252.9 ± 55.75 G2c: 258.6 ± 60.63 G1d: 217.6 ± 67.07 G2d: 203.1 ± 55.43</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Masking: NA</p> <p>Pt selection criteria: +</p> <p>Loss to followup: ++</p> <p>Drop-out rates: -</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: -</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating	
Giannitsas et al., 2004 (continued)	<p>Age, mean ± SD: Total: 56 ± 16.3 Ga: 53 ± 17.2 Gb: 57 ± 16.2 Gc: 57 ± 16.3 Gd: 54 ± 16.6</p> <p>Weight (kg), mean ± SD: Total: 63 ± 5.6 Ga: 63 ± 5.6 Gb: 70 ± 9.1 Gc: 67 ± 8.8 Gd: 69 ± 7.5</p>		<p>Pressure (cmH₂O), first contraction, mean ± SD: Total: 34.8 ± 21.97 Ga: 17.4 ± NR Gb: 37.7 ± 14.03 Gc: 18.5 ± 4.60 Gd: 50.3 ± 25.14</p> <p>Overactivity index, mean ± SD: Total: 36.8 ± 31.36 Ga: 15.3 ± NR Gb: 24.8 ± 19.66 Gc: 26.3 ± 16.14 Gd: 57.0 ± 38.85</p> <p>Cystometric capacity (mL), mean ± SD: Total: 362.8 ± 119.10 Ga: 403 ± NR Gb: 410.0 ± 97.78 Gc: 357.6 ± 127.52 Gd: 331.4 ± 114.17</p>	<p>Bladder volume (mL), first desire void, mean ± SD: G1: 129.0 ± 30.14 G2: 117.9 ± 27.62 G1a: 144 ± NR G2a: 140 ± NR G1b: 153.5 ± 25.72 G2b: 132.0 ± 31.01 G1c: 120.8 ± 25.07 G2c: 113.2 ± 23.16 G1d: 119.4 ± 29.43 G2d: 110.1 ± 26.15 G1/BL: <i>P</i> < 0.05 G2/BL: <i>P</i> < 0.05</p> <p>Bladder volume (mL), first contraction, mean ± SD: G1: 212.9 ± 106.10 G2: 206.9 ± 103.56 G1a: 382 ± NR G2a: 364 ± NR G1b: 355.28 ± 74.79 G1c: 142.4 ± 43.51 G2c: 144.6 ± 48.43 G1d: 173.3 ± 57.37 G2d: 162.7 ± 53.87 G1d/BL: <i>P</i> < 0.01 G2d/BL: <i>P</i> < 0.01</p>		

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Giannitsas et al., 2004 (continued)				<p>Pressure (cmH₂O), first contraction, mean ± SD: G1: 30.9 ± 22.63 G2: 30.9 ± 19.01 G1a: 14.0 ± NR G2a: 17.4 ± NR G1b: 35.6 ± 12.86 G2b: 29.6 ± 13.56 G1c: 17.2 ± 6.80 G2c: 17.9 ± 6.69 G1d: 42.9 ± 28.91 G2d: 44.1 ± 20.75</p> <p>Overactivity index, mean ± SD: G1: 24.4 ± 22.61 G2: 24.7 ± 23.46 G1a: 7.0 ± NR G2a: 9.5 ± NR G1b: 14.1 ± 12.09 G2b: 14.1 ± 12.71 G1c: 18.3 ± 15.89 G2c: 16.9 ± 16.84 G1d: 38.9 ± 26.50 G2d: 40.7 ± 26.58</p> <p>Cystometric capacity (mL), mean ± SD: G1: 419.3 ± 120.86 G2: 415.63 ± 114.06 G1a: 465 ± NR G2a: 453 ± NR G1b: 449.6 ± 106.23 G2b: 459.4 ± 101.17 G1c: 409.9 ± 130.22 G2c: 411.05 ± 132.49 G1d: 401.8 ± 118.34 G2d: 386.7 ± 96.53</p> <p>Dry mouth, n (%): G1: 52 (40.6) G2: 20 (15.6)</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Giannitsas et al., 2004 (continued)				Constipation, n (%): G1: 11 (10.3) G2: 3 (2.8) Discontinued due to AEs, n: Dry mouth: 12 Palpitations: 1	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Gleason et al. 1999</p> <p>Country and setting: US, Community</p> <p>Enrollment period: NR 12 week follow up</p> <p>Funding: Alza Corporation</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Multicenter Open label Single treatment</p> <p>Intervention: Oxybutynin ER</p> <p>Groups: G1: Oxybutynin ER 5- 30 mg/ day (dose adjustment period)</p> <p>N at enrollment: G1: 256</p> <p>N at follow-up: G1: 219 37 missing data 20 from AEs 4 dc'd lack of effectiveness, included in analysis</p> <p>Age, yrs ± SD: NR 38.8% were 65+</p> <p>Race/ethnicity, n (%): White: G1: 235 (91.8)</p> <p>Women, N (%): G1: 234 (91.4)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • ≥ adult men and women • Idiopathic urge incontinence • Mixed incontinence with clinically significant urge component <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Uncontrolled medical condition • PVR > 100 mL • Significant bacteruria • Significant pyuria 	<p>UII episodes/ week, mean ± SD: G1: 18.8 ± 1.2</p> <p>Total incontinent episodes/ week, mean ± SD: G1: 22.2 ± 1.3</p> <p>Voids/ week, mean ± SD: G1: 81.1 ± 1.8</p> <p>PVR, mL, mean ± SD: G1: 17.4 ± 1.3</p> <p>Voided volume, mL, ± SD: G1: 121 ± 79.1</p>	<p>UII episodes/ week, mean ± SD: G1: 2.8 ± 0.6 <i>P</i> < 0.001</p> <p>Total incontinent episodes/ week, mean ± SD: G1: 4.0 ± 0.7 <i>P</i> < 0.001</p> <p>Voids/ week, mean ± SD: G1: 66.8 ± 1.4</p> <p>Reduction Voids/week, mean ± SD: G1: 14.3 ± 1.3 95%CI: 11.7-16.8</p> <p>PVR, mL, mean ± SD: G1: 21.0 ± 2.2 <i>P</i> = NR</p> <p>Voided volume, mL, ± SD: G1: 113.4 ± 84.6 <i>P</i> = NR</p> <p>Participants free UII episodes, %: 55.7</p> <p>Dry Mouth, n (%): G1: 128 (58.6)</p> <p>Discontinuation d/t AE, n (%): 20 (7.8)</p> <p>Nausea (%): 2.3</p> <p>Dry Mouth (%): 1.6</p> <p>Somnolence (%): 1.2</p> <p>Urinary retention, n (%): 2 (0.8)</p> <p>Increased PVR, n, (%): 1 (0.4)</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: NA</p> <p>Masking: NA</p> <p>Pt selection criteria: +</p> <p>Loss to followup: +</p> <p>Drop-out rates: ++</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Goode et al., 2002</p> <p>Country and setting: US, academic health center outpatient geriatric medicine clinic</p> <p>Enrollment period: July 1989 to August 1995</p> <p>Funding: National Institutes on Aging</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT, placebo controlled</p> <p>Computer-generated random numbers using a block size of 6, w/ prior stratification by type and severity of incontinence</p> <p>Intervention: Biofeedback-assisted behavioral vs. drug treatment (oxybutynin chloride; possible range of doses 2.5 mg/d-5.0 mg t.i.d.) vs. placebo</p> <p>All patients had 4 visits over an 8-week period. Patients in G1 had biofeedback added to behavioral training in absence of 50% improvement by session 3.</p> <p>Groups: G1: Behavioral ± biofeedback G2: Pharmacologic G3: Placebo</p> <p>N at enrollment: 468 screened 271 not eligible 197 randomized 105 had pre and post treatment urodynamics</p> <p>G1: 33 G2: 35 G3: 37</p> <p>N at follow-up: NA</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> Community-dwelling women at least age 55 Ambulatory At least 2 urge accidents per week by baseline bladder diary Urge incontinence as predominant pattern Urodynamic evidence of bladder dysfunction <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Continual leakage Postvoid residual urine volume >200mL Uterine prolapse past the introitus Narrow-angle glaucoma Unstable angina Decompensated congestive heart failure History of malignant arrhythmias MMSE <20 (Dementia) 	<p>Voids per day, mean ± SD: G1: 10.0 G2: 10.9 G3: 10.0</p> <p>Cystometric volume (mL), first desire to void, mean ± SD: G1: 97.1 ± 50.7 G2: 101.1 ± 62.1 G3: 124.6 ± 73.7</p> <p>Cystometric volume (mL), strong desire to void mean ± SD: G1: 188.5 ± 93.1 G2: 212.1 ± 86.7 G3: 222.3 ± 87.0</p> <p>Cystometric volume (mL), bladder capacity, mean ± SD: G1: 288.3 ± 117.0 G2: 308.7 ± 93.7 G3: 328.9 ± 107.6</p>	<p>Voids per day, mean ± SD: G1: 8.2 G2: 8.8 G3: 9.7</p> <p>DI on UDS, n (%): +Baseline DI/ +DI post-treatment: G1: 7 (21.2) G2: 1 (2.9) G3: 5 (13.5) +Baseline DI/ -DI post-treatment: G1: 1 (3.0) G2: 7 (20.0) G3: 7 (18.9) -Baseline DI/ +DI post-treatment: G1: 3 (9.1) G2: 3 (8.6) G3: 3 (8.1) -Baseline DI/-DI post-treatment: G1: 22 (66.7) G2: 24 (68.6) G3: 22 (59.5)</p> <p>Cystometric volume (mL), first desire to void, mean ± SD: G1: 115.9 ± 64.9 G2: 145.6 ± 74.0 G3: 133.5 ± 59.6</p> <p>Cystometric volume (mL), strong desire to void mean ± SD: G1: 228.9 ± 106.4 G2: 282.0 ± 93.2 G3: 230.1 ± 78.8</p> <p>Cystometric volume (mL), bladder capacity, mean ± SD: G1: 305.6 ± 117.9 G2: 377.6 ± 92.1 G3: 323.0 ± 109.0</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: + Masking: + Pt selection criteria: + Loss to followup: + Drop-out rates: + Power calculation: - Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: + Baseline OAB status: + Baseline characteristics: ++ Length of followup: + Measurement methods: + Measurement reliability: + Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Goode et al., 2002 (continued)	<p>Age, yrs ± SD: G1: 65.3 ± 4.5 G2: 67.9 ± 7.9 G3: 67.6 ± 7.7</p> <p>Race/ethnicity, %: Black: 2 White: 98</p> <p>Women: 100%</p> <p>Parity mean ± SD: G1: 3.1 ± 1.7 G2: 2.1 ± 1.3 G3: 2.3 ± 1.5</p>			<p>Cystometric, volume (mL), first desire to void, mean change: G1: 18.8 G2: 44.4 G3: 8.9 <i>P</i> = 0.149</p> <p>Cystometric, volume (mL), strong desire to void, mean change: G1: 40.5 G2: 69.9 G3: 7.8 <i>P</i> = 0.018</p> <p>Cystometric, volume (mL), bladder capacity, mean change: G1: 17.3 G2: 68.9 G3: -6.0 <i>P</i> = 0.000</p> <p>Standardized estimates of direct and mediated effects of treatment: G1/G3: Total effect: 0.28* Direct effect: 0.23 Mediated Effect: 0.05 G2/G3: Total Effect: 0.34* Direct Effect: 0.30* Mediated Effect: 0.04 *<i>P</i> < 0.01</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Haab et al., 2005</p> <p>Country and setting: Multinational, Community</p> <p>Enrollment period: NR 52 week open enrollment</p> <p>Funding: Yamanouchi</p> <p>Author industry relationship disclosures: 1 of 4 Yamanouchi (1)</p>	<p>Design: Cohort open-label extension</p> <p>Intervention: Solifenacin 5mg for 4 weeks, then pt-selected dose of either 5 or 10 mg daily for remainder of study</p> <p>Groups: NA</p> <p>N at enrollment: 1,637</p> <p>N at follow-up: 1,329</p> <p>Women, n (%): 1,280 (78)</p> <p>Age, mean ± SD: 56.4 ± 13.5</p> <p>Race/ethnicity, n (%): Black: 8 (0.5) White: 1,602 (98.1) Asian: 13 (0.8) Other: 10 (0.6)</p> <p>Weight (kg), mean ± SD: 74.5 ± 14.6</p> <p>Height (cm), mean ± SD: 165.3 ± 8.4</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 18 • Symptoms of OAB ≥ 3 months • ≥ 8 voids per day • ≥ 1 urgency episode / day • ≥ 1 UUI episodes / day • Completed treatment in previous RCT ≤ 14 days prior to extension study entry <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • BOO • PVR > 200 mL • Persistent or recurrent UTI • Bladder stones • IC • Previous pelvic radiation • Previous or current pelvic organ malignancy • Narrow angle glaucoma • Gastric retention • Urinary retention • Pregnant/nursing • Non-reliable BC in childbearing woman 	<p>Urgency episodes/day, mean ± SD: 5.76 ± 4.46</p> <p>Incontinence episodes/day, mean ± SD: 2.66 ± 2.51</p> <p>Voids/day, mean ± SD: 12.16 ± 3.79</p> <p>Nocturia episodes/day, mean ± SD: 1.95 ± 1.22</p> <p>Voided volume (mL), mean ± SD: 147.6 ± 53.8</p>	<p>Urgency episodes/day, mean ± SD (% change): 2.28 ± 3.59 (-63)</p> <p>Incontinence episodes/day, mean ± SD (% change): 0.93 ± 2.06 (-66)</p> <p>Voids/day, mean ± SD (% change): 9.18 ± 3.33 (-23)</p> <p>Nocturia episodes/day, mean ± SD (% change): 1.25 ± 1.26 (-32)</p> <p>Voided volume (mL), mean ± SD (% change): 187.4 ± 75.4 (31)</p> <p>Dry mouth, n (%): 339 (20.7)</p> <p>Constipation, n (%): 157 (9.6)</p> <p>Blurred vision, n (%): 113 (6.9)</p> <p>Discontinuation rate due to AE, %: 4.7</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: NA</p> <p>Masking: NA</p> <p>Pt selection criteria: +</p> <p>Loss to followup: +</p> <p>Drop-out rates: ++</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: ++</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Haab et al., 2006 [See evidence table for Hill et al., 2007]</p> <p>Country and setting: France, Denmark, Sweden, Australia, Canada, US</p> <p>Enrollment period: NR</p> <p>Funding: Novartis Pharma AG (educational grant)</p> <p>Author industry relationship disclosures: 8 of 9 Astellas (1) Novartis (8) Pfizer (1)</p>	<p>Design: 2-year, non-comparative, open-label extension study</p> <p>Intervention: Darifenacin following 2 feeder studies of darifenacin 3.75/7.5/15 mg qd or placebo x 12 wks</p> <p>Groups: NA</p> <p>N at enrollment: 719</p> <p>N at follow-up (%): 475 (66.3)</p> <p>Women, n (%): 609 (85.1)</p> <p>Age, mean (range): 57.3 (19-89)</p> <p>Age ≥ 65, %: 29.9</p> <p>Race/ethnicity: NR</p> <p>Received treatment ≥ 365 days, n (%): 532 (74.3)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> Successful completion of one of two previous, 12-wk feeder studies with no major protocol violation <p>Exclusion criteria: NR</p>	<p>Urgency episodes/day, median (IQR): 8.1 (5.7-10.3)</p> <p>Urgency severity, VAS score, median (IQR): 54.7 (43.7-66.1)</p> <p>Incontinence episodes/week, median (IQR): 16.8 (10.0-27.0)</p> <p>Significant leaks/week, median (IQR): 7.0 (2.3-14.0)</p> <p>Voids/day, median (IQR): 10.1 (8.9-12.1)</p> <p>Nocturia episodes/week, median (IQR): 11.0 (7.0-16.0)</p> <p>Voided volume (mL), median (IQR): 163 (121-217)</p> <p>Primary diagnosis, n (%) Idiopathic: 674 (94.1) Neurogenic: 42 (5.9)</p> <p>Previous OAB drug treatment, n (%): 163 (22.8)</p>	<p>Urgency episodes/day, median change (median % change): -3.9 (-56.4) <i>P</i> < 0.001</p> <p>Urgency severity, median change (median % change): -15.4 (-28.8) <i>P</i> < 0.001</p> <p>Incontinence episodes/week, median change (median % change): -11.0 (-84.4) <i>P</i> < 0.001</p> <p>Significant leaks/week, median change (median % change): -4.7 (-100.0) <i>P</i> < 0.001</p> <p>Incontinence episodes, responders, reduction, %: ≥ 50%: 77.2 ≥ 70%: 62.3 ≥ 90%: 43.8</p> <p>Voids/day, median change (median % change): -1.4 (-13.9) <i>P</i> < 0.001</p> <p>Nocturia episodes/week, median change (median % change): -1.5 (-14.3) <i>P</i> < 0.001</p> <p>Increased dose to 15 mg at 2 wks and maintained, n (%): 363 (50.7)</p>	<p>Quality: Overall quality score: fair INTERNAL VALIDITY: poor Randomization: NA Masking: NA Pt selection criteria: + Loss to followup: - Drop-out rates: - Power calculation: - Statistical issues: - EXTERNAL VALIDITY: good Age: + Baseline OAB status: + Baseline characteristics: + Length of followup: ++ Measurement methods: + Measurement reliability: + Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Haab et al., 2006 (continued)				<p>Remained on 7.5 mg dose, n (%): 182 (25.4)</p> <p>Dose adjusted at other times, n (%): 171 (23.9)</p> <p>Compliance, \geq 80% of doses, %: >85</p> <p>Voided volume (mL), median change (median % change): 19.0 (11.9) $P < 0.001$</p> <p>Adverse events, all causality, n (%): Any: 572 (79.9) Serious: 84 (11.7) Dry mouth: 167 (23.3) Constipation: 150 (20.9) UTI: 82 (11.5) Respiratory d/o: 76 (10.6) Dyspepsia: 65 (9.1) Accidental injury: 57 (8.0) 'Flu' syndrome: 49 (6.8) HA: 42 (5.9) HTN: 39 (5.4) Arthritis: 36 (5.0) Bronchitis: 34 (4.7) Arthralgia: 33 (4.6) Back pain: 30 (4.2) Increased cough: 26 (3.6) Arthrosis: 25 (3.5) Pain: 23 (3.2) Sinusitis: 22 (3.1)</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Haab et al., 2006 (continued)				<p>Adverse events, treatment related, n (%): Any: 343 (47.9) Serious: 1 (0.1) Dry mouth: 166 (23.2) Constipation: 142 (19.8) UTI: 8 (1.1) Respiratory d/o: 2 (0.3) Dyspepsia: 36 (5.2) Accidental injury: 1 (0.1) 'Flu' syndrome: 0 HA: 14 (2.0) HTN: 3 (0.4) Arthritis: 1 (0.1) Bronchitis: 0 Arthralgia: 1 (0.1) Back pain: 4 (0.6) Increased cough: 1 (0.1) Arthrosis: 0 Pain: 1 (0.1) Sinusitis: 1 (0.1)</p> <p>Days of bowel motions, mean change*: -0.04</p> <p>Days of hard or lumpy bowel motions, mean change*: 0.13</p> <p>Days of straining, mean change*: 0.22</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Haab et al., 2006 (continued)				Days of incomplete bowel openings:* 0.39 Days on which patient felt life was affected by bowel habits:* 0.17 Extent to which life affected by bowel habits in previous 2 wks, mean score:* 0.31	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Halaska et al., 2003</p> <p>Country and setting: Europe and Asia, Academic medical center</p> <p>Enrollment period: May 1996 to May 1999</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Randomized controlled double blind crossover</p> <p>Intervention: Trospium vs. Oxybutynin</p> <p>Groups: G1: Trospium 20 mg b.i.d. G2: Oxybutynin 5 mg b.i.d.</p> <p>N at enrollment: G1: 268 G2: 90</p> <p>N at follow-up: G1: 200 G2: 66</p> <p>Women, n (%): G1: 228 (85) G2: 78 (87)</p> <p>Age, mean (range): G1: 54.2 (19, 89) G2: 52.2 (19, 85)</p> <p>Weight (kg), mean (range): G1: 72.3 (50-120) G2: 70.4 (50-90)</p> <p>Height (cm), mean (range): G1: 164.8 (144-185) G2: 165.5 (145-183)</p> <p>Smokers, n (%): G1: 38 (14) G2: 10 (11)</p> <p>Previous illness, n (%): G1: 184 (69) G2: 66 (73)</p> <p>Previous medication, n (%): G1: 101 (38) G2: 46 (51)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age \geq 18 yrs • Urge syndrome • Urge incontinence • UUI as one component of MUI • UUI due to neurologic condition (detrusor hyperreflexia) <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Absolute tachycardia • Closed angle glaucoma • Myasthenia gravis • Arteriosclerosis of cerebral vessels • SUI • Heart or renal failure • Frequency from diuretics • BOO • Acute UTI • Hiatus hernia with reflux esophagitis • Stenosis of GI tract • Megacolon • Colonic ulceration • Allergy to study medications • Anticholinergics, TCAs, alpha blockers, beta sympathomimetics \leq 7 days • Urological or gynecologic surgery \leq 3 mos • Pregnant or lactating • In another study 	<p>Urgency episodes/day, mean: G1: 10.2 G2: 11.0</p> <p>Incontinence episodes/ day, mean: G1: 1.5 G2: 2.1</p> <p>Voids/day, mean: G1: 11.4 G2: 12.5</p> <p>Max cystometric capacity (mL), mean: G1: 205 G2: 205</p>	<p>Urgency episodes/day, mean: G1: 6.7 G2: 7.4</p> <p>Incontinence episodes/ day, mean: G1: 0.5 G2: 1.1</p> <p>Voids/day, mean: G1: 7.9 G2: 8.3</p> <p>Max cystometric capacity (mL), 26 wks, mean change: G1: 92.0 G2: 117.0 G1/BL: $P \leq 0.001$ G2/BL: $P \leq 0.001$</p> <p>Max cystometric capacity (mL), 52 wks, mean change: G1: 115.0 G2: 119.4 G1/BL: $P \leq 0.001$ G2/BL: $P \leq 0.001$</p> <p>Bladder volume (mL), first contraction, 26 wks, mean change: G1: 63.5 G2: 61.2</p> <p>Bladder volume (mL), first contraction, 52 wks, mean change: G1: 46.1 G2: 36.7</p> <p>Bladder volume (mL), first sensation, 26 wks, mean change: G1: 73.6 G2: 76.93</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: -</p> <p>Masking: -</p> <p>Pt selection criteria: +</p> <p>Loss to followup: ++</p> <p>Drop-out rates: ++</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: ++</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Halaska et al., 2003 (continued)				<p>Bladder volume (mL), first sensation, 26 wks, mean change: G1: 78.6 G2: 70.2</p> <p>Abnormal EKG, n (%): G1: 4 (0.1) G2: 2 (0.2)</p> <p>Any adverse event, n (%): G1: 10 (3.7) G2: 6 (6.7)</p> <p>Poor efficacy, n (%): G1: 8 (3) G2: 2 (2.2)</p> <p>Poor compliance, n (%): G1: 15 (15.6) G2: 6 (6.7)</p> <p>Abdominal pain, n (%): G1: 5 (2) G2: 0 (0)</p> <p>Constipation, n (%): G1: 18 (7) G2: 4 (4)</p> <p>Diarrhea, n (%): G1: 2 (1) G2: 2 (2)</p> <p>Dyspepsia, n (%): G1: 13 (5) G2: 3 (3)</p> <p>Dysphagia, n (%): G1: 9 (3) G2: 3 (3)</p> <p>Dry mouth, n (%): G1: 87 (33) G2: 45 (50) <i>P</i> < 0.001</p> <p>Nausea, n (%): G1: 6 (2) G2: 2 (2)</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Halaska et al., 2003 (continued)				UTI, n (%): G1: 33 (12) G2: 10 (11) Headache, n (%): G1: 11 (4) G2: 8 (9) Visual disturbances, n (%): G1: 9 (3) G2: 5 (6) Virus infection, n (%): G1: 9 (3) G2: 4 (4) Abnormal EKG, n (%): G1: 4 (0.1) G2: 2 (0.2) Sleeplessness, n (%): G1: 10 (4) G2: 2 (2)	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Herschorn et al., 2004</p> <p>Country and setting: Canada, Family medicine and urology clinics</p> <p>Enrollment period: June 2000 to December 2001</p> <p>Funding: Pharmacia Pfizer Canada</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: Health education with tolterodine vs. tolterodine alone</p> <p>Groups: G1: Health education with tolterodine G2: tolterodine alone</p> <p>N at enrollment: G1: 39 G2: 45</p> <p>N at follow-up, 5 weeks: G1: 37 G2: 40</p> <p>N at follow-up, 10 weeks: G1: 35 G2: 32</p> <p>N at follow-up, 16 weeks: G1: 34 G2: 31</p> <p>Age, yrs ± SD: G1: 65.7 ± 14.5 G2: 63.1 ± 15.7</p> <p>Race/ethnicity: NR</p> <p>Women, %: G1: 92.3 G2: 84.4</p> <p>Parity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 50 • Symptoms of OAB • Attend investigators' practice • Normal cognitive function • Able to read English <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Enrollment in another clinical trial • Interstitial cystitis • UTI • Already taking tolterodine 	<p>OAB duration (yrs), mean ± SD: G1: 8.7 ± 11.0 G2: 8.7 ± 10.8</p> <p>Mild bladder problems, n (%): G1: 13 (33.3) G2: 13 (28.9)</p> <p>Moderate bladder problems, n (%): G1: 19 (48.7) G2: 28 (62.2)</p> <p>Severe bladder problems, n (%): G1: 7 (18.0) G2: 4 (8.9)</p> <p>Obtained prescription, n (%): G1: 38 (97.4) G2: 37 (82.2) <i>P</i> < 0.05</p> <p>Intends to fill prescription, (%): G1: 0 (0) G2: 6 (7.5)</p>	<p>Incontinence episodes/week, mean change ± SD: G1: -7.72 ± 21.16 G2: -10.24 ± 19.56†</p> <p>Voids/day, mean change ± SD: G1: -1.82 ± 3.41 G2: -2.18 ± 4.89 <i>P</i> = NR</p> <p>Nocturia episodes/day, mean change ± SD: G1: -0.44 ± 1.13† G2: -0.07 ± 0.91</p> <p>No change in bladder problem severity, n (%): G1: 14 (42.4) G2: 20 (66.7)</p> <p>Improved bladder problem severity, n (%): G1: 15 (45.4)† G2: 6 (20)</p> <p>Worsened bladder problem severity, n (%): G1: 4 (12.1) G2: 4 (13.3)</p> <p>Compliance, 10 weeks, %: G1: 41 G2: 38 <i>P</i> > 0.05</p> <p>Compliance, 16 weeks, %: G1: 39 G2: 31 <i>P</i> > 0.05</p> <p>Continued or started non-drug OAB treatment, 16 weeks, %: G1: 82 G2: 53 <i>P</i> > 0.05</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: -</p> <p>Masking: -</p> <p>Pt selection criteria: +</p> <p>Loss to followup: -</p> <p>Drop-out rates: NR</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: +</p> <p>Baseline OAB status: NR</p> <p>Baseline characteristics: -</p> <p>Length of followup: ++</p> <p>Measurement methods: +</p> <p>Measurement reliability: -</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Herschorn et al., 2004 (continued)				Stopped non-drug OAB treatments, %: G1: 12.8 G2: 28.9	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Hill et al., 2006</p> <p>Country and setting: Multinational-European; Community; 62 centers</p> <p>Enrollment period: NR</p> <p>Funding: Pfizer Novartis Thomson ACUMED</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: Darifenacin vs. placebo</p> <p>Groups: G1: Darifenacin 7.5 mg daily G2: Darifenacin 15 mg daily G3: Darifenacin 30 mg daily G4: placebo</p> <p>N at enrollment: G1: 108 G2: 107 G3: 115 G4: 109</p> <p>N at follow-up: G1: 99 G2: 93 G3: 96 G4: 101</p> <p>Total 50 (11.4%) discontinued study- AE and withdrawal of consent</p> <p>Age, mean (range): G1: 56.1 (23-88) G2: 55.1 (24-82) G3: 54.0 (23-79) G4: 53.7 (21-85)</p> <p>Race/ethnicity: NR</p> <p>Women, N (%): G1: 94 (87) G2: 92 (86) G3: 99 (86) G4: 90 (83)</p> <p>Parity : NR</p> <p>Previous drug therapy, n (%): G1: 21 (19.4) G2: 22 (20.6) G3: 33 (28.7) G4: 24 (22.0)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Male and female • Age ≥18 • ≥ 10 episodes UUI over 14 days • ≥ 8 voids/ day • ≥ 1 episode of urinary urgency • Symptoms OAB ≥ 6 months <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • SUJ • > 2 UTI/ year • BOO • PVR> 200 • IC • Bladder stones • Severe constipation • History of intermittent UTI • Urogenital surgery within previous 6 months • Cystoscopy within previous 30 days • Indwelling catheter • CIC • Clinically significant systemic disease • Bladder training • CI to antimuscarinic therapy • Pregnant/ lactating • Concomitant treatment with anticholinergic/ antispasmodics, opioids, HRT, drugs that inhibit cytochrome P450, 2D6, and 3A4 isoenzymes 	<p>Urgency episodes/day, median (95% CI): G1: 8.5 (7.0, 8.7) G2: 8.6 (7.8, 9.4) G3: 8.4 (7.8, 8.8) G4: 8.1 (7.4, 8.7)</p> <p>Severity of urgency, visual analog scale (0=mild, 100=severe), (range): G1: 53.2 (50.1-56.5) G2: 55.8 (51.5-59.8) G3: 53.5 (50.0-57.9) G4: 53.5 (51.2-57.0)</p> <p>Incontinence episodes/week, median (95% CI): G1: 13.7 (11.8, 17.8) G2: 17.3 (13.5, 21.5) G3: 19.1 (15.8, 22.8) G4: 16.1 (14.0, 19.4)</p> <p>Voids/day, median (95% CI): G1: 10.3 (9.8, 10.9) G2: 11.0 (10.4, 12.1) G3: 10.4 (9.8, 11.4) G4: 10.1 (9.8, 10.8)</p> <p>Number OAB-related nocturnal awakenings/ week, median (95%CI): G1: 12.4 (10.2, 12.9) G2: 11.7 (9.8, 14.5) G3: 12.4 (10.5, 14.0) G4: 10.4 (9.2, 12.9)</p>	<p>Urgency episodes/ day, median change (%): G1: -1.8 (-29.2) G2: -2.3 (-26.9) G3: -3.0 (-33.1) G4: -1.2 (-15.7) G1/G4: <i>P</i> = 0.196 G2/G4: <i>P</i> = 0.013 G3/G4: <i>P</i> < 0.001</p> <p>Severity of urgency, visual analog scale (0=mild, 100=severe), median change (range): G1: -7.0 (-14.2) G2: -7.0 (-11.6) G3: -9.4 (-19.9) G4: -3.9 (-8.0) G1/G4: <i>P</i> = 0.177 G2/G4: <i>P</i> = 0.045 G3/G4: <i>P</i> = 0.011</p> <p>Incontinence episodes/wk, median change (%): G1: -8.1 (-68.7) G2: -10.4 (-76.5) G3: -11.4 (-77.3) G4: -5.9 (-46.0) G1/G4: <i>P</i> = 0.007 G2/G4: <i>P</i> < 0.001 G3/G4: <i>P</i> < 0.001</p> <p>Leaks/week, median change (%): G1: -4.3 (-67.0) G2: -5.3 (-76.7) G3: -6.0 (-73.7) G4: -2.4 (-43.2) G1/G4: <i>P</i> = 0.012 G2/G4: <i>P</i> < 0.001 G3/G4: <i>P</i> < 0.001</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: +</p> <p>Drop-out rates: -</p> <p>Power calculation: +</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Hill et al., 2006 (continued)			Voided volume (mL), median (95%CI): G1: 162 (153, 181) G2: 157 (143, 173) G3: 156 (144, 168) G4: 162 (149, 174)	Patients achieving ≥7 consecutive days “dry”, (%): G1: NR G2: 28 G3: 30 G4: 17 G2/G4: <i>P</i> = 0.29 G3/G4: <i>P</i> = 0.011 voids/day, median change (%): G1: -1.7 (-16.7) G2: -1.9 (-17.9) G3: -2.2 (-21.2) G4: -1.1 (-9.6) G1/G4: <i>P</i> = 0.06 G2/G4: <i>P</i> = 0.033 G3/G4: <i>P</i> < 0.001 Number OAB-related nocturnal awakenings/ week, median change (%): G1: -1.9 (-22.1) G2: -1.7 (-22.7) G3: -2.0 (-19.2) G4: -0.4 (-3.6) G1/G4: <i>P</i> = 0.028 G2/G4: <i>P</i> = 0.022 G3/G4: <i>P</i> = 0.008 Voided volume (mL), median change (%): G1: 17 (10.3) G2: 24 (15.6) G3: 44 (26.0) G4: 7 (4.1) G1/G4: <i>P</i> = 0.079 G2/G4: <i>P</i> = 0.001 G3/G4: <i>P</i> < 0.001 Dry mouth, n (%): G1: 25 (23.1) G2: 43 (40.2) G3: 68 (59.1) G4: 6 (5.5)	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Hill et al., 2006 (continued)				<p>Constipation, n (%): G1: 17 (15.7) G2: 27 (25.2) G3: 32 (27.8) G4: 5 (4.6)</p> <p>Dyspepsia, n (%): G1: 4 (3.7) G2: 9 (8.4) G3: 10 (8.7) G4: 1 (0.9)</p> <p>Headache, n (%): G1: 7 (6.5) G2: 7 (6.5) G3: 7 (6.1) G4: 2 (1.8)</p> <p>Respiratory tract infection, n (%): G1: 4 (3.7) G2: 6 (5.6) G3: 1 (0.9) G4: 6 (5.5)</p> <p>Urinary tract disorder, n (%): G1: 0 (0) G2: 6 (5.6) G3: 1 (0.9) G4: 0</p> <p>UTI, n (%): G1: 3 (2.8) G2: 3 (2.8) G3: 5 (4.3) G4: 2 (1.8)</p> <p>Flu syndrome, n (%): G1: 3 (2.8) G2: 2 (1.9) G3: 0 G4: 5 (4.6)</p> <p>Back pain, n (%): G1: 3 (2.8) G2: 1 (0.9) G3: 5 (4.3) G4: 3 (2.8)</p> <p>Abdominal pain, n (%): G1: 1 (0.9) G2: 3 (2.8) G3: 4 (3.5) G4: 1 (0.9)</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Hill et al., 2006 (continued)				Abnormal vision, n (%): G1: 2 (1.9) G2: 0 G3: 4 (3.5) G4: 0	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Hill et al., 2007</p> <p>[See evidence table for Haab et al., 2006]</p> <p>Country and setting: France, Denmark, Sweden, Australia, Canada, US, Multicenter</p> <p>Enrollment period: NR</p> <p>Funding: Novartis Pharma AG (educational grant)</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: 2-yr, non-comparative, open-label extension study</p> <p>Intervention: Darifenacin following 2 feeder studies of darifenacin 3.75/7.5/15 mg qd or placebo x 12 wks</p> <p>Groups: NA</p> <p>N at enrollment: 214</p> <p>N at follow-up (%): 137 (64.0)</p> <p>Patient years of exposure: 308</p> <p>Women, n (%): 172 (80.4)</p> <p>Age, n (%): 65-74 years: 166 (77.6) 75-89 years: 48 (22.4)</p> <p>Race/ethnicity: NR</p> <p>Menopausal, n (%): 13 (6.1)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 65 • Successful completion of one of 2 previous, 12-wk feeder studies with no major protocol violation <p>Exclusion criteria: NR</p>	<p>Urgency episodes/day, median: 8.1</p> <p>Urgency severity, median VAS score: 51.8</p> <p>Incontinence episodes/week, median: 18.7</p> <p>Voids/day, median: 10.1</p> <p>Voided volume (mL), median: 156</p> <p>Concomitant drug treatment, n (%): 210 (98.1)</p>	<p>Urgency episodes/day, median change (median % change): -3.7 (-52.0) <i>P</i> < 0.05</p> <p>Urgency severity, VAS score, median change (median % change): -12.6 (-23.3) <i>P</i> < 0.05</p> <p>Incontinence episodes/week, median change (median % change): -11.0 (-83.7) <i>P</i> < 0.05</p> <p>Significant leaks/week, median change (median % change): -4.9 (-100.0) <i>P</i> < 0.05</p> <p>Incontinence episodes, responders, % reduction, %: ≥ 50%: 74.1 ≥ 70%: 60.0 ≥ 90%: 44.4</p> <p>Voids/day, median change (median % change): -1.2 (-12.4) <i>P</i> < 0.05</p> <p>Normalization of voiding frequency (< 8 voids/day) after darifenacin treatment, %: 3 months: 34 2 years: 33.8</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: NA</p> <p>Masking: NA</p> <p>Pt selection criteria: +</p> <p>Loss to followup: -</p> <p>Drop-out rates: -</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: +</p> <p>Length of followup: ++</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Hill et al., 2007 (continued)				<p>Nocturia episodes/week, median change (median % change): -1.4 (-10.9) G1/BL: <i>P</i> < 0.05</p> <p>Increased dose to 15 mg at 2 wks and maintained, n (%): 110 (51.4)</p> <p>Remained on 7.5 mg dose, n (%): 55 (25.7)</p> <p>Dose adjusted at other times, n (%): 49 (22.9)</p> <p>Compliance, ≥ 80% of doses, n (%): 214 (84)</p> <p>Voided volume (mL), median change (median % change): 11.1 (6.3)</p> <p>Dry mouth, n (%): 50 (23.4)</p> <p>Constipation, n (%): 48 (22.4)</p> <p>CVD, %: 1.4</p> <p>Peripheral/CNS, %: 3.3</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Homma and Koyama, 2006</p> <p>Country and setting: Japan, Community</p> <p>Enrollment period: NR 9 weeks 1 week wash out 8 week treatment</p> <p>Funding: Sanyko Company</p> <p>Author industry relationship disclosures: 1 of 2 Sanyko (1)</p>	<p>Design: RCT</p> <p>Intervention: Oxybutynin Transdermal (TDS) patch vs placebo</p> <p>Groups: G1: OXY TDS 26 cm² G2: OXY TDS 39 cm² G3: OXY TDS 52 cm² G4: placebo</p> <p>N at enrollment: G1: 160 G2: 164 G3: 152 G4: 161</p> <p>N at follow-up: G1: 160 G2: 164 G3: 152 G4: 161</p> <p>Women, n (%): G1: 122 (76.3) G2: 132 (80.5) G3: 119 (78.3) G4: 117 (72.7)</p> <p>Age, mean ± SD: G1: 63.5 ± 11.5 G2: 62.7 ± 12.4 G3: 62.0 ± 13.6 G4: 62.9 ± 12.3</p> <p>Race/ethnicity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> Adult with OAB syndrome ≥ 1 episodes UUI per day ≥ 8 voids per day <p>Exclusion criteria: NR</p>	<p>Incontinence episodes/week, mean ± SD: G1: 21.1 ± 13.9 G2: 20.8 ± 14.6 G3: 20.4 ± 14.1 G4: 21.1 ± 16.6</p> <p>KHQ, general health, mean ± SD: G1: 36.9 ± 21.0 G2: 38.6 ± 20.3 G3: 36.0 ± 21.4 G4: 36.0 ± 20.5</p> <p>KHQ, incontinence impact, mean ± SD: G1: 54.6 ± 27.1 G2: 54.9 ± 26.7 G3: 57.1 ± 27.4 G4: 54.0 ± 25.5</p> <p>KHQ, role limitation, mean ± SD: G1: 42.5 ± 24.6 G2: 40.7 ± 24.5 G3: 41.8 ± 26.5 G4: 43.0 ± 25.1</p> <p>KHQ, physical limitation, mean ± SD: G1: 48.6 ± 26.0 G2: 45.8 ± 26.0 G3: 48.9 ± 29.6 G4: 46.9 ± 26.2</p> <p>KHQ, social limitation, mean ± SD: G1: 27.4 ± 23.7 G2: 25.2 ± 24.3 G3: 26.1 ± 24.5 G4: 26.3 ± 25.7</p> <p>KHQ, personal relationship, mean ± SD: G1: 17.9 ± 21.6 G2: 13.2 ± 20.7 G3: 16.4 ± 23.4 G4: 17.0 ± 24.7</p>	<p>Incontinence episodes/week, mean ± SD: G1: 10.3 ± 17.8 G2: 6.3 ± 10.4 G3: 8.9 ± 15.8 G4: 11.0 ± 15.4 G2/G4: <i>P</i> = 0.014</p> <p>KHQ domain, general health, mean ± SD: G1: 33.4 ± 20.3 G2: 30.9 ± 22.2 G3: 33.9 ± 21.6 G4: 33.0 ± 22.7 G2/G4: <i>P</i> < 0.05</p> <p>KHQ domain, incontinence impact, mean ± SD: G1: 34.6 ± 23.2 G2: 32.7 ± 23.6 G3: 34.0 ± 24.4 G4: 39.7 ± 26.0 G2/G4: <i>P</i> < 0.05</p> <p>KHQ domain, role limitation, mean ± SD: G1: 24.8 ± 22.0 G2: 22.0 ± 20.3 G3: 26.5 ± 24.7 G4: 31.9 ± 24.1 G2/G4: <i>P</i> < 0.05</p> <p>KHQ domain, physical limitation, mean ± SD: G1: 29.7 ± 25.6 G2: 26.6 ± 22.8 G3: 29.7 ± 27.3 G4: 36.5 ± 27.5 G2/G4: <i>P</i> < 0.05</p> <p>KHQ domain, social limitation, mean ± SD: G1: 16.3 ± 21.3 G2: 13.2 ± 17.1 G3: 18.4 ± 22.8 G4: 21.6 ± 24.2 G2/G4: <i>P</i> < 0.05</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: -</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: ++</p> <p>Drop-out rates: NA</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: +</p> <p>Length of followup: -</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Homma and Koyama, 2006 (continued)			<p>KHQ, emotions, mean ± SD: G1: 45.3 ± 28.9 G2: 46.5 ± 27.3 G3: 46.1 ± 28.1 G4: 46.2 ± 26.2</p> <p>KHQ, sleep/ energy, mean ± SD: G1: 32.0 ± 25.4 G2: 35.3 ± 24.9 G3: 33.4 ± 28.5 G4: 32.5 ± 26.3</p>	<p>KHQ domain, personal relationship, mean ± SD: G1: 10.4 ± 17.3 G2: 8.4 ± 16.8 G3: 11.6 ± 22.1 G4: 12.0 ± 20.2 G2/G4: <i>P</i> < 0.05</p> <p>KHQ domain, emotions, mean ± SD: G1: 28.2 ± 25.8 G2: 24.9 ± 21.6 G3: 29.3 ± 26.7 G4: 35.2 ± 28.4 G2/G4: <i>P</i> < 0.05</p> <p>KHQ domain, sleep/ energy, mean ± SD: G1: 18.2 ± 19.2 G2: 17.9 ± 18.9 G3: 21.1 ± 22.8 G4: 26.0 ± 25.6 G2/G4: <i>P</i> < 0.05</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Jacquetin and Wyndaele, 2001</p> <p>Country and setting: France (22 centers) and Belgium (10 centers)</p> <p>Enrollment period: NR</p> <p>Funding: Pharmacia</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Double-blind, placebo-controlled, parallel-group, multicenter phase III study w/ 2-wk washout period</p> <p>Intervention: Tolterodine vs placebo for 4 weeks</p> <p>Groups: G1: Tolterodine 2mg b.i.d. G2: Tolterodine 1 mg b.i.d. G3: placebo (frequency NR)</p> <p>N at enrollment: NR</p> <p>N at follow-up: G1: 103 G2: 97 G3: 51</p> <p>Women, n (%): G1: 84 (82) G2: 74 (76) G3: 41 (80)</p> <p>Age, mean (range): G1: 58 (21-88) G2: 53 (18-85) G3: 56 (19-89)</p> <p>Race/ethnicity: NR</p> <p>BMI, kg/m² (range): G1: 26.4 (17.7-39.7) G2: 25.5 (16.7-46.3) G3: 24.8 (17.6-36.9)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 18 • UDS-verified detrusor overactivity • ≥ 1 UUI episode/ day • ≥ 8 voids/day <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • SI • Hepatic or renal dz • +UTI or recurrent UTIs • Interstitial cystitis • Hematuria • Clinically significant voiding difficulty • Pts receiving bladder training, electrostimulation therapy • Indwelling catheter • Intermittent cath • Pregnant or nursing • Women of childbearing age w/o reliable contraception 	<p>UUI, n (%) G1: 75 (73) G2: 75 (77) G3: 39 (76)</p> <p>UUI episodes/day, mean (range): G1: 3.2 (0.1-24.0) G2: 2.7 (0.1-24.0) G3: 2.4 (0.1-8.4)</p> <p>Voids/day, mean (range): G1: 10.8 (6.2-34.7) G2: 10.7 (4.9-26.4) G3: 11.7 (6.3-26.3)</p> <p>≥ 8 voids/day, n (%): G1: 96 (93) G2: 89 (92) G3: 49 (96)</p> <p>Urinary symptoms > 5 years, n (%): G1: 46 (45) G2: 42 (43) G3: 17 (33)</p> <p>Previous lower urinary tract surgery, n (%): G1: 36 (35) G2: 29 (30) G3: 16 (31)</p> <p>Previous drug therapy for OAB, n (%): G1: 66 (64) G2: 59 (61) G3: 36 (71)</p> <p>Previous poor efficacy response, n (%): G1: 50 (76) G2: 45 (76) G3: 26 (72)</p>	<p>UUI episodes/day, mean change ± SD: G1: -1.3 ± 1.8 G2: -1.1 ± 2.2 G3: -0.4 ± 1.9 G1/G3: <i>P</i> = 0.0089 G2/G3: <i>P</i> = 0.045</p> <p>Voids/day, mean change ± SD: G1: -1.4 ± 4.3 G2: -1.4 ± 2.8 G3: -1.2 ± 2.7 G1/G3: <i>P</i> = NS G2/G3: <i>P</i> = NS</p> <p>Good efficacy response in current study, previous poor efficacy response, n/N (%) G1: 20/39 (51) G2: 18/37 (49) G3: 7/19 (37) G1/G3: <i>P</i> = NS G2/G3: <i>P</i> = NS</p> <p>Voided volume (mL), mean change ± SD: G1: 19 ± 46 G2: 20 ± 42 G3: 7 ± 40 G1/G3: <i>P</i> = 0.056 G2/G3: <i>P</i> = 0.055</p> <p>Dry mouth, n (%): G1: 35 (34) G2: 20 (21) G3: 3 (6) G1/G2: <i>P</i> < 0.05 G1/G3: <i>P</i> < 0.05 G2/G3: <i>P</i> < 0.05</p> <p>Abdominal pain, n (%): G1: 4 (4) G2: 6 (6) G3: 2 (4)</p> <p>Constipation, n (%): G1: 2 (2) G2: 4 (4) G3: 2 (4)</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: fair</p> <p>Randomization: -</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: NR</p> <p>Drop-out rates: ++</p> <p>Power calculation: +</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: -</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Jacquetin and Wyndaele, 2001 (continued)			Voided volume (mL), mean (range): G1: 158 (43-382) G2: 150 (46-320) G3: 148 (23-284)	Headache, n (%): G1: 3 (3) G2: 3 (3) G3: 2 (4) Total AEs reported, n: G1: 84 G2: 78 G3: 26 Any AE, n (%): G1: 55 (53) G2: 39 (40) G3: 16 (31) G1/G3: $P < 0.05$ Discontinued due to AEs, n (%):* G1: 2 (2) G2: 3 (3) G3: 1 (2)	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Jarvis et al., 1981</p> <p>Country and setting: UK, Academic medical center</p> <p>Enrollment period: NR</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: Inpatient bladder drill vs outpatient drug therapy</p> <p>Groups: G1: inpatient bladder drill G2: flavoxate hydrochloride 200 mg tds and imipramine 25 mg tds x 4 wks</p> <p>N at enrollment: G1: 25 G2: 25</p> <p>N at follow-up: G1: 25 G2: 25</p> <p>Women, %: 100</p> <p>Age, mean ± SD (range): G1: 47 ± 15.4 (17-78) G2: 46 ± 12.8 (17-65)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Women • UDS-diagnosed detrusor instability <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • DM • Neurological abnormalities • UTIs • Taking a drug suspected of affecting lower urinary tract function • Genuine stress incontinence 	<p>UUI, n: G1: 25 G2: 25</p> <p>SUI, n: G1: 19 G2: 17</p> <p>Urgency, n: G1: 25 G2: 25</p> <p>Frequency, n: G1: 25 G2: 25</p> <p>Nocturia, n: G1: 21 G2: 19</p> <p>Duration of symptoms (years), mean ± SD (range): G1: 4.3 ± 2.7 (1-15) G2: 5.4 ± 3.2 (1-20)</p> <p>Bladder volume (mL), first sensation, mean: G1: 87 G2: 79</p> <p>Max cystometric capacity (mL), mean: G1: 381 G2: 353</p>	<p>UUI, n: G1: 4 G2: 11</p> <p>SUI, n: G1: 1 G2: 9</p> <p>Urgency, n: G1: 4 G2: 11</p> <p>Frequency, n: G1: 6 G2: 12</p> <p>Nocturia, n: G1: 4 G2: 13</p> <p>Continent, n (%): G1: 21 (84) G2: 14 (56) <i>P</i> < 0.05</p> <p>Symptom-free, n (%): G1: 19 (76) G2: 12 (48) <i>P</i> < 0.05</p> <p>Bladder volume (mL), first sensation, mean: G1: 152 G2: 140</p> <p>Max cystometric capacity (mL), mean: G1: 470 G2: 446</p> <p>Adverse events, drug therapy, n:* Dizziness: 8 Headache: 6 Dry mouth: 6 Nausea: 4 Drowsiness: 2 Vomiting: 1</p> <p>Discontinued due to AEs, n: Dizziness: 1 Headache: 1 Vomiting: 1</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: -</p> <p>Masking: -</p> <p>Pt selection criteria: -</p> <p>Loss to followup: ++</p> <p>Drop-out rates: ++</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: poor</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: -</p> <p>Measurement reliability: -</p> <p>Intervention description: -</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Jonas et al., 1997</p> <p>Country and setting: Austria, Germany, Sweden, Academic medical center</p> <p>Enrollment period: NR 1-2 week washout-run in to total 4 weeks efficacy/ tolerability</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: Tolterodine 1 mg b.i.d. vs tolterodine 2 mg b.i.d. vs placebo</p> <p>Groups: G1: Tolterodine 1 mg b.i.d. G2: Tolterodine 2mg b.i.d. G3: placebo</p> <p>N at enrollment: G1: 99 G2: 99 G3: 44</p> <p>N at follow-up: G1: 98 G2: 99 G3: 44</p> <p>Women, n (%): G1: 73 (74) G2: 76 (77) G3: 33 (75)</p> <p>Age, mean (range): G1: 59 (21, 81) G2: 57 (20, 83) G3: 57 (23, 92)</p> <p>Race/ethnicity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥18 • Detrusor overactivity (phasic activity of amplitude of ≥ 10 cm H2O or 1 strong detrusor contraction that caused end of infusion) • ≥ 8 voids/ day and ≥ 1 incontinence episode/ day, urinary urgency, or both <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • SUI • Hepatic disease • Renal disease • Any condition contraindicating anticholinergic activity • Indwelling catheter • Recurrent UTI • IC • Uninvestigated hematuria • Voiding difficulty at risk for urinary retention • Patients on anticholinergic therapy • Electrostimulation therapy or bladder training within 14 days prior to enrollment 	<p>Urinary incontinence, n (%): G1: 79 (80) G2: 83 (84) G3: 38 (86)</p> <p>Voids/day, mean (range): G1: 11.7 (0.0, 25.9) G2: 11.2 (4.1, 23.4) G3: 12.2 (7.3, 22.0)</p> <p>≥ 8 voids/day, n (%): G1: 94 (95) G2: 91 (92) G3: 42 (96)</p> <p>Volume (mL), first contraction, mean ± SD: G1: 142 ± 102 G2: 141 ± 102 G3: 140 ± 86</p> <p>Maximal height of wave (cmH₂O), mean ± SD: G1: 41 ± 34 G2: 52 ± 43 G3: 47 ± 34</p> <p>Max cystometric capacity (mL), mean ± SD: G1: 276 ± 142 G2: 272 ± 136 G3: 264 ± 116</p> <p>Postvoid residual volume (mL), mean ± SD: G1: 25 ± 45 G2: 28 ± 47 G3: 38 ± 53</p> <p>DO ≥ 10 cmH₂O, n (%): G1: 99(100) G2: 97 (98) G3: 44 (100)</p>	<p>Volume (mL), first contraction, mean ± SD: G1: 210 ± 163 G2: 230 ± 160 G3: 181 ± 142 G1/G3: <i>P</i> = 0.22 G2/G3: <i>P</i> = 0.030 G1/BL: <i>P</i> < 0.001 G2/BL: <i>P</i> < 0.001 G3/BL: <i>P</i> = 0.11</p> <p>Maximal height of wave (cmH₂O), mean ± SD: G1: 35 ± 36 G2: 37 ± 38 G3: 40 ± 36 G1/G3: <i>P</i> = 0.97 G2/G3: <i>P</i> = 0.17 G1/BL: <i>P</i> = 0.007 G2/BL: <i>P</i> < 0.001 G3/BL: <i>P</i> = 0.11</p> <p>Max cystometric capacity (mL), mean ± SD: G1: 294 ± 151 G2: 316 ± 156 G3: 268 ± 135 G1/G3: <i>P</i> = 0.43 G2/G3: <i>P</i> = 0.034 G1/BL: <i>P</i> = 0.36 G2/BL: <i>P</i> < 0.001 G3/BL: <i>P</i> = 0.80</p> <p>Postvoid residual volume (mL), mean ± SD: G1: 35 ± 52 G2: 46 ± 70 G3: 41 ± 62 G1/G3: <i>P</i> = 0.034 G2/G3: <i>P</i> = 0.042 G1/BL: <i>P</i> = 0.063 G2/BL: <i>P</i> = 0.0076 G3/BL: <i>P</i> = 0.99</p> <p>Dry mouth, n (%): G1: 8 (8) G2: 10 (10) G3: 1 (2)</p> <p>Dry mouth, mild, n (%): G1: 5 (5) G2: 7 (7) G3: 1 (2)</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: -</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: NR</p> <p>Drop-out rates: +</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Jonas et al., 1997 (continued)				<p>Dry mouth, moderate, n (%): G1: 3 (3) G2: 2 (2) G3: 0 (0)</p> <p>Dry mouth, severe, n (%): G1: 0 (0) G2: 1 (1) G3: 0 (0)</p> <p>UTI, n (%): G1: 5 (5) G2: 2 (2) G3: 2 (5)</p> <p>Accomodation abnormal, n (%): G1: 3 (3) G2: 5 (5) G3: NA</p> <p>Constipation, n (%): G1: 2 (2) G2: 3 (3) G3: 2 (5)</p> <p>Headache, n (%): G1: 3 (3) G2: 3 (3) G3: 1 (2)</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Karram et al., 2009</p> <p>Country and setting: USA, multicenter (61 sites)</p> <p>Enrollment period: NR</p> <p>Funding: Astellas Pharma US, Glaxo-SmithKline</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: Solifenacin 5 or 10 mg daily</p> <p>Groups: G1: solifenacin 5 mg daily (with dose adjustment at weeks 4 and 8) G2: placebo</p> <p>N at enrollment: G1: 372 G2: 367</p> <p>N at follow-up (%): G1: 357 G2: 350</p> <p>Age, mean: Total: 57</p> <p>Race/ethnicity, %: White: Total: 83.5</p> <p>Women, %: Total: 84.2</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 18 • OAB symptoms • ≥ 1 Urgency episode/day with or without leakage • >8 voids/day • ± nocturia <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • SUI • MUI with primary stress • UTI • Chronic bladder inflammation • Bladder stones • Bladder cancer • Severe constipation • Narrow angle glaucoma • Urinary retention • Gastric retention • Hypersensitivity to drugs • BOO 	<p>Urgency episodes/day, mean ± SD: G1: 6.15 ± 3.92 G2: 6.03 ± 3.90</p> <p>Warning time (sec), mean ± SD: G1: 136.4 ± 224.2 G2: 161.6 ± 333.9</p> <p>Incontinence episodes/ day, mean ± SD: G1: 2.82 ± 2.70 G2: 2.56 ± 2.72</p> <p>Voids/day, mean ± SD G1: 11.65 ± 3.72 G2: 11.70 ± 3.62</p>	<p>Urgency episodes/day, mean ± SD: G1: 2.24 ± 3.04 G2: 3.30 ± 3.84</p> <p>Urgency episodes/day, mean change ± SD: G1: -3.91 ± 3.54 G2: -2.73 ± 3.84 P < 0.0001</p> <p>Warning time (sec), mean change ± SD: G1: 186.4 ± 600.7 G2: 54.7 ± 393.5 P = 0.008</p> <p>Incontinence episodes/day, mean ± SD: G1: 0.72 ± 1.45 G2: 1.32 ± 2.65</p> <p>Incontinence episodes, mean change ± SD: G1: -2.10 ± 2.39 G2: -1.24 ± 2.30 P < 0.0001</p> <p>Voids/day, mean ± SD: G1: 8.98 ± 3.26 G2: 9.76 ± 3.50</p> <p>Voids/day, mean change ± SD: G1: -2.67 ± 3.31 G2: -1.94 ± 3.30 P = 0.0014</p> <p>Dry Mouth, %: G1: 25.3 G2: 9.0</p> <p>Constipation, %: G1: 14.8 G2: 9.3</p> <p>Blurred vision, %: G1: 3.8 G2: 1.1</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: -</p> <p>Method and blinding: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: +</p> <p>Drop-out rates: +</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Karram et al., 2009 (continued)				Headache, %: G1: 4.6 G2: 5.2 Dizziness, %: G1: 3.2 G2: 1.9 Fatigue, %: G1: 2.7 G2: 1.1	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Kelleher, Reese et al., 2002* Kelleher, Kreder, et al., 2002 [See evidence table for Van Kerrebroeck, et al., 2001]</p> <p>Country and setting: Multinational: Europe North America Australia/New Zealand Russia/Ukraine, Community</p> <p>Enrollment period: NR 12 week 12 month†</p> <p>Funding: Pharmacia Corporation</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Randomized placebo-controlled double blind parallel-group; Open label uncontrolled non-randomized extension†</p> <p>Intervention: Tolterodine ER vs placebo</p> <p>Groups: G1: Tolterodine ER 4mg daily G2: placebo G3: Tolterodine IR 2mg b.i.d.*</p> <p>N at enrollment: G1: 507 G2: 508 G3: 514</p> <p>N at follow-up: G1: 450 G2: 440 G3: 447</p> <p>Women, n (%): Total: 827 (81.5) G1: 417 (82.2) G2: 410 (80.7)* G3: 408 (79.4)*</p> <p>Age, mean ± SD: G1: 60.3 ± 14.4 G2: 61.1 ± 13.9 G3: 60.0 ± 14.3*</p> <p>Race/ethnicity: NR</p> <p>Extension:† Tolterodine ER 4mg daily</p> <p>Groups:† G4: KHQ G5: SF-36</p> <p>N at enrollment:† Total: 1077</p>	<p>Inclusion criteria: NR</p> <ul style="list-style-type: none"> • Age ≥ 18 • ≥ 8 voids/day on 7 day voiding diary • ≥ 5 UUI episodes/ week • OAB symptoms for at least 6 months <p>Exclusion criteria: NR</p>		<p>Patient rating of bladder condition, improvement, %: G1: 43 G2: 58 <i>P</i> = 0.001</p> <p>Patient rating of bladder condition, deterioration, %: G1: 13 G2: 7 <i>P</i> = 0.004</p> <p>KHQ, incontinence impact, mean change ± SD: G1: -15.68 ± 29.36 G2: -8.86 ± 26.65 <i>P</i> = 0.001</p> <p>KHQ, incontinence impact, 12 months, mean change (SE):† G4: -23.1 (1.1) <i>P</i> = 0.002</p> <p>KHQ, role limitations, mean change ± SD: G1: -17.93 ± 30.58 G2: -10.26 ± 29.20 <i>P</i> = 0.001</p> <p>KHQ, role limitations, 12 months, mean change (SE):† G4: -23.8 (1.1) <i>P</i> = 0.002</p> <p>KHQ, physical limitations, mean change ± SD: G1: -15.60 ± 29.98 G2: -8.73 ± 27.90 <i>P</i> = 0.001</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: -</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: +</p> <p>Drop-out rates: -</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: +</p> <p>Baseline OAB status: NR</p> <p>Baseline characteristics: -</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Kelleher, Reese et al., 2002	N at follow-up: † G4: 838 G5: 961			KHQ, physical limitations, 12 months, mean change (SE): † G4: -19.9 (1.1) <i>P</i> = 0.002	
Kelleher, Kreder, et al., 2002 (continued)	Women, n (%): † G4: 678 (80.9) G5: 793 (82.5)			KHQ, social limitations, mean change ± SD: G1: -8.49 ± 23.24 G2: -6.25 ± 22.76 <i>P</i> = 0.062	
	Age, yrs ± SD: † G4: 61.1 ± 13.5 G5: 60.2 ± 13.7			KHQ, social limitations, 12 months, mean change (SE): † G4: -11.5 (0.8) <i>P</i> = 0.002	
	Race/ethnicity, n, (%): † White: G4: 800 (95.5) G5: 927 (96.5) Black: G4: 28 (3.3) G5: 25 (2.5) Asian or Pacific: G4: 6 (0.7) G5: 6 (0.6) Mixed: G4: 4 (0.5) G5: 4 (0.4)			KHQ, personal relationships, mean change ± SD: G1: -5.66 ± 26.74 G2: -3.44 ± 23.15 <i>P</i> = 0.446	
				KHQ, personal relationships, 12 months, mean change (SE): † G4: -8.4 (1.2) <i>P</i> = 0.002	
				KHQ, emotions, mean change ± SD: G1: -9.31 ± 24.85 G2: -6.52 ± 23.98 <i>P</i> = 0.106	
				KHQ, emotions, 12 months, mean change (SE): † G4: -13.8 (0.9) <i>P</i> = 0.002	
				KHQ, sleep and energy, mean change ± SD: G1: -9.82 ± 24.45 G2: -5.08 ± 21.06 <i>P</i> = 0.006	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Kelleher, Reese et al., 2002</p> <p>Kelleher, Kreder, et al., 2002 (continued)</p>				<p>KHQ, sleep and energy, 12 months, mean change (SE):† G4: -11.7 (0.8) <i>P</i> = 0.002</p> <p>KHQ, severity (coping) measures, mean change ± SD: G1: -11.98 ± 22.04 G2: -6.12 ± 20.39 <i>P</i> = 0.001</p> <p>KHQ, severity (coping) measures, 12 months, mean change (SE):† G4: -13.8 (0.8) <i>P</i> = 0.002</p> <p>KHQ, general health perception, mean change ± SD: G1: -0.41 ± 17.55 G2: -0.12 ± 17.49 <i>P</i> = 0.900</p> <p>KHQ, general health perception, 12 months, mean change ± SD:† G4: -0.1 (0.6) <i>P</i> = 0.8302</p> <p>KHQ, symptom severity, mean change ± SD: G1: -2.90 ± 4.10 G2: -1.42 ± 3.99 <i>P</i> = 0.001</p> <p>KHQ, symptom severity, 12 months, mean change (SE):† G4: -11.7 (0.5) <i>P</i> = 0.002</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Kelleher, Reese et al., 2002 Kelleher, Kreder, et al., 2002 (continued)				<p>SF-36, physical summary, mean change \pm SD: G1: 0.97 \pm 7.34 G2: 0.72 \pm 6.57 <i>P</i> = 0.451</p> <p>SF-36, mental summary, mean change \pm SD: G1: 0.67 \pm 8.63 G2: 0.10 \pm 8.43 <i>P</i> = 0.684</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Kelleher et al., 2005</p> <p>[See evidence table for Chapple et al., 2004 and Cardozo et al. 2004]</p> <p>Country and setting: Multinational, multicenter</p> <p>Enrollment period: NR</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: 1 of 5 Pfizer (1) Yamanouchi (1) Lilly (1) Novartis (1) Q Med (1)</p>	<p>Design: Pooled analysis from the 40-wk open-label extension studies of 2 12-wk multinational, multicentre, double-blind RCTs</p> <p>Intervention: Solifenacin 5 mg vs solifenacin 10 mg vs placebo</p> <p>Groups: G1: solifenacin 5 mg qd G2: solifenacin 10 mg qd G2: placebo qd</p> <p>N at enrollment: 1637</p> <p>N at follow-up: 1347</p>	<p>Inclusion criteria: [See evidence table for Chapple et al., 2004 and Cardozo et al. 2004]</p> <p>Exclusion criteria: [See evidence table for Chapple et al., 2004 and Cardozo et al. 2004]</p>	<p>[See evidence table for Chapple et al., 2004 and Cardozo et al. 2004]</p>	<p>KHQ, general health perception, 12 weeks, mean change: G1: -4.3 G2: -4.0 G3: -2.3 G1/G3: $P < 0.001$ G2/G3: $P = 0.031$</p> <p>KHQ, incontinence impact, 12 weeks, mean change: G1: -24.7 G2: -27.3 G3: -18.2 G1/G3: $P < 0.001$ G2/G3: $P < 0.001$</p> <p>KHQ, role limitations, 12 weeks, mean change: G1: -20.6 G2: -22.7 G3: -15.4 G1/G3: $P < 0.001$ G2/G3: $P < 0.001$</p> <p>KHQ, physical limitations, 12 weeks, mean change: G1: -17.7 G2: -20.3 G3: -13.7 G1/G3: $P = 0.002$ G2/G3: $P < 0.001$</p> <p>KHQ, social limitations, 12 weeks, mean change: G1: -11.3 G2: -11.7 G3: -7.8 G1/G3: $P = 0.003$ G2/G3: $P = 0.015$</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: fair</p> <p>Randomization: -</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: ++</p> <p>Drop-out rates: ++</p> <p>Power calculation: +</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Kelleher et al., 2005 (continued)				<p>KHQ, personal relationships, 12 weeks, mean change: G1: -8.7 G2: -9.3 G3: -9.7 G1/G3: $P = 0.650$ G2/G3: $P = 0.747$</p> <p>KHQ, emotions, 12 weeks, mean change: G1: -16.0 G2: -17.7 G3: -12.3 G1/G3: $P < 0.001$ G2/G3: $P < 0.001$</p> <p>KHQ, sleep/energy, 12 weeks, mean change: G1: -13.8 G2: -14.4 G3: -10.0 G1/G3: $P = 0.002$ G2/G3: $P = 0.001$</p> <p>KHQ, severity measures, 12 weeks, mean change: G1: -10.5 G2: -13.2 G3: -7.3 G1/G3: $P < 0.001$ G2/G3: $P < 0.001$</p> <p>KHQ, symptom severity, 12 weeks, mean change: G1: -3.4 G2: -3.6 G3: -2.6 G1/G3: $P < 0.001$ G2/G3: $P < 0.001$</p> <p>KHQ, general health perception, 40 weeks, mean change: G1: -6.0 G2: -7.1 G3: -7.8</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Kelleher et al., 2005 (continued)				<p>KHQ, incontinence impact, 40 weeks, mean change: G1: -33.5 G2: -35.3 G3: -32.9</p> <p>KHQ, role limitations, 40 weeks, mean change: G1: -29.1 G2: -28.9 G3: -30.8</p> <p>KHQ, physical limitations, 40 weeks, mean change: G1: -25.7 G2: -27.0 G3: -26.0</p> <p>KHQ, social limitations, 40 weeks, mean change: G1: -15.6 G2: -18.1 G3: -15.7</p> <p>KHQ, personal relationships, 40 weeks, mean change: G1: -14.1 G2: -15.0 G3: -13.2</p> <p>KHQ, emotions, 40 weeks, mean change: G1: -22.2 G2: -24.8 G3: -21.0</p> <p>KHQ, sleep/energy, 40 weeks, mean change: G1: -17.9 G2: -19.5 G3: -18.8</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Kelleher et al., 2005 (continued)				KHQ, severity measures, 40 weeks, mean change: G1: -15.3 G2: -17.2 G3: -14.7 KHQ, symptom severity, 40 weeks, mean change: G1: -4.4 G2: -4.7 G3: -4.7	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Koonings et al., 1991</p> <p>Country and setting: US, Academic medical center</p> <p>Enrollment period: January 1986 to October 1987</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Cohort</p> <p>Intervention: Oxybutynin chloride 5 mg t.i.d. x 4 wks</p> <p>Groups: NA</p> <p>N at enrollment: 126</p> <p>N at follow-up: 114</p> <p>Women, %: 100</p> <p>Age, mean (range): 39 (21-74)</p> <p>Race/ethnicity: NR</p> <p>Menopausal, n: 44</p> <p>Parity, mean (range): 2 (0-12)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> Confirmed diagnosis of detrusor instability Uninhibited detrusor contraction > 15 cm H₂O on standing provocative urethra cystometry <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Urethritis (on urethroscopy) Cystitis (on cystoscopy) Neurologic findings on screening test of S2-S4 lower voiding center MUI Glaucoma 	<p>Uninhibited detrusor contraction starting prior to urethral pressure change, n (%): 73 (64%)</p> <p>Urethral pressure drop (≥ 20 cmH₂O) prior to detrusor contraction, n (%): 41 (6)</p>	<p>Good response, n (%): 66 (58)</p> <p>Poor response, n (%): 48 (42)</p> <p>Response, women with uninhibited detrusor contraction starting prior to urethral pressure, n (%): Good: 61 (81) Poor: 5 (12) <i>P</i> < 0.01</p> <p>Response, women with urethral pressure drop (≥ 20 cmH₂O) prior to detrusor contraction, n (%): Good: 12 (16) Poor: 36 (88) <i>P</i> < 0.01</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: NA</p> <p>Masking: NA</p> <p>Pt selection criteria: +</p> <p>Loss to followup: ++</p> <p>Drop-out rates: NR</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: -</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Kreder et al., 2002</p> <p>[See evidence table for Van Kerrebroeck, et al., 2001]</p> <p>Country and setting: Multinational; open label extension x 12 months</p> <p>Enrollment period: NR</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Prospective case series Open label, uncontrolled, non-randomized extension study of Phase III</p> <p>Intervention: Tolterodine ER 4 mg daily</p> <p>Groups: Tolterodine ER 4 mg daily</p> <p>N at enrollment: 1077</p> <p>N at follow-up (%): 759 (70.6)</p> <p>Age, mean (range): 60.3 (19.6-93.2)</p> <p>Race/ethnicity: NR</p> <p>Women, N (%): 883 (82)</p>	<p>Inclusion criteria: [See evidence table for Van Kerrebroeck, et al., 2001]</p> <p>Exclusion criteria: [See evidence table for Van Kerrebroeck, et al., 2001]</p>	<p>Frequency ≥ 8 voids/day, n (%): 1041 (96.7)</p> <p>UUI episodes/day ≥ 5, n (%): 1056 (98.1)</p> <p>Voided volume ≤ 200 mL, n (%): 1032 (95.8)</p> <p>Incontinence episodes/week, n (range): 21.6 (0.0-168.0)</p> <p>Voids/day, n (range): 10.9 (2.3-48.6)</p> <p>Voided volume (mL), mean (range): 137.7 (21.1-373.5)</p> <p>Previous treatment for OAB, n (%): 591 (54.9)</p> <p>Previous treatment for OAB with poor efficacy, n (%): 231 (39.1)</p>	<p>Incontinence episodes/week, 3 months, mean change \pm SD: -13.4 \pm 19.3 $P = 0.297$</p> <p>Incontinence episodes/week, 12 months, mean change \pm SD: -12.5 \pm 19.9 $P = 0.5888$</p> <p>Voids/day, 3 months, mean change \pm SD: -2.4 \pm 3.2 $P = 0.0077$</p> <p>Voids/day, 12 months, mean change \pm SD: -2.3 \pm 3.4 $P = 0.0661$</p> <p>Voided volume (mL), 3 months, mean change \pm SD: 42.9 \pm 55.6 $P = 0.6319$</p> <p>Voided volume (mL), 12 months, mean change \pm SD: 43.2 \pm 60.3 $P = 0.6426$</p> <p>Dry mouth, n (%): 139 (12.9)</p> <p>Constipation, n (%): 35 (3.3)</p> <p>Dyspepsia, n (%): 24 (2.2)</p> <p>Upper Respiratory Tract Infection, n (%): 43 (4.0)</p> <p>Bronchitis, n (%): 28 (2.6)</p> <p>UTI, n (%): 44 (4.1)</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: -</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: -</p> <p>Drop-out rates: -</p> <p>Power calculation: +</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: ++</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Kreder et al., 2002 (continued)				<p>Cystitis, n (%): 23 (2.1)</p> <p>Headache, n (%): 26 (2.4)</p> <p>Back pain, n (%): 35 (3.3)</p> <p>Flu-like illness, n (%): 28 (2.6)</p> <p>Serious adverse events related to tolterodine, n: Urinary retention: 1 Aggravated MS: 1 Med error: 1</p> <p>Withdrawal due to AEs (N= 107), n (%): Dry Mouth: 19 (1.8) Headache: 9 (0.8) Abd pain: 9 (0.8) Dizziness: 7 (0.7) UTI: 7 (0.7) Dyspepsia: 6 (0.6) Constipation: 6 (0.6) Xerophthalmia: 5 (0.5) Voiding disorder: 5 (0.5)</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Kreder et al., 2003</p> <p>Country and setting: US and Canada, Community</p> <p>Enrollment period: NR 16 week follow up</p> <p>Funding: Pharmacia Corporation</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Subgroup analysis of multicenter, single blind cohort</p> <p>Intervention: Tolterodine 1-2 mg b.i.d.</p> <p>Groups: G1: MUI with primary urge G2: UUI</p> <p>N at enrollment: G1: 171 G2: 552</p> <p>N at follow-up: G1: 170 G2: 551</p> <p>Women, n (%): G1: 165 (96.5) G2: 464 (84)</p> <p>Age, median (range): G1: 62 (21, 88) G2: 65 (20, 88)</p> <p>Race/ethnicity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 65 • History, PE, UDS consistent with urge incontinence • ≥ 4 episodes UI on 5 day voiding diary • ≥ 8 voids/day • Either urgency or ≥ 1 UI episodes / day <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • MUI with predominate stress component • Contraindications to antimuscarinic therapy • Hepatic/renal disease • Symptomatic or recurrent UTI • Hematuria • IC • Voiding difficulty with risk of retention • Concurrent bladder training • Electrostimulation therapy • Indwelling catheter • CIC • Women with reproduction potential • Pregnant/nursing • Concomitant anticholinergic meds/ treatment 	<p>Incontinence episodes/day, median (range): G1: 3 (1, 19) G2: 3 (1, 15)</p> <p>≥ 2 nocturia episodes/day, n (%): G1: 100 (58.5) G2: 334 (60.5)</p> <p>Using pads, n (%): G1: 122 (71) G2: 342 (62)</p> <p>Voided volume (mL), median (range): G1: 169 (62, 506) G2: 164 (31, 524)</p> <p>Duration of symptoms > 5 years, n (%): G1: 91 (53) G2: 259 (47)</p> <p>Previous drug therapy for OAB, n (%): G1: 85 (50) G2: 275 (50)</p>	<p>Incontinence episodes/day, median % change: G1: -67 G2: -75 <i>P</i> = NS</p> <p>Cure rate, achieved dryness, n (%): G1: 66 (39) G2: 243 (44) <i>P</i> = NS</p> <p>Voids/day, median % change: G1: -15 G2: -17 <i>P</i> = NS</p> <p>Cure rate, ≤ 8 voids/day dryness, n (%): G1: 40 (23.5) G2: 130 (24) <i>P</i> = NS</p> <p>Nocturia episodes/day, median % change: G1: -50 G2: -33 <i>P</i> = NS</p> <p>Cure rate, achieving ≤ 2 nocturnal voiding episodes/day, n (%): G1: 83 (83) G2: 254 (76) <i>P</i> = NS</p> <p>Cure rate, achieved no pad usage, n (%): G1: 26 (21) G2: 93 (27) <i>P</i> = NS</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: NA</p> <p>Masking: NA</p> <p>Pt selection criteria: +</p> <p>Loss to followup: -</p> <p>Drop-out rates: NR</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Kreder et al., 2003 (continued)				Voided volume (mL), median change (range): G1: 26.5 (-261, 195) G2: 27 (-345, 389) G1/BL: $P < 0.001$ G2/BL: $P < 0.001$	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Landis et al., 2004</p> <p>[See evidence table for van Kerrebroeck et al., 2001]</p> <p>Country and setting: Multinational, Community</p> <p>Enrollment period: NR 12 weeks</p> <p>Funding: Pharmacia</p> <p>Author industry relationship disclosures: 2 of 4 ALZA (1) Bristol-Myers Squibb (1) Pharmacia (2)</p>	<p>Design: RCT</p> <p>Intervention: Tolterodine ER vs placebo 1 week run-in period with 12 weeks treatment</p> <p>Groups: G1: Tolterodine ER 4 mg daily G2: placebo</p> <p>Stratified: a: 5-20 UUI episodes/week b: ≥ 21 UUI episodes/week</p> <p>N at enrollment: G1: 492 G1a: 321 G1b: 171 G2: 494 G2a: 284 G2b: 210</p> <p>N at follow-up: G1a: 321 G1b: 171 G2a: 284 G2b: 210</p> <p>Women, N (%): G1a: 262 (81.6) G1b: 143 (83.6) G2a: 223 (78.5) G2b: 178 (84.8)</p> <p>Age, mean ± SD: G1a: 60.86 ± 14.45 G1b: 60.04 ± 14.10 G2a: 60.61 ± 13.59 G2b: 61.84 ± 13.85</p> <p>Race/ethnicity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 18 • ≥ 5 UUI episodes/ week • ≥ 8 voids/day • Symptoms of OAB x 5 months <p>Exclusion criteria: NR</p>	<p>UUI episodes/ week, median (range): G1: 16 (5-168) G1a: 11 (5-20) G1b: 34 (21-168) G2: 18 (5-168) G2a: 11 (5-20) G2b: 31.5 (21-168)</p> <p>Voids/day, median (range): G1a: 9.57 (2.57-34.85) G1b: 10.71 (2.28-51.28) G2a: 10.28 (2.28-28.42) G2b: 11.14 (2-37.42)</p> <p>Voided volume (mL), median (range): G1a: 142.72 (36.19-338) G1b: 135.12 (51.45-270.52) G2a: 136.25 (32.51-237.5) G2b: 126.94 (21.16-373.46)</p> <p>Number prior Tx for OAB, n (%): G1a: 162 (50.6) G1b: 100 (58.5) G2a: 139 (49.1) G2b: 117 (55.7)</p> <p>Number prior Tx for OAB w/ good efficacy, n (%): G1a: 85 (52.8) G1b: 63 (63.0) G2a: 84 (60.4) G2b: 68 (58.6)</p>	<p>UUI episodes/ week, median change: G1: -9.0 G1a: -6.0 G1b: -21 G2: -5.0 G2a: -4.0 G2b: -9.5 G1/G2: $P < 0.0001$ G1a/G2a: $P < 0.0001$ G1b/G2b: $P < 0.0001$</p> <p>UUI episodes/ week, median % change: G1a: -71.42 G1b: -67.56 G2a: -38.46 G2b: -29.81 G1a/G2a: $P = 0.0264$ (adjusted) G1b/G2b: $P = 0.0221$ (adjusted)</p> <p>Voids/day, median change: G1a: -1.22 G1b: -1.9 G2a: -0.85 G2b: -0.4 G1a/G2a: $P < 0.04$ G1b/G2b: $P < 0.02$</p> <p>Voided volume (mL), median change: G1a: 24.0 G1b: 27.0 G2a: 4.0 G2b: 2.9 G1a/G2a: $P < 0.0001$ G1a/G2a: $P < 0.0001$</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: +</p> <p>Drop-out rates: -</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Lauti et al., 2008</p> <p>Country and setting: New Zealand, Academic</p> <p>Enrollment period: February 2003 to July 2003</p> <p>Funding: University of Otago</p> <p>Author industry relationship disclosures: None</p>	<p>Design: RCT pilot study, unmasked</p> <p>Intervention: Oxybutynin vs. bladder retraining vs. combination therapy</p> <p>Groups: G1: Oxybutynin 2.5 mg/day (daily dose could be increased by 2.5 mg every 5 days to a maximum of 15 mg/day) G2: Bladder retraining G3: Combination therapy</p> <p>N screened: 120</p> <p>N at enrollment: G1: 21 G2: 16 G3: 19</p> <p>N at 3 month follow-up: G1: 18 G2: 16 G3: 12</p> <p>N at 12 month follow-up: G1: 16 G2: 14 G3: 12</p> <p>Age, mean ± SD: G1: 53.8 ± 14.8 G2: 63.9 ± 17.2 G3: 47.6 ± 16.3</p> <p>Race/ethnicity: NR</p> <p>Women, %: Total: 100</p> <p>Parous, %: G1: 81 G2: 62.5 G3: 73.7</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age > 18 • Predominant UUI <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Predominant SUI • Contraindications to anticholinergic drugs • Current UTI • Neurological disease • Psychiatric disorder • Untreated co-existing pelvic organ prolapse below the hymenal ring • Obstructed voiding • Functional-reversible cause of incontinence • Inability to toilet independently • Limited fluency of written/spoken English • Current or recent use of any of the trial interventions 	<p>Urgency episodes/day, mean ± SD: G1: 3.8 ± 2.7 G2: 3.1 ± 2.2 G3: 3.5 ± 2.0</p> <p>Incontinence episodes/day, mean ± SD: G1: 2.2 ± 1.5 G2: 1.0 ± 1.1 G3: 1.8 ± 1.6</p> <p>Nocturia episodes/day, mean ± SD: G1: 1.1 ± 1.0 G2: 1.4 ± 1.0 G3: 0.8 ± 0.7</p> <p>Voids per day, mean ± SD: G1: 7.8 ± 2.8 G2: 8.0 ± 1.7 G3: 8.4 ± 2.5</p> <p>OAB-q total HRQL, mean ± SD: G1: 73.1 ± 17.4 G2: 69.5 ± 24.6 G3: 71.6 ± 21.5</p> <p>OAB-q severity, mean ± SD: G1: 47.0 ± 16.2 G2: 42.3 ± 17.7 G3: 45.9 ± 18.7</p> <p>OAB-q coping, mean ± SD: G1: 72.0 ± 21.6 G2: 66.2 ± 31.7 G3: 73.8 ± 26.2</p> <p>OAB-q concern, mean ± SD: G1: 68.2 ± 19.0 G2: 68.8 ± 27.6 G3: 63.8 ± 29.2</p> <p>OAB-q sleep, mean ± SD: G1: 63.1 ± 28.7 G2: 59.8 ± 29.9 G3: 55.1 ± 27.6</p>	<p>Urgency episodes/day, 3 mos, mean ± SD: G1: 2.2 ± 1.8 G2: 1.5 ± 2.1 G3: 1.7 ± 1.8</p> <p>Urgency episodes/day, 12 mos, mean ± SD: G1: 2.3 ± 2.5 G2: 1.9 ± 2.1 G3: 2.0 ± 1.1</p> <p>Incontinence episodes/day, 3 mos, mean ± SD: G1: 0.8 ± 0.8 G2: 0.1 ± 0.3 G3: 0.6 ± 0.8</p> <p>Incontinence episodes/day, 12 mos, mean ± SD: G1: 0.9 ± 0.0 G2: 0.9 ± 1.0 G3: 0.8 ± 0.7</p> <p>Nocturia episodes/day, 3 mos, mean ± SD: G1: 1.0 ± 0.5 G2: 0.8 ± 0.7 G3: 0.6 ± 0.5</p> <p>Nocturia episodes/day, 12 mos, mean ± SD: G1: 1.0 ± 0.9 G2: 1.2 ± 0.6 G3: 0.7 ± 0.7</p> <p>Voids/day, 3 mos, mean ± SD: G1: 6.7 ± 1.8 G2: 6.3 ± 1.6 G3: 6.7 ± 2.2</p> <p>Voids/day, 12 mos, mean ± SD: G1: 7.2 ± 1.1 G2: 6.8 ± 1.4 G3: 7.6 ± 1.5</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Method and blinding: -</p> <p>Pt selection criteria: +</p> <p>Loss to followup: +</p> <p>Drop-out rates: +</p> <p>Power calculation: +</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: ++</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Lauti et al., 2008 (continued)			<p>OAB-q social, mean ± SD: G1: 92.5 ± 14.6 G2: 85.4 ± 19.9 G3: 92.8 ± 18.6</p> <p>SF-12 quality of life, physical, mean ± SD: G1: 49.0 ± 9.6 G2: 41.7 ± 11.5 G3: 46.2 ± 10.6</p> <p>SF-12 quality of life, mental, mean ± SD: G1: 49.1 ± 9.3 G2: 53.1 ± 8.8 G3: 46.3 ± 8.3</p>	<p>OAB-q total HRQL, 3 mos, mean ± SD: G1: 82.3 ± 16.1 G2: 89.6 ± 9.4 G3: 91.8 ± 7.4</p> <p>OAB-q total HRQL, 12 mos, mean ± SD: G1: 87.9 ± 11.6 G2: 81.6 ± 19.3 G3: 88.9 ± 9.9</p> <p>OAB-q severity, 3 mos, mean ± SD: G1: 37.2 ± 22.0 G2: 16.8 ± 12.0 G3: 21.6 ± 10.9</p> <p>OAB-q severity, 12 mos, mean ± SD: G1: 24.6 ± 10.6 G2: 33.1 ± 16.6 G3: 21.9 ± 14.8</p> <p>OAB-q coping, 3 mos, mean ± SD: G1: 79.2 ± 22.1 G2: 91.6 ± 9.5 G3: 92.7 ± 9.4</p> <p>OAB-q coping, 12 mos, mean ± SD: G1: 89.2 ± 13.7 G2: 81.5 ± 23.7 G3: 90.5 ± 10.0</p> <p>OAB-q concern, 3 mos, mean ± SD: G1: 78.6 ± 18.0 G2: 87.7 ± 14.5 G3: 90.2 ± 12.4</p> <p>OAB-q concern, 12 mos, mean ± SD: G1: 85.3 ± 15.5 G2: 81.7 ± 19.7 G3: 85.2 ± 13.4</p> <p>OAB-q sleep, 3 mos mean ± SD: G1: 77.7 ± 24.9 G2: 81.3 ± 14.6 G3: 85.0 ± 19.6</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Lauti et al., 2008 (continued)				<p>OAB-q sleep, 12 mos, mean ± SD: G1: 79.9 ± 18.3 G2: 72.0 ± 24.5 G3: 83.2 ± 18.4</p> <p>OAB-q social, 3 mos, mean ± SD: G1: 96.4 ± 9.7 G2: 95.6 ± 7.0 G3: 98.9 ± 1.9</p> <p>OAB-q social, 12 mos, mean ± SD: G1: 97.3 ± 7.1 G2: 91.9 ± 14.2 G3: 97.3 ± 6.9</p> <p>SF-12 quality of life, physical G1: 50.6 ± 8.0 G2: 42.1 ± 12.7 G3: 48.4 ± 10.8</p> <p>SF-12 quality of life, physical, 12 mos, mean ± SD: G1: 50.0 ± 7.3 G2: 45.1 ± 13.9 G3: 45.3 ± 13.4</p> <p>SF-12 quality of life, mental, 3 mos, mean ± SD: G1: 50.4 ± 9.6 G2: 51.2 ± 9.5 G3: 46.7 ± 7.6</p> <p>SF-12 quality of life, mental, 12 mos, mean ± SD: G1: 49.6 ± 7.5 G2: 50.1 ± 10.7 G3: 50.6 ± 8.4</p> <p>Dry mouth, n (%): G1: 3 (21) G2: 5 (46) G3: 5 (42)</p> <p>Headaches, n (%): G1: 6 (43) G2: 1 (11) G3: 7 (58)</p> <p>Dizziness, n (%): G1: 4 (29) G2: 2 (20) G3: 3 (25)</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Lauti et al., 2008 (continued)				Constipation, n (%): G1: 3 (21) G2: 3 (27) G3: 3 (27) Fatigue, n (%): G1: 9 (64) G2: 5 (46) G3: 7 (64)	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Layton et al., 2001</p> <p>Country and setting: England, Community</p> <p>Enrollment period: November 1998 to May 1999</p> <p>Funding: Pharmacia Upjohn</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Prospective observational cohort</p> <p>Intervention: Tolterodine prescription event monitoring in UK</p> <p>Groups: NA</p> <p>N at enrollment: 35,295 had commenced treatment</p> <p>26,991 green forms mailed out</p> <p>14,526 returned forms</p> <p>Response rate: 53.8%</p> <p>Age, mean ± SD: 62.7 ± 16.4</p> <p>Women, N (%): 9965 (68.6)</p> <p>Parity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Patients prescribed tolterodine in general practice in UK <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • None 		<p>Dry mouth, n: 250</p> <p>Unspecified adverse effects, n: 168</p> <p>Headache, n: 123</p> <p>Constipation, n: 78</p> <p>General malaise, n: 78</p> <p>Hallucinations, n: 23</p> <p>Palpitations/tachycardia, n: 42</p> <p>Other cardiac arrhythmias, n: 29</p> <p>Chest pains, n: 87</p>	<p>Quality: Overall quality score: good</p> <p>INTERNAL VALIDITY: good</p> <p>Randomization: NA</p> <p>Masking: NA</p> <p>Pt selection criteria: +</p> <p>Loss to followup: NA</p> <p>Drop-out rates: NA</p> <p>Power calculation: +</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: NR</p> <p>Baseline characteristics: +</p> <p>Length of followup: ++</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: -</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Lee et al., 2002</p> <p>Country and setting: South Korea, University</p> <p>Enrollment period: NR</p> <p>Funding: Pharmacia Corp</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: Tolterodine 2mg b.i.d. vs Oxybutynin 5mg b.i.d.</p> <p>Groups: G1: Tolterodine 2mg b.i.d. G2: Oxybutynin 5mg b.i.d.</p> <p>N at enrollment: G1: 112 G2: 116</p> <p>N at follow-up: G1: 97 G2: 90</p> <p>Women, n (%): G1: 84 (74) G2: 92 (79)</p> <p>Age, mean (range): G1: 52 (27, 82) G2: 52 (20, 86)</p> <p>Race/ethnicity (%): Asian: G1: 100 G2: 100</p> <p>BMI, kg/m² (range): G1: 23 (17, 32.5) G2: 23.5 (16, 38)</p> <p>Previous drug therapy: N (%) G1: 36 (32) G2: 26 (22)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 18 • OAB symptoms > 6 mos • ≥ 8 voids/day, with or without incontinence (measured by diary) <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • SUI • Women not using reliable contraception • Pregnant or nursing • Prior treatment with anticholinergic < 2 wks • Renal or hepatic disease • Narrow angle glaucoma • Urinary retention • Gastric retention • Hypersensitivity to drugs • UTI • IC • Hematuria • BOO • Concomitant bladder training, e-stim treatment • Indwelling catheter • CIC • Concomitant treatment for OAB ≤ 2 mos 	<p>Incontinence episodes/day, mean (range): G1: 2.6 (0.3, 9.3) G2: 2.4 (3.0, 14.7)</p> <p>Patients with incontinence episodes, n (%): G1: 46 (41) G2: 42 (36)</p> <p>Voids/day, mean (range): G1: 12.2 (8.0, 23.7) G2: 12.4 (7.7, 29.7)</p>	<p>Incontinence episodes/day, mean change ± SD (% change): G1: -2.2 ± 2.3 (-85) G2: -1.4 ± 1.8 (-58)</p> <p>G1/BL: P = 0.0001 G1/BL: P = 0.0001 G1/G2: P = 0.10</p> <p>Voids/ day, mean change ± SD (% change): G1: -2.6 ± 2.9 (-21) G2: -1.8 ± 4.2 (-15)</p> <p>G1/BL: P = 0.0001 G1/BL: P = 0.0001 G1/G2: P = 0.14</p> <p>Benefit of Tx, %: G1: 45 G2: 46 G1/G2: P = NS</p> <p>Patients reporting adverse events, n (%): G1: 62 (55) G2: 94 (82)</p> <p>Discontinued due to AEs, n (%): G1: 11 G2: 18</p> <p>Dry mouth, n (%): G1: 39 (35) G2: 72 (63) G1/G2: P = 0.001</p> <p>Dry mouth, mild, n (%): G1: 29 (26) G2: 40 (35)</p> <p>Dry mouth, moderate, n (%): G1: 9 (8) G2: 26 (23)</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: -</p> <p>Drop-out rates: -</p> <p>Power calculation: +</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: -</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Lee et al., 2002 (continued)				<p>Dry mouth, severe, n (%): G1: 1 (1) G2: 6 (5)</p> <p>Voiding disorder, n (%): G1: 10 (9) G2: 16 (14)</p> <p>Dyspepsia, n (%): G1: 8 (7) G2: 6 (5)</p> <p>Abdominal pain, n (%): G1: 6 (5) G2: 6 (5)</p> <p>Headache, n (%): G1: 4 (4) G2: 6 (5)</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Leung et al., 2002</p> <p>Country and setting: Hong Kong, Academic, uro-gynecology centers</p> <p>Enrollment period: April 2000 to December 2000</p> <p>Funding: Pharmacia Limited</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: oral tolterodine 2 mg vs. oxybutynin 5 mg b.i.d. for 10 weeks</p> <p>Groups: G1: tolterodine 2 mg G2: oxybutynin 5 mg b.i.d.</p> <p>N at enrollment: G1: 53 G2: 53</p> <p>N at follow-up: G1: 44 G2: 45</p> <p>Women, %: 100</p> <p>Age, median (IQR): G1: 51 (44, 67) G2: 49 (43, 57)</p> <p>Race/ethnicity, %: Hong Kong Chinese: 100</p> <p>Menopausal, %: G1: 66.0 G2: 45.3 G1/G2: $P = 0.073$</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • UDS-confirmed detrusor instability • Age ≥ 18 • Diagnosis of OAB (phasic detrusor contraction with an amplitude ≥ 15 cmH₂O by ICS criteria • ≥ 8 voids/day • ≥ 1 incontinent episodes/day <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Genuine SUI • Voiding difficulty (max flow rate < 10 mL/s with a residual volume of > 200 mL) • Recurrent or acute UTIs • Intermittent or indwelling catheter • Hematuria or bladder cancer • Currently on treatment for OAB or on anticholinergic medications • Psychiatric disease or cognitive impairment (by history or MMSE) • Cardiac, hepatic, renal, or hematologic disorder • Contraindications for antimuscarinic agents • Pregnant or lactating • Women not using reliable contraception 	<p>Overall severity, VAS score, wk 0, median (IQR): G1: 5.2 (5.0, 7.85) G2: 5.7 (5.0, 8.0)</p> <p>Peak flow rate (mL/s), median (IQR): G1: 13.1 (10.0, 17.4) G2: 13.9 (11.2, 21.2)</p> <p>Max cystometric capacity (mL), median (IQR): G1: 333 (263.3, 400.8) G2: 373 (293, 416.8)</p>	<p>Overall severity, VAS score, wk 4, median (IQR): G1: 5.0 (5.0, 7.0) G2: 5.0 (3.0, 6.0)</p> <p>Overall severity, VAS score, wk 10, median (IQR): G1: 5.0 (3.5, 6.0) G2: 5.0 (4.0, 6.0)</p> <p>$P < 0.005$ for change with duration of treatment, $P < 0.005$ for linear pattern</p> <p>Perceived change in symptoms, double-sided VAS score, wk 4, median (IQR): G1: 1.0 (0.0, 2.5) G2: 2.0 (0.0, 3.0)</p> <p>Perceived change in symptoms, double-sided VAS score, wk 10, median (IQR): G1: 1.0 (0.0, 2.0) G2: 2.0 (0.0, 3.0) G1/G2: $P = 0.053$</p> <p>ITT analysis G1/G2: $P = 0.047$ per-protocol analysis</p> <p>Urinary pad weight (g), ITT analysis, median change (IQR): G1: -5.0 (-54.75, 0) G2: 0.0 (-4.5, -0.75) G1/G2: $P = 0.019$</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: +</p> <p>Drop-out rates: -</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: -</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Leung et al., 2002 (continued)				<p>Urinary pad weight (g), per-protocol analysis, median change (IQR): G1: -5.0 (-54.75, 0) G2: 0.0 (-5.0, 0) G1/G2: $P = 0.031$</p> <p>Drug compliance, %: G1: 75 G2: 87.5</p> <p>Significance of variables, change with duration of treatment, ITT analysis: XQ overall dryness: $P < 0.001$ XQ uncomfortable: $P < 0.001$ XQ sleep: $P = 0.020$ XQ speak: $P = 0.042$ XQ swallow: $P = 0.002$ XQ liquid: $P = 0.005$ XQ dentures: $P = 0.361$ Diary frequency: $P < 0.001$ Diary urgency episodes: $P = 0.109$ Diary incontinence episodes: $P = 0.203$ Diary pad use: $P = 0.593$</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Leung et al., 2002 (continued)				<p>Significance of variables, between groups, ITT analysis: XQ overall dryness: $P = 0.062$ XQ uncomfortable: $P = 0.285$ XQ sleep: $P = 0.626$ XQ speak: $P = 0.652$ XQ swallow: $P = 0.197$ XQ liquid: $P = 0.451$ XQ dentures: $P = 0.480$ Diary frequency: $P = 0.965$ Diary urgency episodes: $P = 0.672$ Diary incontinence episodes: $P = 0.993$ Diary pad use: $P = 0.665$</p> <p>Drug compliance rate (IQR): G1: 75.0 (8.9, 98.8) G2: 87.5 (11.4, 99.3)</p> <p>Withdrawal rate, %: G1: 17.0 G2: 15.1</p> <p>Side effects, %: G1: 60.4 G2: 49.1</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Lose et al., 2000</p> <p>Country and setting: Denmark, 26 clinics in the Dept of Obstetrics & Gynaecology Outpatient Clinic</p> <p>Enrollment period: September 1994 to April 1996</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT parallel group with active control, sequential numbers</p> <p>Intervention: oestradiol-releasing vaginal ring vs. oestriol vaginal pessaries (DANUGA study)</p> <p>Groups: G1: 7.5 mg oestradiol per 24 h oestradiol-releasing vaginal ring x 12 wks G2: 1-0.5 mg oestriol vaginal pessaries per day x 3 wks followed by 1 every 2 days for 21 wks</p> <p>N at enrollment: G1: 134 G2: 117</p> <p>N at follow-up: G1: 129 G2: 114</p> <p>Age, yrs ± SD: G1: 65.6±9.5 G2: 66.8 ± 9.1</p> <p>Race/ethnicity: NR</p> <p>Women, N (%): G1: 134 (100) G2: 117 (100)</p> <p>Height (cm) mean ± SD: G1: 164 ± 5.9 G2: 164 ± 6.4</p> <p>Body weight (kg) mean ± SD: G1: 69.8 ± 12.7 G2: 68.1 ± 12.4</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> Women who reported at least one bothersome lower urinary tract symptom appearing at least two years after spontaneous or surgical postmenopause <p>Exclusion criteria:</p> <ul style="list-style-type: none"> known or suspected oestrogen-dependent neoplasia or mammary, ovarian endometrioid) or corpus uteri malignancies vaginal bleeding of unknown origin clinically significant liver disease acute or intermittent porphyria, uterovaginal prolapse of grade II or III sex hormone treatment within the last 6 mos previous participation in clinical trials within 3 mos prior to inclusion 	<p>Persistence of symptoms (months), mean ± SD: G1: 74 ± 88 G2: 80 ± 101</p> <p>Urgency %: G1: 84 G2: 91</p> <p>Frequency %: G1: 75 G2: 73</p> <p>Urge Incontinence %: G1: 66 G2: 64</p> <p>Stress Incontinence %: G1: 67 G2: 58</p> <p>Nocturia %: G1: 60 G2: 66</p> <p>Dysuria %: G1: 25 G2: 23</p> <p>Mucosal atrophy, n (%): G1: 127 (95) G2: 115 (98)</p> <p>Vaginal pH, mean: G1: 6.0 G2: 6.0</p> <p>Urinary symptoms measured by VAS, mean: G1: 21.1 G2: 23.2</p> <p>Concomitant medication n (%): G1: 54 (41) G2: 50 (42)</p>	<p>Responder rate, 24 weeks, %: Urgency : G1: 51 G2: 56 Frequency : G1: 61 G2: 58 UUI: G1: 58 G2: 58 SUI: G1: 53 G2: 59 Nocturia: G1: 51 G2: 54 Dysuria: G1: 76 G2: 63 G1/G2: <i>P</i> = 0.282 Vaginal dryness: G1: 77 G2: 89 Dyspareunia: G1: 75 G2: 78</p>	<p>Quality: Overall quality score: fair INTERNAL VALIDITY: fair Randomization: + Masking: - Pt selection criteria: + Loss to followup: ++ Drop-out rates: + Power calculation: + Statistical issues: + EXTERNAL VALIDITY: fair Age: + Baseline OAB status: + Baseline characteristics: ++ Length of followup: ++ Measurement methods: + Measurement reliability: - Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Lose et al., 2000 (continued)		<ul style="list-style-type: none"> signs of vaginal irritation other than atrophy derived or signs of vaginal ulceration 		<p>Symptom free rate, 24 weeks, %: Urgency : G1: 27 G2: 33 Frequency : G1: 34 G2: 44 G1/G2: $P = 0.203$ UUI: G1: 33 G2: 34 SUI: G1: 34 G2: 41 Nocturia : G1: 31 G2: 35 Dysuria : G1: 67 G2: 52 G1/G2: $P = 0.244$ Vaginal dryness: G1: 57 G2: 83 G1/G2: $P = 0.019$ Dyspareunia : G1: 61 G2: 71</p> <p>Vaginal pH at 24 weeks, mean: G1: 4.8 G2: 4.8</p> <p>Subjective judgement of administration form, %:</p> <p>Excellent: G1: 60 G2: 14</p> <p>Good: G1: 27 G2: 34</p> <p>Acceptable: G1: 6 G2: 48</p> <p>Bad: G1: 2 G2: 3</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Lose et al., 2000 (continued)				<p>Unacceptable: G1: 3 G2: 2 <i>P</i> = 0.0001</p> <p>At least 1 adverse events, n (%): 49 (19.5) difference between groups = NS</p> <p>Vaginal discomfort, n : G1: 6 G2: 13</p> <p>Leukorrhea, n : G1: 7 G2: 7</p> <p>Vaginal itching, n: G1: 1 G2: 4</p> <p>Breast pain, n: G1: 2 G2: 3</p> <p>Vaginal haemorrhage, n: G1: 2 G2: 2</p> <p>Weight increase, n: G1: 2 G2: 3</p> <p>Breast discharge, n: G1: 1 G2: 3</p> <p>Abdominal pain, n: G1: 2 G2: 0</p> <p>Nausea, n: G1: 2 G2: 0</p> <p>Headache, n: G1: 1 G2: 1</p> <p>Pruritus, n: G1: 1 G2: 1</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Lose et al., 2000 (continued)				<p>Flushing, n: G1: 1 G2: 0</p> <p>Urinary urgency, n: G1: 1 G2: 0</p> <p>Change in body smell, n: G1: 0 G2: 1</p> <p>Malaise, n: G1: 0 G2: 1</p> <p>Hypertrichosis, n: G1: 0 G2: 1</p> <p>Stranguria, n: G1: 1 G2: 0</p> <p>Oedema, n: G1: 1 G2: 0</p> <p>Palpitation, n: G1: 1 G2: 0</p> <p>Increased sweating, n: G1: 1 G2: 0</p> <p>Uterine discomfort, n: G1: 0 G2: 1</p> <p>Amnesia, n: G1: 0 G2: 1</p> <p>Pelvic inflammation, n: G1: 1 G2: 0</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Macaulay et al., 1987</p> <p>Country and setting: UK, Specialty treatment center</p> <p>Enrollment period: NR</p> <p>Funding: Wellcome Trust, trustees of St. George's Hospital</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT (randomization not specified)</p> <p>Intervention: Brief eclectic psychotherapy, bladder training or medication</p> <p>Groups: G1: psychotherapy G2: bladder drill G3: Propantheline</p> <p>N at enrollment: G1: 19 G2: 16 G3: 15</p> <p>N at follow-up: G1: 18 G2: 15 G3: 14</p> <p>Women, %: 100</p> <p>Age: NR</p> <p>Race/ethnicity: NR</p> <p>Follow-up: 3 months</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Previous completion of survey on psych conditions w/ OAB • Detrusor instability on UDS or sensory urgency <p>Exclusion criteria: NR</p>	<p>Detrusor instability, n: G1: 10 G2: 8 G3: 8</p> <p>Sensory urgency, n: G1: 9 G2: 8 G3: 7</p> <p>Urinary frequency, n: G1: NR G2: NR G3: 10.8</p> <p>Bladder capacity (mL), mean: G1: 393 G2: NR G3: 323</p> <p>Bladder volume (mL), first sensation, mean: G1: 107 G2: 110 G3: 70</p> <p>Detrusor pressure rise (cm H₂O), mean: G1: NR G2: 45.5 G3: NR</p>	<p>Urinary frequency, mean: G1: NR G2: NR G3: 8.3 G3/BL: $P < 0.005$</p> <p>Bladder capacity (mL), mean: G1: 414 G2: NR G3: 368 G1/BL: $P = NS$ G3/BL: $P = NS$</p> <p>Bladder volume (mL), first sensation, mean: G1: 142 G2: 150 G3: 137 G1/BL: $P = NS$ G2/BL: $P < 0.05$ G3/BL: $P = 0.06$</p> <p>Detrusor pressure rise (cm H₂O), mean: G1: NR G2: 29.5 G3: NR G2/BL: $P < 0.05$</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: -</p> <p>Masking: -</p> <p>Pt selection criteria: -</p> <p>Loss to followup: ++</p> <p>Drop-out rates: ++</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: -, NR</p> <p>Baseline OAB status: NR</p> <p>Baseline characteristics: -</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Malone-Lee et al., 2003</p> <p>Country and setting: UK, Hospital</p> <p>Enrollment period: 1993-1999</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Prospective observational</p> <p>Intervention: Oxybutynin 2.5mg b.i.d. and bladder retraining</p> <p>Groups: G1: urinary frequency and urgency w/ detrusor instability G2: urinary frequency and urgency w/o detrusor instability</p> <p>N at enrollment: G1: 266 G2: 86</p> <p>N at follow-up: Total: 347 G1: NR G2: NR</p> <p>Age, mean (range): G1: 54.8 (20, 90) G2: 51.8 (21, 88)</p> <p>Race/ethnicity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Women • Age ≥ 18 • ≥ 8 voids/day <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Neurological disease • Significant stress incontinence • Symptomatic UTI • Interstitial cystitis 	<p>Incontinence episodes/day, median (95% CI): G1: 2 (0, 7) G2: 2 (0, 6.3)</p> <p>Void/day median (95% CI): G1: 14 (8, 24) G2: 12 (8, 22)</p>	<p>Incontinence episodes/day, median change (95% CI): G1: 0 (2, 6) G2: 0 (2, 6) G1/G2: <i>P</i> = 0.73</p> <p>Void/day, median change (95% CI): G1: -5 (1.9-14) G2: -5 (1-12.3) G1/G2: <i>P</i> = 0.61</p> <p>Dry mouth, n (%): G1: 219 (84) G2: 69 (70)</p> <p>Constipation, n (%): G1: 84 (32) G2: 19 (22)</p> <p>Heartburn, n (%): G1: 71 (27) G2: 20 (23)</p> <p>Dry skin, n (%): G1: 46 (18) G2: 4 (5) G1/G2: <i>P</i> = 0.01</p> <p>Headache, n (%): G1: 25 (10) G2: 3 (3.5)</p> <p>Dry eyes, n (%): G1: 12 (5) G2: 1 (1)</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: fair</p> <p>Randomization: NA</p> <p>Masking: NA</p> <p>Pt selection criteria: +</p> <p>Loss to followup: ++</p> <p>Drop-out rates: NR</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: +</p> <p>Length of followup: -</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Mattiason et al., 2003</p> <p>Country and setting: Sweden, Denmark, Norway (Tolterodine Scandinavian Study Group)</p> <p>Enrollment period: October 1999 to December 2000</p> <p>Funding: Pharmacia Corp</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT, single-blinded (balanced blocks of 4, computerized randomization list)</p> <p>Intervention: Pharmacologic ± behavioral therapy; Tolterodine 2mg b.i.d. ± bladder training (BT); Tolterodine dosage could be decreased to 1mg po b.i.d. during the first 2 wks if intolerable SE; BT taught with a written handout</p> <p>Groups: G1: Tolterodine + BT x 24 wks G2: Tolterodine x 24 wks</p> <p>N at enrollment: Total: 505</p> <p>N at follow-up: G1: 244 G2: 257</p> <p>N Completed treatment: G1: 77% G2: 79% Total: 74%</p> <p>Women, n (%) G1: 177 (73) G2: 201 (78)</p> <p>Age, median (range): G1: 62 (19, 86) G2: 63 (22, 86)</p> <p>Race/ethnicity: NR</p> <p>Follow-up: 24 weeks</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 18 • ≥ 8 voids/day and urinary urgency (± UUI) as determined by 1 wk bladder diary • With or without UUI • Women of reproductive age had to be using reliable birth control <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Contraindication to antimuscarinic therapy • Use of electrostimulation therapy or BT within prior 3 mo • Indwelling catheter or intermittent catheterization • Pregnancy or lactation • Use of anticholinergic agents or concomitant treatment for OAB (estrogen permitted) 	<p>Urgency episodes/day, mean (range): G1: 6.0 (0, 23.0) G2: 6.6 (0, 34.3)</p> <p>Incontinence episodes/day, mean (range): G1: 2 (0.3, 20.3) G2: 2.3 (0.3, 16.3)</p> <p>Void/day, mean (range): G1: 10.3 (7.3, 27.6) G2: 10.6 (7.7, 24.6)</p> <p>Duration of symptoms > 5 years, n (%): G1: 120 (49) G2: 124 (48)</p> <p>Previous drug therapy for OAB, n (%): G1: 40 (16) G2: 35 (14)</p> <p>Previous surgery affecting lower urinary tract, n (%): G1: 28 (11.5) G2: 30 (12)</p> <p>Symptoms caused at least moderate problems, %: G1: 92 G2: 93</p>	<p>Urgency episodes/day, median % change (IQR): G1: -38 (-76.7, -14.1) G2: -38 (-68.7, -8.0) G1/G2: P = 0.75</p> <p>Incontinence episodes/day, median IQR% change; n=301: G1: -87 (-100, -20) G2: -81 (-100, -41.8) G1/G2: P = 0.28</p> <p>Void/day, median % change (IQR): G1: -33 (-42.3, 21.3) G2: -25 (-38.8, -13.0) G1/G2: P < 0.001</p> <p>Voided volume (mL), median % change (IQR): G1: 31.5 (13.3, 56.2) G2: 20 (3.1, 45.4) G1/G2: P < 0.001</p> <p>Symptoms are "minor or less", %: G1: 66.5 G2: 61.5</p> <p>Overall improvement in symptoms, %: G1: 76 G2: 71</p> <p>Worsening of symptoms, %: G1: 3 G2: 5</p> <p>Dry mouth, n (%): G1: 76 (31) G2: 90 (35)</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: + Masking: + Pt selection criteria: + Loss to followup: - Drop-out rates: - Power calculation: + Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: + Baseline OAB status: + Baseline characteristics: ++ Length of followup: ++ Measurement methods: + Measurement reliability: + Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Mattiason et al., 2003 (continued)				<p>Headache, n (%): G1: 15 (6) G2: 21 (8)</p> <p>Constipation, n (%): G1: 7 (3) G2: 14 (5)</p> <p>≥ 1 SE, n (%): G1: 158 (65) G2: 177 (69) G1/G2: <i>P</i> = NS</p> <p>Withdrawal due to, %: AE: 15</p> <p>Withdrawal due to lack of efficacy: 3</p> <p>Withdrawal of consent : 2</p> <p>Protocol violations: 1</p> <p>Completed treatment: G1: 77% G2: 79%</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Messelink, 1999</p> <p>Country and setting: NR</p> <p>Enrollment period: NR</p> <p>Funding: Pharmacia & Upjohn</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Compilation of findings from four Phase II and eight Phase III studies with open label follow-up; not all trials have been previously published*</p> <p>Intervention: Tolterodine 2mg b.i.d.</p> <p>Groups: G1: tolterodine 2mg b.i.d. previously in 4 week placebo controlled trial; continued for 12 months open label G2: tolterodine 2mg b.i.d. previously in 12 week placebo controlled trial continued for 9 months open label</p> <p>N at enrollment: G1: 135 G2: 121</p> <p>N at follow-up: G1: 135 G2: 121</p> <p>Age, yrs (range): G1: 60 (18, 92) G2: 61 (18, 88)</p> <p>Race/ethnicity: NR</p> <p>Women, %: G1: 68 G2: 76</p> <p>Parity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Participants in prior trials (details NR) • Proceeded into long term open label study <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • NR 	<p>Incontinence episodes/day, mean: G1: 2.9 G2: 3.5</p> <p>Voids/day, mean: G1: 10.9 G2: 11.4</p> <p>Voided volume (mL), mean: G1: 159 G2: 159</p>	<p>Incontinence episodes/day, 3 months, mean: G1: ND G2: 1.4</p> <p>Incontinence episodes/day, 6 months, mean: G1: 1.4 G2: ND</p> <p>Incontinence episodes/day, 9 months, mean: G1: ND G2: 1.5</p> <p>Incontinence episodes/day, 12 months, mean: G1: 1.6 G2: ND</p> <p>Voids/day, 3 months, mean: G1: ND G2: 8.8</p> <p>Voids/day, 6 months, mean: G1: 8.6 G2: ND</p> <p>Voids/day, 9 months, mean: G1: ND G2: 8.9</p> <p>Voids/day, 12 months, mean: G1: 8.6 G2: ND</p> <p>Voided volume (mL), 3 months, mean: G1: ND G2: 201</p> <p>Voided volume, (mL), 6 months, mean: G1: 193 G2: ND</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: -</p> <p>Masking: -</p> <p>Pt selection criteria: +</p> <p>Loss to followup: NR</p> <p>Drop-out rates: NR</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: +</p> <p>Length of followup: ++</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Messelink, 1999 (continued)				<p>Voided volume (mL), 9 months, mean: G1: ND G2: 199</p> <p>Voided volume (mL), 12 months, mean: G1: 190 G2: ND</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Michel et al., 2002</p> <p>Country and setting: Germany, Community</p> <p>Enrollment period: NR 12 week follow up</p> <p>Funding: Pharmacia GmbH</p> <p>Author industry relationship disclosures: 1 of 4 Pharmacia(1) Sanofi-Synthelabo(1)</p>	<p>Design: Open-label observational post-marketing surveillance</p> <p>Intervention: Tolterodine</p> <p>Groups: Tolterodine Median dose: 2 mg daily</p> <p>Dose (mg), mean ± SD: 3.81 ± 1.16</p> <p>N at enrollment: 2,250</p> <p>N at follow-up: 1,979</p> <p>Women, n (%): 1,730 (76.9)</p> <p>Age, mean ± SD: 61.1 ± 13.8</p> <p>Race/ethnicity: NR</p> <p>BMI, kg/m² ± SD: 73.2 ± 11.3</p>	<p>Inclusion criteria: • Physician medical judgement</p> <p>Exclusion criteria: NR</p>	<p>Urgency episodes/day, mean ± SD: 8.4 ± 5.1</p> <p>Incontinence episodes/day, mean ± SD: 3.4 ± 4.2</p> <p>Voids/day, mean ± SD: 12.4 ± 4.3</p>	<p>Urgency episodes/day, mean ± SD: 2.0 ± 3.0 <i>P</i> = NR</p> <p>Incontinence episodes/day, mean ± SD: 0.8 ± 2.0 <i>P</i> = NR</p> <p>Voids/day, mean ± SD: 7.7 ± 2.7 <i>P</i> = NR</p> <p>Urgency, successful treatment, OR (95% CI): Gender, M/F: 0.764 (0.583, 1.001) <i>P</i> = 0.0508 Age, years: 0.981 (0.973, 0.990) <i>P</i> < 0.001 Frequency, BL episodes/day: 1.038 (0.997- 1.080) <i>P</i> = 0.0724 Urgency, BL episodes/day: 0.851 (0.823, 0.880) <i>P</i> < 0.001 Incontinence, BL episodes/day: 0.979 (0.950, 1.000) <i>P</i> = 0.1735 Tolterodine dose, mg/day: 0.913 (0.830, 1.005) <i>P</i> = 0.0623</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: NA</p> <p>Masking: NA</p> <p>Pt selection criteria: -</p> <p>Loss to followup: +</p> <p>Drop-out rates: -</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Michel et al., 2002 (continued)				<p>Incontinence, successful treatment, OR (95% CI): Gender, M/F: 1.453 (1.062, 1.990) <i>P</i> = 0.0196 Age, years: 0.978 (0.968, 0.987) <i>P</i> < 0.001 Frequency, BL episodes/day: 1.034 (0.993, 1.076) <i>P</i> = 0.1036 Urgency, BL episodes/day: 1.053 (1.018, 1.088) <i>P</i> = 0.0027 Incontinence, BL baseline episodes/day: 0.744 (0.716, 0.774) <i>P</i> < 0.001 Tolterodine dose, mg/day: 0.866 (0.784, 0.956) <i>P</i> = 0.0043</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Michel et al., 2002 (continued)				Frequency, successful treatment, OR (95% CI): Gender, M/F: 0.745 (0.552, 1.004) P = 0.0532 Age, years: 0.981 (0.971, 0.991) P = 0.001 Frequency, BL episodes/day: 0.735 (0.701, 0.771) P < 0.001 Urgency, BL episodes/day: 1.008 (0.975, 1.041) P = 0.6526 Incontinence, BL episodes/day: 0.969 (0.937, 1.002) P = 0.0644 Tolterodine dose, mg/day: 1.070 (0.957, 1.196) P = 0.2335	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Michel et al., 2002 (continued)				<p>Effect on Global efficacy, OR (95% CI): Gender, M/F: 0.656 (0.526, 0.818) <i>P</i> = 0.0002 Age, years: 0.986 (0.980, 0.993) <i>P</i> < 0.001 Frequency, BL epsidoes/day: 1.002 (0.972, 1.033) <i>P</i> = 0.8896 Urgency, BL episodes/day: 1.009 (0.984, 1.033) <i>P</i> = 0.4906 Incontinence, BL episodes/day: 0.963 (0.939, 0.987) <i>P</i> < 0.0026 Tolterodine dose, mg/day: 1.000 (0.926, 1.080) <i>P</i> = 0.9971</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Michel et al., 2002 (continued)				<p>Effect on global tolerability of Tolterodine, OR (95% CI): Gender, M/F: 0.993 (0.790, 1.249) <i>P</i> = 0.9519 Age, years: 0.995 (0.988, 1.002) <i>P</i> = 0.1915 Frequency, BL episodes/day: 1.002 (0.971, 1.034) <i>P</i> = 0.8995 Urgency, BL episodes/day: 1.022 (0.997, 1.048) <i>P</i> = 0.08 Incontinence, BL episodes/day: 0.990 (0.965, 1.016) <i>P</i> < 0.459 Tolterodine dose, mg/day: 1.114 (1.028, 1.206) <i>P</i> = 0.008589</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Michel et al., 2007*</p> <p>Michel et al., 2005</p> <p>Country and setting: Germany, NR</p> <p>Enrollment period: November 2001 to June 2003</p> <p>Funding: Pharmacia (became Pfizer before publication)</p> <p>Author industry relationship disclosures: 5 of 5* 4SC (1) Astellas (1) Boehringer Ingelheim (2) Eli Lilly (2) Pfizer (4) Theravance (1) Anformed (1)</p>	<p>Design: Retrospective cohort, open label</p> <p>Intervention: Tolterodine ER</p> <p>Groups: G1: Incontinent G2: Continent</p> <p>N at enrollment: Total: 3,824 G1: 2,571 G2: 1,253</p> <p>N at follow-up: Total: 3,416</p> <p>Women, %: Total: 75.8 G1: 81.7 G2: 62.6 P < 0.001</p> <p>Age, mean ± SD: Total: 64.8 ± 13.3 G1: 66.3 ± 12.6 G2: 61.4 ± 14.1 P < 0.001</p> <p>Race/ethnicity: NR</p> <p>Follow-up: 9 months</p>	<p>Inclusion criteria: NR</p> <p>Exclusion criteria: NR</p>	<p>Urgency episodes/day, mean ± SD: Total: 7.9 ± 5.2 G1: 8.0 ± 5.2 G2: 7.7 ± 5.2 P = 0.1937</p> <p>Incontinence episodes/day, mean ± SD: G1: 4.8 ± 3.7 G2: NA</p> <p>Voids/day, mean ± SD: Total: 14.0 ± 4.6 G1: 14.1 ± 4.6 G2: 13.5 ± 4.4 P < 0.001</p> <p>Daytime voids/day, mean ± SD: Total: 10.7 ± 3.7 G1: 10.8 ± 3.7 G2: 10.4 ± 3.4 P = 0.0128</p> <p>Frequency ≥ 8 voids/day, %: G1: 95.7 G2: 95.2</p> <p>Nocturia episodes/day, mean ± SD: Total: 3.4 ± 1.7 G1: 3.5 ± 1.8 G2: 3.2 ± 1.6 P < 0.001</p> <p>Pad use/day, mean ± SD: G1: 3.4 ± 2.8 G2: 0.1 ± 0.6 P < 0.001</p> <p>Duration of symptoms (mos), mean ± SD: G1: 50 ± 53 G2: 40 ± 46 P < 0.001</p>	<p>Incontinence episodes/day, mean change ± SD: G1: -3.8 ± 3.5 G2: NA</p> <p>Urgency episodes/day, mean ± SD: Total: 1.6 ± 2.8</p> <p>No urgency, %: G1: 53.4 G2: 63.2</p> <p>Voids/day, mean ± SD: Total: 7.5 ± 3.0</p> <p>Daytime voids/day, mean ± SD: Total: 6.2 ± 2.3</p> <p>Frequency ≥ 8 voids/day, %: G1: 43.7 G2: 39.1</p> <p>Nocturia episodes/day, mean ± SD: Total: 1.4 ± 1.1</p> <p>Pad use/day, mean change ± SD: G1: -2.4 ± 2.5</p> <p>Limitation of daily activities, score change ± SD: G1: 4.49 ± 2.65 G2: 4.10 ± 2.51</p> <p>Limitations in daily life caused by bladder problems, score ± SD: G1: 7.59 ± 1.65 G2: 6.66 ± 1.69 P < 0.001</p> <p>Total adverse events, n (%): Total: 496 (13.0)</p> <p>Dry mouth, n (%): 299 (7.8)</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: NA</p> <p>Masking: NA</p> <p>Pt selection criteria: -</p> <p>Loss to followup: NR</p> <p>Drop-out rates: NA</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: +</p> <p>Length of followup: ++</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Michel et al., 2007*			Previous treatment, %:	Withdrew: 11%	
Michel et al., 2005 (continued)			G1: 50.8 G2: 44.8 <i>P</i> = 0.0007	Unable to tolerate medication: 2.8% Administrative reasons: 2.6% Lack of efficacy : 2.4% Patient request: 1.2%	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Michel et al., 2008</p> <p>Country and setting: Germany, 50% of office-based urologists</p> <p>Enrollment period: September 2004 to November 2005</p> <p>Funding: Astellas Pharma GmbH</p> <p>Author industry relationship disclosures: 3 of 4 Astellas (3) Bayer (1) Boehringer (1) Elbion (1) Eli Lilly (1) Pfizer (1)</p>	<p>Design: Case series</p> <p>Intervention: Solifenacin 5-10 mg qd x 12 wks (initial dose could be adjusted at first and second follow-up visits)</p> <p>Groups: NA</p> <p>N at enrollment: 4,450</p> <p>N at follow-up: 4,146</p> <p>Age, mean ± SD: 63.6 ± 13.1</p> <p>Weight (kg), mean ± SD: 75.5 ± 12.3</p> <p>Height (cm), mean ± SD: 167 ± 7</p> <p>Race/ethnicity: "the vast majority of OAB patients in Germany are Caucasian"</p> <p>Women, %: 83.5</p> <p>Follow-up: 12 weeks</p>	<p>Inclusion criteria: • Age ≥ 18</p> <p>Exclusion criteria: NR</p>	<p>Duration of OAB (years), mean ± SD: 2.9 ± 3.7</p> <p>Duration of OAB (years), median (range): 1.5 (0-34.6)</p> <p>Previous OAB drug treatment, %: Herbal drugs: 28 Oxybutynin: 23 Trospium chloride: 20 Tolterodine: 13 Topical estrogens: 12 No treatment: 30</p> <p>Previous non-pharma OAB treatment, %: Absorbent products: 51 Pelvic floor exercises: 41 Bladder training: 22 No treatment: 30</p> <p>Initial solifenacin dose, %: 5 mg: 93.4 10 mg: 6.1</p> <p>Final solifenacin dose, %: 5 mg: 78.7 10 mg: 20.8</p> <p>Heart rate (beats/min), mean ± SD: 75.2 ± 8.2</p> <p>Blood pressure (mm Hg), mean ± SD: 137 ± 15 / 82 ± 7</p>	<p>Mult regression analysis of the presence of trt-emergent AEs, OR (95% CI): Male sex: 0.840 (0.559, 1.265) <i>P</i> = 0.4042 Age: NA <i>P</i> = 0.0019 Age 41-50: 1.144 (0.363, 3.608) <i>P</i> = 0.8180 Age 51-60: 1.773 (0.614, 5.119) <i>P</i> = 0.2900 Age 61-70: 1.941 (0.682, 5.524) <i>P</i> = 0.2137 Age 71-80: 1.816 (0.627, 5.262) <i>P</i> = 0.2715 Age > 80: 3.902 (1.327, 11.472) <i>P</i> = 0.0133 BMI (kg/m²): NA <i>P</i> = 0.9052 Comedication present: 1.768 (1.219, 2.564) <i>P</i> = 0.0027 CHD/MI present: 0.997 (0.667, 1.491) <i>P</i> = 0.9892 CHF present: 1.243 (0.774, 1.997) <i>P</i> = 0.3684 Diabetes mellitus present: 1.083 (0.762, 1.541) <i>P</i> = 0.6557</p> <p>Heart rate (beats/min), mean ± SD: 74.5 ± 7.6</p> <p>Blood pressure (mm Hg), mean ± SD: 134 ± 13 / 81 ± 8</p> <p>Treatment-emergent AEs, n (%): 215 (4.8)</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: fair</p> <p>Randomization: NA</p> <p>Method and blinding: NA</p> <p>Pt selection criteria: +</p> <p>Loss to followup: ++</p> <p>Drop-out rates: ++</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: +</p> <p>Baseline OAB status: NR</p> <p>Baseline characteristics: +</p> <p>Length of followup: ++</p> <p>Measurement methods: +</p> <p>Measurement reliability: -</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Michel et al., 2008 (continued)				<p>Leukocyturia, n (%): 48 (1.1)</p> <p>Dry mouth, n (%): 31 (0.7)</p> <p>Haematuria, n (%): 26 (0.6)</p> <p>Constipation, n (%): 20 (0.4)</p> <p>Proteinuria, n (%): 17 (0.4)</p> <p>UTI, n (%): 15 (0.3)</p> <p>Nausea, n (%): 14 (0.3)</p> <p>Nitrate present in urine, n (%): 13 (0.3)</p> <p>Bacteriuria, n (%): 9 (0.2)</p> <p>Glycosuria, n (%): 7 (0.2)</p> <p>Discontinued dt AEs, n (%): 62 (1.4)</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Millard et al., 1999</p> <p>Country and setting: Australia & US, Academic medical centers</p> <p>Enrollment period: NR</p> <p>Funding: Pharmacia and Upjohn AB</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: Tolterodine 1 mg vs. tolterodine 2 mg vs. placebo</p> <p>Groups: G1: tolterodine 1 mg x 12 wks G2: tolterodine 2 mg x 12 wks G3: placebo x 12 wks</p> <p>N at enrollment: G1: 129 G2: 123 G3: 64</p> <p>N at follow-up: G1: 114 G2: 116 G3: 61</p> <p>Women, %: G1: 78 G2: 77 G3: 66</p> <p>Age, mean (range): G1: 60.1 (24-89) G2: 60.2 (24-83) G2: 60.5 (25-84)</p> <p>BMI, kg/m² (range): G1: 28.0 (17.2-47.2) G2: 27.3 (16.6-51.4) G2: 26.8 (19.9-47.1)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 18 • Cystometrically proved DO • Average urinary frequency ≥8 voids/day • UI (≥ 1 incontinence episodes/day) and/or urinary urgency • Adequate contraception <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • SUJ • Voiding difficulty (max flow rate < 10 mL/second with post-void residual volume > 200 mL) • Recurrent UTI • Interstitial cystitis • Hematuria • Bladder cancer • Intermittent catheterization or indwelling catheter • Hepatic or renal disease • Narrow angle glaucoma • Electro-stimulation for bladder training • Started primarily anticholinergic drug ≤ 14 days • Unstable dose of any treatment with anticholinergic side effects • Average total voided volume >3 L/24 h 	<p>Urgency, %: G1: 100 G2: 100 G3: 98</p> <p>Incontinence, %: G1: 88 G2: 90 G3: 86</p> <p>Incontinence episodes/day, mean ± SD: G1: 3.9 ± 4.0 G2: 3.6 ± 4.0 G3: 3.5 ± 3.2</p> <p>Incontinence episodes/day, mean (range): G1: 3.9 (0.1-24.0) G2: 3.6 (0.3-24.0) G3: 3.5 (0.1-18.4)</p> <p>≥ 8 voids/day, %: G1: 99 G2: 98 G3: 98</p> <p>Voids/day, mean ± SD: G1: 11.5 ± 3.7 G2: 11.2 ± 3.1 G3: 11.3 ± 3.4</p> <p>Voids/day, mean (range): G1: 11.5 (7.0-26.3) G2: 11.2 (6.3-22.0) G3: 11.3 (7.1-21.7)</p> <p>Duration of symptoms > 5 yrs, %: G1: 44 G2: 49 G3: 47</p> <p>Detrusor instability, %: G1: 87 G2: 88 G3: 91</p>	<p>Incontinence episodes/day, mean change ± SD (mean % change): G1: -1.7 ± 2.8 (-43) G2: -1.7 ± 2.5 (-50) G3: -1.3 ± 2.5 G1/G3: P = 0.27 G2/G3: P < 0.19</p> <p>Voids/day, mean change ± SD: G1: -2.3 ± 3.0 (-20) G2: -2.3 ± 2.1 (-25) G3: -1.4 ± 2.3 G1/G3: P = 0.0029 G2/G3: P = 0.045</p> <p>Patients achieving normalized voiding frequency, %: G1: 26 G2: 43 G3: 26 G2/G3: P = 0.012 G2/G1: P = 0.022</p> <p>Complete cure, %: G1: 19 G2: 11 G3: 10</p> <p>Compliance (12 of 14 wks), %: G1: 87 G2: 89</p> <p>Perceived improvement, %: G1: 41 G2: 59 G3: 38 G2/G3: P = 0.015 G2/G1: P = 0.018</p>	<p>Quality: Overall quality score: fair INTERNAL VALIDITY: poor Randomization: - Masking: + Pt selection criteria: + Loss to followup: ++ Drop-out rates: + Power calculation: + Statistical issues: - EXTERNAL VALIDITY: good Age: + Baseline OAB status: + Baseline characteristics: ++ Length of followup: + Measurement methods: + Measurement reliability: + Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Millard et al., 1999 (continued)		<ul style="list-style-type: none"> Treatment with any investigational drug during or 2 mos before study 	<p>Hyper reflexia, %: G1: 13 G2: 12 G3: 9</p> <p>Severe problems, patient perception, %: G1: 42 G2: 50 G3: 36</p> <p>Previous therapy for UI, %: G1: 50 G2: 47 G3: 48</p> <p>Previous therapy for UI, good efficacy response, %: G1: 39 G2: 48 G3: 45</p> <p>Previous lower urinary tract surgery, %: G1: 35 G2: 33 G3: 30</p> <p>Voided volume (mL), mean ± SD: G1: 151 ± 56 G2: 155 ± 52 G3: 158 ± 53</p> <p>Voided volume (mL), mean (range): G1: 151 (34-334) G2: 155 (32-304) G3: 158 (47-299)</p> <p>Bladder volume (mL), initial contraction, mean (range): G1: 167 (4-600) G2: 188 (7-520) G3: 212 (6-600)</p>	<p>Voided volume (mL), mean change ± SD: G1: 27 ± 41 G2: 36 ± 50 G3: 10 ± 47 G1/G3: <i>P</i> = 0.0059 G2/G3: <i>P</i> < 0.0001</p> <p>Minor noncholinergic and cholinergic AEs, %: G1: 74 G2: 73 G3: 78</p> <p>AEs per patient, mean: G1: 1.9 G2: 2.2 G3: 2.0</p> <p>Autonomic nervous system disorders, %: G1: 29 G2: 43 G3: 17</p> <p>Dry mouth, %: G1: 24 G2: 39 G3: 13</p> <p>Dry mouth, severe, %: G1: 1 G2: 2 G3: 2</p> <p>Dry eyes, %: G1: 2 G2: 6 G3: 2</p> <p>Serious AEs, n (%): G1: 5 (4) G2: 7 (5) G3: 5 (4) G3: 2</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Millard et al., 1999 (continued)			<p>Post voiding residual volume (mL), mean (range): G1: 28 (0-180) G2: 25 (0-178) G3: 33 (0-200)</p> <p>Max flow rate, (mL/sec), mean (range): G1: 18 (2-56) G2: 20 (3-70) G3: 19 (5-57)</p> <p>Detrusor overactivity \geq 10 cm, %: G1: 98 G2: 98 G3: 98</p> <p>Max wave height (cm), mean (range): G1: 44 (2-144) G2: 45 (2-165) G3: 52 (4-188)</p>	<p>Severe AEs possibly indicating cardiac dysfunction, %: G1: 0 G2: 2</p> <p>Nonserious cardiac AEs, %: G1: 3 G2: 3 G3: 6</p> <p>Discontinued due to AEs, n (%): G1: 2 (2) G2: 8 (6) G3: 0</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Millard, 2004</p> <p>Country and setting: International, Multicenter study, 54 sites</p> <p>Enrollment period: NR</p> <p>Funding: Pharmacia Corp</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: Tolterodine ± simplified pelvic-floor exercise regimen for 24 weeks</p> <p>Groups: G1: tolterodine 2 mg b.i.d. and simple pelvic floor muscle exercise (PFME) program G2: tolterodine 2 mg b.i.d. daily</p> <p>N at enrollment: G1: 227 G2: 253</p> <p>N at follow-up, 12 weeks, n (%): G1: 181 (79.7) G2: 205 (81.0)</p> <p>N at follow-up, 24 weeks, n (%): G1: 164 (72.2) G2: 190 (75.1)</p> <p>Women, n (%): G1: 169 (75.4) G2: 190 (75.4)</p> <p>Age, mean ± SD: G1: 53.6 ± 16.9 G2: 53.2 ± 17.4</p> <p>Race/ethnicity: Asian: G1: 176 (78.6) G2: 203 (80.6) White/mixed: G1: 48 (21.4) G2: 49 (19.4)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age 18-90 • ≥ 8 voids/day • Urgency • ≥ 1 UI episode/d • sx for ≥ 6 months <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Symptomatic SI • Significant PVR volume • Neuropathy • Glaucoma • UTI • + urine cytology • Use of concomitant anticholinergic therapy w/in 14 days of randomization 	<p>Urgency episodes/day, mean ± SD: G1: 4.2 ± 3.6 G2: 4.1 ± 4.0</p> <p>Urgency episodes/day, median (IQR): G1: 3.6 (1.3-6.0) G2: 3.0 (1.3-6.0)</p> <p>Incontinence episodes/day, mean ± SD: G1: 3.44 ± 3.4 G2: 3.21 ± 3.4</p> <p>Incontinence episodes/day, median (IQR): G1: 2.3 (1.3-4.0) G2: 2.9 (1.3-3.7)</p> <p>Voids/day, mean ± SD: G1: 11.87 ± 4.3 G2: 12.78 ± 5.6</p> <p>Voids/day, median (IQR): G1: 10.7 (9.0-13.7) G2: 11.3 (9.0-15.0)</p> <p>Voided volume (mL), mean ± SD: G1: 146.1 ± 67.7 G2: 146.0 ± 83.3</p> <p>Voided volume (mL), median (IQR): G1: 137 (98-186) G2: 132 (99-189)</p> <p>Duration of symptoms, n (%): ≤ 5 years: G1: 166 (74.1) G2: 173 (68.7) > 5 years: G1: 56 (25) G2: 78 (31)</p>	<p>Urgency episodes/day, wk 12, mean change ± SD (% change): G1: -1.9 ± 4.0 (-64.5) G2: -2.2 ± 3.6 (-69.8) G1/BL: <i>P</i> = 0.001 G1/BL: <i>P</i> = 0.001 G1/G2: <i>P</i> = NS</p> <p>Urgency episodes/day, wk 12, median change: G1: -1.6 G2: -1.3 G1/G2: <i>P</i> = 0.7658</p> <p>Urgency episodes/day, wk 24, mean change ± SD (% change): G1: -2.2 ± 3.6 (-78.7) G2: -2.7 ± 3.5 (-83) G1/BL: <i>P</i> = 0.001 G1/BL: <i>P</i> = 0.001 G1/G2: <i>P</i> = NS</p> <p>Urgency episodes/day, wk 24, median change: G1: -1.9 G2: -2.0 G1/G2: <i>P</i> = 0.3029</p> <p>Incontinence episodes/day, wk 12, mean change ± SD: G1: -2.15 ± 3.0 G2: -2.15 ± 2.7 G1/BL: <i>P</i> = 0.001 G1/BL: <i>P</i> = 0.001 G1/G2: <i>P</i> = 0.2215</p>	<p>Quality: Overall quality score: fair INTERNAL VALIDITY: poor Randomization: - Masking: - Pt selection criteria: + Loss to followup: ++ Drop-out rates: - Power calculation: + Statistical issues: - EXTERNAL VALIDITY: good Age: + Baseline OAB status: + Baseline characteristics: ++ Length of followup: ++ Measurement methods: + Measurement reliability: + Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Millard, 2004 (continued)			<p>Previous UT surgery, n (%): G1: 44 (19.6) G2: 43 (17.1)</p> <p>Prev anticholinergics, n (%): G1: 19 (8.5) G2: 14 (5.6)</p> <p>Prev UUI drug, n (%): G1: 101 (45.1) G2: 91 (36.1)</p> <p>Patient subjective rating of UI symptoms as “severe” or “many severe”, %: G1: 54.0 G2: 56.8</p>	<p>Incontinence episodes/day, wk 12, median change: G1: -1.6 G2: -1.6 G1/G2: <i>P</i> = 0.8251</p> <p>Incontinence episodes/day, wk 24, mean change ± SD (% change): G1: -2.23 ± 3.0 (-64) G2: -2.26 ± 3.0 (-70) G1/BL: <i>P</i> = 0.001 G1/BL: <i>P</i> = 0.001 G1/G2: <i>P</i> = NS</p> <p>Incontinence episodes/day, wk 24, median change: G1: -1.6 G2: -1.6 G1/G2: <i>P</i> = 0.8341</p> <p>Voids/day, wk 12, mean change ± SD (% change): G1: -2.68 ± 3.8 (-22) G2: -3.42 ± 4.6 (-26) G1/BL: <i>P</i> = 0.001 G1/BL: <i>P</i> = 0.001 G1/G2: <i>P</i> = 0.9478</p> <p>Voids/day, wk 24, mean change ± SD (% change): G1: -2.58 ± 5.0 (-22) G2: -3.58 ± 5.2 (-26) G1/BL: <i>P</i> = 0.001 G1/BL: <i>P</i> = 0.001 G1/G2: <i>P</i> = 0.3549</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Millard, 2004 (continued)				<p>Voided volume (mL), wk 12, median change (% change): G1: 20.4 (17.2) G2: 17.5 (15.8)</p> <p>Voided volume (mL), wk 12, median change (% change): G1: 21.1 (18.1) G2: 19.1 (15.4)</p> <p>Patient subjective report of improvement in bladder condition, wk 12, %: G1: 82.6 G2: 83.9</p> <p>Patient subjective report of improvement in bladder condition, wk 24, %: G1: 81.7 G2: 85.9</p> <p>Adverse events, n (%): G1: 22 (9.7) G2: 23 (9.1)</p> <p>Mild dry mouth, %: G1: 18.1 G2: 21.3</p> <p>Moderate dry mouth, %: G1: 7.5 G2: 5.1</p> <p>Severe dry mouth, %: G1: 4.0 G2: 3.2</p> <p>Headache, %: 6</p> <p>Constipation, %: 4.8</p> <p>Nausea, %: 2.7</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Millard, 2004 (continued)				Dry eyes, %: 2.5 Dizziness, %: 2.4	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Minassian et al., 2007</p> <p>Country and setting: Canada, Tertiary care center</p> <p>Enrollment period: February 2003 to December 2005</p> <p>Funding: Janssen-Ortho Inc.</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: Oxybutynin XL vs oxybutynin IR</p> <p>Groups: G1: oxybutynin XL 5 mg qd after 4 weeks non responders to oxybutynin XL 10 mg qd G2: oxybutynin IR 2.5 mg t.i.d. after 4 weeks non responders to oxybutynin 5 mg t.i.d.</p> <p>N at enrollment: G1: 39 G2: 33</p> <p>N at follow-up: G1: 35 G2: 27</p> <p>Age, mean ± SD: G1: 75 ± 6 G2: 73 ± 5</p> <p>Women, %: G1: 100 G2: 100</p> <p>Race/ethnicity: NR</p> <p>Height, cm ± SD: G1: 158 ± 6 G2: 159 ± 5</p> <p>Weight, kg ± SD: G1: 73 ± 14 G2: 76 ± 14</p> <p>BMI, kg/m² ± SD: G1: 29 ± 5 G2: 30 ± 5</p> <p>Parous, %: G1: 85 G2: 88</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Symptoms of OAB (ICS) • Female • Age > 65 • Community-dwelling • OAB main presenting symptom, 2-week washout if on anticholinergic medication <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Bedridden • Permanent indwelling catheter • MMSE score <24 • Glaucoma • Gastric retention or bowel obstruction • History of allergy to anticholinergic medications • Taking tricyclic antidepressants or anticholinesterase inhibitors • Post-void residual bladder volume >100 mL • History of neurologic disorder 	<p>UUI, %: G1: 95 G2: 94</p> <p>SUI, %: G1: 67 G2: 52</p> <p>Urgency, %: G1: 100 G2: 97</p> <p>Incontinence episodes/day, median (IQR): G1: 2 (0, 4) G2: 1 (0, 3)</p> <p>Frequency episodes/day, median (IQR): G1: 9 (7, 11) G2: 10 (8, 12)</p> <p>Voids/day, median (IQR): G1: 13 (10, 16) G2: 14 (11, 16)</p> <p>Nocturia episodes/day, median (IQR): G1: 2 (2, 3) G2: 3 (2, 4)</p> <p>Pads/day, median (IQR): G1: 1 (0, 3) G2: 1 (0, 3)</p> <p>Mini-mental state exam, median score (IQR): G1: 29 (29, 30) G2: 30 (28, 30)</p> <p>U-IIQ, activities score ± SD: G1: 2.7 ± 0.9 G2: 3.1 ± 1.1</p> <p>U-IIQ, travel score ± SD: G1: 2.4 ± 1.3 G2: 2.8 ± 1.6</p>	<p>Incontinence episodes/day, 12 wks, median (IQR): G1: 1 (0, 2) G2: 0 (0, 1) P = 0.05</p> <p>Frequency episodes/day, wk 1-12, median change (IQR): G1: -1.4 (-3.3, 0.4) G2: -1.3 (-4.1, 0.3) P = 0.65</p> <p>Voids/day, wk 12, median (IQR): G1: 11 (9, 13) G2: 11 (9, 14) P = 0.35</p> <p>Pads/day, 12 wks, median (IQR): G1: 0 (0, 2) G2: 0 (0, 1) P = 0.53</p> <p>U-IIQ, activities score ± SD: G1: 2.2 ± 1.0 G2: 2.1 ± 1.2 P = 0.73</p> <p>U-IIQ, travel score ± SD: G1: 2.0 ± 1.1 G2: 1.9 ± 1.2 P = 0.79</p> <p>U-IIQ, physical activities score ± SD: G1: 2.3 ± 1.3 G2: 1.9 ± 1.2 P = 0.45</p> <p>U-IIQ, feelings score ± SD: G1: 2.0 ± 1.1 G2: 1.9 ± 1.3</p> <p>U-IIQ, relationships score ± SD: G1: 1.4 ± 0.9 G2: 1.5 ± 1.0</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Masking: -</p> <p>Pt selection criteria: +</p> <p>Loss to followup: +</p> <p>Drop-out rates: -</p> <p>Power calculation: +</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Minassian et al., 2007 (continued)			U-IIQ, physical activities score ± SD: G1: 2.4 ± 1.2 G2: 2.6 ± 1.5	U-UDI, mean score ± SD: G1: 2.1 ± 1.0 G2: 1.7 ± 1.0 <i>P</i> = 0.10	
			U-IIQ, feelings score ± SD: G1: 2.5 ± 1.2 G2: 2.7 ± 1.5	Voided volume (mL), 12 wks, median (IQR): G1: 164 (129, 187) G2: 161 (114, 109) <i>P</i> = 0.78	
			U-IIQ, relationships score ± SD: G1: 1.7 ± 0.6 G2: 2.0 ± 1.4	Postvoid residual volume (mL), 12 wks, median (IQR): G1: 0 (0, 29) G2: 4 (0, 87) <i>P</i> = 0.33	
			U-UDI, mean score ± SD: G1: 2.9 ± 0.6 G2: 2.7 ± 0.8	Remained on med, 12 wks, %: G1: 70 G2: 61 <i>P</i> = 0.42	
			Voided volume (mL), median (IQR): G1: 142 (109, 192) G2: 138 (108, 165)	Experienced side effects, %: G1: 51 G2: 57	
			Postvoid residual volume (mL), median (IQR): G1: 0 (0, 23) G2: 0 (0, 22)	Dry mouth, n: G1: 14 G2: 16	
			Urine culture positive, % G1: 8 G2: 9	Gastro-intestinal, n: G1: 6 G2: 2	
			Previous prolapse/incontinence surgery, %: G1: 13 G2: 49		
			Positive cough test, lying, %: G1: 26 G2: 15		

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Minassian et al., 2007 (continued)			Positive cough test, standing, %: G1: 33 G2: 15		

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Molander et al., 1991</p> <p>Country and setting: Sweden, Academic hospital</p> <p>Enrollment period: NA</p> <p>Funding: Goteborg Medical Society, University of Goteborg</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Case series</p> <p>Intervention: 3mg estriol qd for 1 mo, then 1-2 mg estriol qd for 2 mos</p> <p>Groups: G1: Intervention among SUI G2: Intervention among UUI G3: Intervention among MUI</p> <p>N at enrollment: G1: 91 G2: 113 G3: 142</p> <p>N at follow-up: G1: 55 G2: 69 G3: 82</p> <p>Women, %: 100</p> <p>Age, mean (SE): G1: 70.1 (1.9) G2: 73.5 (1.2) G3: 72.6 (1.3)</p> <p>Race/ethnicity: NR</p> <p>Parity, mean (SE): G1: 2.0 (0.2) G2: 1.7 (0.1) G3: 1.9 (0.1)</p> <p>Menopausal age, mean (SE): G1: 49.6 (0.4) G2: 49.1 (0.4) G3: 48.8 (0.5)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> Self-reported UI (with UUI defined as involuntary urine loss preceded by the urge to void or uncontrollable voiding with little or no warning; SUI defined as involuntary urinary loss precipitated by coughing, sneezing, or physical exertion; and MUI defined as a combination of UUI+SUI) <p>Exclusion criteria: NR</p>	<p>Total leakage/48 hours (mL), mean (SEM): G1: 25.2 (6.1) G2: 44.4 (9.8) G3: 65.1 (11.5)</p> <p>Max single leakage/48 hours, mean (SE): G1: 14.0 (3.1) G2: 21.7 (3.6) G3: 31.5 (3.1)</p> <p>Voids/day, mean (SE): G1: 6.9 (0.3) G2: 8.4 (0.3) G3: 7.6 (0.3)</p> <p>Total voided volume/day (mL), mean (SE): G1: 1405 (68) G2: 1515 (71) G3: 1497 (68)</p> <p>Max voided volume (mL)/day, mean (SE): G1: 401 (22) G2: 396 (23) G3: 383 (18)</p>	<p>Total leakage/48 hours (mL), mean (SEM): G1: 19.7+/-6.1 G2: 37.3+/-11.9 G3: 26.2+/-6.8 G3/BL: <i>P</i> < 0.01</p> <p>Max single leakage/48 hours, mean (SE): G1: 11.3 (3.1) G2: 19.2 (4.6) G3: 16.1 (4.2) G3/BL: <i>P</i> < 0.05</p> <p>Voids/day, mean (SE): G1: 7.1 (0.4) G2: 7.9 (0.4) G3: 7.4 (0.4) G2/BL: <i>P</i> < 0.05</p> <p>Total voided volume/day (mL), mean (SE): G1: 1539 (67) G2: 1481 (74) G3: 1483 (88)</p> <p>Max voided volume (mL)/day, mean (SE): G1: 438 (24) G2: 399 (27) G3: 392 (26)</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: NA</p> <p>Masking: NA</p> <p>Pt selection criteria: +</p> <p>Loss to followup: -</p> <p>Drop-out rates: +</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Moore et al., 1990</p> <p>Moore and Sutherst, 1990*</p> <p>Country and setting: UK, academic health center</p> <p>Enrollment period: NR</p> <p>Funding: Tillots Laboratories NR*</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT with crossover</p> <p>Intervention: oxybutynin hydrochloride vs placebo</p> <p>Groups: G1:oxybutynin hydrochloride 3 mg t.i.d. G2:placebo</p> <p>N at enrollment: 53</p> <p>N at follow-up: G1: 27 G2: 22</p> <p>Age, mean ± SD: G1: 46.0 ± 11.7 G2: 46.4 ± 12.4</p> <p>Race/ethnicity: NR</p> <p>Parity, mean: G1: 2.3 G2: 2.2</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • UDS-defined idiopathic detrusor instability • Involuntary detrusor contractions > 30 cm H2O during filling phase of cystometry <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Neurological disorder • Urologic disorder • Age > 75 years • Genuine SI • Low-compliance bladder • Bacterial or interstitial cystitis • Previous treatment with oxybutynin 	<p>Voids/day, mean ± SD: G1: 11.1 ± 7.9 G2: 12.1 ± 5.7</p> <p>Volume at first desire to void mean mL ± SD: G1: 94.2 ± 66 G2: 107.4 ± 101</p> <p>Max detrusor filling pressure mean cm H2O, ±SD: G1: 55.3 ± 23 G2: 59.2 ± 26</p> <p>Max cystometric capacity (mL) G1: 275 ± 164 G2: 290 ± 168</p> <p>Residual urine volume mean mL ± SD: G1: 23.1 ± 34.2 G2: 22.7 ± 34.0</p> <p>Duration of OAB, mean yrs ± SD): G1: 7.5 ± 7.9 G2: 8.9 ± 9.0</p>	<p>Change in volume at first desire to void, period 1, mean mL ± SD (95% CI) G1: 70.0 ± 103 G2: 7.7 ± 76 G1/G2: P = 0.02 95% CI: (10, 113)</p> <p>Change in maximum detrusor filling pressure, mean ± SD: G1:-16.9 ± 19.5 G2:1.0 ± 28.7 G1/G2: P = 0.02 95% CI: (-32.5, -3.2)</p> <p>Change in maximum cystometric capacity, mean ± SD: G1:104.0 ± 131 G2:7.0 ± 103 G1/G2: P = 0.006 95% CI: (29, 165)</p> <p>Change in residual urine Volume, period 1: G1: 22.4 (79.3) G2: 7.7 (45.7) P = 0.42</p> <p>Change in residual urine Volume, period 2: G1: 10.0 ± 27.1 G2: 25.7 ± 55.6 P = 0.33</p> <p>Dry mouth on oxybutynin, %: 88</p> <p>Dry mouth on placebo, %: 33</p> <p>Mouth ulcers on oxybutynin, %: 16</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: -</p> <p>Drop-out rates: -</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: ++</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Moore et al., 1990 Moore and Sutherst, 1990 (continued)				<p>Mouth ulcers on placebo, %: 0</p> <p>Constipation on oxybutynin, % 12.5</p> <p>Constipation on placebo, % 0</p> <p>Drowsiness on oxybutynin, %: 12.5</p> <p>Drowsiness on placebo, %: 7</p> <p>Nausea on oxybutynin, %: 8.3</p> <p>Nausea on placebo, %: 2.3</p> <p>Initial hesitancy on oxybutynin, %: 4.2</p> <p>Initial hesitancy on placebo, %: 2.3</p> <p>Dizziness on oxybutynin, %: 4.2</p> <p>Dizziness on placebo, %: 7.0</p> <p>Metallic taste on oxybutynin, %: 2.4</p> <p>Metallic taste on placebo, %: 2.3</p> <p>Crown Crisp Experimental Index, >50% improvement:*</p> <p>Free floating anxiety: 6.4</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Moore et al., 1990 Moore and Sutherst, 1990 (continued)				Phobic anxiety: 5.3 Obsessionalism: 6.2 Somatic complaints: 6.2 Depression: 4.0 Hysteria: 2.6 Total: 30.7 Crown Crisp Experimental Index, <49% improvement:* Free floating anxiety: 8.8 Phobic anxiety: 6.8 Obsessionalism: 8.1 Somatic complaints: 8.9	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Nitti et al.,2007</p> <p>Country and setting: US, 83 center</p> <p>Enrollment period: October 2003 to February 2005</p> <p>Funding: Schwarz BioSciences GmbH and Pfizer, Inc</p> <p>Author industry relationship disclosures: 9 of 9 Allergan (3) Astellas (3) Gynecare (1) Novartis (2) Pfizer (5) Boston Scientific (1) Espirito (4) Ortho (1) Watson (1) Indevus (1) GlaxoSmithKline (1) Schwarz Biosciences (3)</p>	<p>Design: RCT</p> <p>Intervention: Placebo, Fesoteorine 4 or 8 mg, bladder diary before randomization, 2, 8, 12 wks</p> <p>Groups: G1: Placebo G2: Fesoterodine 4 mg/d G3: Fesoterodine 8 mg/d</p> <p>N at enrollment: Total: 836 G1: 274 G2: 283 G3: 279</p> <p>N at follow-up: G1: 266 G2: 267 G3: 267</p> <p>Age, mean (range): Total: 59 (21, 91) G1: 59 (24, 88) G2: 59 (21, 85) G3: 59 (23, 91)</p> <p>Women, %: Total: 76 G1: 74 G2: 76 G3: 78</p> <p>Race/ethnicity: White: Total: 82% G1: 80% G2: 82% G3: 84% Black: G1: 10% G2: 9% G3: 8% Asian: G1: 2% G2: 1% G3: 1%</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 18 • Urinary frequency ≥8 voids/day • Urgency ≥ 6 for 3 days or UUI ≥ 3 episodes/3 day diary • At least moderate bladder problems on a Likert scale <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Pregnant • Women of reproductive age not on adequate contraception • Lower urinary tract pathology (SUI, urolithiasis, interstitial cystitis, urothelial tumors, ≥ grade 3 pelvic organ prolapse, BOO, postvoid residual > 100 mL, polyuria >3 L/24 h, rec UTI) • Current antimuscarinic • Neurogenic cause for OAB • Arrhythmia • Unstable angina or corrected QT interval > 500 ms • Electrostimulation or bladder training in past 4 weeks or currently 	<p>UUI episodes/day, mean ± SD: G1: 3.7 ± 3.3 (n=205) G2: 3.9 ± 3.5 (n=228) G3: 3.9 ± 3.3 (n=218)</p> <p>Urgency episodes/day, mean ± SD: G1: 11.4 ± 3.8 G2: 12.5 ± 4.1 G3: 11.6 ± 3.7</p> <p>Incontinence, n (%): G1: 205 (77) G2: 228 (85) G3: 218 (82)</p> <p>Continent days/wk, mean ± SD: G1: 0.6 ± 1.3 G2: 0.7 ± 1.5 G3: 0.7 ± 1.4</p> <p>Voids/day, mean ± SD: G1: 12.2 ± 3.7 G2: 12.9 ± 3.9 G3: 12.0 ± 3.3</p> <p>Daytime voids/day, mean ± SD: G1: 10.2 ± 3.3 G2: 10.7 ± 3.4 G3: 10.1 ± 2.9</p> <p>Nocturia episodes/day, mean ± SD: G1: 2.0 ± 1.3 G2: 2.2 ± 1.6 G3: 1.9 ± 1.4</p> <p>Voided volume (mL), mean ± SD: G1: 159 ± 69 G2: 152 ± 60 G3: 156 ± 58</p>	<p>UUI episodes/day, LS mean change (SE): G1: -0.96 (0.17) G2: -1.65 (0.16)* G3: -2.28 (0.16)*</p> <p>UUI episodes/day, median % change: G1: -40.0 (n=205) G2: -67.4* (n=228) G3: -81.8* (n=218)</p> <p>Urgency episodes/day, LS mean change (SE): G1: -0.79 (0.20) G2: -1.91 (0.20) G3: -2.30 (0.20)</p> <p>Urgency episodes/day, median % change: G1: -3.3 G2: -16.3* G3: -18.4*</p> <p>Continent days/wk, LS mean change (SE): G1: 1.31 (0.20) G2: 2.33 (0.19)* G3: 2.80 (0.19)*</p> <p>Voids/day, LS mean change (SE): G1: -1.08 (0.18) G2: -1.61 (0.18)* G3: -2.09 (0.18)*</p> <p>Voids/day, mean % change: G1: -6.9 G2: -14.9* G3: -16.0*</p> <p>Daytime voids/day, LS mean change (SE): G1: -0.69 (0.16) G2: -1.04 (0.16) (P = 0.107) G3: -1.54 (0.16)*</p>	<p>Quality: Overall quality score: fair INTERNAL VALIDITY: poor Randomization: + Masking: + Pt selection criteria: + Loss to followup: ++ Drop-out rates: - Power calculation: - Statistical issues: + EXTERNAL VALIDITY: good Age: + Baseline OAB status: + Baseline characteristics: ++ Length of followup: + Measurement methods: + Measurement reliability: + Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating	
Nitti et al., 2007 (continued)			<p>Years since OAB diagnosis, mean \pm SD: G1: 9.8 \pm 10.3 G2: 9.1 \pm 10.3 G3: 10.1 \pm 11.5</p> <p>Previous tolterodine use, n (%): G1: 90 (33) G2: 88 (31) G3: 95 (34)</p> <p>Previous oxybutynin use, n (%): G1: 101 (37) G2: 94 (33) G3: 97 (35)</p> <p>Previous trospium chloride use, n (%): G1: 7 (2.6) G2: 6 (2.1) G3: 3 (1.1)</p>	<p>Daytime voids/median % change: G1: -5.9 G2: -11.1 (<i>P</i> = 0.008) G3: -15.6*</p> <p>Nocturia episodes/day, LS mean change (SE): G1: -0.39 (0.07) G2: -0.58 (0.17) (<i>P</i> = 0.042) G3: -0.55 (0.17) (<i>P</i> = 0.09)</p> <p>Nocturia episodes/day, median % change: G1: -25.0 G2: -33.3 (<i>P</i> = 0.013) G3: -25.0 (<i>P</i> = 0.267)</p> <p>Treatment response, n (% yes) G1: 266 (45) G2: 267 (64)* G3: 267 (74)*</p> <p>Voided volume (mL), LS mean change (SE): G1: 8.38 (4.06) G2: 16.5 (4.00) (<i>P</i> = 0.15) G3: 33.6 (4.04)*</p> <p>Side effects, n (%) G1: 149 (55) G2: 171 (61) G3: 193 (69)</p> <p>Dry mouth, n (%): G1: 19 (7) G2: 45 (16) G3: 99 (36)</p> <p>Constipation, n (%) G1: 7 (3) G2: 14 (5) G3: 21 (8)</p>		

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Nitti et al., 2007 (continued)				<p>UTI, n (%): G1: 11 (4) G2: 10 (4) G3: 15 (5)</p> <p>Headache, n (%): G1: 9 (3) G2: 12 (4) G3: 8 (3)</p> <p>Withdrew, %: Total: 19 G1: 41 G2: 58 G3: 56</p> <p>Withdrew due to AE, %: G1: 4 G2: 6 G3: 9 *<i>P</i> < 0.001</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Preik et al., 2004</p> <p>[See evidence table for Anderson et al. 1999]</p> <p>Country and setting: Germany & US, Multiple academic hospitals</p> <p>Enrollment period: NR</p> <p>Funding: ALZA Corporation</p> <p>Author industry relationship disclosures: 4 of 5 ALZA Corp (2) Ditropan XL (1) Janssen-Cilag GmbH (1) Johnson and Johnson (1) Ortho-McNeil (1)</p>	<p>Design: RCT</p> <p>Intervention: Oxybutynin</p> <p>Groups: daily dose increased in 5 mg increments each 4-7 days until reaching the minimum effective dose, maximum tolerated dose, or maximum allowable dose. Once participant reached titrated dose, maintained for 1 week confirmation period and 1 week maintenance period.</p> <p>G1: Controlled-release oxybutynin (OROS-O) G2: Immediate release oxybutynin (IR-O)</p> <p>N at enrollment: 105</p> <p>N at follow-up: Total: 93 G1: 46 G2: 47</p> <p>Women, n (%): G1: 50 (94.3) G2: 47 (90.4)</p> <p>Age, mean (SE): G1: 59.2 (10.6) G2: 59.6 (10.0)</p> <p>Race/ethnicity: NR</p> <p>Weight (kg), mean (SE): G1: 78.0 (18.3) G2: 80 (20.2)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> Community-dwelling men & women age 40-75 Urge or mixed UI ≥ 6 UUI episodes per week recorded on the run-in diary after washout of anticholinergic Previous response to anticholinergic Urge predominant SUI <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Known treatable GU disorders that could cause incontinence Men who had had prostate surgery < 9 mos before study enrollment or with a PSA level of >10 ng/mL PVR of > 100mL Estimated Cr clearance of < 50 mL/min Glaucoma or untreated anterior chamber angles Hgb < 100g/L Known hypersensitivity to oxybutynin History of drug or alcohol abuse Positive urine drug screen Pregnant or breastfeeding Myasthenia gravis 	<p>UUI episodes/wk, mean (SE): G1: 27.6 (24.0) G2: 23.4 (16.3)</p> <p>Duration of incontinence, n (%): 1-5 years: G1: 26 (49) G2: 22 (42) > 5 years: G1: 23 (43) G2: 22 (42)</p>	<p>Dose-titration endpoint, MED, n/N (%): G1: 29/46 (63) G2: 26/47 (55)</p> <p>Dry mouth, MED, n/N (%): G1: 3/29 (10) G2: 7/26 (27)</p> <p>Dose-titration endpoint, MTD, n/N (%): G1: 11/46 (24) G2: 19/47 (41)</p> <p>Dry mouth, MTD, n/N (%): G1: 7/11 (64) G2: 13/19 (68)</p> <p>Dose-titration endpoint, MAD, n/N (%): G1: 6/46 (13) G2: 2/47 (4)</p> <p>Dry mouth, MAD, n/N (%): G1: 1/6 (17) G2: 1/2 (50)</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: -</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: +</p> <p>Drop-out rates: -</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: -</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Rentzhog et al., 1993</p> <p>Country and setting: Sweden & UK, 17 academic hospitals</p> <p>Enrollment period: NR</p> <p>Funding: Pharmacia & Upjohn AB</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT, double blind, placebo controlled</p> <p>Intervention: tolterodine vs placebo</p> <p>Groups: G1: tolterodine 1mg/d G2: tolterodine 2 mg/d G3: tolterodine 4mg/d G4: tolterodine 8mg/d G5: placebo</p> <p>N at enrollment: G1: 21 G2: 16 G3: 14 G4: 16 G5: 13</p> <p>N at follow-up: G1: 17 G2: 15 G3: 11 G4: 11 G5: 10</p> <p>Women, n: G1: 14 G2: 10 G3: 12 G4: 15 G5: 10</p> <p>Age, mean: G1: 56 G2: 59 G3: 56 G4: 58 G5: 58</p> <p>Race/ethnicity: NR</p> <p>Body weight, kg, mean: G1: 73 G2: 76 G3: 72 G4: 76 G5: 73</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • age 18-75 • urinary urgency, 8+ micturitions/d, and/or UI (1+ episode/d) • UDS-confirmed detrusor instability (phasic increase in detrusor pressure in the presence of typical sx) • max urinary flow rate of >15 mL/s • sterile or clinically insignificant bacteriuria • normal routine laboratory tests <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • antimuscarinic agents or drugs w/ antimuscarinic side effects • other drugs for UI (except estrogens) • clinically significant cardiac, hepatic, renal, or heme disorders • contraindications to antimuscarinic agents • pregnant or lactating • women not using reliable contraception 	<p>Frequency of micturitions/d: G1: 10.3 G2: 11.2 G3: 12.8 G4: 10.2 G5: 10.2</p> <p>Incontinence episodes/d: G1: 2.1 G2: 1.7 G3: 1.8 G4: 2.7 G5: 4.1</p> <p>Previous drug therapy: G1: 9 G2: 10 G3: 9 G4: 9 G5: 4</p> <p>Frequency, n: G1: 7 G2: 7 G3: 4 G4: 5 G5: 4</p> <p>Leakage, n: G1: 2 G2: - G3: 2 G4: 1 G5: 3</p> <p>Leakage and frequency, n: G1: 11 G2: 9 G3: 7 G4: 9 G5: 6</p> <p>Volume at first contraction (mL), mean mL ± SD G1: 243 ± 220 G2: 274 ± 129 G3: 171 ± 109 G4: 297 ± 123 G5: 295 ± 16723 (</p>	<p>Volume at first contraction (mL), mean change (% change): G1: 74 (63) G2: 17 (17) G3: 89 (114) G4: 115 (119) G5: -23 (-4)</p> <p>Linear regression of change: $P = 0.0459$</p> <p>Residual urinary volume (mL), mean change (% change): G1: 6 (2) G2: 10 (57) G3: 30 (122) G4: 143 (225) G5: 4 (126)</p> <p>Linear regression of change: $P = 0.0003$</p> <p>Max cystometric capacity (mL), mean change (% change): G1: 15 (20) G2: -25 (5) G3: 2 (10) G4: 76 (30) G5: -17 (3)</p> <p>Linear regression of change: $P = 0.0921$</p> <p>Bladder compliance (mL/cmH2O), mean change (% change): G1: -29 (-9) G2: 10 (164) G3: 2 (72) G4: 18 (355) G5: -6 (11)</p> <p>Linear regression of change: $P = 0.1646$</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: -</p> <p>Methods and blinding: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: -</p> <p>Drop-out rates: -</p> <p>Power calculation: +</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Rentzhog et al., 1993 (continued)			<p>Residual urinary volume, mean mL ± SD: G1: 24 ± 31 G2: 51 ± 94 G3: 25 ± 18 G4: 52 ± 130 G5: 26 ± 35</p> <p>Max cystometric capacity, mean mL ± SD: G1: 381 ± 200 G2: 397 ± 197 G3: 284 ± 195 G4: 335 ± 145) G5: 377 ± 181)</p> <p>Bladder compliance, mean mL/cmH2O ± SD: G1: 65 ± 105 G2: 50 ± 44 G3: 36 ± 43 G4: 44 ± 46 G5: 44 ± 28</p> <p># detrusor contractions >10 cmH20, mean ± SD: G1: 4.1 ± 3.3 G2: 2.3 ± 1.8 G3: 3.4 ± 2.5 G4: 3.3 ± 2.7 G5: 3.1 ± 3.0</p>	<p>Detrusor contractions >10 cmH20, mean change (% change): G1: -1.6 (-34) G2: -1.2 (-38) G3: -2.0 (-33) G4: -0.3 (-32) G5: 0.0 (48)</p> <p>Linear regression of change: <i>P</i> = 0.7284</p> <p>Max urinary flow (mL/s), mean change (% change): G1: 1.2 (9) G2: 1.3 (18) G3: -2.4 (-3) G4: -0.3 (-32) G5: -2.0 (28)</p> <p>Linear regression of change: <i>P</i> = 0.3072</p> <p>Volume at normal desire to void (mL), mean change (% change): G1: -1 (2) G2: -18 (1) G3: 32 (25) G4: 66 (45) G5: -18 (15)</p> <p>Linear regression of change: <i>P</i> = 0.0492</p> <p>Patients reporting 1+ adverse event, n: G1: 8 G2: 6 G3: 7 G4: 12 G5: 6</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Rentzhog et al., 1993 (continued)			Max urinary flow, mean mL/s ± SD: G1: 21.0 ± 7.5 G2: 2.33 ± 9.9 G3: 19.2 ± 9.8 G4: 20.5 ± 8.2 G5: -2.0 ± 28	Total AE reported: G1: 10 G2: 14 G3: 10 G4: 25 G5: 10	
			Volume at normal desire to void, mean mL ± SD: G1: 235 ± 128 G2: 273 ± 121 G3: 143 ± 83 G4: 211 ± 128 G5: 255 ± 157	Dry mouth G1: 2 G2: 2 G3: 5 G4: 9 G5: 2	
			Concomitant dz, n: G1: 11 G2: 11 G3: 7 G4: 11 G5: 9	Constipation G1: 1 G2: 3 G3: 1 G4: 2 G5: 0 Abnormal vision G1: 0 G2: 3 G3: 1 G4: 1 G5: 1	
				Urinary retention G1: 0 G2: 0 G3: 0 G4: 1 G5: 0	
				Other G1: 7 G2: 6 G3: 3 G4: 12 G5: 7	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Robinson et al., 2007</p> <p>Country and setting: 12 countries in Europe, 39 study sites</p> <p>Enrollment period: July 2002 to December 2002</p> <p>Funding: Astellas</p> <p>Author industry relationship disclosures: 2 of 4 Astellas (2)</p>	<p>Design: RCT</p> <p>Intervention: tamsulosin OCAS (0.25, 0.5, 1.0, 1.5 mg) qd vs tolterodine ER qd vs placebo qd</p> <p>Groups: G1: 0.25 mg tamsulosin OCAS qd G2: 0.5 mg tamsulosin OCAS qd G3: 1.0 mg tamsulosin OCAS qd G4: 1.5 mg tamsulosin OCAS qd G5: tolterodine ER qd G6: placebo qd</p> <p>N at enrollment: G1: 58 G2: 65 G2: 63 G4: 60 G5: 61 G6: 61</p> <p>N at follow-up: G1: 52 G2: 61 G3: 55 G4: 46 G5: 53 G6: 59</p> <p>Women, %: 100</p> <p>Age, range: 18-70</p> <p>Race/ethnicity: NR</p>	<p>Inclusion criteria: • Symptoms of OAB ≥ 3 months</p> <p>Exclusion criteria: • SUI or MUI where stress symptoms predominant and neurogenic</p>	<p>Urgency episodes/day, mean: G1: 5.83 G2: 5.58 G3: 5.82 G4: 6.28 G5: 6.78 G6: 5.58</p> <p>Incontinence episodes/day, mean: G1: 3.22 G2: 2.17 G3: 3.33 G4: 2.98 G5: 2.33 G6: 2.90</p> <p>Voids/day, mean: G1: 12.20 G2: 11.09 G3: 12.96 G4: 12.46 G5: 13.22 G6: 11.90</p> <p>Nocturia episodes/day, mean: G1: 1.97 G2: 1.76 G3: 1.82 G4: 2.10 G5: 2.34 G6: 1.87</p> <p>Voided volume (mL), mean: G1: 135.5 G2: 148.6 G3: 135.1 G4: 131.4 G5: 148.9 G6: 145.5</p>	<p>Urgency episodes/day, mean change: G1: -1.51 G2: -1.66 G3: -1.83 G4: -1.91 G5: -2.49 G6: -1.86 G4/G6: <i>P</i> = 0.901 G5/G6: <i>P</i> = 0.632</p> <p>Incontinence episodes/day, mean change: G1: -1.40 G2: -0.93 G3: 0.02 G4: -0.66 G5: -1.66 G6: -0.66 G4/G6: <i>P</i> = 0.945 G5/G6: <i>P</i> = 0.025</p> <p>Voids/day, mean change: G1: -1.60 G2: -1.01 G3: -1.38 G4: -1.18 G5: -2.59 G6: -1.81 G4/G6: <i>P</i> = 0.189 G5/G6: <i>P</i> = 0.353</p> <p>Nocturia episodes/day, mean change: G1: -0.35 G2: -0.36 G3: -0.33 G4: -0.34 G5: -0.74 G6: -0.41 G4/G6: <i>P</i> = 0.495 G5/G6: <i>P</i> = 0.353</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: -</p> <p>Drop-out rates: NR</p> <p>Power calculation: +</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: -, NR</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: -</p> <p>Length of followup: -</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Robinson et al., 2007 (continued)				Void volume (mL), mean change: G1: 13.9 G2: 9.7 G3: 12.6 G4: 12.9 G5: 24.3 G6: 11.4 G4/G6: $P = 0.992$ G5/G6: $P = 0.092$	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Rogers et al., 2008</p> <p>Country and setting: USA, multicenter (54 outpatient sites)</p> <p>Enrollment period: NR</p> <p>Funding: Pfizer</p> <p>Author industry relationship disclosures: 7 of 7 Bayer (1) Boeringer Ingelheim(1) Duramed (1) J&J (1) P&G (1) Novartis (1) Novo-Nordisk (1) Pfizer (7) Wyeth (1)</p>	<p>Design: Multicenter randomized placebo-controlled double-blind</p> <p>Intervention: Tolterodine 4 mg ER vs placebo within 4 hours of bedtime</p> <p>Groups: G1: Tolterodine 4mg ER daily G2: Placebo</p> <p>N at enrollment: G1: 202 G2: 211</p> <p>N at follow-up: G1: 163 G2: 167</p> <p>Women, %: Total: 100</p> <p>Age, yrs ± SD: G1: 49 ± 12 G2: 47 ± 12</p> <p>Race/ethnicity, n (%): White: G1: 139 (69) G2: 138 (66) Black: G1: 37 (18) G2: 45 (21) Asian: G1: 3 (2) G2: 4 (2) Other: G1: 22 (11) G2: 23 (11)</p> <p>Parous, %: G1: 86 G2: 91</p> <p>Follow-up: 12 weeks</p> <p>Premenopausal, n (%): G1: 71 (35) G2: 95 (45)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 18 • Women • ≥ 0.6 UUI episodes/day • ≥ 8 voids/day • ≥ 3 OAB voids/day (urgency-associated voids) • “some moderate problems” on PPBC • OAB symptoms ≥ 3 months • Stable, sexually active relationship with man ≥ 6 months <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • POP Stage 3 or greater • History of lower urinary tract surgery • Lifelong sexual dysfunction unrelated to lifelong UUI • Predominate SUI 	<p>UUI episodes/day, mean ± SD: G1: 2.5 ± 2.1 G2: 2.2 ± 1.8</p> <p>Voids/ day, mean ± SD: G1: 13.0 ± 4.1 G2: 12.5 ± 3.9</p> <p>OAB voids/ day, mean ± SD: G1: 10.6 ± 4.4 G2: 9.9 ± 4.2</p> <p>Pads/ day, mean ± SD: G1: 2.9 ± 2.7 G2: 2.9 ± 3.1</p> <p>SQOL-F total score, mean± SD: G1: 69.6 ± 23.1 G2: 69.2 ± 23.0</p> <p>PISQ total score, mean ± SD: G1: 88.7 ± 13.9 G2: 88.9 ± 14.2</p> <p>PISQ Behavioral/emotive score, mean ± SD: G1: 38.0 ± 8.9 G2: 38.0 ± 9.0</p> <p>PISQ Physical score, mean ± SD: G1: 31.7 ± 5.9 G2: 31.8 ± 6.0</p> <p>PISQ partner related score, mean ± SD: G1: 19.0 ± 2.8 G2: 19.0 ± 2.5</p> <p>HAD Anxiety score, mean ± SD: G1: 8.3 ± 4.0 G2: 8.2 ± 4.0</p> <p>HAD Depression score, mean ± SD: G1: 5.1 ± 3.3 G2: 5.4 ± 3.4</p>	<p>UUI episodes/day, mean change (SE): G1: 1.8 (0.1) G2: 1.4 (0.1) <i>P</i> = 0.0029 (<i>P</i> = 0.0525 when a single statistical outlier is excluded)</p> <p>UUI episodes/day, median % change: G1: -100.0 G2: -82.5 <i>P</i> = 0.0003</p> <p>No UUI events, week 12, %: G1: 57 G2: 42 <i>P</i> < 0.004</p> <p>OAB voids/day, mean change (SE): G1: -4.7 (0.3) G2: -3.0 (0.3) <i>P</i> < 0.0001</p> <p>Pads/day, mean change (SE): G1: -1.5 (0.1) G2: -1.0 (0.1) <i>P</i> = 0.0024</p> <p>Voids/day, mean change (SE): G1: -3.3 (0.2) G2: -2.3 (0.2) <i>P</i> = 0.0006</p> <p>SQOL-F score, mean change (SE): G1: 6.4 (1.2) G2: 1.5 (1.2) <i>P</i> = 0.004</p> <p>PISQ total score, mean change (SE): G1: 4.9 (0.7) G2: 2.3 (0.7) <i>P</i> = 0.009</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: fair</p> <p>Randomization: +</p> <p>Method and blinding: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: -</p> <p>Drop-out rates: +</p> <p>Power calculation: +</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Rogers et al., 2008 (continued)	<p>Perimenopausal, n (%): G1: 18 (9) G2: 16 (8)</p> <p>Postmenopausal, n (%): G1: 112 (56) G2: 99 (47)</p>			<p>PISQ behavioral/emotive score, mean change (SE): G1: 1.6 (0.4) G2: 0.5 (0.4) <i>P</i> = NS</p> <p>PISQ physical score, mean change (SE): G1: 2.6 (0.3) G2: 1.6 (0.3) <i>P</i> ≤ 0.01</p> <p>PISQ partner related score, mean change (SE): G1: 0.6 (0.1) G2: 0.3 (0.1) <i>P</i> = NS</p> <p>HAD anxiety score, mean change (SE): G1: -1.9 (0.3) G2: -1.1 (0.3) <i>P</i> = 0.03</p> <p>HAD depression score, mean change (SE): G1: -1.2 (0.2) G2: -0.8 (0.2) <i>P</i> = NS</p> <p>Subjects with adverse events, n (%): G1: 114 (57) G2: 111 (53)</p> <p>Withdrawn due to AEs, n (%): G1: 9 (5) G2: 6 (3)</p> <p>Dry mouth, n (%): G1: 26 (13) G2: 19 (9)</p> <p>Constipation, n (%): G1: 7 (4) G2: 8 (4)</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Rogers et al., 2008 (continued)				Nasopharyngitis, n (%): G1: 9 (5) G2: 10 (5) Sinusitis, n (%): G1: 8 (4) G2: 9 (4) URI, n (%): G1: 9 (4) G2: 9 (5) UTI, n (%): G1: 12 (6) G2: 5 (2) Headache, n (%): G1: 7 (4) G2: 6 (3) Depression, n (%): G1: 5 (3) G2: 0 (0) Insomnia, n (%): G1: 5 (3) G2: 0 (0)	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Rudy et al., 2006</p> <p>Rudy et al., 2006*</p> <p>Staskin and Harnett, 2004†</p> <p>Country and setting: US, 52 sites</p> <p>Enrollment period: NR</p> <p>Funding: Indevus NR†</p> <p>Author industry relationship disclosures: 5 of 5 Indevus (5) NR†</p>	<p>Design: RCT</p> <p>Intervention: Trospium chloride vs placebo x 12 wks</p> <p>Groups: G1: trospium chloride 20 mg b.i.d. G2: matching placebo</p> <p>N at enrollment: G1: 329 G2: 329</p> <p>N at follow-up: G1: 323 G2: 325</p> <p>Age, mean (SE): G1: 61.1 (0.69) G2: 61.0 (0.70)</p> <p>Women, N (%): G1: 269 (81.8) G2: 267 (81.2)</p> <p>Race/ethnicity, n (%): Black: G1: 26 (7.9) G2: 21 (6.4) White: G1: 284 (86.3) G2: 300 (91.2) Hispanic: G1: 13 (4.0) G2: 5 (1.5) Asian: G1: 5 (1.5) G2: 3 (0.9)</p> <p>Parity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥18 • ≥ 6 mos of OAB sx • ≥10 voids/day • Sx of urgency • ≥ 7 UUI episodes/wk <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Predominantly SUI, insensate UI, or overflow UI • Neurogenic bladder d/o's • Significant renal dz • Uninvestigated hematuria • UTI at washout or more than 2x during the prior 12 mos • PVR >100mL • Use of any anticholinergic drug or other drug therapy for OAB w/in 21 days before randomization • Bladder surgery w/in 6 mos before randomization • Cancer • Interstitial cystitis • PSA >10ng/mL • Diuretic use • Estrogen therapy • Nonmedical bladder therapy not part of a stable, long-term program (ie, timed voids and straight caths; Kegels were allowed) • Pregnancy 	<p>UUI episodes/day, median: G1: 2.86 G2: 2.86 <i>P</i> = 0.9849</p> <p>Urgency severity score associated with toilet voids, median: G1: 1.79 G2: 1.75 <i>P</i> = 0.4100</p> <p>Voids/day, mean: G1: 12.94 G2: 13.17 <i>P</i> = 0.3169</p> <p>Nocturia episodes/day, median: G1: 2.00 G2: 2.00 <i>P</i> = 0.9048</p> <p>Nocturnal urgency severity score associated with toilet voids, mean: G1: 2.03 G2: 2.01 <i>P</i> = 0.6863</p> <p>OAB-SCS, median: G1: 36.56 G2: 36.88 <i>P</i> = 0.7176</p> <p>Stanford Sleepiness Scale, mean: G1: 1.98 G2: 1.94</p> <p>Stanford Sleepiness Scale, age group < 65 years: G1: 1.99 G2: 2.09</p>	<p>UUI episodes/day, day 1, median change: G1: -1.00 G2: -0.57 <i>P</i> = 0.042</p> <p>UUI episodes/day, day 2, median change: G1: -1.29 G2: -0.86 <i>P</i> = 0.0022</p> <p>UUI episodes/day, day 3, median change: G1: -1.57 G2: -0.86 <i>P</i> < 0.0001</p> <p>UUI episodes/day, day 4, median change: G1: -1.57 G2: -1.00 <i>P</i> < 0.0001</p> <p>UUI episodes/day, day 5, median change: G1: -1.57 G2: -1.00 <i>P</i> < 0.0001</p> <p>UUI episodes/day, day 6, median change: G1: -1.71 G2: -1.00 <i>P</i> < 0.0001</p> <p>UUI episodes/day, day 7, median change: G1: -1.71 G2: -1.00 <i>P</i> < 0.0001</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: NA Masking: NA Pt selection criteria: + Loss to followup: - Drop-out rates: NR Power calculation: - Statistical issues: - EXTERNAL VALIDITY: fair Age: + Baseline OAB status: + Baseline characteristics: + Length of followup: + Measurement methods: + Measurement reliability: - Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Rudy et al., 2006 Rudy et al., 2006 Staskin and Harnett, 2004 (continued)		<ul style="list-style-type: none"> Contraindication to antimuscarinic therapy 	Stanford Sleepiness Scale, age group > 65 years: G1: 1.96 G2: 1.71 Stanford Sleepiness Scale, age group < 75 years: G1: 2 G2: 1.97 Stanford Sleepiness Scale, age group >75 years: G1: 1.86 G2: 1.79 Voided volume (mL), mean: G1: 154.80 ± NR G2: 154.64 ± NR <i>P</i> = 0.9667 Prior OAB medications, n (%): G1: 162 (49.2) G2: 169 (51.4) Prior hx of pelvic-floor training, n (%): G1: 62 (18.8) G2: 76 (23.1) Currently practice undergarment change d/t incontinence, n (%): G1: 227 (69.0) G2: 239 (72.6)	UUI episodes/day, week 1, median change:* G1: -1.43 G2: -0.86 <i>P</i> < 0.0001 UUI episodes/day, week 4, median change:* G1: -1.71 G2: -1.14 <i>P</i> < 0.0001 UUI episodes/day, week 12, median change:* G1: -1.86 G2: -1.29 <i>P</i> < 0.0001 Urgency severity score, day 1, mean change: G1: 0.00 G2: 0.02 <i>P</i> = 0.47 Urgency severity score, day 2, mean change: G1: -0.08 G2: -0.03 <i>P</i> = 0.13 Urgency severity score, day 3, mean change: G1: -0.11 G2: -0.04 <i>P</i> = 0.015 Urgency severity score, day 4, mean change: G1: -0.12 G2: -0.05 <i>P</i> = 0.031 Urgency severity score, day 5, mean change: G1: -0.13 G2: -0.05 <i>P</i> = 0.021	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Rudy et al., 2006 Rudy et al., 2006 Staskin and Harnett, 2004 (continued)				<p>Urgency severity score, day 6, mean change: G1: -0.08 G2: 0.04 <i>P</i> < 0.0001</p> <p>Urgency severity score, day 7, mean change: G1: -0.09 G2: 0.03 <i>P</i> < 0.0001</p> <p>Urgency severity score, week 1, mean change:* G1: -0.09 G2: -0.01 <i>P</i> = 0.0023</p> <p>Urgency severity score, week 4, mean change:* G1: -0.19 G2: -0.04 <i>P</i> < 0.0001</p> <p>Urgency severity score, week 12, mean change:* G1: -0.21 G2: -0.02 <i>P</i> < 0.0001</p> <p>Voids/day, day 1, mean change: G1: -0.66 G2: -0.35 <i>P</i> = 0.14</p> <p>Voids/day, day 2, mean change: G1: -1.11 G2: -0.75 <i>P</i> = 0.09</p> <p>Voids/day, day 3, mean change: G1: -1.30 G2: -0.77 <i>P</i> = 0.012</p> <p>Voids/day, day 4, mean change: G1: -1.33 G2: -0.95 <i>P</i> = 0.086</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Rudy et al., 2006 Rudy et al., 2006 Staskin and Harnett, 2004 (continued)				<p>Voids/day, day 5, mean change: G1: -1.73 G2: -1.12 <i>P</i> = 0.0037</p> <p>Voids/day, day 6, mean change: G1: -1.80 G2: -1.28 <i>P</i> = 0.017</p> <p>Voids/day, day 7, mean change: G1: -1.99 G2: -1.44 <i>P</i> = 0.011</p> <p>Voids/day, week 1, mean change:* G1: -1.42 G2: -0.96 <i>P</i> = 0.0039</p> <p>Voids/day, week 4, mean change:* G1: -2.34 G2: -1.55 <i>P</i> < 0.0001</p> <p>Voids/day, week 12, mean change:* G1: -2.67 G2: -1.76 <i>P</i> < 0.0001</p> <p>Nocturia episodes/day, week 1, median change:* G1: -0.29 G2: -0.29 <i>P</i> = 0.8454</p> <p>Nocturia episodes/day, week 4, median change:* G1: -0.43 G2: -0.29 <i>P</i> = 0.0299</p> <p>Nocturia episodes/day, week 12, median change:* G1: -0.57 G2: -0.29 <i>P</i> = 0.0026</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Rudy et al., 2006 Rudy et al., 2006 Staskin and Harnett, 2004 (continued)				<p>Nocturnal urgency severity score, week 1, mean change:* G1: -0.05 G2: 0.03 <i>P</i> = 0.0442</p> <p>Nocturnal urgency severity score, week 4, mean change:* G1: -0.13 G2: =0.01 <i>P</i> = 0.0062</p> <p>Nocturnal urgency severity score, week 12, mean change:* G1: -0.17 G2: 0.01 <i>P</i> = 0.0005</p> <p>OAB-SCS, day 1, median change: G1: -2.50 G2: -0.14 <i>P</i> = 0.014</p> <p>OAB-SCS, day 2 median change: G1: -3.86 G2: -2.14 <i>P</i> = 0.015</p> <p>OAB-SCS, day 3, median change: G1: -4.71 G2: -2.71 <i>P</i> = 0.0019</p> <p>OAB-SCS, day 4, median change: G1: -4.86 G2: -3.36 <i>P</i> = 0.011</p> <p>OAB-SCS, day 5, median change: G1: -6.43 G2: -3.36 <i>P</i> < 0.0001</p> <p>OAB-SCS, day 6, median change: G1: -5.43 G2: -2.93 <i>P</i> < 0.0001</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Rudy et al., 2006 Rudy et al., 2006 Staskin and Harnett, 2004 (continued)				<p>OAB-SCS, day 7, median change: G1: -6.29 G2: -3.00 <i>P</i> < 0.0001</p> <p>OAB-SCS, week 1, median change:* G1: -4.71 G2: -2.29 <i>P</i> < 0.0001</p> <p>OAB-SCS, week 4, median change:* G1: -8.14 G2: -3.86 <i>P</i> < 0.0001</p> <p>OAB-SCS, week 12, median change:* G1: -8.43 G2: -4.62 <i>P</i> < 0.0001</p> <p>Stanford Sleepiness Scale, week 1, mean change:† G1: -0.2 G2: -0.12</p> <p>Stanford Sleepiness Scale, week 4, mean change:† G1: -0.17 G2: -0.14</p> <p>Stanford Sleepiness Scale, week 12, mean change:† G1: -0.16 G2: -0.11</p> <p>Stanford Sleepiness Scale, including T-max time point values only (G1=93, G2=97), week 1, mean change:† G1: -0.42 G2: -0.2</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Rudy et al., 2006 Rudy et al., 2006 Staskin and Harnett, 2004 (continued)				<p>Stanford Sleepiness Scale, including T-max time point values only (G1=144, G2=148), week 4, mean change:† G1: -0.2 G2: -0.19</p>	
				<p>Stanford Sleepiness Scale, including T-max time point values only (G1=182, G2=179), week 12, mean change:† G1: -0.17 G2: -0.18</p>	
				<p>Mean Stanford Sleepiness Scale, age group < 65 years, week 1, mean change:† G1: -0.27 G2: -0.19</p>	
				<p>Mean Stanford Sleepiness Scale, age group < 65 years, week 4, mean change:† G1: -0.14 G2: -0.27</p>	
				<p>Mean Stanford Sleepiness Scale, age group < 65 years, week 12, mean change:† G1: -0.14 G2: -0.22</p>	
				<p>Mean Stanford Sleepiness Scale, age group >65 years, week 1, mean change:† G1: -0.04 G2: -0.03</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Rudy et al., 2006 Rudy et al., 2006 Staskin and Harnett, 2004 (continued)				<p>Mean Stanford Sleepiness Scale, age group >65 years, week 4, mean change:† G1: -0.23 G2: 0.03</p>	
				<p>Mean Stanford Sleepiness Scale, age group >65 years, week 12, mean change:† G1: -0.22 G2: 0.03</p>	
				<p>Mean Stanford Sleepiness Scale, age group < 75 years, change from baseline to week 1:† G1: -0.21 G2: -0.14</p>	
				<p>Mean Stanford Sleepiness Scale, age group < 75 years, change from baseline to week 4:† G1: 0.14 G2: -0.16</p>	
				<p>Mean Stanford Sleepiness Scale, age group < 75 years, week 12, mean change:† G1: 0.15 G2: -0.14</p>	
				<p>Mean Stanford Sleepiness Scale, age group > 75 years, change from baseline to week 1:† G1: -0.02 G2: 0</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Rudy et al., 2006 Rudy et al., 2006 Staskin and Harnett, 2004 (continued)				<p>Mean Stanford Sleepiness Scale, age group > 75 years, change from baseline to week 4:† G1: -0.49 G2: -0.09</p> <p>Mean Stanford Sleepiness Scale, age group > 75 years, week 12, mean change:† G1: -0.33 G2: 0.02</p> <p>Clinically significant increase (≥ 3 points) from baseline to week 12 in SSS score, n (%):† G1: 5 (1.5) G2: 8 (2.5)</p> <p>Voided volume (mL), week 1, mean change:* G1: 29.23 G2: 6.05 <i>P</i> < 0.0001</p> <p>Voided volume (mL), week 4, mean change:* G1: 39.50 G2: 9.45 <i>P</i> < 0.0001</p> <p>Voided volume (mL), week 12, mean change:* G1: 35.59 G2: 9.44 <i>P</i> < 0.0001</p> <p>Adverse events Any, n (%) G1: 196 (59.6) G2: 153 (46.5)</p> <p>Dry mouth: G1: 65 (19.8) G2: 17 (5.2)</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Rudy et al., 2006 Rudy et al., 2006 Staskin and Harnett, 2004 (continued)				<p>Constipation: G1: 36 (10.9) G2: 19 (5.8)</p> <p>Headache not otherwise specified: G1: 18 (5.5) G2: 15 (4.6)</p> <p>UTI not otherwise specified: G1: 16 (4.9) G2: 8 (2.4)</p> <p>Nasopharyngitis: G1: 13 (4.0) G2: 12 (3.6)</p> <p>Cough: G1: 8 (2.4) G2: 1 (0.3)</p> <p>Diarrhea: G1: 7 (2.1) G2: 13 (4.0)</p> <p>AEs leading to treatment discontinuation, %: G1: 7.3 G2: 4.6</p> <p>Most common AEs leading to discontinuation:</p> <p>Constipation: G1: 1.8% G2: 0.6%</p> <p>Dry mouth: G1: 1.5% G2: 0.0%</p> <p>At least one CNS event: G1: 5.8% G2: 5.2%</p> <p>Somnolence G1: 0.3% G2: 0.6%</p> <p>Sedation: G1: 0 G2: 0.3%</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Salvatore et al., 2005</p> <p>Country and setting: UK, Urogynecology specialty</p> <p>Enrollment period: 1 year with phone questionnaire at 2 years</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Prospective randomized phone questionnaire</p> <p>Intervention: Oxybutynin with incremental increase in dosage to 5mg t.i.d. over 6 weeks at 14 day intervals</p> <p>Groups: G1: Oxybutynin 2.5 mg b.i.d. G2: Oxybutynin 5 mg at night</p> <p>N at enrollment: G1: 42 G2: 44</p> <p>N at follow-up: G1: 27 G2: 39</p> <p>Women, %: 100</p> <p>Age, yrs ± SD: G1: 59.2 ± 12.6 G2: 53.3 ± 14.4</p> <p>Race/ethnicity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Women • Video-urodynamic diagnosis of DO or low bladder capacity • DO- ≥ 1 contractions on filling cystometry associated with urgency • Low bladder compliance = linear detrusor pressure rise during filling cystometry > 15 cmH2O at 500mL associated with urgency or leakage <p>Exclusion criteria: NR</p>	<p>Stopped treatment, n (%) G1: 16 (59.3) G2: 28 (71.8)</p>	<p>SUI episodes/day, mean ± SD: G1: 0.85 ± 0.78 G2: 1.27 ± 1.17 <i>P</i> = 0.04</p> <p>Nocturia episodes/ day, mean ± SD: G1: 1.21 ± 1.17 G2: 2.04 ± 1.45 <i>P</i> = 0.01</p> <p>Oxybutynin compliance, 2 yrs, n (%) : G1: 11 (40.7) G2: 11 (28.2)</p> <p>Time on drug, n: 1 month: G1: 6 G2: 12 1-2 months: G1: 6 G2: 11 1-2 months: G1: 2 G2: 3 3-4 months: G1: 2 G2: 2</p> <p>Duration of treatment: G1: NR G2: NR <i>P</i> = 0.08</p> <p>General perception of improvement: <i>P</i> = 0.58</p> <p>Max dose of oxybutynin reached: <i>P</i> = 0.31</p> <p>Present dose of oxybutynin: <i>P</i> = 0.45</p> <p>Present overall severity of symptom: <i>P</i> = 0.12</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: -</p> <p>Masking: NA</p> <p>Pt selection criteria: -</p> <p>Loss to followup: -</p> <p>Drop-out rates: NR</p> <p>Power calculation: +</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: +</p> <p>Baseline OAB status: NR</p> <p>Baseline characteristics: -</p> <p>Length of followup: ++</p> <p>Measurement methods: +</p> <p>Measurement reliability: -</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Salvatore et al., 2005 (continued)				<p>Daytime frequency: <i>P</i> = 0.78</p> <p>Urgency: <i>P</i> = 0.78</p> <p>Urge incontinence: <i>P</i> = 0.18</p> <p>Pad usage: <i>P</i> = 0.24</p> <p>Side effects, n: G1: 3 G2: 4</p> <p>Dry Mouth, n: G1: 1 G2: 4</p> <p>Dry Throat, n: G1: 2 G2: 0</p> <p>Dry nose, n: G1: 0 G2: 1</p> <p>Dry eyes, n: G1: 0 G2: 1</p> <p>Nausea, n: G1: 0 G2: 4</p> <p>Tachycardia, n: G1: 0 G2: 1</p> <p>Dizziness, n: G1: 0 G2: 2</p> <p>Constipation, n: G1: 0 G2: 0</p> <p>Headache. N: G1: 0 G2: 1</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Salvatore et al., 2007</p> <p>Country and setting: Italy, Urogynecology outpatient clinic</p> <p>Enrollment period: January 2004 to October 2005</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Cohort</p> <p>Intervention: Tolterodine SR 4 mg qd</p> <p>Groups: G1: vaginal profile stage 0a or Ia G2: anterior prolapse ≥ stage IIa</p> <p>N at enrollment: G1: 184 G2: 51</p> <p>N at follow-up: NR</p> <p>Women, %: 100</p> <p>Age, median (range): G1: 59 (20, 85) G2: 59 (35, 82)</p> <p>Race/ethnicity: NR</p> <p>Parity, mean (range): G1: 2 (0-6) G2: 2 (0-6)</p> <p>Menopausal, n (%): G1: 132 (72) G2: 7 (18)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Women • Urodynamically proven OAB who had no prolapse or pure anterior vaginal prolapse • Proven detrusor overactivity <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • All symptoms or signs related to voiding difficulties • Prolapse 	<p>Previous surgery, n (%): G1: 52 (28) G2: 11 (22)</p> <p>HRT, n (%): G1: 35 (26) G2: 7 (18)</p>	<p>Improvement in condition or cured, n (%): G1: 158 (85.9) G2: 31 (60.8)* <i>P</i> = 0.0002</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: NA</p> <p>Masking: NA</p> <p>Pt selection criteria: +</p> <p>Loss to followup: NR</p> <p>Drop-out rates: NR</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: poor</p> <p>Age: +</p> <p>Baseline OAB status: NR</p> <p>Baseline characteristics: -</p> <p>Length of followup: +</p> <p>Measurement methods: -</p> <p>Measurement reliability: -</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Salvatore et al., 2008</p> <p>Country and setting: Italy, academic health center</p> <p>Enrollment period: January 2005 to August 2005</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: None</p>	<p>Design: prospective cohort</p> <p>Intervention: tolterodine ER 4 mg qd for 12 wks</p> <p>Groups: G1: Involuntary detrusor contractions during filling G2: Involuntary detrusor contractions after p</p> <p>N at enrollment: 111 N at follow-up: 109 G1: 84 G2: 25</p> <p>Age, mean yrs: G1: 54.5 G2: 52</p> <p>BMI G1: 25.5 (19-36) G2: 25 (18-38)</p>	<p>Inclusion criteria: • UDS+ detrusor overactivity</p> <p>Exclusion criteria: • UTI • DM • neurological dz • genital prolapse >/= POP-Q stage II • previous anti-incontinence or prolapse surgery • previous trt w/ antimuscarinics</p> <p>Indications, N:</p>	<p>Menopause, n(%) G1: 53 (63) G2: 16 (64)</p> <p>HRT, n(%) G1: 13 (24.5) G2: 6 (37.5)</p> <p>Reported in relation to urodynamic findings</p> <p>1. women who experienced involuntary detrusor contractions during the filling phase spontaneously N=84</p> <p>2. women who experienced involuntary detrusor contractions after provocative maneuvers such as listening to running water or washing hands in cold water N=25</p>	<p>Non-response to trt, proportion (%) G1: 12/84 (14) G2: 12/25 (48) G1<G2 (P = 0.0008)</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: -</p> <p>Masking: -</p> <p>Pt selection criteria: -</p> <p>Loss to followup: ++</p> <p>Drop-out rates: ++</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: -</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Sand et al., 2004</p> <p>Country and setting: US, Specialty clinic</p> <p>Enrollment period: Subanalysis of OBJECT trial, women only</p> <p>Funding: ALZA Corp</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: 10mg ER Oxybutynin daily vs 4mg Tolterodine (2mg b.i.d.)</p> <p>Groups: G1: 10mg ER Oxybutynin daily x 12 wks G2: 4mg Tolterodine (2mg b.i.d.) x 12 wks</p> <p>Stratified by age: a: Age ≤ 64 b: Age 65-74 c: Age ≥ 75</p> <p>N at enrollment: G1: 152 G2: 163</p> <p>N at follow-up: G1: 132 G2: 146</p> <p>Women, %: 100</p> <p>Age, yrs ± SD: G1: 58.4 G2: 58.8</p> <p>Race/ethnicity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • ≥ 7 and ≤ 50 UUI episodes/week • ≥ 10 voids/day • MUI included if predominant UUI <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • UTI • IC • Urethral diverticulum • Bladder tumor • Bladder stone • Delivery within 6 mos • Pelvic, bladder, vaginal surgery in ≤ 6 mos • PVR ≥ 150 mL • Cardiovascular, renal, pulmonary, gastrointestinal, endocrine, neurologic, autoimmune, hematological, urological, psychiatric, or hepatic disease • Hematuria • Positive urine culture • Narrow angle glaucoma • Obstructive uropathy • Myasthenia gravis • POP to hymenal ring • Gastrointestinal obstruction • Decreased GI motility • GI narrowing • GI retention 	<p>UUI episodes/week, mean: G1: 25.2 G2: 25.1</p> <p>Incontinence episodes/week, mean: G1: 28.1 G2: 28.9</p> <p>Voids/week, mean: G1: 91.7 G2: 91.6</p> <p>Naïve to anticholinergics, %: G1: 60.5 G2: 60.7</p>	<p>UUI episodes/week, %: G1: 6.2 G2: 8.5 G1a: 5.0 G2a: 8.4 G1b: 5.5 G2b: 7.5 G1c: 8.5 G2c: 11.1 G1/G2: <i>P</i> = 0.038 G1a/G2a: <i>P</i> = 0.005 G1b/G2b: <i>P</i> = 0.337 G1c/G2c: <i>P</i> = 0.568</p> <p>Incontinence episodes/week, mean: G1: 7.3 G2: 10.1 G1a: 5.8 G2a: 10.0 G1b: 6.1 G2b: 9.2 G1c: 10.5 G2c: 12.5 G1/G2: <i>P</i> = 0.030 G1a/G2a: <i>P</i> = 0.005 G1b/G2b: <i>P</i> = 0.164 G1c/G2c: <i>P</i> = 0.714</p> <p>Voids/week, mean: G1: 68.0 G2: 71.2 G1a: 63.7 G2a: 71.2 G1b: 73.8 G2b: 71.9 G1c: 66.8 G2c: 65.5 G1/G2: <i>P</i> = 0.272 G1a/G2a: <i>P</i> = 0.024 G1b/G2b: <i>P</i> = 0.706 G1c/G2c: <i>P</i> = 0.838</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: fair</p> <p>Randomization: +</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: +</p> <p>Drop-out rates: +</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Sand et al., 2004 (continued)		<ul style="list-style-type: none"> • Investigational drugs within 1 month of screening • Hypersensitivity to drugs • Current drug/ EtOH abuse • Pregnant • Breastfeeding • Inability to follow protocol 		<p>Dry mouth, %: G1: 28.3 G2: 33.7</p> <p>Constipation, %: G1: 8.6 G2: 6.7</p> <p>Retention, %: G1: 4.0 G2: 1.2</p> <p>Blurred vision, %: G1: 2.6 G2: 0.6</p> <p>Dizziness, %: G1: 3.9 G2: 4.3</p> <p>Insomnia, %: G1: 0.7 G2: 1.8</p> <p>Somnolence, %: G1: 3.3 G2: 1.8</p> <p>Nervousness, %: G1: 0 G2: 1.2</p> <p>Headache, %: G1: 9.2 G2: 10.4</p> <p>Dyspepsia, %: G1: 5.3 G2: 6.1</p> <p>Nausea, %: G1: 3.3 G2: 1.8</p> <p>Vomiting, %: G1: 2.0 G2: 1.8</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Sand et al., 2006</p> <p>[See evidence table for Sand et al., 2007]</p> <p>Country and setting: US, 327 centers</p> <p>Enrollment period: NR</p> <p>Funding: Watson Laboratories, Inc</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: Oxybutynin transdermal system 3.9 mg/day, twice weekly patch for up to 6 months with "standard instruction" or with "educational intervention". Evaluate changes in HRQoL using Kings Health Questionnaire (KHQ) and Beck Depression Inventory II (BDI-II)</p> <p>Groups: G1: Standard instructions G2: Educational intervention (educational booklet, OAB newsletters, dosing reminders, calendar reminders, bladder diary)</p> <p>N at enrollment: Total: 2878 G1: G2:1596</p> <p>N at follow-up: NR</p> <p>Age, mean yrs ± SD: 62.5 ± 14.8</p> <p>Women, %: 87.2</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • ≥18 years old • UUI, urinary urgency or frequency • Willing to discontinue all prescription and OTC medications for OAB • Capable of completing QOL questionnaires without assistance • Negative pregnancy test & medically acceptable contraceptive <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Contra-indications to oxybutynin • Reversible etiologies for OAB • Prior treatment with Oxytrol • Long-term care facilities and nursing homes 	<p>History of OAB, years, %: <1 yr: 12.0% 1-2 yr: 18.5% 2-4 yr: 23.1% ≥ 4 yr: 46.4%</p> <p>Sexually active: Yes: 1500 (59.2%) No: 1034 (40.8%)</p> <p>Coital incontinence, n (%): Omitted or NA: 1924 (77.2) A little: 324 (13) Moderately: 137 (5.5) A lot: 108 (4.3) Total with incontinence: 569 (22.8%)</p> <p>KHQ, embarrassed by bladder problems: Yes: (1845/2571) 71.8%</p> <p>OAB affects sex life: 586 (39% of 1500 who answered; 23.1% of total population)</p> <p>OAB affects relationship with partner: Yes: 622 (37.3% of 1666 who answered, 24.4% of total population)</p>	<p>Coital incontinence, n (%): Omitted or NA: 1831 (80.7) A little: 260 (11.5) Moderately: 116 (5.1) A lot: 62 (2.7) Total with incontinence: 438 (19.3%)</p> <p>Coital incontinence, n (%): Improved: 277 (12.6%) Worsened: 165 (7.5%) <i>P</i> < 0.0001</p> <p>Embarrassed by OAB: Improved: 828 (35.5%) Worsened: 251 (11.1%) <i>P</i> < 0.0001</p> <p>Effect of OAB on sex life: Improved: 429 (19.1%) Worsened 251 (11.1%) <i>P</i> < 0.0001</p> <p>Effect of OAB on relationship with partners: Improved: 444 (19.6%) Worsened: 271 (11.9%) <i>P</i> < 0.0001</p> <p>Interest in sex: Improved: 472 (23.4%) Worsened: 246 (12.2%) <i>P</i> < 0.0001</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Masking: -</p> <p>Pt selection criteria: +</p> <p>Loss to followup: +</p> <p>Drop-out rates: -</p> <p>Power calculation: +</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: ++</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Sand et al., 2006 (continued)	Race/ethnicity, %: White: 83.6 African American: 9.9 Hispanic: 4.8 Asian: 1.2 Other: 0.5 Follow-up: 6 months		Decreased interest in sex: Total: 1219 (52.1%) Less interest: 589 (25.2%) Much less interest: 228 (9.7%) Complete loss of interest: 402 (17.2%)	Improvement in all KHQ domains and BDI-II summary scores (<i>P</i> < 0.0001) No difference between G1 and G2 in baseline sexual symptoms or improvement	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Sand et al., 2007</p> <p>Country and setting: US, Multicenter, 327 sites (141 Urology 141, 96 Primary care, 43 Ob-Gyn, 17 Geriatric Med)</p> <p>Enrollment period: NR</p> <p>Funding: Watson Laboratories, Inc</p> <p>Author industry relationship disclosures: 7 of 7 Allergan (1) Astellas (1) Esprit (1) GlaxoSmithKline (2) GSK (1) GTx (1) Indevus-Esprit (1) Lilly (1) Novartis (3) Ortho-McNeil (1) Ortho Urology (1) Pfizer (4) Sanofi (1) USB (1) Watson (6)</p>	<p>Design: RCT, open-label (case series for drug, RCT for educational materials)</p> <p>Intervention: Oxybutynin transdermal system 3.9 mg/day, twice weekly patch for up to 6 months with "standard instruction" or with "educational intervention". Evaluate changes in HRQoL using KHQ</p> <p>Groups: G1: Standard instructions G2: Educational intervention (educational booklet, OAB newsletters, dosing reminders, calendar reminders, bladder diary)</p> <p>N at enrollment: Total: 2,878 G1: 1,282 G2: 1,596</p> <p>Women, %: 87.2</p> <p>Age, yrs ± SD: 62.5 ± 14.8</p> <p>Race/ethnicity, %: White: 83.6 African American: 9.9 Hispanic: 4.8 Asian: 1.2 Other: 0.5</p> <p>Follow-up: 6 months</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥18 • UUI, urinary urgency or frequency • Willing to discontinue all prescription and OTC medications for OAB • Capable of completing QOL questionnaires without assistance • Negative pregnancy test & medically acceptable contraceptive <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Contraindications to oxybutynin • Reversible etiologies for OAB • Prior treatment with Oxytrol • Long-term care facilities and nursing homes 	<p>KHQ, UUI severity item score, %: A lot: 38.1 Moderately: 31.3 A little: 20.5 Omitted, N/A: 10.1</p> <p>KHQ, urgency severity item score, %: A lot: 49.2 Moderately: 30.3 A little: 14.8 Omitted, N/A: 5.6</p> <p>KHQ, frequency severity item score, %: A lot: 59.3 Moderately: 32.5 A little: 7.2 Omitted, N/A: 1.0</p> <p>KHQ, nocturia severity item score, %: A lot: 47.3 Moderately: 33.5 A little: 16.7 Omitted, N/A: 2.4</p> <p>History of OAB, years, %: <1 year: 12.0 1-2 years: 18.5 2-4 years: 23.1 ≥ 4 years: 46.4</p> <p>OAB Severity: No problem: 1.8 Very minor: 4.6 Minor: 15.5 Moderate: 33.0 Severe: 28.4 Many severe problems: 16.7</p> <p>Previous OAB treatments, n (%): 0: 43.4 1: 38.7 2: 13.1 3: 2.9 4: 1.9</p>	<p>KHQ, UUI severity item score, %: A lot: 21.3 Moderately: 29.5 A little: 31.6 Omitted, N/A: 17.7</p> <p>KHQ, urgency severity item score, %: A lot: 27.3 Moderately: 33.8 A little: 27.3 Omitted, N/A: 11.5</p> <p>KHQ, frequency severity item score, %: A lot: 33.5 Moderately: 45.2 A little: 18.7 Omitted, N/A: 2.6</p> <p>KHQ, nocturia severity item score, %: A lot: 27.5 Moderately: 35.4 A little: 31.3 Omitted, N/A: 5.7</p> <p>KHQ, general health perception, mean ± SD (% improvement): -1.2 ± 17.4 (-4.3)</p> <p>KHQ, incontinence Impact, mean ± SD (% improvement): -13.5 ± 29.5 (-19.5)</p> <p>KHQ, symptom severity, mean ± SD (% improvement): -12.4 ± 24.8 (-22.2)***</p>	<p>Quality: Overall quality score: fair INTERNAL VALIDITY: poor Randomization: - Masking: NA Pt selection criteria: + Loss to followup: - Drop-out rates: - Power calculation: + Statistical issues: + EXTERNAL VALIDITY: good Age: + Baseline OAB status: + Baseline characteristics: ++ Length of followup: ++ Measurement methods: + Measurement reliability: + Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Sand et al., 2007 (continued)			KHQ, general health perception, mean \pm SD:	KHQ, role limitations, mean \pm SD (% improvement):	
			28.2 \pm 19.8	-13.3 \pm 29.2 (-29.5)	
			KHQ, incontinence impact, mean \pm SD:	KHQ, physical limitations, mean \pm SD (% improvement):	
			69.3 \pm 27.4	-11.7 \pm 29.9 (-25.1)	
			KHQ, symptom severity, mean \pm SD:	KHQ, social limitations, mean \pm SD (% improvement):	
			55.9 \pm 20.5*	-6.7 \pm 23.7 (-26.2)	
			KHQ, role limitations, mean \pm SD:	KHQ, emotions, mean \pm SD (% improvement):	
			45.1 \pm 31.0	-8.8 \pm 25.4 (-29.3)	
			KHQ, physical limitations, mean \pm SD:	KHQ, personal relationships, mean \pm SD (% improvement):	
			46.7 \pm 31.6	-6.0 \pm 23.5 (-29.1)***	
			KHQ, social limitations, mean \pm SD:	KHQ, sleep/energy, mean \pm SD (% improvement):	
			25.6 \pm 28.3	-11.2 \pm 24.1 (-20.7)	
			KHQ, emotions, mean \pm SD:	KHQ, severity (coping) measures, mean \pm SD (% improvement):	
30.0 \pm 29.2	-8.6 \pm 21.3 (-18.0)				
KHQ, personal relationships, mean \pm SD:	Side effects, application site, %:				
20.6 \pm 29.5***	Total: 14.0				
KHQ, sleep/energy, mean \pm SD:	Pruritis: 4.9				
54.2 \pm 27.3	Erythema: 4.6				
KHQ, severity (coping) measures, mean \pm SD:	Dermatitis: 4.4				
47.9 \pm 26.4	Irritation: 3.2				
	Other: 2.0				
	Side effects, %:				
	Rash: 3.0				
	Dry mouth: 2.6				
	Pruritis: 2.6				
	Skin irritation: 2.1				

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Sand et al., 2007 (continued)				<p>Withdrawn due to AEs, %: 21.3</p> <p>Withdrawn: 1452 (50.5%)</p> <p>Adverse events 21.3%</p> <p>Withdrawn consent: 7.5%</p> <p>Requirement for alternative therapy 7.4%</p> <p>Loss to follow-up 7.2%</p> <p>Noncompliance 5.6%</p> <p>Administrative decision 0.7%</p> <p>Ineligible: 0.4%</p> <p>Death 0.1%</p> <p>No reason given 0.3%</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Song et al., 2006</p> <p>Country and setting: Korea, Medical Center</p> <p>Enrollment period: May 2001 to April 2002</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: bladder training (BT) vs. Tolterodine vs. BT + Tolterodine</p> <p>Groups: G1: BT x 12 wks G2: Tolterodine 2 mg b.i.d. x 12 wks G3: Tolterodine 2 mg b.i.d. + BT x 12 wks</p> <p>N at enrollment: G1: 46 G2: 47 G3: 46</p> <p>N at follow-up: G1: 26 G2: 32 G3: 31</p> <p>Women, %: 100</p> <p>Age, mean ± SD: G1: 45.73 ± 12.68 G2: 48.41 ± 9.38 G3: 45.42 ± 9.54</p> <p>Race/ethnicity, %: Korean: 100</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 18 • ≥ 8 voids/day • Urge with or without incontinence • Symptom duration ≥ 3 months • No prior history of treatment for OAB <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Active urinary tract infection • Clinically significant SUI • Bladder outlet obstruction • Interstitial cystitis • Glaucoma • Megacolon • Maximal urine flow rate of < 10 mL/sec • Postvoid residual volume > 30% of the total amount voided on uroflowmetry 	<p>Voids/day, mean ± SD: G1: 10.93 ± 2.14 G2: 11.63 ± 2.57 G3: 11.90 ± 1.51</p> <p>Nocturia episodes/day, mean ± SD: G1: 1.45 ± 1.14 G2: 1.72 ± 1.04 G3: 1.96 ± 1.49</p> <p>Urgency, mean score ± SD: G1: 2.58 ± 1.30 G2: 2.81 ± 0.74 G3: 3.00 ± 1.10</p> <p>Maximum flow rate (mL/s), mean ± SD: G1: 20.35 ± 8.44 G2: 22.56 ± 4.94 G3: 21.19 ± 4.96</p> <p>Residual urine (mL), mean ± SD: G1: 9.08 ± 22.56 G2: 7.59 ± 12.39 G3: 6.42 ± 10.16</p> <p>Symptom duration (years), mean ± SD: G1: 6.44 ± 6.84 G2: 4.54 ± 5.15 G3: 4.10 ± 3.99</p>	<p>Voids/day, mean (% decrease) G1: 8.1 (25.9%)* G2: 8.1 (30.2%)* G3: 7.9 (33.5%)* G3/G1: <i>P</i> < 0.05</p> <p>Nocturia episodes/day, mean (% reduction): G1: 0.6 (56.1%)* G2: 0.6 (65.4%)* G3: 0.6 (66.3%)*</p> <p>Urgency, mean score (% reduction): G1: 1.4 (44.8%)* G2: 1.1 (62.2%)* G3: 1.2 (60.2%)* G3/G1: <i>P</i> = 0.021 G2/G1: <i>P</i> = 0.017 G2/G3: <i>P</i> = NS</p> <p>Satisfaction, mean score (% improved): G1: 1.5 (53.9) G2: 1.4 (63.0) G3: 1.3 (71.0)</p> <p>Dry mouth, n (%): G1: 0 (0.0) G2: 7 (21.9) G3: 9 (28.9)</p> <p>Hesitancy, n (%): G1: 0 (0.0) G2: 3 (9.4) G3: 2 (6.5)</p> <p>Decreased appetite/constipation, n (%): G1: 0 (0.0) G2: 2 (6.3) G3: 2 (6.5)</p> <p>Headache, n (%): G1: 0 (0.0) G2: 1 (3.1) G3: 0 (0.0)</p>	<p>Quality: Overall quality score: fair INTERNAL VALIDITY: poor Randomization: - Masking: - Pt selection criteria: + Loss to followup: - Drop-out rates: + Power calculation: - Statistical issues: - EXTERNAL VALIDITY: good Age: + Baseline OAB status: + Baseline characteristics: ++ Length of followup: + Measurement methods: + Measurement reliability: + Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Staskin et al., 2007</p> <p>Country and setting: US, Multicenter</p> <p>Enrollment period: August 2005 to May 2006</p> <p>Funding: Esprit Pharma, Indevus Pharmaceuticals</p> <p>Author industry relationship disclosures: 4 of 4 Allergan (3) Astellas (3) Eli Lilly (1) Esprit (4) GlaxoSmithKline (2) Indevus (3) Medtronic (1) Novartis (3) Ortho (1) Pfizer (2) Schwartz (1) Watson (4)</p>	<p>Design: RCT</p> <p>Intervention: Trospium chloride ER</p> <p>Groups: G1: trospium chloride ER 60mg qd G2: placebo</p> <p>N at enrollment: G1: 298 G2: 303</p> <p>N at follow-up: G1: 263 G2: 273</p> <p>Women, n (%): G1: 254 (85.2) G2: 256 (84.5)</p> <p>Age, mean (SE): G1: 59.6 (0.77) G2: 59.3 (0.70)</p> <p>Race/ethnicity, %: White: G1: 86.6 G2: 85.5 Black: G1: 8.7 G2: 9.9 Hispanic: G1: 3.0 G2: 2.3 Asian: G1: 1.0 G2: 1.3 Other G1: 0.7 G2: 1.0</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥18 • OAB symptoms for ≥ 6 mos • ≥ 1 “severe” urgency severity rating per 3 days, measured Indevus Urgency Severity Scale • ≥ 30 voids per 3 days • ≥ 1 UUI episodes/day • Average total volume voided < 3L per day and < 250 per void <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • SUI • Insensate incontinence or overflow incontinence • Neurogenic bladder • Significant renal disease • Hematuria • Current UTI • ≥ 3 UTIs during previous year • Significant bladder outlet obstruction • Indwelling catheter • Active IBD • Interstitial cystitis • Bladder cancer • PSA > 4 ng/ml • Prostate cancer or chronic prostatitis • Undergoing or likely to undergo bladder retraining or drill program • Diuretic estrogen use outside of a long-term stable program 	<p>Prior OAB medications, n (%): G1: 159 (53.4) G2: 151 (49.8)</p>	<p>UUI episodes/day, wk 1, mean change ± SD: G1: -1.86 ± 0.13 G2: -1.24 ± 0.14 P = .0003</p> <p>UUI episodes/day, wk 4, mean change ± SD: G1: -2.36 ± 0.17 G2: -1.75 ± 0.15 P = 0.0051</p> <p>UUI episodes/day, wk 12, mean change ± SD: G1: -2.48 ± 0.17 G2: -1.93 ± 0.16 P = 0.0022</p> <p>UUI episodes/wk, wk 1, mean change ± SD: G1: -13.03 ± 0.91 G2: -8.66 ± 0.95 P = 0.0003</p> <p>UUI episodes/wk, wk 4, mean change ± SD: G1: -16.50 ± 1.17 G2: -12.24 ± 1.07 P = 0.0054</p> <p>UUI episodes/wk, wk 12, mean change ± SD: G1: -17.34 ± 1.18 G2: -13.49 ± 1.09 P = 0.0024</p> <p>Urgency-associated voids/day, wk 1, mean change ± SD: G1: -1.90 ± 0.16 G2: -1.34 ± 0.16 P = 0.0033</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: + Masking: + Pt selection criteria: - Loss to followup: + Drop-out rates: + Power calculation: + Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: + Baseline OAB status: + Baseline characteristics: ++ Length of followup: + Measurement methods: + Measurement reliability: + Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Staskin et al., 2007 (continued)				<p>Urgency-associated voids/day, wk 4, mean change ± SD: G1: -2.66 ± 0.17 G2: -1.71 ± 0.18 <i>P</i> = 0.0003</p> <p>Urgency-associated voids/day, wk 12, mean change ± SD: G1: -3.11 ± 0.17 G2: -2.12 ± 0.19 <i>P</i> < 0.0001</p> <p>Voids/day, wk 1, mean change ± SD: G1: -1.66 ± 0.14 G2: -1.24 ± 0.13 <i>P</i> = 0.0092</p> <p>Voids/day, wk 4, mean change ± SD: G1: -2.44 ± 0.15 G2: -1.58 ± 0.15 <i>P</i> < 0.001</p> <p>Voids/day, wk 12, mean change ± SD: G1: -2.81 ± 0.15 G2: -1.99 ± 0.16 <i>P</i> < 0.001</p> <p>Urgency severity score/void, wk 1, mean change ± SD: G1: -0.21 ± 0.02 G2: -0.10 ± 0.02 <i>P</i> = 0.0002</p> <p>Urgency severity score/void, wk 4, mean change ± SD: G1: -0.26 ± 0.03 G2: -0.13 ± 0.03 <i>P</i> = 0.0008</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Staskin et al., 2007 (continued)				<p>Urgency severity score/void, wk 12, mean change ± SD: G1: -0.32 ± 0.03 G2: -0.18 ± 0.03 <i>P</i> = 0.0004</p> <p>OAB-SCS, wk 1, mean change ± SD: G1: -6.96 ± 0.51 G2: -4.81 ± 0.55 <i>P</i> = 0.0002</p> <p>OAB-SCS, wk 4, mean change ± SD: G1: -9.67 ± 0.56 G2: -6.13 ± 0.64 <i>P</i> < 0.0001</p> <p>OAB-SCS, wk 12, mean change ± SD: G1: -11.20 ± 0.55 G2: -7.80 ± 0.67 <i>P</i> < 0.0001</p> <p>Voided volume (mL), wk 1, mean change ± SD: G1: 21.61 ± 2.76 G2: 12.07 ± 2.11 <i>P</i> = 0.0036</p> <p>Voided volume (mL), wk 4, mean change ± SD: G1: 30.00 ± 3.14 G2: 17.24 ± 2.47 <i>P</i> = 0.0007</p> <p>Voided volume (mL), wk 12, mean change ± SD: G1: 29.77 ± 3.16 G2: 18.89 ± 2.79 <i>P</i> = 0.0039</p> <p>Subjects who achieved normalization, %: G1: 20.5 G2: 11.3 <i>P</i> < 0.01</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Staskin et al., 2007 (continued)				<p>At least 1 treatment emergent AE, n (%): G1: 80 (26.8) G2: 53 (17.5)</p> <p>Dry mouth, n (%): G1: 26 (8.7) G2: 9 (3.0)</p> <p>Constipation, n (%): G1: 28 (9.4) G2: 4 (1.3)</p> <p>Headache, n (%): G1: 3 (1.0) G2: 8 (2.6)</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Steers et al., 2005</p> <p>Country and setting: US (49) and Canada (15), Multicenter</p> <p>Enrollment period: NR</p> <p>Funding: Pfizer</p> <p>Author industry relationship disclosures: 3 of 4 Pfizer (3)</p>	<p>Design: RCT</p> <p>Intervention: Darifenacin CR vs. placebo</p> <p>Groups: G1: Darifenacin CR 7.5-15 mg qd G2: Placebo G3: Increased dose at 2 wks G4: No increased dose at 2 wks</p> <p>N at enrollment: G1: 269 G2: 129</p> <p>N at follow-up: G1: 261 G2: 123</p> <p>Women, N (%): G1: 227 (84.7) G2: 106 (83.5)</p> <p>Age, yrs (range): G1: 57.5 (27-89) G2: 58.5 (22-86)</p> <p>Aged < 65 years, n (%) G1: 183 (68.3) G2: 81 (63.8)</p> <p>Race/ethnicity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 18 • OAB symptoms for ≥ 6 mos • Capable of independent toileting • ≥ 5 episodes of UI per wk • ≥ 8 voids/day • Strong desire to void ≥ 1 times per day • Adequate method of contraception <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • BPH on a non-stable dose of finasteride for < 6mos • Contra-indications to anticholinergic medication • Clinically significant SU1 • BOO • PVR of > 200 mL • Pregnant or breastfeeding • GU conditions that could cause urinary symptoms • Fecal impaction or severe constipation (< 3 BM/wk) • Urogenital surgery in past 6 mos • Bladder biopsy in last 30 days • Indwelling catheter • Intermittent self-catherization • Intention to start a bladder-training program during study 	<p>Urgency episodes/day, median (IQR): G1: 8.3 (6.4, 10.6) G2: 8.0 (5.6, 10.3)</p> <p>Severity of urgency, VAS score, median (IQR): G1: 53 (41-66) G2: 53 (39-66)</p> <p>Incontinence episodes/wk, median (IQR): G1: 16.3 (9.1, 29.6) G2: 14.0 (9.0, 26.0)</p> <p>Significant leaks/week, median (IQR): G1: 7.0 (2.0, 13.1) G2: 5.3 (1.0, 11.2)</p> <p>Voids/day, median (IQR): G1: 9.9 (8.6, 11.9) G2: 10.4 (8.3, 11.6)</p> <p>Nocturia episodes/week, median (IQR): G1: 12.0 (7.0-16) G2: 11.2 (6.1-16)</p> <p>Voided volume (mL), median (IQR): G1: 175 (125, 253) G2: 177 (120, 229)</p> <p>Idiopathic OAB, n (%): G1: 253 (94.4) G2: 123 (96.8)</p>	<p>Urgency episodes/day, median change (%): G1: -2.3 (-28.2) G2: -0.9 (-11.0) G1/G2: <i>P</i> < 0.001</p> <p>Severity of urgency, VAS score, median change (%): G1: -9.1 (-16.8) G2: -3.2 (-5.6) G1/G2: <i>P</i> < 0.05</p> <p>Significant leaks/week, median change (%): G1: -3.0 (-67.3) G2: -1.8 (-42.9) G1/G2: <i>P</i> < 0.01</p> <p>Voids/day, median change (%): G1: -1.9 (-18.9) G2: -1.0 (-10.0) G1/G2: <i>P</i> < 0.001</p> <p>Nocturia episodes/week, median change (%): G1: -2.0 (-18.0) G2: -1.0 (-12.5) G1/G2: <i>P</i> = NS</p> <p>Voided volume (mL), median change (%): G1: 19 (10.5) G2: 7 (5.4) G1/G2: <i>P</i> < 0.05</p> <p>Constipation, n (%): G3: 24 (22.2) G4: 32 (20.0)</p> <p>Dry Mouth, n (%): G3: 22 (20.4) G4: 28 (17.5)</p> <p>Headache, n (%): G3: 5 (4.6) G4: 13 (8.1)</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: NA</p> <p>Masking: NA</p> <p>Pt selection criteria: +</p> <p>Loss to followup: -</p> <p>Drop-out rates: -</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: +</p> <p>Length of followup: ++</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Steers et al., 2005 (continued)		<ul style="list-style-type: none"> Concomitant use anticholinergics, antispasmodics, opioids, other drugs known to cause constipation, HRT (unless taken for ≥ 2 months), and P450 3A4 inhibitors 	<p>Previous drug therapy for frequency, enuresis, or incontinence, n (%): G1: 62 (23.1) G2: 32 (25.2)</p>		

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Steers et al., 2007</p> <p>Country and setting: Australia, Canada, US; multicenter, 30 centers</p> <p>Enrollment period: July 2001 to September 2003</p> <p>Funding: Eli Lilly, Boehringer Ingelheim GmbH.</p> <p>Author industry relationship disclosures: 7 of 7 Eli Lilly (7)</p>	<p>Design: RCT, placebo-controlled, double-blind, stratified by urodynamic observation</p> <p>Intervention: duloxetine vs placebo</p> <p>Groups: G1: placebo x 2 wks, duloxetine 40 mg b.i.d. x 4 wks, duloxetine 60 mg b.i.d. x 8 wks G2: placebo b.i.d. x 14 wks</p> <p>N at enrollment: G1: 153 G2: 153</p> <p>N at follow-up, N (%): G1: 90 (59) G2: 120 (78)</p> <p>Age, mean ± SD: G1: 56.0 ± 14.8 G2: 53.3 ± 14.1</p> <p>Women, N (%): 306 (100)</p> <p>BMI, mean kg/m2 ± SD: G1: 29.2 (7.4) G2: 30.2 (7.8)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> women >18 yoa predominant symptoms of OAB ≥ 3 months defined as bothersome urinary urgency and/or urge, abnormal voiding frequency (≤2 h mean daytime voiding interval) UDS observation of either detrusor OA or urgency that limited bladder capacity to <400mL <p>Exclusion criteria:</p> <ul style="list-style-type: none"> -SUI PVR >100 mL Mean 24 h total volume voided of ≥ 3L Urine culture ≥4 UTIs in past yr Regular use of meds for OAB w/in 4wks of enrollment Previous use of duloxetine Continence surgery ≤ 6 mos Any major surgery ≤ 3 mos Pelvic organ prolapse > ICS Stage II Non-pharm intervention ≤ 3 mos 	<p>Incontinence episodes, all assessable pts, mean: G1: 1.70 G2: 1.44</p> <p>Urinary incontinence episode (wet OAB): G1: 2.34 G2: 2.07</p> <p>Pads/week, mean: G1: 7.81 G2: 7.05</p> <p>Voids/day, mean: G1: 10.76 G2: 10.49</p> <p>Daytime voiding interval, mean: G1: 113.58 G2: 119.63</p> <p>Nocturia episodes/day, mean: G1: 1.47 G2: 1.63</p> <p>Voided volume (mL), mean: G1: 175.41 G2: 183.40</p> <p>QoL Instrument score, mean ± SD: I-QOL: G1: 56.65 ± 24.80 G2: 57.11 ± 23.3 U-IIQ: G1: 2.44 ± 1.18 G2: 2.39 ± 1.15 U-UDI: G1: 3.18 ± 0.58 G2: 3.15 ± 0.71</p> <p>Bladder compliance (mL/cmH₂O) G1: 71.8 ± 85.7 G2: 79.3 ± 91.0</p>	<p>Incontinence episodes, all assessable pts, mean change: G1: -0.74 G2: -0.14 <i>P</i> = 0.006</p> <p>Incontinence episodes, wet OAB pts, mean change: G1: -1.03 G2: -0.24 <i>P</i> = 0.032</p> <p>Pads/week, mean change: G1: -1.41 G2: -0.53 <i>P</i> = 0.143</p> <p>Voids/day, mean change: G1: -1.81 G2: -0.62 <i>P</i> < 0.001</p> <p>Daytime voiding interval, mean change: G1: 29.46 G2: 6.51 <i>P</i> < 0.001</p> <p>Nocturia episodes/day, mean change: G1: 0.05 G2: -0.18 <i>P</i> = 0.004</p> <p>Voided volume (mL), mean change: G1: 8.85 G2: 5.07 <i>P</i> = 0.171</p> <p>Condition improved, 80 mg/day, %: G1: 59.9 G2: 42.9 <i>P</i> = 0.005</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: -</p> <p>Drop-out rates: ++</p> <p>Power calculation: +</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Steers et al., 2007 (continued)		<ul style="list-style-type: none"> • Pelvic-floor-muscle training that had not been stable for 3 mos or would not remain stable during the trial 	<p>SUI: G1: 0 G2: 0</p> <p>Maximum cystometric capacity mL G1: 318.3 ± 152.4 G2: 330.4 ± 135.4</p> <p>Volume threshold for first detrusor contraction mL G1: 226.4 ± 141.8 G2: 250.6 ± 146.0</p> <p>Moderate or severe bladder condition from PGI-I Scale (%): G1: 84.9 G2: 88.8</p> <p>I-QOL total score, mean ± SD: G1: 56.6 (24.9) G2: 57.0 (23.2)</p> <p>UIE , mean ± SD: G1: 1.55 (2.08) G2: 1.41 (2.00)</p> <p>VE24 , mean ± SD: G1: 10.8 (3.3) G2: 10.6 (3.6)</p> <p>Symptom of bothersome urgency, %: G1: 99.3 G2: 99.3</p> <p>Symptoms of urge UI, %: G1: 88.2 G2: 88.9</p> <p>Symptoms of SUI: G1: 42.5 G2: 50.3</p> <p>Urodynamic DOA G1: 42.5 G2: 41.8</p>	<p>Condition improved, 120 mg/day, %: G1: 61.6 G2: 42.1 <i>P</i> = 0.008</p> <p>Condition improved, overall, %: G1: 53.4 G2: 41.9 <i>P</i> = 0.052</p> <p>Headache: G1: 13 (8.5) G2: 8 (5.2) <i>P</i> = 0.336</p> <p>Diarrhea: G1: 10 (6.5) G2: 5 (3.3) <i>P</i> = 0.289</p> <p>Cough: G1: 7 (4.6) G2: 6 (3.9) <i>p</i>>0.999</p> <p>Change in QOL Instrument score, mean ±: I-QOL G1: 8.37 ± 15.89 G2: 4.87 ± 15.27 G1>G2 (<i>P</i> = .035)</p> <p>U-IIQ G1: -0.44 ± 0.76 G2: -0.22 ± 0.87 <i>P</i> = 0.018</p> <p>U-UDI G1: -0.26± 0.67 G2: -0.19 ± 0.59 <i>P</i> = 0.440</p> <p>Bladder compliance (mL/cmH20) G1: 16.5 ±131.3 G2: 1.0 ± 120.8 <i>P</i> = 0.655</p> <p>Maximum cystometric capacity mL G1: 22.1 ± 116.4 G2: 23.8 ± 108.5 <i>P</i> = 0.619</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Steers et al., 2007 (continued)			<p>Sensory urgency: G1: 57.5 G2: 58.2</p> <p>Previous continence surgery, %: G1: 13.1 G2: 11.8</p> <p>Current pelvic-floor muscle training, %: G1: 9.2 G2: 11.8</p> <p>Previous behavioral therapy, %: G1: 3.3 G2: 2.6</p> <p>Previous tolterodine therapy, %: G1: 10.5 G2: 7.2</p> <p>Previous oxybutynin therapy, %: G1: 13.0 G2: 9.2</p>	<p>Volume threshold for first detrusor contraction mL G1: 90.0 ± 200.6 G2 : 13.9 ± 112.8 <i>P</i> = 0.254</p> <p>PGI-I score Better 4 wks (80 mg/day) G1: 59.9 G2: 42.9 <i>P</i> = 0.005</p> <p>TEAEs reported, % G1: 79.1 G2: 55.6 <i>P</i> < 0.001</p> <p>Appetite decreased G1: 6 (3.9) G2: 0 G1>G2, <i>P</i> = 0.030</p> <p>Arthralgia G1: 6 (3.9) G2: 3 (2.0) <i>P</i> = 0.501</p> <p>Somnolence: G1: 6 (3.9) G2: 0 <i>P</i> = 0.30</p> <p>Sweating increased: G1: 6 (3.9) G2: 2 (1.3) <i>P</i> = 0.283</p> <p>UTI: G1: 6 (3.9) G2: 6 (3.9) <i>p</i>>0.999</p> <p>Anorgasmia G1: 5 (3.3) G2: 0 G1>G2, <i>P</i> = 0.060</p> <p>Anxiety: G1: 5 (3.3) G2: 0 G1>G2, <i>P</i> = 0.060</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Steers et al., 2007 (continued)				<p>Tremor: G1: 5 (3.3) G2: 0 G1>G2, P = 0.060</p> <p>Upper respiratory infection: G1: 5 (3.3) G2: 4 (2.6) <i>P</i> > 0.999</p> <p>Vomiting: G1: 5 (3.3) G2: 3 (2.0) <i>P</i> = 0.723</p> <p>Abdominal pain: G1: 4 (2.6) G2: 1 (0.7) <i>P</i> = 0.371</p> <p>Back pain: G1: 4 (2.6) G2: 1 (0.7) <i>P</i> = 0.371</p> <p>Note: mean change consistent at 4 wks (duloxetine 80 mg/d) and 8 wks (duloxetine 120 mg/d)</p> <p>Change in I-QOL total score, mean ± SD: Discontinuation due to TEAEs, % G1: 28.1 G2: 5.2 <i>P</i> < 0.001</p> <p>Treatment emergent adverse events, n (%):</p> <p>Nausea: G1: 47 (30.7) G2: 7 (4.6) G1>G2, P < 0.001</p> <p>Dry mouth: G1: 25 (16.3) G2: 2 (1.3) G1>G2, P < 0.001</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Steers et al., 2007 (continued)				<p>Dizziness: G1: 22 (14.4) G2: 1 (0.07) G1>G2, <i>P</i> < 0.001</p> <p>Constipation: G1: 21 (13.7) G2: 5 (3.3) G1>G2, <i>P</i> = 0.002</p> <p>Insomnia: G1: 20 (13.1) G2: 5 (3.3) G1>G2, <i>P</i> = 0.003</p> <p>Fatigue: G1: 16 (10.5) G2: 3 (2.0) G1>G2, <i>P</i> = 0.003</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Sussman and Garely, 2002</p> <p>Country and setting: US, 2 sites</p> <p>Enrollment period: NR</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Multicenter RCT</p> <p>Intervention: Tolterodine ER vs. Oxybutynin ER</p> <p>Groups: G1: Tolterodine ER 2 mg qd x 8 wks G2: Tolterodine ER 4 mg qd x 8 wks G3: Oxybutynin ER 5 mg qd x 8 wks G4: Oxybutynin ER 10 mg qd x 8 wks</p> <p>N at enrollment: G1: 333 G2: 336 G3: 313 G4: 307</p> <p>N at follow-up: G1: 313 (86) G2: 316 (88) G3: 286 (81) G4: 285 (79)</p> <p>Age, yrs ± SD: G1: 63.8 ± 15.7 G2: 63.4 ± 16.6 G3: 59.8 ± 16.5 G4: 63.2 ± 15.9</p> <p>Women, N (%): G1: 243 (73) G2: 254 (76) G3: 245 (78) G4: 222 (72)</p> <p>Race/ethnicity, n (%): White: G1: 278 (84) G2: 296 (88) G3: 256 (82) G4: 247 (81) Black: G1: 28 (8) G2: 23 (7) G3: 41 (13) G4: 42 (14)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • 18+ years old • OAB (urinary frequency + urgency) w/ or w/o UI <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Pure SI • Urinary retention • Gastric retention • Uncontrolled narrow-angle glaucoma • Significant hepatic or renal dysfunction • Symptomatic or recurrent UTI • Electrostimulation • Bladder training • Pelvic floor exercise w/in 1 wk of first study visit or expected to start during study • Indwelling catheter • Intermittent self-cath • Contraindication to antimuscarinic trt • Estrogen therapy started less than 2 mos prior to first visit • Any trt for UI w/in 1 wk of first visit • Use of anticholinergic drug or potent inhibitors of cytochrome P450 3A4 • Pregnant • Lactating 	<p>Presence of UI, n (%): G1: 195 (59) G2: 214 (64) G3: 199 (64) G4: 195 (64)</p> <p>Duration of sx, n (%): < 6 months: G1: 38 (11) G2: 48 (14) G3: 62 (20) G4: 43 (14) 6 months to 5 years: G1: 226 (68) G2: 224 (67) G3: 183 (59) G4: 205 (67) > 5 years: G1: 68 (20) G2: 64 (19) G3: 67 (21) G4: 59 (19)</p> <p>Severity of bladder condition, % No problems: G1: 1.3 G2: 0.3 G3: 1.0 G4: 0.7 Some very minor problems: G1: 6 G2: 5 G3: 5 G4: 5 Some minor problems: G1: 18 G2: 16 G3: 21 G4: 15 Moderate problems: G1: 45 G2: 50 G3: 41 G4: 47</p>	<p>Improvement in bladder condition, 8 wks, overall, %: G1: 60 G2: 70 G3: 59 G4: 60 G2/G3: $P < 0.01$ G2/G4: $P < 0.01$</p> <p>Improvement in bladder condition, 8 wks, moderate or severe bladder condition, %: G2: 77 G4: 65 G2/G4: $P < 0.01$</p> <p>Improvement in bladder condition, 8 wks, treatment naïve, %: G1: 60 G2: 69 G3: 60 G4: 61 $P = 0.11$ for improvement rates $P > 0.05$ for overall difference btw trt arms</p> <p>Improvement in bladder condition, 8 wks, treatment experienced, %: G1: 57 G2: 75 G3: 59 G4: 54 $P = 0.11$ for improvement rates $P > 0.05$ for overall difference btw trt arms</p> <p>Withdrawal due to AE, (%): G2: 6 G4: 13 $P = 0.001$</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Masking: -</p> <p>Pt selection criteria: +</p> <p>Loss to followup: -</p> <p>Drop-out rates: -</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: -</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Sussman and Garely, 2002 (continued)	<p>Hispanic: G1: 19 (6) G2: 15 (5) G3: 15 (5) G4: 16 (5)</p> <p>Other, n (%): G1: 8 (2) G2: 2 (<1) G3: 1 (<1) G4: 2 (<1)</p> <p>Parity: NR</p>	<ul style="list-style-type: none"> • Childbearing potential w/o adequate contraception 	<p>Severe problems: G1: 26 G2: 21 G3: 23 G4: 25</p> <p>Many severe problems: G1: 4 G2: 7 G3: 8 G4: 8</p>	<p>Mean change in severity of dry mouth (visual analogue scale): G1: 2.3 G2: 6.0 G3: 6.3 G4: 11.3 G1 vs G2: <i>P</i> = NS G3 vs G4: <i>P</i> = 0.05 G2 vs G4: <i>P</i> = 0.03</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Swift et al., 2003</p> <p>Country and setting: Europe (167 centers), North America (74 centers), Australia and New Zealand (4 centers), University</p> <p>Enrollment period: February 1999 to October 1999</p> <p>Funding: Pharmacia Corp</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT double blind placebo-controlled double dummy, random permuted blocks of 6</p> <p>Intervention: Tolterodine ER vs. Tolterodine IR</p> <p>Groups: G1: Tolterodine ER 4 mg daily G2: Tolterodine IR 2 mg b.i.d. G3: placebo</p> <p>N at enrollment: G1: 417 G2: 408 G3: 410</p> <p>N at follow-up: Total: 1092</p> <p>Women, %: 100</p> <p>Age, yrs ± SD: G1: 59 ± 14 G2: 59 ± 14 G3: 60 ± 14</p> <p>Race/ethnicity, n (%): White: G1: 396 (95) G2: 389 (95) G3: 383 (93) Black: G1: 15 (4) G2: 12 (3) G3: 20 (5) Asian/Pacific: G1: 5 (1) G2: 4 (1) G3: 2 (1) Mixed: G1: 0 G2: 3 (1) G3: 5 (1) Unknown: G1: 1 (<1) G2: 0 G3: 0</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Women • Age ≥ 18 • ≥ 8 voids/day • > 5 UUI/ week • Symptoms x ≥ 6 months (per voiding diary) <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • SUI • Total daily urine volume > 3 liters • Hepatic/ renal disease • UTI • IC • Hematuria • BOO • Current e-stim • Current bladder training • Indwelling catheter • CIC • Pregnant/ nursing • Childbearing age without BC • Anticholinergic meds • Meds inhibit cytochrome450 	<p>Incontinence episodes/week, mean ± SD: G1: 22.1 ± 22.5 G2: 22.9 ± 21.9 G3: 23.9 ± 21.2</p> <p>Pads/day, mean ± SD: G1: 1.6 ± 2.1 G2: 1.5 ± 2.0 G3: 1.7 ± 2.4</p> <p>Voids/day, mean ± SD: G1: 10.8 ± 4.2 G2: 11.1 ± 3.7 G3: 11.2 ± 3.9</p> <p>Voided volume (mL), mean ± SD: G1: 141.2 ± 43.1 G2: 137.2 ± 41.2 G3: 135.7 ± 43.2</p> <p>Previous drug therapy: n (%): G1: 235 (56) G2: 222 (54) G3: 225 (55)</p>	<p>Incontinence episodes/week, mean ± SD: G1: 10.3 ± 17.2 G2: 12.8 ± 19.8 G3: 16.7 ± 19.7 G1/G3: P = 0.001 G2/G3: P = 0.001</p> <p>Pads/day, mean ± SD: G1: 1.0 ± 1.8 G2: 1.0 ± 1.5 G3: 1.5 ± 2.2 G1/G3: P = 0.001 G2/G3: P = 0.001</p> <p>Voids/day, mean ± SD: G1: 9.0 ± 3.2 G2: 9.3 ± 4.0 G3: 9.9 ± 3.8 G1/G3: P = 0.001 G2/G3: P = 0.005</p> <p>Voided volume (mL), mean ± SD: G1: 179.1 ± 66.6 G2: 169.7 ± 65.6 G3: 149.0 ± 56.3 G1/G3: P = 0.001 G2/G3: P = 0.001</p> <p>Clinical effectiveness*, dry mouth: G1: 0.53 G2: 0.39 G3: 0.30</p> <p>Dry mouth, n (%): G1: 105 (25.3) G2: 127 (31.2) G3: 33 (8.0) G1/G3: P < 0.01 G2/G3: P < 0.01</p> <p>Abdominal pain, n (%): G1: 18 (4.3) G2: 12 (2.9) G3: 7 (1.7) G1/G3: P = 0.03</p> <p>Constipation, n (%): G1: 27 (6.5) G2: 27 (6.6) G3: 14 (3.4)</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: fair</p> <p>Randomization: +</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: ++</p> <p>Drop-out rates: ++</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Swift et al., 2003 (continued)	BMI, kg/m² ± SD: G1: 28.8 ± 13.8 G2: 29.0 ± 11.0 G3: 28.8 ± 6.7			Dyspepsia, n (%): G1: 11 (2.7) G2: 14 (3.4) G3: 6 (1.5) Nausea, n (%): G1: 7 (1.7) G2: 9 (2.2) G3: 9 (2.2) Diarrhea, n (%): G1: 10 (2.4) G2: 14 (3.4) G3: 9 (2.2) Flatulence, n (%): G1: 8 (1.9) G2: 11 (2.7) G3: 6 (1.5) Xerophthalmia, n (%): G1: 16 (3.9) G2: 8 (2.0) G3: 8 (2.0) Abnormal vision, n (%): G1: 5 (1.2) G2: 4 (1.0) G3: 2 (0.5) Headache, n (%): G1: 29 (7.0) G2: 14 (3.4) G3: 19 (4.6) UTI, n (%): G1: 15 (3.6) G2: 11 (2.7) G3: 19 (4.6) Insomnia, n (%): G1: 7 (1.7) G2: 2 (0.5) G3: 9 (2.2) Somnolence, n (%): G1: 12 (2.9) G2: 11 (2.7) G3: 8 (2.0) Dizziness, n (%): G1: 7 (1.7) G2: 7 (1.7) G3: 4 (1.0)	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Swift et al., 2003 (continued)				<p>Hypertension, n (%): G1: 6 (1.4) G2: 4 (1.0) G3: 4 (1.0)</p> <p>Sinusitis, n (%): G1: 8 (1.9) G2: 2 (0.5) G3: 3 (0.7)</p> <p>Arthritis, n (%): G1: 1 (0.2) G2: 5 (1.2) G3: 1 (0.2)</p> <p>Dry skin, n (%): G1: 2 (0.5) G2: 5 (1.2) G3: 1 (0.2)</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics*	Outcomes	Quality Rating
<p>Author: Szonyi et al., 1995</p> <p>Country and setting: UK</p> <p>Enrollment period: NR</p> <p>Funding: Smith & Nephew Pharmaceuticals Ltd</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: Oxybutynin plus bladder training vs placebo plus bladder training</p> <p>Groups: G1: Oxybutynin 2.5 mg b.i.d. with dose titration on days 29 and 43 plus bladder training G2: placebo + bladder training</p> <p>N at enrollment: G1: 30 G2: 30</p> <p>N at follow-up: G1: 16 G2: 23</p> <p>Women, n (%): 56 (93)</p> <p>Age, mean ± SD: 82.2 ± 6.06</p> <p>Race/ethnicity: NR</p> <p>Weight (kg), mean ± SD: 67.4 ± 14.92</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age > 70 • Frequency, urgency and UUI • Mobile • Able to keep diary <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • UTI • Hepatic or renal disease • Glaucoma • Uncontrolled diabetes • Taking imipramine or propantheline 		<p>Voids/2 weeks, median change (95% CI): G1/G2: 577 (-27.0, 6.0) <i>P</i> = 0.0025</p> <p>Nocturia episodes/2 weeks, median change (95% CI): G1 vs. G2: -9.5 (-11.0, 3.0)</p> <p>Nocturia episodes/2 weeks, median change (95% CI): G1/G2: -1.0 (-3.0, 2.0)</p> <p>Patient assessment of benefit, %: 29 days: G1: 86 G2: 55 <i>P</i> = 0.02 43 days: G1: 71 G2: 59 <i>P</i> = 0.41 57 days: G1: 79 G2: 55 <i>P</i> = 0.09</p> <p>Patient response, 29 days, n: Cure: G1: 1 G2: 0 Significant improvement: G1: 15 G2: 8 Marginal improvement: G1: 7 G2: 8</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: -</p> <p>Drop-out rates: -</p> <p>Power calculation: +</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: +</p> <p>Length of followup: -</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics*	Outcomes	Quality Rating
Szonyi et al., 1995 (continued)				<p>No change: G1: 5 G2: 13</p> <p>Patient response, 57 days, n: Cure: G1: 4 G2: 3 Significant improvement: G1: 14 G2: 8 Marginal improvement: G1: 3 G2: 4 No change: G1: 7 G2: 14</p> <p>Dry mouth, %: G1: 93 G2: 86</p> <p>Blurred vision, %: G1: 50 G2: 59</p> <p>Heartburn, %: G1: 57 G2: 45</p> <p>Constipation, %: G1: 50 G2: 45</p> <p>Dry skin, %: G1: 50 G2: 59</p> <p>Poor compliance (< 75% of tablets), %: G1: 20 G2: 20</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Tseng et al., 2009</p> <p>Country and setting: Taiwan, University</p> <p>Enrollment period: January 2005 to November 2005</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Prospective cohort randomized</p> <p>Intervention: Tolterodine 2 mg b.i.d. vs. tolterodine 2 mg b.i.d. + conjugated equine estrogen 0.625 mg twice per week</p> <p>Groups: G1: tolterodine 2 mg b.i.d. G2: tolterodine 2 mg b.i.d. + conjugated equine estrogen 0.625 mg twice per week</p> <p>N at enrollment: G1: 40 G2: 40</p> <p>N at follow-up: G1: 40 G2: 40</p> <p>Age, mean ± SD: G1: 64.5 ± 7.4 G2: 66.2 ± 6.8</p> <p>Race/ethnicity: NR</p> <p>Women, N (%): G1: 40 (100) G2: 40 (100)</p> <p>BMI, kg/m² ± SD: G1: 24.5 ± 3.9 G2: 25.3 ± 3.8</p> <p>Previous antimuscarinic Rx, n (%): G1: 5 (12.5) G2: 6 (15)</p> <p>Prior hysterectomy, n (%): G1: 12 (30) G2: 14 (35)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 18 • OAB symptoms • Amenorrheic <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Advanced POP > Stage 2 • Women with storage and voiding dysfunction • undiagnosed • Severe constipation • Elevated PVR • Neurological deficit • Renal/ hepatic disease • Narrow angle glaucoma • Urinary retention • Gastric retention • Hypersensitivity to drugs • BOO • Cardiac conduction disorders • Myasthenia gravis • History of CVA • History of VTE • Gallbladder disease • Known or suspected breast carcinoma • Undiagnosed genital bleeding • HRT within 3 months of study enrollment 	<p>UUI episodes/day, mean ± SD: G1: 1.8 ± 0.7 G2: 2.1 ± 1.1</p> <p>Urgency episodes/ day, mean ± SD: G1: 4.5 ± 0.8 G2: 4.3 ± 0.7</p> <p>Nocturia episodes/ day, mean ± SD: G1: 3.5 ± 0.8 G2: 3.3 ± 0.8</p> <p>Voids/day, mean ± SD: G1: 14.1 ± 1.3 G2: 14.8 ± 1.5</p> <p>UDI-6 score, mean ± SD: G1: 9.5 ± 3.9 G2: 8.6 ± 3.8</p> <p>IIQ-7 score, mean ± SD: G1: 10.2 ± 4.5 G2: 9.4 ± 3.6</p> <p>Voided volume (mL), mean ± SD: G1: 108.5 ± 14.0 G2: 115.8 ± 15.1</p>	<p>UUI episodes/day, mean ± SD: G1: 1.5 ± 0.5 G2: 1.5 ± 0.5 <i>P</i> = NS</p> <p>Urgency episodes/ day, mean ± SD: G1: 3.5 ± 0.5 G2: 3.3 ± 0.6</p> <p>Nocturia episodes/ day, mean ± SD: G1: 2.9 ± 0.6 G2: 2.6 ± 0.7</p> <p>Voids/day, mean ± SD: G1: 6.4 ± 1.9 G2: 5.8 ± 0.9 <i>P</i> = 0.001</p> <p>UDI-6 score, mean ± SD: G1: 7.2 ± 2.9 G2: 6.9 ± 2.7 <i>P</i> < 0.001</p> <p>IIQ-7 score, mean ± SD: G1: 6.5 ± 2.7 G2: 6.1 ± 2.5 <i>P</i> < 0.001</p> <p>Voided volume (mL), mean ± SD: G1: 134.5 ± 15.8 G2: 141.9 ± 16.1 <i>P</i> = 0.007</p> <p>Adverse events: None</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: fair</p> <p>Randomization: +</p> <p>Method and blinding: -</p> <p>Pt selection criteria: +</p> <p>Loss to followup: ++</p> <p>Drop-out rates: ++</p> <p>Power calculation: +</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Van Kerrebroeck et al., 2001</p> <p>Freeman et al., 2003*</p> <p>Country and setting: North America (74 centers), Australasia (4 centers), Europe (89 centers)</p> <p>Enrollment period: NR</p> <p>Funding: Pharmacia Corporation</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: Tolterodine ER vs Tolterodine IR vs placebo</p> <p>Groups: G1: tolterodine ER 4 mg qd G2: tolterodine IR 2 mg b.i.d. G3: placebo</p> <p>N at enrollment: G1: 507 G2: 514 G3: 508</p> <p>N at follow-up: Total: 1442 G1: 398 G3: 374</p> <p>Women, n (%): G1: 417 (82) G2: 408 (79) G3: 410 (81)</p> <p>Age, mean (range): G1: 60 (20, 89) G2: 60 (22, 92) G3: 61 (22, 93)</p> <p>Race/ethnicity, %:* White: G1: 95.7 G3: 94.7 Black: G1: 3.0 G3: 3.5 Asian/Pacific Islander: G1: 1.0 G3: 0.8 Other: G1: 0.3 G3: 1.1</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 18 • Urinary frequency (≥ 8 voids/day) <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • SUI • Total daily urine volume > 3 L • Contra-indications to antimuscarinic treatment • Hepatic or renal disease • UITs • Interstitial cystitis • Hematuria • BOO • Current electrostimulation or bladder training therapy • Indwelling catheter or intermittent self-catheterization • Pregnant or nursing • Women not using reliable contraception • Being treated for OAB with other anticholinergic drugs or drugs that inhibit cytochrome P450 3A4 isoenzymes • Estrogen therapy < 2 months • Treatment w/ investigational drug < 2 months 	<p>Incontinence episodes/week, mean (range): G1: 22.1 (0, 168.0) G2: 23.2 (0, 168.0) G3: 23.3 (0, 168.0)</p> <p>≥ 5 incontinence episodes/week, n (%): G1: 492 (97) G2: 498 (97) G3: 494 (97)</p> <p>Pads/day, mean (range): G1: 1.4 (0-18) G2: 1.4 (0-25) G3: 1.5 (0-22)</p> <p>Voids/day, mean (range): G1: 10.9 (2.3, 51.3) G2: 11.1 (2.0, 48.6) G3: 11.3 (2.0, 37.4)</p> <p>≥ 8 voids/day, n (%): G1: 458 (90) G2: 469 (91) G3: 467 (92)</p> <p>Previous drug therapy, n (%): G1: 270 (53) G2: 276 (54) G3: 263 (52)</p> <p>Poor efficacy, %: G1: 43 G2: 38.4 G3: 40.7</p> <p>Able to finish tasks before visiting a toilet, %:* G1: 5 G3: 6</p>	<p>Urgency, subjective assessment, 12 wks, n (%):* Improvement: G1: 173 (44) G3: 118 (32) G1/G3: <i>P</i> < 0.001 No change: G1: 201 (51) G3: 212 (57) Deterioration: G1: 22 (6) G3: 44 (12) G1/G3: <i>P</i> < 0.002</p> <p>Urgency, improvement, 12 wks, women only, %:* G1: 46.6 G3: 26.6 G1/G3: <i>P</i> = 0.001 OR 1.81 (95% CI: 1.31, 2.49)</p> <p>Not able to hold urine, 12 wks, %:* G1: 58 G3: 32 G1/G3: <i>P</i> < 0.001</p> <p>Incontinence episodes/week, mean change ± SD (%) G1: -11.8 ± 17.8 G2: -10.6 ± 16.9 G3: -6.9 ± 15.4 G1/G3: <i>P</i> = 0.00001 G2/G3: <i>P</i> = 0.0005 G1/G2: <i>P</i> < 0.05</p> <p>Incontinence episodes/week, median % change: G1: -71 G2: -60 G3: -30 G1/G2: <i>P</i> < 0.05</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: + Masking: + Pt selection criteria: + Loss to followup: + Drop-out rates: - Power calculation: - Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: + Baseline OAB status: + Baseline characteristics: ++ Length of followup: + Measurement methods: + Measurement reliability: + Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Van Kerrebroeck et al., 2001 Freeman et al., 2003* (continued)			Voided volume (mL), mean (range): G1: 141 (36, 338) G2: 137 (38, 283) G3: 136 (31, 374)	Pads/day, mean ± SD: G1: -0.5 ± 1.4 G2: -0.5 ± 1.8 G3: -0.2 ± 1.4 G1/G3: <i>P</i> = 0.0145 G2/G3: <i>P</i> = 0.0035 Voids/day, mean ± SD: G1: -3.5 ± 4.9 G2: -3.3 ± 4.4 G3: -2.2 ± 4.0 G1/G3: <i>P</i> = 0.00001 G2/G3: <i>P</i> = 0.0002 Voluntary voids/day, mean ± SD: G1: -1.8 ± 3.4 G2: -1.7 ± 3.3 G3: -1.2 ± 2.9 G1 vs G3 G1/G3: <i>P</i> = 0.00047 G2/G3: <i>P</i> = 0.0079 Bladder symptoms, improvement, 12 wks, women only, %:* G1: 62.8 G3: 48.4 G1/G3: <i>P</i> = 0.001 OR 1.78 (95% CI: 0.34, 2.37) Treatment benefit, 12 wks, n (%):* Much benefit: G1: 172 (43.2) G3: 88 (23.5) G1/G3: <i>P</i> < 0.001 Little benefit G1: 138 (34.7) G3: 118 (31.6) No benefit G1: 88 (22.1) G3: 168 (44.9)	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Van Kerrebroeck et al., 2001</p> <p>Freeman et al., 2003* (continued)</p>				<p>Able to finish tasks before visiting a toilet, 12 wks, %:* G1: 33 G3: 18</p> <p>G1/G3: $P < 0.001$</p> <p>Voided volume (mL), mean change \pm SD: G1: $+34 \pm 51$ G2: $+29 \pm 47$ G3: $+14 \pm 41$ G1/G3: $P = 0.00001$ G2/G3: $P = 0.0001$</p> <p>Discontinued due to AEs, n (%): G1: 27 (5) G2: 28 (5) G3: 33 (6)</p> <p>Reported serious adverse events, n: G1: 7 G2: 12 G3: 18</p> <p>Parasympathetic Dry mouth, n (%): G1: 118 (23) G2: 156 (30) G3: 39 (8)</p> <p>Xerophthalmia, n (%): G1: 17 (3) G2: 12 (2) G3: 10 (2)</p> <p>Abnormal vision, n (%): G1: 6 (1) G2: 4 (1) G3: 2 (0.5)</p> <p>Dry skin, n (%): G1: 2 (0.5) G2: 6 (1) G3: 1 (0.5)</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Van Kerrebroeck et al., 2001 Freeman et al., 2003* (continued)				<p>Gastrointestinal Constipation, n (%): G1: 30 (6) G2: 35 (7) G3: 22 (4)</p> <p>Dyspepsia, n (%): G1: 15 (3) G2: 16 (3) G3: 7 (1)</p> <p>Abdominal pain, n (%): G1: 19 (4) G2: 13 (3) G3: 8 (2)</p> <p>Diarrhea, n (%): G1: 10 (2) G2: 16 (3) G3: 11 (2)</p> <p>Flatulence, n (%): G1: 10 (2) G2: 14 (3) G3: 9 (2)</p> <p>Nausea, n (%): G1: 7 (1) G2: 10 (2) G3: 10 (2)</p> <p>Headache, n (%): G1: 32 (6) G2: 19 (4) G3: 23 (5)</p> <p>Somnolence, n (%): G1: 14 (3) G2: 13 (3) G3: 9 (2)</p> <p>Dizziness, n (%): G1: 11 (2) G2: 9 (2) G3: 5 (1)</p> <p>Fatigue, n (%): G1: 11 (2) G2: 6 (1) G3: 4 (1)</p> <p>Insomnia, n (%): G1: 7 (1) G2: 2 (0.5) G3: 9 (2)</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Van Kerrebroeck et al., 2001				Urinary tract infection, n (%): G1: 16 (3) G2: 13 (3) G3: 20 (4)	
Freeman et al., 2003* (continued)				Dysuria, n (%): G1: 5 (1) G2: 8 (2) G3: 1 (0.5)	
				Peripheral edema, n (%): G1: 7 (1) G2: 7 (1) G3: 4 (1)	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Versi et al., 2000</p> <p>Country and setting: US, 20 Academic health centers</p> <p>Enrollment period: NR</p> <p>Funding: ALZA Corp</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: Oxybutynin CR vs Oxybutynin IR</p> <p>Doses were increased/decreased in 5 mg/d increments until optimal dose identified. Once identified the dose was maintained for 1 wk and urinary diary completed</p> <p>Groups: G1: Oxybutynin CR 5-20 mg/d G2: Oxybutynin IR 5-20 mg/d</p> <p>N at enrollment: G1: 111 G2: 115</p> <p>N at follow-up: G1: 107 G2: 106</p> <p>Age, yrs ± SD: G1: 58.8 G2: 59.6</p> <p>Race/ethnicity, %: White: G1: 86.5 G2: 90.4 Black: G1: 5.4 G2: 3.5 Hispanic G1: 5.4 G2: 5.2 Asian: G1: 0.9 G2: 0 Native American G1: 0 G2: 0.9</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> Known to be responsive to anticholinergic trt 7-45 UI episodes per wk Adult Community-dwelling 4+ days of incontinence/wk <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Clinically significant medical problems PVR>100mL Contraindication to oxybutynin 	<p>UUI episodes/week, mean: G1: 18.6 G2: 19.8</p> <p>Incontinence episodes/week, mean: G1: 20.2 G2: 22.4</p>	<p>UUI episodes/week, mean (%): G1: 2.9 (83) G2: 4.4 (76) P value for change from baseline: G1: $P < 0.001$ G2: $P < 0.001$ $P = 0.36$ for % reduction</p> <p>Incontinence episodes/week, mean (%): G1: 3.5 (81) G2: 5.4 (75) P value for change from baseline: G1: $P < 0.001$ G2: $P < 0.001$ $P = 0.85$ for % reduction</p> <p>Rates for Dry Mouth (%): G1: 47.7 G2: 59.1 $P = 0.09$</p> <p>Pts reporting any dry mouth, cumulative %: Dose of 5mg/day: G1: 18.9 G2: 35.7 Dose of 10mg/day: G1: 39.5 G2: 61.2 Dose of 15 mg/day: G1: 56.8 G2: 74.1 Dose of 20 mg/day: G1: 80.1 G2: 82.7 G1/G2: $P = 0.003$ for all comparisons</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: -</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: +</p> <p>Drop-out rates: +</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: -</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Versi et al., 2000 (continued)	<p>Other G1: 2 G2: 0</p> <p>Weight, mean kg: G1: 78.0 G2: 79.0</p> <p>Height, mean cm: G1: 163.7 G2: 165.1</p> <p>Women, %: G1: 88.3 G2: 90.4</p> <p>Parity mean ± SD: NR</p>				

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Wang et al., 2002</p> <p>Country and setting: US, Health care database</p> <p>Enrollment period: January 1991 to June 1995</p> <p>Funding: Pharmacia</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Healthcare database utilization claims analysis (retrospective cohort)</p> <p>Intervention: Urinary antispasmodics including flavoxate, oxybutynin, hyocyanine and hyocyanine sulfate</p> <p>Groups: G1: Prescribed any urinary antispasmodic in interval G2: No prescriptions for urinary antispasmodic agents</p> <p>N in database: G1: 3,898 G2: 10,740</p> <p>N at follow-up: NA</p> <p>Women, %: G1: 75.4 G2: 67.0</p> <p>Age, mean: G1: 78.0 G2: 80.3</p> <p>Race/ethnicity, %: White: G1: 82.9 G2: 82.5 Black: G1: 12.5 G2: 13.0 Other: G1: 4.6 G2: 4.5</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> New Jersey residents in Medicaid/Medicare or Pharmacy Assistance for the Aged and Disabled with comprehensive prescription programs Covered by programs from January 1990 to June 1995 Age ≥ 65 Diagnosis of urinary incontinence Care utilization or prescription for any indication in the six months before and after the index date of the diagnosis of incontinence <p>Exclusion criteria:</p> <ul style="list-style-type: none"> No record of care utilization or prescriptions filled in the study interval 	<p>Charlson comorbidity index, mean score: G1: 1.8 G2: 2.6 <i>P</i> < 0.001</p> <p>Hospital days in prior 180 days, mean: G1: 6.8 G2: 12.5 <i>P</i> < 0.001</p> <p>Physician visits in prior 180 days, mean: G1: 7.3 G2: 5.4 <i>P</i> < 0.001</p> <p>Nursing home days in prior 180 days, mean: G1: 17.0 G2: 36.8 <i>P</i> < 0.001</p> <p>SES: Medicaid, %: G1: 51.5 G2: 65.7 <i>P</i> < 0.001</p> <p>SES: PAAD, %: G1: 48.5 G2: 34.3 <i>P</i> < 0.001</p> <p>Dx of CVD, %: G1: 18.2 G2: 25.2 <i>P</i> < 0.001</p> <p>Dx of CHF, %: G1: 16.9 G2: 25.1 <i>P</i> < 0.001</p> <p>Dx of conduction disturbance, %: G1: 3.0 G2: 3.9 <i>P</i> = 0.008</p>	<p>Adjusted relative risk of ventricular arrhythmia, G1/G2 (95% CI): Urinary antispasmodic use: 1.23 (0.87, 1.75) Nonsedating antihistamine use: 0.95 (0.23, 3.84) Cytochrome P450 3A4 inhibitor use: 1.12 (0.60, 2.08) Concurrent nonsedating antihistamine/cytochrome inhibitor use: 5.47 (1.34, 22.26)</p> <p>Adjusted relative risk of sudden death, G1/G2 (95% CI): Urinary antispasmodic use: 0.70 (0.28, 1.74) Nonsedating antihistamine use: 1.43 (0.20, 10.35) Cytochrome P450 3A4 inhibitor use: 1.80 (0.86, 3.76) Concurrent nonsedating antihistamine/cytochrome inhibitor use: 21.50 (5.23, 88.37)</p>	<p>Quality: Overall quality score: fair INTERNAL VALIDITY: fair Randomization: NA Masking: NA Pt selection criteria: + Loss to followup: NA Drop-out rates: NA Power calculation: - Statistical issues: + EXTERNAL VALIDITY: fair Age: + Baseline OAB status: NR Baseline characteristics: - Length of followup: NA Measurement methods: + Measurement reliability: + Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Wang et al., 2002 (continued)			<p>Dx of supra-ventricular dysrhythmia, %: G1: 7.8 G2: 12.2 <i>P</i> < 0.001</p> <p>Dx of HTN, %: G1: 44.8 G2: 43.8 <i>P</i> = 0.255</p> <p>Dx of Ischemic heart disease, %: G1: 31.0 G2: 36.7 <i>P</i> < 0.001</p> <p>Dx of other CVD, %: G1: 28.0 G2: 35.0 <i>P</i> < 0.001</p> <p>Dx of electrolyte abnormalities, %: G1: 6.4 G2: 11.8 <i>P</i> < 0.001</p>	<p>Incidence rate of ventricular arrhythmia with medication use, (95% CI): Urinary antispasmodics: 1.08 (0.85, 1.37) Nonsedating antihistamines: 1.00 (0.50, 1.99) Cytochrome P450 3A4 inhibitors: 1.02 (0.77, 1.34) Concurrent nonsedating antihistamine/cytochrome inhibitor use: 3.45 (1.11, 10.69)</p> <p>Incidence rate of sudden death with medication use, (95% CI): Urinary antispasmodics: 0.69 (0.29, 1.66) Nonsedating antihistamines: 1.14 (0.16, 8.06) Cytochrome P450 3A4 inhibitors: 1.51 (0.76, 3.02) Concurrent nonsedating antihistamine/cytochrome inhibitor use: 24.35 (6.09, 97.35)</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Wang et al., 2006</p> <p>Country and setting: Taiwan; Academic medical center</p> <p>Enrollment period: July 2004 to November 2005</p> <p>Funding: National Science Council, Taiwan</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: Electric Stimulation (ES) vs Oxybutynin vs Placebo for 12 weeks</p> <p>Groups: G1: ES: intravaginal electrode; biphasic symmetric pulsed current w/ a 10-Hz frequency, 400-millisecond pulse width, 10/5 duty cycle, and varying intensity, 20 min/session, twice weekly G2: Oxybutynin, 2.5 mg 3 times daily G3: Placebo pill 3 times daily</p> <p>N at enrollment: G1: 25 G2: 26 G3: 23</p> <p>N at follow-up: G1: 24 G2: 23 G3: 21</p> <p>Age: NR</p> <p>Race/ethnicity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> Female OAB for ≥ 6 mos Age 16-80 Urgency ≥ 4 times per day <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Use of anticholinergics or TCAs Previous treatment with pelvic-floor muscle training, bladder training Pelvic prolapse repair Pregnancy Neurologic disorders DM Demand cardiac pacemaker IUD use Genital prolapse greater than ICS Stage II PVR>100 mL Overt SI Anti-incontinence surgery UTI 	<p>UUI episodes/day, median (range): G1: 1 (0-2) G2: 0 (0-2) G3: 1 (0-2)</p> <p>Urgency episodes/day, median (range): G1: 11.4 (4.0-16.0) G2: 12.1 (5-18) G3: 9.8 (4.2-15.5)</p> <p>Voids/day, median (range): G1: 12.8 (8.8-16.0) G2: 11.5 (4.3-19.5) G3: 11.5 (6.5-22.8)</p> <p>Nocturia episodes/day, median (range): G1: 1.75 (0-6.5) G2: 0 (0-2.5) G3: 0.65 (0-3.1)</p> <p>Pads/day, median (range): G1: 1 (0-4.1) G2: 0 (0-3) G3: 1 (0-4)</p> <p>Warning time (sec), median (range): G1: 41.5 (8-105) G2: 44 (2-215) G3: 65 (26-265)</p> <p>Max voided volume (mL), median (range): G1: 340 (120-450) G2: 310 (130-800) G3: 350 (120-600)</p>	<p>UUI episodes/day, median (range): G1: 0.5 (0-2) G2: 0 (0-2) G3: 1 (0-2) G1/BL: <i>P</i> = 0.814 G2/BL: <i>P</i> = 0.083 G3/BL: <i>P</i> = 0.854 G1-G3: <i>P</i> = 0.413</p> <p>Urgency episodes/day, median (range): G1: 1.0 (0.0-12.3) G2: 6 (0.5-13) G3: 7.4 (3.9-13.4) G1/BL: <i>P</i> < 0.001 G2/BL: <i>P</i> < 0.001 G3/BL: <i>P</i> = 0.003 G1-G3: <i>P</i> < 0.001</p> <p>Voids/day, median (range): G1: 7.8 (1.8-13.0) G2: 7.4 (2-14) G3: 10 (3.9-13.4) G1/BL: <i>P</i> < 0.001 G2/BL: <i>P</i> < 0.001 G3/BL: <i>P</i> = 0.070 G1-G3: <i>P</i> = 0.002</p> <p>Nocturia episodes/day, median (range): G1: 0 (0-3.0) G2: 0 (0-2.5) G3: 1 (0-3.6) G1/BL: <i>P</i> = 0.001 G2/BL: <i>P</i> = 0.394 G3/BL: <i>P</i> = 0.176 G1-G3: <i>P</i> = 0.002</p> <p>Pads/day, median (range): G1: 0 (0-2) G2: 0 (0-2.5) G3: 1 (0-3) G1/BL: <i>P</i> = 0.010 G2/BL: <i>P</i> = 0.662 G3/BL: <i>P</i> = 0.501 G1-G3: <i>P</i> = 0.012</p>	<p>Quality: Overall quality score: fair INTERNAL VALIDITY: poor Randomization: + Masking: - Pt selection criteria: + Loss to followup: ++ Drop-out rates: + Power calculation: + Statistical issues: - EXTERNAL VALIDITY: good Age: + Baseline OAB status: + Baseline characteristics: ++ Length of followup: + Measurement methods: + Measurement reliability: + Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Wang et al., 2006 (continued)			Total voided volume/day (mL), median (range): G1: 2160 (1010-2950) G2: 2106 (1560-3153) G3: 2305 (1305-3300)	Warning time (sec), median (range): G1: 72 (32-633) G2: 54.5 (138-850) G3: 66.5 (26-219) G1/BL: <i>P</i> = 0.002 G2/BL: <i>P</i> = 0.001 G3/BL: <i>P</i> = 0.532 G1-G3: <i>P</i> < 0.001	
				Max voided volume (mL), median (range): G1: 355 (150-550) G2: 336.5 (138-850) G3: 340 (160-600) G1/BL: <i>P</i> = 0.018 G2/BL: <i>P</i> = 0.004 G3/BL: <i>P</i> = 0.979 G1-G3: <i>P</i> = 0.035	
				Total voided volume/day (mL), median (range): G1: 2270 (1210-3106) G2: 2100 (1619-3200) G3: 2305 (1351-3221) G1/BL: <i>P</i> = 0.024 G2/BL: <i>P</i> = 0.728 G3/BL: <i>P</i> = 0.627 G1-G3: <i>P</i> = 0.050	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Yamaguchi et al., 2007</p> <p>Country and setting: Japan, 155 academic health centers</p> <p>Enrollment period: June 2003 to January 2004</p> <p>Funding: Astellas Pharma Inc.</p> <p>Author industry relationship disclosures: 2 of 12 Astellas (2)</p>	<p>Design: RCT, double-blind, double-dummy, placebo controlled</p> <p>Intervention: solifenacin vs propiverine vs placebo</p> <p>Groups: G1: solifenacin 5 mg qd x 12 wks G2: solifenacin 10 mg qd x 12 wks G3: propiverine 20 mg qd x 12 wks G4: placebo qd x 12 wks</p> <p>N at enrollment: G1: 398 G2: 381 G3: 400 G4: 405</p> <p>N at follow-up: G1: 364 G2: 349 G3: 364 G4: 371</p> <p>Women, n (%): G1: 318 (83.0) G2: 318 (85.7) G3: 321 (83.6) G4: 333 (84.3)</p> <p>Age, mean ± SD, (range): G1: 60.4 ± 13.3 (20.0-89.0) G2: 59.9 ± 13.0 (20.0-86.0) G3: 59.6 ± 13.6 (23.0-94.0) G4: 60.8 ± 12.5 (28.0-90.0)</p> <p>Age ≥ 65, N (%): G1: 161 (42.0) G2: 157 (42.3) G3: 166 (43.2) G4: 163 (41.3)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 20 • OAB sx for ≥ 6 mos • ≥ 8 voids/day • ≥ 3 episodes of urgency or ≥ 3 episodes of urgency incontinence over 72 hours <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • PVR >100 mL • Significant BOO • Urinary retention • SI • Bladder stones • UTI • Interstitial cystitis • Previous or current malignant dz of the pelvic organs • Previous pelvic radiation • Current trt w/ anticholinergic meds • Known or suspected hypersensitivity to anticholinergic meds or lactose 	<p>UUI episodes/day, mean ± SD: G1: 1.95 ± 2.14 G2: 1.89 ± 1.91 G3: 1.82 ± 1.94 G4: 1.67 ± 1.95</p> <p>Urgency episodes, n: G1: 383 G2: 371 G3: 384 G4: 395</p> <p>Urgency episodes/day, mean ± SD: G1: 4.40 ± 3.38 G2: 4.47 ± 3.30 G3: 4.07 ± 3.19 G4: 4.04 ± 3.11</p> <p>Incontinent, n: G1: 274 G2: 270 G3: 295 G4: 283</p> <p>Incontinence type, n (%): Urge: G1: 235 (61.4) G2: 213 (57.4) G3: 218 (56.8) G4: 244 (61.8) Mixed: G1: 130 (33.9) G2: 143 (38.5) G3: 144 (37.5) G4: 135 (34.2) None: G1: 18 (4.7) G2: 15 (4.0) G3: 22 (5.7) G4: 16 (4.1)</p> <p>Incontinence episodes/day, mean ± SD: G1: 2.35 ± 2.45 G2: 2.19 ± 2.04 G3: 2.15 ± 2.31 G4: 1.99 ± 2.11</p>	<p>UUI episodes/day, mean change ± SD: G1: -1.45 ± 1.89 G2: -1.52 ± 1.77 G3: -1.19 ± 2.20 G4: -0.69 ± 2.00 G1/G4: <i>P</i> < 0.001 G2/G4: <i>P</i> < 0.001 G3/G4: <i>P</i> < 0.025</p> <p>Urgency free, n (%): G1: 126 (32.9) G2: 138 (37.2) G3: 128 (33.3) G4: 82 (20.8)</p> <p>Urgency episodes/day, mean change ± SD: G1: -2.41 ± 2.88 G2: -2.78 ± 2.82 G3: -2.30 ± 3.08 G4: -1.28 ± 2.90 G1/G4: <i>P</i> < 0.001 G2/G4: <i>P</i> < 0.001 G2/G3: <i>P</i> = 0.012</p> <p>Continent, n (%): G1: 154 (56.2) G2: 161 (59.6) G3: 165 (55.9) G4: 105 (37.1)</p> <p>Incontinence episodes/day, mean change ± SD: G1: -1.59 ± 2.12 G2: -1.60 ± 1.81 G3: -1.25 ± 2.79 G4: -0.72 ± 1.95 G1/G4: <i>P</i> < 0.001 G2/G4: <i>P</i> < 0.001 G3/G4: <i>P</i> < 0.025</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: -</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: ++</p> <p>Drop-out rates: -</p> <p>Power calculation: +</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating	
Yamaguchi et al., 2007 (continued)	<p>Race/ethnicity: NR</p> <p>Weight (kg), mean ± SD (range): G1: 56.3 ± 9.7 (30.5-123.0) G2: 57.1 ± 10.5 (35.0-117.4) G3: 56.6 ± 10.0 (33.0-103.0) G4: 56.2 ± 9.4 (36.0-90.0)</p>		<p>Voids/day, mean ± SD G1: 11.44 ± 2.94 G2: 11.15 ± 2.76 G3: 11.37 ± 2.71 G4: 11.25 ± 2.73</p> <p>≥ 8 voids/day, n: G1: 383 G2: 371 G3: 384 G4: 395</p> <p>Nocturia episodes/day, mean ± SD: G1: 1.91 ± 1.22 G2: 1.78 ± 1.04 G3: 1.95 ± 1.29 G4: 1.84 ± 1.10</p> <p>Duration of symptoms (years), n (%): ≥ 0.5 to < 1: G1: 46 (12.0) G2: 36 (9.7) G3: 38 (9.9) G4: 39 (9.9) ≥ 1 to < 3: G1: 89 (23.2) G2: 97 (26.1) G3: 85 (22.1) G4: 81 (20.5) ≥ 3 to < 5: G1: 34 (8.9) G2: 45 (12.1) G3: 38 (9.9) G4: 40 (10.1) ≥ 5 to < 10: G1: 24 (6.3) G2: 23 (6.2) G3: 22 (5.7) G4: 37 (9.4) ≥ 10: G1: 23 (6.0) G2: 14 (3.8) G3: 20 (5.2) G4: 24 (6.1) Unknown: G1: 167 (43.6) G2: 36 (9.7) G3: 38 (9.9) G4: 39 (9.9)</p>	<p>Voids/day, mean change ± SD: G1: -1.93 ± 1.97 G2: -2.19 ± 2.09 G3: -1.87 ± 2.70 G4: -0.94 ± 2.29</p> <p>G1/G4: <i>P</i> < 0.001 G2/G4: <i>P</i> < 0.001 G3/G4: <i>P</i> < 0.001 G1/G3: <i>P</i> = NS G2/G3: <i>P</i> = NS</p> <p>≥ 8 voids/day, n (%): G1: 109 (28.5) G2: 137 (36.9) G3: 101 (26.3) G4: 83 (21.0)</p> <p>Nocturia episodes/day, mean change ± SD: G1: -0.41 ± 0.96 G2: -0.46 ± 0.90 G3: -0.43 ± 1.21 G4: -0.30 ± 0.91 G1/G4: <i>P</i> = NS G2/G4: <i>P</i> < 0.025 G3/G4: <i>P</i> = NS</p> <p>KHQ, general health:* G1/G4: <i>P</i> < 0.05 G3/G4: <i>P</i> < 0.05</p> <p>KHQ, incontinence impact:* G1/G4: <i>P</i> < 0.05 G2/G4: <i>P</i> < 0.05 G3/G4: <i>P</i> < 0.05</p> <p>KHQ, role limitations:* G1/G4: <i>P</i> < 0.05 G2/G4: <i>P</i> < 0.05 G3/G4: <i>P</i> < 0.05</p> <p>KHQ, physical limitations:* G1/G4: <i>P</i> < 0.05 G2/G4: <i>P</i> < 0.05 G3/G4: <i>P</i> < 0.05</p>		

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Yamaguchi et al., 2007 (continued)			Volume voided, mean mL ± SD: G1: 153.4 ± 52.91 G2: 154.4 ± 50.21 G3: 151.2 ± 49.38 G4: 152.8 ± 44.28	KHQ, social limitations:* G1/G4: <i>P</i> < 0.05 G2/G4: <i>P</i> < 0.05 G3/G4: <i>P</i> < 0.05 KHQ, personal relationships:* G2/G4: <i>P</i> < 0.05 G3/G4: <i>P</i> < 0.05 KHQ, emotions:* G1/G4: <i>P</i> < 0.05 G2/G4: <i>P</i> < 0.05 G3/G4: <i>P</i> < 0.05 KHQ, sleep/energy:* G1/G4: <i>P</i> < 0.05 G2/G4: <i>P</i> < 0.05 G3/G4: <i>P</i> < 0.05 KHQ, severity:* G1/G4: <i>P</i> < 0.05 G2/G4: <i>P</i> < 0.05 G3/G4: <i>P</i> < 0.05 G2/G3: <i>P</i> < 0.05 Voided volume (mL), mean change ± SD: G1: 35.78 ± 43.49 G2: 43.59 ± 44.52 G3: 36.62 ± 37.99 G4: 11.67 ± 33.74 G1/G4: <i>P</i> < 0.001 G2/G4: <i>P</i> < 0.001 G3/G4: <i>P</i> < 0.001 G2/G3: <i>P</i> = 0.009 Discontinued due to AEs, n (%): G1: 20 (5.1) G2: 26 (6.8) G3: 26 (6.5) G4: 11 (2.7) Dry mouth, n (%): G1: 67 (16.9) G2: 130 (34.1) G3: 103 (25.8) G4: 23 (5.7) G1/G3: <i>P</i> = 0.003 G2/G3: <i>P</i> = 0.012	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Yamaguchi et al., 2007 (continued)				<p>Dry mouth, mild, n (%): G1: 64 (16.2) G2: 124 (32.5) G3: 100 (25.0) G4: 23 (5.7)</p> <p>Dry mouth, moderate, n (%): G1: 3 (0.8) G2: 6 (1.6) G3: 3 (0.8) G4: 1 (0.2)</p> <p>Constipation, n (%): G1: 42 (10.6) G2: 72 (18.9) G3: 45 (11.3) G4: 16 (4.0) G2/G3: <i>P</i> = 0.004</p> <p>Blurred vision, n (%): G1: 7 (1.8) G2: 14 (3.7) G3: 15 (3.8) G4: 8 (2.0)</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Zinner et al., 2002</p> <p>[See evidence table for Van Kerrebroeck et al., 2001]</p> <p>Country and setting: North America (74 centers), Australasia (4 centers), Europe (89 centers)</p> <p>Enrollment period: NR</p> <p>Funding: Pharmacia Corporation</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: Tolterodine ER vs placebo</p> <p>Groups: G1: tolterodine ER 4 mg qd < 65 years of age G2: placebo < 65 years of age G3: tolterodine ER 4 mg qd ≥ 65 years of age G4: placebo ≥ 65 years of age</p> <p>N at enrollment: G1: 293 G2: 285 G3: 214 G4: 223</p> <p>N at follow-up: G1: 257 G2: 246 G3: 193 G4: 194</p> <p>Women, n (%): G1: 255 (87)** G2: 247 (87)** G3: 162 (76) G4: 163 (73)</p> <p>Age, mean ± SD: G1: 51 ± 10.5 G2: 51 ± 10 G3: 74 ± 6 G2: 74 ± 6</p> <p>Race/ethnicity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 18 • Urinary frequency (≥ 8 voids/day) <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Demonstrable SUI • Total daily urine volume > 3 L • Contra-indications to antimuscarinic treatment • Hepatic or renal disease • UTI • Interstitial cystitis • Hematuria • BOO • Current electrostimulation or bladder training therapy • Indwelling catheter or intermittent self-catheterization • Pregnant or nursing • Women not using reliable contraception • Being treated for OAB with other anticholinergic drugs or drugs that inhibit cytochrome P450 3A4 isoenzymes • Estrogen therapy < 2 months • Treatment w/ investigational drug < 2 months 	<p>Not able to hold urine upon experiencing urgency, %: G1: 24.9 G2: 29.1 G3: 33.6 G4: 34.5</p> <p>Able to complete tasks before toilet visit in response to urgency, %: G1: 6.5 G2: 7.7 G3: 5.1 G4: 4.9</p> <p>Incontinence episodes/week, mean ± SD: G1: 21.4 ± 22.1 G2: 23.2 ± 22.0 G3: 23.2 ± 22.7 G4: 23.4 ± 18.9</p> <p>Voids/day, mean ± SD: G1: 11.0 ± 3.9 G2: 11.4 ± 4.2 G3: 10.8 ± 4.5 G4: 11.0 ± 3.2</p> <p>Voided volume (mL), mean ± SD: G1: 140 ± 41 G2: 137 ± 45 G3: 141 ± 45 G4: 134 ± 39</p> <p>Previous drug therapy, n (%): G1: 148 (50.5) G2: 146 (51.2) G3: 121 (56.5) G4: 117 (52.5)</p> <p>Percentage with poor efficacy, %: G1: 74 (50.0)** G2: 60 (41.1)** G3: 40 (33.1) G4: 46 (39.3)</p>	<p>Not able to hold urine upon experiencing urgency, %: G1: 11.3 G2: 21.1 G3: 15.9 G4: 25.6 G1/G2: <i>P</i> = 0.003 G3/G4: <i>P</i> = 0.007</p> <p>No age-related difference</p> <p>Able to complete tasks before toilet visit in response to urgency, %: G1: 32.8 G2: 16.8 G3: 26.2 G4: 14.8 G1/G2: <i>P</i> = 0.001 G3/G4: <i>P</i> = 0.003</p> <p>No age-related difference</p> <p>Incontinence episodes/week, mean ± SD G1: -12.0 ± 17.6 G2: -7.4 ± 15.6 G3: -11.5 ± 18.2 G4: -6.3 ± 15.0 G1/G2: <i>P</i> = 0.001 G3/G4: <i>P</i> < 0.001</p> <p>No age-related difference</p> <p>Voids/day, mean ± SD G1: -2.0 ± 3.1 G2: -1.4 ± 3.1 G3: -1.4 ± 3.7 G4: -0.9 ± 2.6 G1/G2: <i>P</i> = 0.26 G3/G4: <i>P</i> = 0.92</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: +</p> <p>Drop-out rates: NR</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: -</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Zinner et al., 2002 (continued)				<p>Perception of bladder condition, improved, %: G1: 60.1 G2: 51.9 G3: 54.2 G4: 31.4 G1/G2: $P < 0.05$ G3/G4: $P < 0.0001$</p> <p>Perception of bladder condition, no change, %: G1: 32.4 G2: 38.9 G3: 38.3 G4: 51.1 G4/G2: $P < 0.0001$</p> <p>Perception of bladder condition, deterioration, %: G1: 7.2 G2: 9.1 G3: 7.5 G4: 17.5</p> <p>Treatment beneficial, %: G1: 78.3 G2: 58.3 G3: 69.8 G4: 46.9 G1/G2: $P = 0.001$ G3/G4: $P = 0.001$ No age-related difference</p> <p>Voided volume (mL), mean \pm SD: G1: 35 \pm 53 G2: 13 \pm 41 G3: 33 \pm 47 G4: 16 \pm 41 G1/G2: $P < 0.001$ G3/G4: $P < 0.001$ No age-related difference</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Zinner et al., 2002 (continued)				Adverse events, %: G1: 50.7 G2: 50.5 G3: 54.2 G4: 46.0 Dry mouth, severe, % G1-G2: 1.7 G3-G4: 1.9 Dry mouth, moderate, % G1-G2: 7.6 G3-G4: 6.5 Dry mouth, mild, % G1-G2: 13.4 G3-G4: 15.9 No dry mouth, % G1-G2: 77.3 G3-G4: 75.7	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Zinner et al., 2004</p> <p>Country and setting: US, 51 sites</p> <p>Enrollment period: NR</p> <p>Funding: Indevus Corporation</p> <p>Author industry relationship disclosures: 4 of 5 Alza (1) Indevus (4) Kyowa (1) Lilly (2) Merck (1) Ortho-McNeil (1) Pfizer (2) Pharma (1) Pharmacia (1) Schwarz (1) Watson (2) Yamanouchi (1)</p>	<p>Design: RCT with 2 wk washout period</p> <p>Intervention: Tropium chloride vs placebo</p> <p>Groups: G1: Tropium chloride 20 mg b.i.d. G2: placebo</p> <p>N at enrollment: G1: 262 G2: 261</p> <p>N at follow-up: G1: 256 G2: 256</p> <p>Women, n (%): G1: 203 (77.5) G2: 186 (71.3)</p> <p>Age, mean ± SD: G1: 63 ± 0.8 G2: 61.5 ± 0.8</p> <p>Race/ethnicity, n (%): White: G1: 222 (84.7) G2: 225 (86.2) Black: G1: 26 (9.9) G2: 20 (7.7) Hispanic: G1: 10 (3.8) G2: 13 (5.0) Other: G1: 4 (1.6) G2: 3 (1.1)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥18 years • OAB symptoms ≥ 6 mos • Urinary urgency • ≥ 70 voids per week • ≥ 7 UUI episodes/week <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Stress predominant MUI • Insensate or overflow • Neurogenic bladder disorders • Significant renal disease • Hematuria hematuria and UTI at washout or ≥ twice in the prior year • BOO with PVR ≥ 100 mL • Concurrent use of any anticholinergic drug or other drug therapy for OAB within 21 days before randomization • Bladder surgery ≤ 6 mos • Bladder cancer • Interstitial cystitis • PSA ≥ 10 ng/mL • Diuretic use, estrogen therapy and nonmedical bladder therapy that was not part of a stable, long-term program. 	<p>UUI episodes/day, mean: G1: 3.9 G2: 4.3</p> <p>Urgency episodes/day, mean: G1: 11.29 G2: 11.72</p> <p>Urgency severity score, mean: G1: 1.8 G2: 1.8</p> <p>Voids/day, mean: G1: 12.7 G2: 12.9</p> <p>Daytime voids/day, mean: G1: 10.6 G2: 10.9</p> <p>Nocturia episodes/day, mean: G1: 2.1 G2: 2.0</p> <p>Voided volume (mL), mean: G1: 155.1 G2: 156.1</p> <p>IIQ, mean score (SE): G1: 183.5 (5.6) G2: 195.4 (5.6)</p> <p>IIQ,, women only, mean score (SE): G1: 191.2 (6.3) G2: 201.9 (6.6)</p> <p>Incontinence Impact Questionnaire, travel subscale, score (SE): G1: 52.4 (1.7) G2: 55.6 (1.7)</p>	<p>UUI episodes/day, week 1, mean % change: G1: -40.1 G2: -27.5 G1/G2: $P \leq 0.001$</p> <p>UUI episodes/day, week 4, mean % change: G1: -58.5 G2: -40.9 G1/G2: $P \leq 0.0001$</p> <p>UUI episodes/day, week 12, mean % change: G1: -59.0 G2: -44.2 G1/G2: $P \leq 0.0001$</p> <p>Urgency episodes/day, week 1, mean change: G1: -1.13 G2: -1.08</p> <p>Urgency episodes/day, week 4, mean change: G1: -2.10 G2: -1.00 G1/G2: $P \leq 0.0001$</p> <p>Urgency episodes/day, week 12, mean change: G1: -2.30 G2: -1.08 G1/G2: $P \leq 0.0001$</p> <p>Urgency severity score, week 1, mean change: G1: -0.11 G2: -0.01 G1/G2: $P \leq 0.01$</p>	<p>Quality: Overall quality score: fair INTERNAL VALIDITY: fair Randomization: + Masking: - Pt selection criteria: + Loss to followup: ++ Drop-out rates: + Power calculation: - Statistical issues: + EXTERNAL VALIDITY: good Age: + Baseline OAB status: + Baseline characteristics: ++ Length of followup: + Measurement methods: + Measurement reliability: + Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Zinner et al., 2004 (continued)			IIQ, social relationships, score (SE): G1: 37.8 (1.5) G2: 40.3 (1.5)	Urgency severity score, week 4, mean change: G1: -0.18 G2: -0.06 G1/G2: $P \leq 0.01$	
			IIQ, emotional health, mean score (SE): G1: 47.1 (1.6) G2: 49.6 (1.6)	Urgency severity score, week 12, mean change: G1: -0.22 G2: -0.04 G1/G2: $P \leq 0.001$	
			IIQ,, physical activity, mean score (SE): G1: 46.1 (1.6) G2: 50.2 (1.6)	Voids/day, week 1, mean change: G1: -1.18 G2: -0.81 G1/G2: $P \leq 0.05$	
			Prior OAB med, n (%): G1: 135 (51.5) G2: 142 (54.5)	Voids/day, week 4, mean change: G1: -2.20 G2: -1.07 G1/G2: $P \leq 0.0001$	
				Voids/day, week 12, mean change: G1: -2.37 G2: -1.29 G1/G2: $P \leq 0.0001$	
				Daytime voids/ day, week 1, mean change: G1: -1.00 G2: -0.68 G1/G2: $P \leq 0.05$	
				Daytime voids/ day, week 4, mean change: G1: -1.77 G2: -0.89 G1/G2: $P \leq 0.0001$	
				Daytime voids/ day, week 12, mean change: G1: -1.90 G2: -0.98 G1/G2: $P \leq 0.0001$	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Zinner et al., 2004 (continued)				Nocturia episodes/day, week 1, mean change: G1: -0.18 G2: -0.15	
				Nocturia episodes/day, week 4, mean change: G1: -0.43 G2: -0.17 G1/G2: $P \leq 0.001$	
				Nocturia episodes/day, week 12, mean change: G1: -0.47 G2: -0.29 G1/G2: $P \leq 0.05$	
				Voided volume (mL), week 1, mean change: G1: 19.9 G2: 6.6 G1/G2: $P \leq 0.0001$	
				Voided volume (mL), week 4, mean change: G1: 30.0 G2: 8.5 G1/G2: $P \leq 0.0001$	
				Voided volume (mL), week 12, mean change: G1: 32.1 G2: 7.7 G1/G2: $P \leq 0.0001$	
				IIQ, week 12, LS mean change (SE): G1: -54.0 (5.6) G2: -36.0 (5.6) G1/G2: $P \leq 0.05$	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Zinner et al., 2004 (continued)				<p>IIQ, women only, week 12, LS mean change (SE): G1: -59.1 (6.6) G2: -35.7 (6.9) G1/G2: $P \leq 0.05$</p> <p>IIQ, travel subscale, week 12, LS mean change (SE): G1: -14.9 (1.7) G2: -9.9 (1.7) G1/G2: $P < \leq 0.05$</p> <p>IIQ, social relationships, week 12, LS mean change (SE): G1: -10.8 (1.4) G2: -6.3 (1.4) G1/G2: $P \leq 0.05$</p> <p>IIQ, emotional health, 12 weeks, LS mean change (SE): G1: 14.1 (1.5) G2: -9.2 (1.5) G1/G2: $P \leq 0.05$</p> <p>IIQ, physical activity, 12 weeks, LS mean change (SE): G1: -13.5 (1.7) G2: -11.0 (1.7)</p> <p>Dry mouth, n (%): G1: 57 (21.8) G2: 17 (6.5)</p> <p>Constipation, n (%): G1: 25 (9.5) G2: 10 (3.8)</p> <p>Headache, n (%): G1: 17 (6.5) G2: 12 (4.6)</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Zinner et al., 2004 (continued)				Diarrhea, n (%): G1: 8 (3.1) G2: 14 (5.4) Abdominal pain, n (%): G1: 8 (3.1) G2: 3 (1.1) Discontinuation due to AE, %: G1: 8.8 G2: 5.7	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Zinner et al., 2005</p> <p>Country and setting: US, Specialty treatment center</p> <p>Enrollment period: NR</p> <p>Funding: Pfizer Novartis Pharma Thomson ACUMED</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Randomized double-blind placebo-controlled four way crossover</p> <p>Intervention: Darifenacin 15 mg qd vs darifenacin 30 mg qd vs. Oxybutynin vs. placebo</p> <p>Groups:* G1: Darifenacin 15mg daily G2: Darifenacin 30mg daily G3: Oxybutynin 5 mg t.i.d. G4: placebo</p> <p>N at enrollment: G1-G4: 76</p> <p>N included, tolerability: G1-G4: 61</p> <p>N included, efficacy: G1-G4: 58</p> <p>Women, n (%): G1-G4: 71 (93.4)</p> <p>Age, mean (range): G1-G4: 59.9 (33-84)</p> <p>Race/ethnicity: NR</p> <p>Weight (kg), mean (range): G1-G4: 75.7 (42-157)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age 18-85 • ≥ 4 UUI episodes/week • ≥ 8 voids/day (from 14 day run in placebo voiding diary) <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Neurogenic bladder • SUI • Contraindication to antimuscarinic treatment • UTI • Bladder symptoms • Bladder stones • BOO • Concomitant disease • Concurrent bladder training • Concurrent thyroid or HRT • Other bladder meds • Pregnant or breastfeeding 	<p>Urgency episodes/day, mean \pm SD: G1-G 4: 9.3 \pm 3.4</p> <p>Urgency severity, mean \pm SD: G1-G4: 2.0 \pm 0.4</p> <p>Incontinence episodes/week, mean \pm SD: G1-G4: 20.4 \pm 17.7</p> <p>Voids/ day, mean \pm SD: G1-G4: 10.4 \pm 3.0</p>	<p>Urgency episodes/day, mean: G1: 7.95 G2: 7.59 G3: 8.12 G4: 8.71 G1-G3/G4: $P < 0.05$</p> <p>Urgency severity, mean: G1: 1.93 G2: 1.84 G3: 1.89 G4: 2.03 G1-G3/G4: $P < 0.05$</p> <p>Incontinence episodes/week, mean: G1: 10.93 G2: 8.82 G3: 9.45 G4: 14.64 G1-G3/G4: $P < 0.05$</p> <p>Voids/day, mean \pm SD: G1: 9.33 G2: 8.85 G3: 9.24 G4: 9.62 G2/G4: $P < 0.05$</p> <p>Dry mouth, (%): G1: 13.1 G2: 34.4 G3: 36.1 G4: 4.9 G1/G3: $P < 0.05$ G2/G3: $P < 0.05$ G2/G4: $P < 0.05$ G3/G4: $P < 0.05$</p> <p>Constipation, (%): G1: 9.8 G2: 21.3 G3: 8.2 G4: 3.3 G2/G3: $P < 0.05$ G2/G4: $P < 0.05$</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: -</p> <p>Drop-out rates: NR</p> <p>Power calculation: +</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: -, NR</p> <p>Baseline OAB status: NR</p> <p>Baseline characteristics: -</p> <p>Length of followup: -</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Zinner et al., 2005 (continued)				Blurred vision, (%): G1: 0 G2: 0 G3: 3.3 G4: 0 Dizziness, (%): G1: 0 G2: 0 G3: 1.6 G4: 0	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Zinner et al., 2006</p> <p>Country and setting: NR, Multicenter</p> <p>Enrollment period: NR</p> <p>Funding: Novartis</p> <p>Author industry relationship disclosures: 5 of 6 Alza (1) Eli Lilly (2) Astellas (1) GSK (1) Indevus (1) Madaus (1) Novartis (5) Pfizer (2) Sanofi (1) Watson (2) Yamanouchi (1)</p>	<p>Design: RCT</p> <p>Intervention: Darifenacin 15 mg controlled release once daily vs. placebo</p> <p>Groups: G1: Darifenacin 15 mg controlled release once daily G2: Placebo</p> <p>N at enrollment: G1: 214 G2: 225</p> <p>N at follow-up: G1: 185 G2: 188</p> <p>Women, n (%): G1: 185 (86.4) G2: 198 (88.0)</p> <p>Age, mean (range): G1: 59.1 (20-93) G2: 59.1 (18-89)</p> <p>Race/ethnicity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> Adults Age ≥18 History of OAB for ≥ 6 mos ≥ 1 urge incontinence episodes/day ≥ 8 voids/day ≥ 4 urgency episodes/day Mean warning time of ≤ 15 min during 12 hrs <p>Exclusion criteria:</p> <ul style="list-style-type: none"> SUI Marked cystocele Pelvic prolapse Use of anticholinergics, antispasmodics, cholinergic agonists, potent cytochrome P450 3A4 inhibitors, or opioids 2 wks prior to screening visit Contraindications to drugs that cause significant constipation, clinically significant bladder outlet obstruction Intention to start bladder training program In-dwelling catheter Intermittent self-catheterization 	<p>UUI episodes/wk, median: G1: 18.73* G2: 21.00^</p> <p>Urgency episodes/wk, median: G1: 82.60* G2: 85.40^</p> <p>Warning time (min), mean: G1: 4.38* G2: 4.74*</p> <p>Warning time (min), median: G1: 2.57* G2: 3.04*</p> <p>Void/day, median: G1: 11.00* G2: 11.20^</p> <p>Duration of OAB (years), mean (range): G1: 7.8 (0, 45) G2: 8.3 (0, 58)</p> <p>Previous OAB therapy, n (%): G1: 112 (52.3) G2: 124 (55.1)</p> <p>Voided volume (mL), median: G1: 153.50* G2: 144.62^</p>	<p>UUI episodes/wk, wk 2, median change: G1: -9.1 G2: -7.0 <i>P</i> < 0.01</p> <p>UUI episodes/wk, wk 6, median change: G1: -11.2 G2: -8.4 <i>P</i> = 0.031</p> <p>UUI episodes/wk, wk 12, median change: G1: -12.60 G2: -9.80 <i>P</i> = 0.035</p> <p>Urgency episodes/wk, wk 12, median change: G1: -18.20 G2: -15.58 <i>P</i> = 0.18</p> <p>Urgency-free time, (min), median: G1: 34.3 G2: 25.0 <i>P</i> = 0.003</p> <p>Warning time, median % change: G1: 41.8 G2: 18.4</p> <p>Void/day, wk 12, median change: G1: -2.20 G2: -1.80 <i>P</i> = 0.176</p> <p>Voided volume (mL), wk 12, mean change: G1: 22.62 G2: 11.31 <i>P</i> = 0.002</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: - Masking: + Pt selection criteria: + Loss to followup: + Drop-out rates: - Power calculation: + Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: + Baseline OAB status: + Baseline characteristics: ++ Length of followup: + Measurement methods: + Measurement reliability: + Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Zinner et al., 2006 (continued)				<p>Total OAB-q score, wk 6, mean change: G1: 24.5 G2: 18.7 <i>P</i> = 0.035</p> <p>Total OAB-q score, wk 12, mean change: G1: 26.4G2: 19.1 <i>P</i> < 0.001</p> <p>Incontinence Impact, wk 12, mean change: G1: -24.7 G2: -17.8 <i>P</i> = 0.022</p> <p>Severity measures, wk 12, mean change: G1: -24.3 G2: -15.6 <i>P</i> < 0.001</p> <p>Adverse events, %: G1: 63.6 G2: 48.9</p> <p>Discontinuation, n (%): G1: 17 (7.9) G2: 9 (4)</p> <p>Dry mouth and constipation, n (%): G1: 29.0 (17.8) G2: 5.8 (4.9)</p> <p>Discontinuation due to dry mouth, %: G1: 2.3 G2: 0</p> <p>Discontinuation due to constipation, %: G1: 2.3 G2: 0</p> <p>Headaches, %: G1: 6.1 G2: 2.2</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Zinner et al., 2006 (continued)				Cardiac AEs, %: G1: 0.5 G2: 0.9	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Zinner et al. 2008</p> <p>Country and setting: US, 89 sites</p> <p>Enrollment period: NR</p> <p>Funding: Astellas Pharma US, GlaxoSmithKline</p> <p>Author industry relationship disclosures: 5 of 5 Astellas (5) Esprit (2) Watson (3) Allergan (2) Medtronics (1) GlaxoSmithKline (3) Indevus (1)</p>	<p>Design: open label, non-comparative, flexible-dosing</p> <p>Intervention: solifenacin 5 or 10 mg qd for 12 wks after washout from tolterodine, with dose adjustments at wk 4 and wk 8</p> <p>Groups: NA</p> <p>N at enrollment: 440</p> <p>N at follow-up: 390</p> <p>Age, yrs ± SD: 61.5 ± 13.69</p> <p>Age < 65, n (%): 246 (55.9)</p> <p>Age ≥ 65-74, n (%): 108 (24.5)</p> <p>Age ≥ 75, n (%): 86 (19.5)</p> <p>Race/ethnicity, n (%): Caucasian 391 (88.9) Black or African 37 (8.4) Asian 1 (0.2) Other 11 (2.5)</p> <p>Women, N (%): 388 (88.2)</p> <p>Women < 65 yrs, n (%): 227 (58.5)</p> <p>Women Age ≥ 65-74, n (%): 92 (23.7)</p> <p>Women Age ≥ 75, n (%): 69 (17.8)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 18 • Symptoms of OAB ≥ 3 mos • Ambulatory and able to use toilet without difficulty • Treated with tolterodine ER 4 mg/day at least 4 wks preceding study but failed to achieve satisfactory improvement in urgency episodes • ≥3 urgency episodes/24 hrs documented in 3-day voiding diary while receiving tolterodine ER 4 mg/d • Prior drug or other treatment for OAB with ≥ 4 wks washout before screening <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Prior treatment with darifenacin • SUI • Stress dominant MUI • UTI • Interstitial cystitis • Bladder stones • Outflow obstruction due to benign prostatic hyperplasia • Uncontrolled narrow-angle glaucoma • Urinary or gastric retention • Severe renal or hepatic impairment 	<p>Urgency episodes/day, mean: 6.03</p> <p>Voids/day, mean: 10.59</p> <p>Incontinence episodes/day, mean: 3.04</p> <p>Nocturia episodes/day, mean: 1.83</p> <p>Nocturnal voids/day, mean: 2.32</p> <p>Pads or diapers/week, mean: 10.7</p>	<p>Urgency episodes/day, mean change (95% CI): -3.41 (-3.79, -3.04)*</p> <p>Voids/day, mean change (95% CI): -1.57 (-1.86, -1.28)*</p> <p>Incontinence episodes/day, mean change (95% CI): -1.86 (-2.19, -1.53)*</p> <p>Nocturia episodes/day, mean change (95% CI): -0.72 (-0.84, -0.59)*</p> <p>Nocturnal voids/day, mean change (95% CI): -0.79 (-0.95, -0.62)*</p> <p>Pads or diapers/week, mean change: 7.9 <i>P</i> = 0.0009</p> <p>Improvement in HRQL, mean (95% CI): 21.1 (19.0, 23.1)*</p> <p>Change in symptom bother score, mean (95% CI):[^] -27.4 (-29.7, -25.1)*</p> <p>Change in coping, score, mean (95% CI)[^]: 23.1 (20.6, 25.6)*</p> <p>Change in concern score, mean (95% CI)[^]: 25.2 (22.8, 27.5)*</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: NA</p> <p>Masking: NA</p> <p>Pt selection criteria: +</p> <p>Loss to followup: +</p> <p>Drop-out rates: NR</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Zinner et al. 2008 (continued)	<p>Height, mean cm ± SD: 164.1 ± 8.39</p> <p>Weight, mean kg ± SD: 81.1 ± 20.56</p> <p>BMI, mean kg/m² ± SD: 30.1 ± 7.12</p>	<ul style="list-style-type: none"> • Chronic severe constipation • Gastrointestinal obstructive disease • Bladder cancer • Pregnant or breast-feeding • Not using a reliable birth control method • Known hypersensitivity to solifenacin or to anticholinergic medicines • Participating in any clinical trial of any investigational drug within 30 days 		<p>Change in sleep score, mean (95% CI)[^]: 21.9 (19.5, 24.3)*</p> <p>Change in social interaction score, mean (95% CI)[^]: 11.1 (9.4, 12.9)*</p> <p>Change in % work time missed, mean ± SD[†]: -1.84 ± 0.57 <i>P</i> = 0.0017</p> <p>Change in % impairment while working, mean ± SD[†]: -11.64 ± 1.64 <i>P</i> < 0.0001</p> <p>Change in % overall work impairment, mean ± SD[†]: -12.14 ± 1.79 <i>P</i> < 0.0001</p> <p>Change in % activity impairment, mean ± SD[†]: -13.21 ± 1.31 <i>P</i> < 0.001</p> <p>Change in office visits, mean ± SD: -1.00 ± 0.08 <i>P</i> < 0.001</p> <p>Change in UTI, mean ± SD: -0.12 ± 0.03 <i>P</i> < 0.001</p> <p>Change in skin rash episodes, mean ± SD: -0.46 ± 0.34 <i>P</i> = 0.1801</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Zinner et al. 2008 (continued)				<p>Change in number of falls, mean ± SD: 0.04 ± 0.07 P = 0.5561</p> <p>Change in number of pads/diapers/ wk, mean ± SD: -2.79 ± 0.83 P = 0.0009</p>	

Evidence Table 3. KQ2 Procedural and surgical treatments

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Enzelsberger et al. 1995</p> <p>Country and setting: Austria, Academic gynecology/ urology outpatient clinic</p> <p>Enrollment period: NR</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: Instillation of oxybutynin or sterile water into the bladder via a disposable catheter, administered over a period of 12 days</p> <p>Groups: G1: instillation of 20 mg oxybutynin in 40 mL sterile water G2: instillation of 40 mL sterile water</p> <p>N at enrollment: G1: 26 G2: 26</p> <p>N at follow-up: G1: 23 G2: 20</p> <p>Women, %: 100</p> <p>Age, median (range): G1: 61 (55, 66) G2: 59 (56, 64)</p> <p>Race/ethnicity: NR</p> <p>Weight (kg), median (range): G1: 68 (58, 74) G2: 70 (59, 77)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Women • Presenting complaint of frequency (more than 5 times/ 12hr), nocturia (more than twice per night), and urgency • Confirmed idiopathic detrusor instability <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Urodynamically assessed genuine stress incontinence • Neurologic disorders 	<p>Daytime voids /day, median (range): G1: 7.5 (5, 15) G2: 8.2 (6, 12)</p> <p>Nocturia episodes/day, median (range): G1: 5.1 (3, 7) G2: 4.6 (3, 8)</p> <p>Voided volume (mL), first desire to void, median (range): G1: 95 (30, 140) G2: 90 (25, 170)</p> <p>Cystometric capacity (mL), median (range): G1: 205 (50, 340) G2: 195 (40, 260)</p> <p>Max pressure rise during filling (cmH₂O), median (range): G1: 16 (10, 22) G2: 14 (9, 19)</p> <p>Max contraction (cmH₂O), median (range): G1: 32 (20, 44) G2: 29 (18, 42)</p> <p>Peak flow rate (mL/s), median (range): G1: 27 (15, 45) G2: 30 (17, 40)</p> <p>Residual urinary volume (mL), median (range): G1: 40 (10, 50) G2: 35 (12, 50)</p> <p>Duration of symptoms (months), median (range): G1: 7 (2, 9) G2: 8 (3, 9)</p>	<p>Daytime voids/day, median (range): G1: 4.0 (2, 7) G2: 6.9 (5, 10) <i>P</i> < 0.05</p> <p>Nocturia episodes/day, median (range): G1: 1.8 (1, 3) G2: 3.5 (3, 6) <i>P</i> < 0.05</p> <p>Voided volume (mL), first desire to void, median (range): G1: 150 (80, 180) G2: 110 (30, 140) <i>P</i> < 0.05</p> <p>Cystometric capacity (mL), median (range): G1: 310 (100, 390) G2: 207 (43, 240) <i>P</i> < 0.05</p> <p>Max pressure rise during filling (cmH₂O), median (range): G1: 8 (5, 10) G2: 11 (6, 12) <i>P</i> < 0.05</p> <p>Max contraction (cmH₂O), median (range): G1: 14 (10, 20) G2: 24 (11, 42) <i>P</i> < 0.05</p> <p>Peak flow rate (mL/s), median (range): G1: 22 (15, 40) G2: 32 (15, 45)</p> <p>Residual urinary volume (mL), median (range): G1: 81 (40, 140) G2: 38 (15, 60) <i>P</i> < 0.05</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: +</p> <p>Drop-out rates: -</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: -</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 3. KQ2 Procedural and surgical treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Enzelsberger et al. 1995 (continued)			Previous operations for UI, n: G1: 11 G2: 10	Dry mouth, n (%): G1: 4 (17) G2: 2 (10) Blurred vision, n (%): G1: 2 (8) G2: 1 (5) Nausea, n (%): G1: 0 G2: 0 Constipation, n (%): G1: 2 (8) G2: 3 (15) Dizziness, n (%): G1: 3 (13) G2: 0 Headache, n (%): G1: 1 (4) G2: 1 (5) UTI, n (%): G1: 4 (17) G2: 2 (10)	

Evidence Table 3. KQ2 Procedural and surgical treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Fossberg et al., 1990</p> <p>Country and setting: Norway, Denmark, Finland, Sweden, Academic medical center</p> <p>Enrollment period: NR</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Prospective case series</p> <p>Intervention: Treatment with a stimulation device including an anal and a vaginal plug electrode; stimulation provided at cyclic sweep within 5-10 Hz, administered 12 times for 20 minutes each (frequency of treatment not indicated)</p> <p>Groups: NA</p> <p>N at enrollment: 91</p> <p>N at follow-up: 74</p> <p>Women, %: 88</p> <p>Age, mean (range): 53 (20-78)</p> <p>Race/ethnicity: NR</p> <p>Menopausal, %: 50</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Cystometrically proven detrusor instability, frequency and UUI <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • UTI • Drugs affecting the lower urinary tract • Discontinued anticholinergic drugs < 6 weeks before study 	<p>Voids/day, mean ± SD (range): 9.1 ± 3.5 (4-20)</p> <p>Nocturia episodes/day, mean ± SD (range): 1.6 ± 1.1 (0-6)</p> <p>Bladder volume (mL), mean ± SD (range): 169 ± 68 (50-425)</p> <p>Bladder volume (mL), first sensation, mean ± SD (range): 139 ± 81 (33-300)</p> <p>Max cystometric capacity (mL), mean ± SD (range): 270 ± 155 (25-750)</p> <p>Previously treated with drugs for UUI, % 67</p>	<p>Voids/day, before last treatment, mean ± SD (range): 7.9 ± 2.9 (3-18) <i>P</i> = 0.001</p> <p>Voids/day, 6 weeks after last treatment, mean ± SD (range): 8.0 ± 2.8 (3-19) <i>P</i> = 0.003</p> <p>Nocturia epsides/day, before last treatment, mean ± SD (range): 1.2 ± 1.2 (0-6) <i>P</i> = 0.002</p> <p>Nocturia epsides/day, 6 weeks after last treatment, mean ± SD (range): 1.1 ± 1.1 (0-4) <i>P</i> < 0.001</p> <p>Bladder volume (mL), before last treatment, mean ± SD (range): 189 ± 82 (50-490) <i>P</i> = 0.002</p> <p>Bladder volume (mL), 6 weeks after last treatment, mean ± SD (range): 194 ± 93 (50-500) <i>P</i> < 0.001</p> <p>Bladder volume (mL), first sensation, before last treatment, mean ± SD (range): 164 ± 93 (35-410) <i>P</i> = 0.008</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: NA</p> <p>Masking: NA</p> <p>Pt selection criteria: +</p> <p>Loss to followup: +</p> <p>Drop-out rates: -</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: ++</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 3. KQ2 Procedural and surgical treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Fossberg et al., 1990 (continued)				<p>Bladder volume (mL), first sensation, 6 weeks after last treatment, mean \pm SD (range): 175 ± 114 (25-470) $P = 0.003$</p> <p>Max cystometric capacity (mL), before last treatment, mean \pm SD (range): 307 ± 159 (50-725) $P = 0.009$</p> <p>Max cystometric capacity (mL), before last treatment, mean \pm SD (range): 299 ± 174 (25-690) $P = 0.12$</p> <p>Patient evaluation of effect, before last treatment, n: Cured: 6 Improved: 45 Unchanged: 23 Worse: 0 $P < 0.001$</p> <p>Patient evaluation of effect, 6 weeks after last treatment, n: Cured: 8 Improved: 32 Unchanged: 34 Worse: 0 $P < 0.001$</p> <p>Adverse effects, n:* 11</p>	

Data analysis indicated no differences between males and females in any data recorded

Evidence Table 3. KQ2 Procedural and surgical treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Groenendijk et al. 2007</p> <p>[See evidence table for Hassouna et al. 2000]</p> <p>Country and setting: US, Canada, and Europe;16 centers</p> <p>Enrollment period: NR</p> <p>Funding: Medtronic Inc.</p> <p>Author conflict of interest: Medtronic (6)</p>	<p>Design: Cohort</p> <p>Intervention: implantable Interstim system stimulating the nerve ramus in preoperatively tested sacral nerve foramen</p> <p>Groups: G1: urge incontinence and detrusor overactivity (DO) G2: urge incontinence without detrusor overactivity</p> <p>N at enrollment: G1: 67 G2: 44</p> <p>N at follow-up: G1: 51 G2: 32</p> <p>Age: NR</p> <p>Race/ethnicity: NR</p> <p>Women, n: G1: 59 G2: 40</p> <p>Follow-up: 6 months</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • >16 years of age • refractory to standard medical therapy • Bladder capacity of at least 100mL • -Normal upper urinary tract <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Treatable etiologies for urinary/frequency symptoms • Neurologic conditions • -Primary stress incontinence • Pelvic pain symptoms 	<p>First sensation of filling, mL ± SD: Overall: 82.8 ± 64.7 G1: NR G2: 122.2 ±78.8</p> <p>Bladder volume at first detrusor contraction, mL ± SD: Overall: 104.7 ± 111.0</p> <p>Bladder capacity, mL ± SD: Overall: 254 ± 138 G1: NR G2: 313 ± 117</p> <p>Detrusor pressure at capacity, cm water ± SD: Overall: 27.7 ± 21.6</p>	<p>First sensation of filling, mL ± SD: Overall: 167.4 ± 109.3, p<0.001 G1: NR G2: 192.4 ± 118.4 p=0.001</p> <p>Bladder volume at first detrusor contraction, mL ± SD: Overall:133.7 ± 125.9, p =0.30</p> <p>Bladder capacity, mL ± SD: Overall: 328 ± 148, p=0.001 G1: NR G2: 365 ± 115, p=0.02</p> <p>Detrusor pressure at capacity, cm water ± SD: Overall: 17.7 ± 16.5</p> <p>DO at 6 months, n (%): G1: -25 (49%) -stimulation was clinically successful in 19 (40%) -failed in 3 (6%) - 26 (51%) were negative for DO -stimulation was clinically successful in 20 (42%) - failed in 6 (12%) G2: -3 (9%)were positive for DO -stimulation was clinically successful in 2 (7%) - failed in 1 (4%) - 29 remained negative for DO -stimulation was successful in 23 (82%) -failed in 2 (7%)</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: NA</p> <p>Masking: NA</p> <p>Pt selection criteria: +</p> <p>Loss to followup: -</p> <p>Drop-out rates: NR</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: -, NR</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: -</p> <p>Length of followup: ++</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 3. KQ2 Procedural and surgical treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Hassouna et al., 2000</p> <p>Country and setting: US, Canada, and Europe, 12centers</p> <p>Enrollment period: Ended June 1998</p> <p>Funding: Medtronic Inc.</p> <p>Author conflict of interest: Medtronic</p>	<p>Design: Prospective cohort</p> <p>Intervention: implantable Interstim system stimulating the nerve ramus in preoperatively tested sacral nerve foramen</p> <p>Groups: G1: Neurostimulation G2: control</p> <p>N at enrollment: G1: 25 G2: 26</p> <p>N at follow-up: 51 patients with urgency/frequency</p> <p>Age, mean yrs ± SD: Total: 39.0 ±11.8</p> <p>Race/ethnicity: NR</p> <p>Women, %: 90</p> <p>Follow-up: 6 months</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • >16 years of age • refractory to standard medical therapy • Bladder capacity of at least 100mL • -Normal upper urinary tract <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Treatable etiologies for urinary/frequency symptoms • Neurologic conditions • -Primary stress incontinence • Pelvic pain symptoms 	<p>Mean voids/day: 16.0 ±8.2</p> <p>Mean mL voided/void: G1: 118±74 G2: 124±66</p> <p>Mean max. voided volume: 288 ±156</p> <p>Mean total voided volume/day: 1693± 866</p> <p>Mean % indicating they felt empty: 44 ±39</p> <p>Mean duration of urinary symptoms before enrollment: 8.1 ± 9.2 years</p> <p>Mean pelvic/bladder discomfort (0 none-3 severe): 2.1 ±0.8</p> <p>Degree urgency before voiding: G1: 2.2±0.6 G2: 2.4±0.5</p>	<p>Mean mL voided/void: G1: 226±124 G2: 123±75 P< 0.001</p> <p>Degree urgency before voiding: G1: 1.6±0.9 G2: 2.3±0.5 P = 0.01</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: -</p> <p>Masking: -</p> <p>Pt selection criteria: +</p> <p>Loss to followup: NR</p> <p>Drop-out rates: NR</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: ++</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 3. KQ2 Procedural and surgical treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Evidence Table 3. KQ2 Procedural and surgical treatments (continued)					
<p>Author: Janknegt et al., 2001</p> <p>Country and setting: 15 investigators in US, Canada, and Europe</p> <p>Enrollment period: December 1993-September 1999</p> <p>Funding: NR.</p> <p>Author conflict of interest: NR</p>	<p>Design: Prospective case series</p> <p>Intervention: Implantable Interstim system stimulating the nerve ramus in preoperatively tested sacral nerve foramen</p> <p>Groups: NA</p> <p>N at enrollment: NA</p> <p>N at follow-up: 96 patients with urge incontinence</p> <p>Age: NR</p> <p>Women, n (%): 85 (89)</p> <p>Race/ethnicity: NR</p> <p>Follow-up: ≥12 months Mean follow-up 30.8 months (range 12-60)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • age > 16 years • refractory to standard medical therapy for UUI • 100 mL bladder capacity with normal upper urinary tract <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • neurological conditions • stress urinary incontinence • primary pelvic pain 	<p>Duration of symptoms, mean yrs ± SD: 9.1 ± 7.0.</p> <p>Leaking episodes/day: 10.9 ± 6.5</p> <p>Leak severity ranking: 2.0 ± 0.6</p> <p>Absorbent pad/diapers replaced/day: 6.6 ± 5.2</p> <p>Pads/diapers used daily, mean ± SD: 7.1 ± 5.1</p> <p>Leaking episodes per day, mean ± SD: 8.0 ± 4.8</p> <p>Heavy leaks per day, mean ± SD: 4.7 ± 4.2</p>	<p>Leaking episodes/day: 4.2 ± 4.9 p<0.0001</p> <p>Completely dry, n (%): 25 (26)</p> <p>> 50% reduction in frequency of incontinent episodes, n (%): 35 (36)</p> <p>Leak severity ranking: 1.7 ± 0.6 P<0.0001</p> <p>Pads/diapers used daily mean ± SD: 2.9 ± 3.8 P<0.0001</p> <p>Stopped using pads, n (%): 30 (33)</p> <p>Reduced pad use ≥ 50%, n (%): 25 (28)</p> <p>Heavy leaks per day, mean ± SD: 2.8 ± 4.0 P<0.0001</p> <p>Eliminated both moderate and heavy leaking episodes, n (%): 38 (43)</p> <p>≥ 50% reduction in moderate/heavy leaking, n (%): 18 (20)</p> <p>11 patients explanted due to lack of efficacy</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: fair</p> <p>Randomization: NA</p> <p>Masking: NA</p> <p>Pt selection criteria: +</p> <p>Loss to followup: ++</p> <p>Drop-out rates: +</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: ++</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 3. KQ2 Procedural and surgical treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
				(n=9), chronic leg pain (n=1), or bowel dysfunction (n=1)	

Evidence Table 3. KQ2 Procedural and surgical treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
--------------------------	--	--------------------------------------	--------------------------------	-----------------	-----------------------

Evidence Table 3. KQ2 Procedural and surgical treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Kessler et al., 2007</p> <p>Country and setting: Switzerland, Swiss Registry database</p> <p>Enrollment period: July 2000 to December 2005</p> <p>Funding: not described</p> <p>Author conflict of interest: Authors Kessler, Buchser, and Burkhard have consultancy agreements with Medtronic; author Zrehen is a Medtronic employee; Metronic Switzerland was assigned to monitor and maintain the Swiss Registry.</p>	<p>Design: Prospective case series</p> <p>Intervention: SNM testing and IPG implantation</p> <p>Groups: A portion of this series is composed of patients with urge incontinence (not further defined).</p> <p>N at enrollment: 209 evaluated: 181 female, 28 male.</p> <p>N at follow-up: 209: all evaluated were considered for adverse events of SNM testing</p> <p>102 successful tests</p> <p>91 had implantation of implantable pulse generator (IPG). 71 of these had urge incontinence.</p> <p>Age, median: 58 yrs</p> <p>Race/ethnicity: NR</p> <p>Follow-up, mean: 24 months for UUI</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Tested for urge incontinence, nonobstructive chronic urinary retention and chronic pain syndrome • Refractory to conventional therapies • Underwent urologic evaluation with medical history, physical exam, bladder and/or pain diary, urine analysis, urine culture and urodynamic studies before SNM. <p>Exclusion criteria: NR</p>	<p>Urge incontinence, n: Female: 137 Male: 16</p> <p>Leakages per 24 hrs, median: 5</p> <p>Pads used per 24 hrs, median: 4</p> <p>Voids per 24 hrs, median: 10</p> <p>Of 91 who underwent IPG implantation (11 of the 102 with successful testing refused implantation): 71 urge incontinence; 13 nonobstructive chronic urinary retention and 7 chronic pelvic pain syndrome.</p>	<p>Median follow-up was 24 months.</p> <p>Primary outcome: subjective symptom improvement, changes in bladder/pain diary variables.</p> <p>SNM was considered successful in 64 of 91 implanted patients, including 50 with urge incontinence. This group at last follow-up had a median of 0 leakages per 24 hours, 1 pad use per 24 hours, 6 voids per 24 hours, and 80% subjective symptom improvement.</p> <p>Secondary outcome: Incidence and treatment of adverse events due to SNM:</p> <p>Adverse events were reported in 12 of 209 patients who were initially tested, including lead migration in 9, wound infection in 2, and refractory pain at tined lead site in 1. These were not broken down separately for urge incontinence group.</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: NA</p> <p>Masking: NA</p> <p>Pt selection criteria: +</p> <p>Loss to followup: ++</p> <p>Drop-out rates: NR</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: ++</p> <p>Measurement methods: -</p> <p>Measurement reliability: -</p> <p>Intervention description: +</p>

Evidence Table 3. KQ2 Procedural and surgical treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Kessler et al., 2007 (continued)				Adverse events with IPG implantation were reported in 10 of 91 patients, including lead migration (2) broken lead (1), wound infection at IPG site (2), pain at IPG site (3), IPG migration (1), and IPG malfunction after MIR (1). 6 of these patients required surgical revision. These were not broken down separately for urge incontinence group.	

Evidence Table 3. KQ2 Procedural and surgical treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Evidence Table 3. KQ2 Procedural and surgical treatments (continued)					
<p>Author: Koldewijn et al. 1994</p> <p>Country and setting: The Netherlands</p> <p>Enrollment period: June 1990 to August 1993</p> <p>Funding: NR</p>	<p>Design: Prospective case series</p> <p>Intervention: percutaneous stimulation of the sacral root S3 on the left and right sides</p> <p>Groups: NA</p> <p>N at enrollment: NR</p> <p>N at follow-up: 100</p> <p>Age (mean): 42 years</p> <p>Women, n: 86</p> <p>Race/ethnicity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> failed previous treatment with medication, behavioral therapy, surgery, or combination therapy <p>Exclusion criteria:</p> <ul style="list-style-type: none"> No other lower urinary tract conditions 	<p>Urge incontinence, n: 46</p> <p>Incomplete bladder emptying, n: 23</p> <p>Urgency and frequency, n: 13</p> <p>Mixed incontinence, n: 10</p> <p>Dysuria or pelvic pain, n: 5</p> <p>Stress incontinence, n: 3</p> <p>Detrusor overactivity, n: 52</p> <p>Detrusor underactivity, n: 24</p> <p>Urethral instability, n: 5</p> <p>No abnormalities by UDS, n: 19</p> <p>No neuro-physiological abnormalities, n: 18</p> <p>Central neuro-physiological abnormality, n: 10</p> <p>Peripheral neuro-physiological abnormality, n: 16</p>	<p>Perfect improvement (>90%), n: 22</p> <p>Moderate improvement (50-90 %), n : 8</p> <p>Slight improvement (10-50%), n : 10</p> <p>No response, n: 17</p> <p>Pain in the area of puncture during stimulation, n: 7</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: fair</p> <p>Randomization: NA</p> <p>Masking: NA</p> <p>Pt selection criteria: +</p> <p>Loss to followup: NR</p> <p>Drop-out rates: NR</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: -</p> <p>Measurement methods: +</p> <p>Measurement reliability: -</p> <p>Intervention description: +</p>

Evidence Table 3. KQ2 Procedural and surgical treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
--------------------------	--	--------------------------------------	--------------------------------	-----------------	-----------------------

Evidence Table 3. KQ2 Procedural and surgical treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
--------------------------	--	--------------------------------------	--------------------------------	-----------------	-----------------------

Evidence Table 3. KQ2 Procedural and surgical treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Korda et al., 1987</p> <p>Country and setting: Australia</p> <p>Enrollment period: 1983 - 1984</p> <p>Funding: NR</p>	<p>Design: Prospective case series</p> <p>Intervention: Bladder distention</p> <p>Groups: NA</p> <p>N at enrollment: 50</p> <p>N at follow-up: 50</p> <p>Age: NR</p> <p>Race/ethnicity: NR</p> <p>Gender: 100% F</p> <p>Follow-up: 3, 6 and 12 mo</p>	<p>Inclusion criteria: NR</p> <ul style="list-style-type: none"> • Diagnosis with detrusor instability and failure to respond to anticholinergic treatment <p>Exclusion criteria: NR</p>	NR	<p>Extraperitoneal rupture: 2 patients</p> <p>At 3 mos: Symptom free: 10 Improved: 25 Unchanged: 15</p> <p>At 12 mos: Symptom free: 9 Improved: 15 Unchanged: 26</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: NA</p> <p>Masking: NA</p> <p>Pt selection criteria: -</p> <p>Loss to followup: ++</p> <p>Drop-out rates: +</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: poor</p> <p>Age: -, NR</p> <p>Baseline OAB status: NR</p> <p>Baseline characteristics: -</p> <p>Length of followup: ++</p> <p>Measurement methods: -</p> <p>Measurement reliability: -</p> <p>Intervention description: +</p>

Evidence Table 3. KQ2 Procedural and surgical treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Evidence Table 3. KQ2 Procedural and surgical treatments (continued)					
<p>Author: Mundy 1983</p> <p>Country and setting: UK</p> <p>Enrollment period: NR</p> <p>Funding: NR</p>	<p>Design: retrospective case series</p> <p>Intervention: bladder transection</p> <p>Groups, N: G1: 23; circumferential transection of the bladder wall just above the ureteric orifices G2: 81; transection of the posterior and posterolateral bladder wall from 1 cm lateral to the right ureteric orifice to 1 cm lateral on the left side ureteric orifice</p> <p>N at enrollment: 104</p> <p>Duration of follow-up: at least 12 months (range 12-59 months; mean 35 months)</p> <p>N at follow-up: 104 at year 1, 82 at year 2, 49 at year 3, 25 at year 4, and 6 at year 5</p> <p>Age mean yrs (range): 37 (19-81)</p> <p>Women, n (%): 91(87.5%)</p> <p>Race/ethnicity: NR</p> <p>No separate analysis by</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> urodynamically proven detrusor instability frequency, urgency and urge incontinence for at least 3 years failure to respond to standard conservative treatment (e.g. bladder drill, probanthine therapy, imipramine therapy, emepromium bromide therapy, hydrostatic bladder dilatation, oxybutynin) <p>Exclusion criteria: NR</p>	<p>NR</p>	<p>Initial post-operative evaluation (3-6 months after operation): 74% cured, 14% improved; 12% failures; authors note same results at year 1</p> <p>At last evaluation (in patients followed for 2-5 years): 53 (65% still cured), 16 (19%) improved; 13 (16%) failures</p> <p>9 participants with residual symptoms after operation later responded to other therapies</p> <p>Among 68 patients with at least one postoperative urodynamic study: 50 symptomatically cured; 14 of these reverted to stability and 26 showed a shift to the right of the urodynamic pattern</p> <p>28 were failures or showed partial response; 8 were no longer unstable, and 14 remained the same as before their operation; authors note 6 with genuine stress incontinence were successfully managed by appropriate surgical treatment</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: NA</p> <p>Masking: NA</p> <p>Pt selection criteria: -</p> <p>Loss to followup: ++</p> <p>Drop-out rates: +</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: poor</p> <p>Age: +</p> <p>Baseline OAB status: NR</p> <p>Baseline characteristics: -</p> <p>Length of followup: ++</p> <p>Measurement methods: -</p> <p>Measurement reliability: -</p> <p>Intervention description: -</p>

Evidence Table 3. KQ2 Procedural and surgical treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
--------------------------	--	--------------------------------------	--------------------------------	-----------------	-----------------------

gender

Evidence Table 3. KQ2 Procedural and surgical treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Mundy 1983 (continued)				<p>Adverse events: 1 patient developed persistent urine leak from her bladder requiring reoperation.</p> <ul style="list-style-type: none"> • Urodynamic evidence of vesicoureteric reflux postoperatively that had not been observed pre-operatively in 14 patients; none were symptomatic or developed acute pyelonephritis 	

Evidence Table 3. KQ2 Procedural and surgical treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: O'Reilly et al., 2008</p> <p>Country and setting: Australia, Academic health centers</p> <p>Enrollment period: NR</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: Trans-sacral nerve (S3-S4) stimulation vs sham</p> <p>Groups: G1: transsacral nerve magnetic stimulator of the S3 and S4 nerve roots x 20 min/day x 12 wk G2: sham x 20 min/day x 12 wk</p> <p>N at enrollment: G1: 33 G2: 30</p> <p>N at follow-up: G1: 33 G2: 30</p> <p>Women, %: 100</p> <p>Age, mean ± SD: G1: 62.6 ± 13.45 G2: 59.8 ± 14.19</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Women • Age ≥ 18 • Ability to give informed consent • Ability to comply w/ instructions to use the device • Ability to perform treatment over an 84-d period and attend for f/u • OAB symptoms for ≥ 6 mos • Unresponsive to behavioral measures or anticholinergic medication <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Pregnancy • Actively trying to conceive • Waking voiding frequency < 7 in 12 hrs • Nocturia < 2 • SUL symptoms • Benign or malignant bladder tumor • UTI • Previous irradiation or drug-induced cystitis • Symptomatic diverticulum • Vaginal infection • DM • Peripheral neuropathy • Cardiovascular disease • Cancer of the genital tract, urinary tract calculi • Cardiac arrhythmia 	<p>Voids/day, mean ± SD G1: 10 ± 3.2 G2: 9 ± 3.3</p> <p>Symptom impact, median VAS score (IQR): G1: 80 (50-90) G2: 80 (70-90)</p> <p>AQoL, overall mean score ± SD G1: 0.67 ± 0.22 G2: 0.67 ± 0.26</p> <p>AQoL, illness, median score (IQR): G1: 0.42 (0.14-0.87) G2: 0.40 (0.17-0.87)</p> <p>AQoL, independent living, median score (IQR): G1: 1 (0.87-1) G2: 1 (0.88-1)</p> <p>AQoL, social relationships, median score (IQR): G1: 1 (0.90-1) G2: 0.97 (0.79-1)</p> <p>AQoL, physical senses, median score (IQR): G1: 0.94 (0.86-1) G2: 0.97 (0.92-1)</p> <p>AQoL, psychological well-being, median score (IQR): G1: 0.85 (0.85-0.91) G2: 0.87 (0.85-0.93)</p> <p>KHQ, general health, median</p>	<p>Voids/day, mean ± SD: G1: 9 ± 2.7 G2: 9 ± 3.2</p> <p>Symptom impact, median VAS score (IQR): G1: 70 (40-90) G2: 75 (50-80)</p> <p>AQoL, overall mean score ± SD: G1: 0.69 ± 0.22 G2: 0.72 ± 0.25</p> <p>AQoL, illness, median score (IQR): G1: 0.52 (0.19-0.81) G2: 0.42 (0.17-0.87)</p> <p>AQoL, independent living, median score (IQR): G1: 1 (0.9-1) G2: 1 (0.83-1)</p> <p>AQoL, social relationships, median score (IQR): G1: 0.94 (0.88-1) G2: 0.94 (0.86-1)</p> <p>AQoL, physical senses, median score (IQR): G1: 0.94 (0.86-1) G2: 0.94 (0.92-1)</p> <p>AQoL, psychological well-being, median score (IQR): G1: 0.85 (0.83-0.93) G2: 0.90 (0.85-0.98)</p> <p>KHQ, general health, median</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: fair</p> <p>Randomization: +</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: ++</p> <p>Drop-out rates: +</p> <p>Power calculation: +</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 3. KQ2 Procedural and surgical treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes score (IQR): G1: 25 (0-50) G2: 25 (0-50)	Quality Rating
--------------------------	--	--------------------------------------	--------------------------------	--	-----------------------

Evidence Table 3. KQ2 Procedural and surgical treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
O'Reilly et al., 2008 (continued)			KHQ, general health, median score (IQR): G1: 25 (0-25) G2: 25 (25-50)	KHQ, incontinence Impact, median score (IQR): G1: 100 (33.3-100) G2: 100 (33.3-100)	
			KHQ, incontinence Impact, median score (IQR): G1: 100 (66.6-100) G2: 100 (66.6-100)	KHQ, role limitations, median score (IQR): G1: 33.3 (16.6-83.3) G2: 33.3 (0-66.6)	
			KHQ, role limitations, median score (IQR): G1: 66.6 (50-83.3) G2: 58 (33.3-83.3)	KHQ, physical limitations, median score (IQR): G1: 41.6 (16.6-83.3) G2: 50 (16.6-66.6)	
			KHQ, physical limitations, median score (IQR): G1: 50 (16.6-83.3) G2: 50 (16.6-66.6)	KHQ, social limitations, median score (IQR): G1: 25 (10-60) G2: 20 (10-30)	
			KHQ, social limitations, median score (IQR): G1: 30 (10-70) G2: 30 (10-60)	KHQ, personal relationships, median score (IQR): G1: 0 (0-33.3) G2: 0 (0-33.3)	
			KHQ, personal relationships, median score (IQR): G1: 16.6 (0-66.6) G2: 16.6 (0-50)	KHQ, emotions, median score (IQR): G1: 44.4 (22.2-88.8) G2: 44.4 (22.2-77.7)	
			KHQ, emotions, median score (IQR): G1: 66.6 (33-100) G2: 55.2 (22.2-88.8)	KHQ, sleep and energy, median score (IQR): G1: 50 (33.3-66.6) G2: 50 (33.3-66.6)	
			KHQ, severity measures, median score (IQR): G1: 50 (33.3-83.3)	KHQ, severity measures, median score (IQR): G1: 53.3 (33.3-73.3) G2: 46.6 (40-66.6)	

Evidence Table 3. KQ2 Procedural and surgical treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
O'Reilly et al., 2008 (continued)			KHQ, sleep and energy, median score (IQR): G1: 66.6 (33.3-83.3)	Voided volume (mL), median (IQR): G1: 425 (250-550) G2: 375 (300-500)	
			KHQ, severity measures, median score (IQR): G1: 60 (46.6-76.6) G2: 50 (40-66.6)	Total voided volume/ day (mL), mean ± SD: G1: 1780 ± 630 G2: 1790 ± 710	
			Voided volume (mL), median (IQR): G1: 350 (250-500) G2: 400 (300-550)		
			Total voided volume/day (mL), mean ± SD: G1: 1850 ± 715 G2: 1800 ± 750		

Evidence Table 3. KQ2 Procedural and surgical treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Evidence Table 3. KQ2 Procedural and surgical treatments (continued)					
Author: Rios et al., 2007	Design: RCT	Inclusion criteria: <ul style="list-style-type: none"> • Urinary frequency, daily urgency & UUI, nocturia for > 6 mos prior to study • Involuntary detrusor contractions on cystometrogram 	Incontinence episodes/day, per-protocol analysis, mean ± SD: G1: 3.11 ± 2.50 G2: 5.76 ± 3.39 <i>P</i> = 0.002	Incontinence episodes/day, per-protocol analysis, mean ± SD: G1: 2.68 ± 3.08 G2: 4.50 ± 3.83 <i>P</i> = 0.012	Quality: Overall quality score: poor INTERNAL VALIDITY: poor Randomization: - Masking: + Pt selection criteria: - Loss to followup: ++ Drop-out rates: NR Power calculation: - Statistical issues: +
Country and setting: Brazil, academic hospital	Intervention: intravesical resiniferatoxin (RTX)	Exclusion criteria: <ul style="list-style-type: none"> • Use of anticholinergics or tricyclic antidepressants in last 2 mos • Neurologic conditions • UTI • Pelvic prolapse > grade 2 • History of pelvic radiation or bladder tumor • Poor bladder wall compliance • Detrusor underactivity 	Incontinence episodes/day, ITT analysis, mean ± SD: G1: 3.00 ± 2.46 G2: 5.06 ± 3.37 <i>P</i> = 0.015	Incontinence episodes/day, ITT analysis, mean ± SD: G1: 2.72 ± 3.04 G2: 4.20 ± 3.57 <i>P</i> = 0.027	EXTERNAL VALIDITY: fair Age: + Baseline OAB status: + Baseline characteristics: ++ Length of followup: - Measurement methods: + Measurement reliability: + Intervention description: +
Enrollment period: NR	Groups: G1: single dose 100 ml 50 nM RTX G2: placebo		Voids/day, per-protocol analysis, mean ± SD: G1: 9.68 ± 2.92 G2: 9.94 ± 2.76 <i>P</i> = 0.756	Voids/day, per-protocol analysis, mean ± SD: G1: 9.03 ± 2.77 G2: 9.10 ± 2.68 <i>P</i> = 0.814	
Funding: Departments of Urology of the Federal University of São Paulo, Paulista School of Medicine and Hospital do Servidor Público Estadual de São Paulo	N at enrollment: G1: 34 G2: 24 N at follow-up: G1: 33 G2: 23 Women, %: 100 Age, mean ± SD: 56 ± 12.09 Race/ethnicity: NR		Voids/day, ITT analysis, mean ± SD: G1: 9.66 ± 2.88 G2: 9.90 ± 2.58 <i>P</i> = 0.746	Voids/day, ITT analysis, mean ± SD: G1: 9.03 ± 2.73 G2: 9.21 ± 2.53 <i>P</i> = 0.738	
Author industry relationship disclosures: None			Enuretic episodes/wk, per-protocol analysis, mean ± SD: G1: 0.33 ± 0.53 G2: 0.59 ± 0.80 <i>P</i> = 0.340	Enuretic episodes/wk, per-protocol analysis, mean ± SD: G1: 0.15 ± 0.34 G2: 0.65 ± 1.06 <i>P</i> = 0.140	
			Enuretic episodes/wk, ITT analysis, mean ± SD: G1: 0.36 ± 0.53 G2: 0.46 ± 0.72 <i>P</i> = 0.864	Enuretic episodes/wk, ITT analysis, mean ± SD: G1: 0.17 ± 0.36 G2: 0.52 ± 0.98 <i>P</i> = 0.321	
			Nocturia episodes/day, per-protocol analysis, mean ± SD: G1: 2.22 ± 1.02	Nocturia episodes/day, per-protocol analysis, mean ± SD: G1: 1.73 ± 1.14 G2: 1.82 ± 1.16 <i>P</i> = 0.794	

Evidence Table 3. KQ2 Procedural and surgical treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
			G2: 2.29 ± 1.26 P = 0.827		

Evidence Table 3. KQ2 Procedural and surgical treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Rios et al., 2007 (continued)			<p>Nocturia episodes/day, ITT analysis, mean ± SD: G1: 2.21 ± 1.02 G2: 2.28 ± 1.19 <i>P</i> = 0.796</p> <p>KHQ, general health perception, mean score ± SD: G1: 37.88 ± 14.14 G2: 42.70 ± 21.61</p> <p>KHQ, incontinence impact, mean score ± SD: G1: 77.78 ± 28.46 G2: 83.33 ± 22.94</p> <p>KHQ, role limitations, mean score ± SD: G1: 62.63 ± 36.33 G2: 70.00 ± 35.29</p> <p>KHQ, physical limitations, mean score ± SD: G1: 59.60 ± 34.37 G2: 60.00 ± 31.71</p> <p>KHQ, social limitations, mean score ± SD: G1: 27.44 ± 25.23 G2: 29.90 ± 28.64 G2/G1: <i>P</i> < 0.05</p> <p>KHQ, personal relationship, mean score ± SD: G1: 34.92 ± 41.47 G2: 26.47 ± 34.89</p> <p>KHQ, emotions, mean score ± SD: G1: 57.24 ± 35.59 G2: 66.11 ± 29.39</p>	<p>Nocturia episodes/day, ITT analysis, mean ± SD: G1: 1.77 ± 1.14 G2: 1.89 ± 1.12 <i>P</i> = 0.769</p> <p>KHQ, general health perception, mean score ± SD: G1: 34.85 ± 13.89 G2: 43.75 ± 24.16</p> <p>KHQ, incontinence impact, mean score ± SD: G1: 60.61 ± 33.80 G2: 65.00 ± 38.20 G1/BL: <i>P</i> < 0.05 G2/BL: <i>P</i> < 0.05</p> <p>KHQ, role limitations, mean score ± SD: G1: 48.99 ± 35.09 G2: 49.12 ± 37.46 G2/BL: <i>P</i> < 0.05</p> <p>KHQ, physical limitations, mean score ± SD: G1: 45.45 ± 36.39** G2: 44.17 ± 39.84</p> <p>KHQ, social limitations, mean score ± SD: G1: 21.55 ± 25.30 G2: 38.50 ± 31.91</p> <p>KHQ, personal relationship, mean score ± SD: G1: 27.08 ± 38.33 G2: 37.50 ± 41.51</p> <p>KHQ, emotions, mean score ± SD: G1: 42.76 ± 35.80 G2: 54.45 ± 37.10 G1/BL: <i>P</i> < 0.05</p> <p>KHQ, sleep and energy, mean score ± SD: G1: 28.79 ± 24.03 G2: 38.33 ± 31.11 G1/BL: <i>P</i> < 0.05</p>	

Evidence Table 3. KQ2 Procedural and surgical treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Rios et al., 2007 (continued)			<p>KHQ, sleep and energy, mean score ± SD: G1: 43.43 ± 26.66 G2: 37.50 ± 24.71</p> <p>KHQ, symptom severity, mean score ± SD: G1: 8.08 ± 7.59 G2: 10.88 ± 13.55</p> <p>Cystometric capacity, per-protocol analysis, mean ± SD: G1: 112.94 ± 68.22 G2: 124.15 ± 76.05 <i>P</i> = 0.581</p> <p>Cystometric capacity, ITT analysis, mean ± SD: G1: 114.62 ± 67.89 G2: 127.42 ± 72.99 <i>P</i> = 0.496</p> <p>Max cystometric capacity, per-protocol analysis, mean ± SD: G1: 285.8 ± 183.20 G2: 246.50 ± 92.87 <i>P</i> = 0.306</p> <p>Max cystometric capacity, ITT analysis, mean ± SD: G1: 287.47 ± 180.64 G2: 240.63 ± 85.77 <i>P</i> = 0.194</p>	<p>KHQ, symptom severity, mean score ± SD: G1: 13.94 ± 9.91 G2: 10.88 ± 11.80 G1/BL: <i>P</i> < 0.05</p> <p>Cystometric capacity, per-protocol analysis, mean ± SD: G1: 144.70 ± 93.41 G2: 172.75 ± 117.83 <i>P</i> = 0.310</p> <p>Cystometric capacity, ITT analysis, mean ± SD: G1: 145.44 ± 92.08 G2: 167.92 ± 109.93 <i>P</i> = 0.322</p> <p>Max cystometric capacity, per-protocol analysis, mean ± SD: G1: 298.03 ± 161.48 G2: 270.50 ± 109.57 <i>P</i> = 0.402</p> <p>Max cystometric capacity, ITT analysis, mean ± SD: G1: 299.26 ± 159.18 G2: 260.63 ± 102.37 <i>P</i> = 0.237</p> <p>Bladder volume, 1st involuntary contraction, per-protocol analysis, mean ± SD: G1: 120.17 ± 68.98 G2: 140.83 ± 84.34 <i>P</i> = 0.093</p>	

Evidence Table 3. KQ2 Procedural and surgical treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Rios et al., 2007 (continued)			<p>Bladder volume, first involuntary contraction, per-protocol analysis, mean ± SD: G1: 89.31 ± 44.58 G2: 125.56 ± 79.92 <i>P</i> = 0.263</p> <p>Bladder volume, 1st involuntary contraction, ITT analysis, mean ± SD: G1: 103.82 ± 59.51 G2: 124.38 ± 73.01 <i>P</i> = 0.243</p> <p>Max detrusor voiding pressure, per-protocol analysis, mean ± SD: G1: 35.90 ± 12.05 G2: 34.06 ± 12.86 <i>P</i> = 0.807</p> <p>Max detrusor voiding pressure, ITT, analysis, mean ± SD: G1: 33.59 ± 12.59 G2: 35.00 ± 11.67 <i>P</i> = 0.666</p>	<p>Bladder volume, 1st involuntary contraction, ITT analysis, mean ± SD: G1: 121.83 ± 68.39 G2: 139.09 ± 79.83 <i>P</i> = 0.102</p> <p>Max detrusor voiding pressure, per-protocol analysis, mean ± SD: G1: 33.83 ± 15.78 G2: 32.33 ± 12.29 <i>P</i> = 0.625</p> <p>Max detrusor voiding pressure, ITT analysis, mean ± SD: G1: 33.33 ± 15.74 G2: 32.91 ± 11.50 <i>P</i> = 0.819</p>	

Evidence Table 3. KQ2 Procedural and surgical treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Schmid et al. 2006</p> <p>Country and setting: Switzerland; 3 clinics</p> <p>Enrollment period: January 2003 to October 2004</p> <p>Funding: International Institute for Research in Paraplegia; Swiss National Science Foundation</p> <p>Author conflict of interest: NR</p>	<p>Design: prospective case series</p> <p>Intervention: 1000 U Botox-A in 10 mL 0.9% normal saline injected into the detrusor muscle 1-2 mm under the urothelium layer at 30 sites on the inner surface of the bladder wall (approx. 0.3 mL each injection)</p> <p>Groups: NA</p> <p>N at enrollment: 100</p> <p>N at follow-up: 100 at 1 month; 80 at 3 months; 20 at 9 months</p> <p>Age: mean 63 years (range 24-89)</p> <p>Race/ethnicity: NR</p> <p>Follow-up: patients followed 4, 12, and 36 weeks after treatment Mean follow-up: 20 weeks (range 4-36)</p> <p>No separate analysis by gender</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> OAB syndrome (ICS) Urodynamically demonstrable nonneurogenic detrusor overactivity or hypersensitive bladder with premature filling sensation ≥ 8 voids per 24 hours (with or without incontinence) Not effectively treated with maximal doses of anticholinergics (or unable to tolerate those drugs) <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Renal dysfunction Myasthenia gravis Neurogenic bladder dysfunction Serious illness Pregnant or breastfeeding Interstitial cystitis Radiocystitis Bladder tumor Chronic pelvic pain syndrome Prostate hyperplasia Infravesical obstruction PVR > 150 mL Hypocontractile/ acontractile detrusor Prior SUI surgery and de novo 	<p>Maximal cystometric bladder capacity (mL; mean ± SD): 246 ±15</p> <p>Detrusor compliance (mL/cm H2O; mean ± SD): 24.5 ± 2.3</p> <p>Reflex volume (mL; mean ± SD): 169 ±14 (n=54)</p> <p>First desire to void (mL mean ± SD): 125 ±13</p> <p>Urgency to void (mL; mean ± SD): 214 ±16</p> <p>Postvoid residual (mL; mean ± SD): 21 ±8</p> <p>Incontinence, n: 84</p> <p>Voiding frequency/24 hrs, n: 9-15 times: 37 16-23 times:41 >24 times: 24</p>	<p>Voiding frequency/24 hrs: 4-8 times: 69 8-15 time : 24</p> <p>Maximal cystometric bladder capacity (mL; mean ± SD): 1 mo.: 381 ±19 3 mos.: 384 ±19 9 mos.: 325 ±55</p> <p>Detrusor compliance (mL/cm H2O; mean ± SD): 1 mo.: 40.5 ±2.9 3 mos.: 45.1 ±2.9 9 mos.: 37.1 ±8.4</p> <p>Reflex volume (mL; mean ± SD): 1 mo.: 229 ±25 (n=14) 3 mos.: 222 ±35 (n=14) 9 mos.: 195 ±57 (n=6)</p> <p>First desire to void (mL mean ± SD): 1 mo.: 212 ±16 3 mos.: 208 ±16 9 mos.: 178 ±40</p> <p>Urgency to void (mL; mean ± SD): 1 mo.: 309 ±20 3 mos.: 332 ±20 9 mos: 210 ±57</p> <p>Postvoid residual (mL; mean ± SD): 1 mo: 75 ±10 3 mos: 85 ±10 9 mos: 14 ±28</p> <ul style="list-style-type: none"> Urgency disappeared in 72 patients at the 4 week assessment and in 66% at the 12 	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: NA</p> <p>Masking: NA</p> <p>Pt selection criteria: +</p> <p>Loss to followup: +</p> <p>Drop-out rates: NR</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 3. KQ2 Procedural and surgical treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
		urge • Mixed incontinence		week followup	

Evidence Table 3. KQ2 Procedural and surgical treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Schmid et al., 2006 (continued)				<ul style="list-style-type: none"> • Incontinence disappeared in 74% at 4 week follow-up and 80% at 12 week followup • No improvement in 8 patients • 90% of patients experienced improvement in at least 1 quality of life category, increasing at 3 month follow-up and decreasing at 9 months <p>Adverse events:</p> <ul style="list-style-type: none"> • 4 patients developed temporary (4 week) urinary retention and postvoid residual > 400 mL at 1-2 weeks after injection, requiring temporary intermittent clean self-catheterization; PVR returned to baseline within 6-9 months • 15 patients reported mild difficulty in voiding during the first moth with PVR 150-200 mL; declined ICSC • 10 patients developed bladder infection successfully treated by antibiotics 	

Evidence Table 3. KQ2 Procedural and surgical treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Evidence Table 3. KQ2 Procedural and surgical treatments (continued)					
<p>Author: Schmidt et al., 1999</p> <p>Country and setting: 16 centers in US, Canada, and Europe</p> <p>Enrollment period: 1993- April 1997</p> <p>Funding: Supported by Medtronic Inc.</p> <p>Author conflict of interest: Schmidt, Jonas, Oleson, Janknegt, Siegel: financial interest and/or other relationship with Medtronic Inc.</p>	<p>Design: randomized controlled trial</p> <p>Intervention: implantable Interstim system stimulating the nerve ramus in preoperatively tested sacral nerve foramen</p> <p>Groups: G1: implantation with immediate stimulation (n=34 for efficacy analysis) G2: control group receiving standard medical therapy for 6 months; allowed to cross over at 6 months if remained eligible for implantation (n=42 for efficacy analysis)</p> <p>N at enrollment: 155</p> <p>N randomized: 98 (the remaining 57 did not respond to test stimulation so were not randomized)</p> <p>N at follow-up: 76 at 6 months</p> <p>Age: mean 46.6 ±13.0 (range 20.2 – 78.9)</p> <p>Gender: 125 F (80.6%), 30 M (19.4%)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • age > 16 years • refractory to standard medical therapy • ≥ 100 mL bladder capacity with normal upper urinary tract • good surgical candidate • able to complete study documentation and return for follow-up evaluation <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • neurological conditions • stress urinary incontinence • primary pelvic pain 	<p>Duration of urinary symptoms before enrollment: mean 9.0±7.4 (range 0.6-35.4)</p> <p>Mean number of daily incontinence episodes: 8.9± 5.9 G1: 9.7 ± 6.3 G2: 9.3 ± 4.8</p> <p>Mean leak severity rating: 1.9± 0.6 G1: 2.0 ± 0.7 G2: 1.8 ± 0.6</p> <p>Mean absorbent pads/diapers replaced daily: 4.8± 4.8 G1: 6.2 ± 5.0 G2: 5.0 ± 3.7</p> <p>Mean heavy incontinence episodes per day: G1: 3.4 ±3.8 G2: 2.6 ±3.5</p> <p>3-day diary data: 90.3% of patients had at least 1 moderate/heavy and 74.8% had at least 1 heavy incontinence episode</p> <p>Previous treatment for urinary problems: 153 (89.7%), including pharmacological in 144 (92.5%, nonsurgical in 55 (35.5%), and surgical in 88 (56.8%)</p>	<p>At 6 months:</p> <p>Mean number of daily incontinence episodes: G1: 2.6± 5.1 (p<0.0001) G2: 11.3 ± 5.9 (p=0.002)</p> <p>Mean leak severity rating: G1: 0.8 ± 0.9 (p<0.0001) G2: 2.0± 0.6 (p=0.006)</p> <p>Mean absorbent pads/diapers replaced daily G1: 1.1 ± 2.0 (p<0.0001) G2: 6.3 ± 3.6 (p=0.003)</p> <p>Mean heavy incontinence episodes per day: G1: 0.3 ±0.9 (P<0.0001) G2: 3.9 ±3.8</p> <p>G1: 47% completely dry at 6 months; 3 patients had no reduction in incontinence after 6 months (one underwent device explantation and 2 had increased frequency of urination and incontinence)</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: -</p> <p>Masking: -</p> <p>Pt selection criteria: +</p> <p>Loss to followup: -</p> <p>Drop-out rates: NR</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: ++</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 3. KQ2 Procedural and surgical treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
--------------------------	--	--------------------------------------	--------------------------------	-----------------	-----------------------

Race/ethnicity:
NR

Evidence Table 3. KQ2 Procedural and surgical treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Schmidt et al., 1999 (continued)				<p>Safety data (pooled for 157 patients): adverse events requiring surgical repositioning or replacement of implant devices document in 51 (32.5%)</p> <p>168 post-implant events reported by 83 patients, including pain at the neurostimulator site in 15.9%, pain at implant site in 19.1, and lead migration in 7.0%</p> <p>Infection or skin irritation led to device explantation in 2 patients and temporary explant in 2 patients</p> <p>No permanent injuries or nerve damage reported.</p>	

Evidence Table 3. KQ2 Procedural and surgical treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Evidence Table 3. KQ2 Procedural and surgical treatments (continued)					
<p>Author: Siegel et al. 2000</p> <p>Country and setting: US, Canada, and Europe, 12 centers in</p> <p>Enrollment period: 1993- 999</p> <p>Funding: partially funded through restricted research grants from Medtronic Inc.</p> <p>Author conflict of interest: NR</p>	<p>Design: case series</p> <p>Intervention: implantable Interstim system stimulating the nerve ramus in preoperatively tested sacral nerve foramen</p> <p>N at enrollment: 219 undergoing stimulator placement (from pool of 581 patients evaluated)*</p> <p>UUI: urge incontinence (n=41)</p> <p>Urgency/Frequency: (n=29)</p> <p>N at follow-up: total=112; 23 additional patients had undergone device explantation and an additional 84 were not yet due for long-term follow-up</p> <p>Age, mean (range): 43 (17-81)</p> <p>Women, %: 78%</p> <p>Race/ethnicity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> patients who received implanted Interstim and were due for long-term follow-up <p>Exclusion criteria: NA</p> <p>Indications, N: Urge incontinence: 41 Urgency/frequency: 29 Retention: 42</p>	<p>UUI episodes/day, mean ± SD: 11.6 ± 6.6</p> <p>Heavy episodes per day, mean ± SD: 3.6 ± 4.0</p> <p>Pads per day, mean ± SD: 6.7 ± 4.6</p> <p>Voids per day: 17.7 ± 8.6</p> <p>Average volume per void: 132.5 ± 93.6</p> <p>Average degree of urgency before voiding (scale 1 mild – 3 severe): 2.2. ± 0.7</p> <p>Void volume per void, mean mL ± SD: 343 ± 167</p>	<p>UUI episodes/day, mean ± SD: 5.0 ± 6.1 P<0.0001</p> <p>Heavy episodes per day, mean ± SD: 1.3 ± 3.5 P<0.0001</p> <p>Pads per day, mean ± SD: 3.4 ± 4.9 P<0.0001</p> <p>Voids per day: 10.6 ± 6.6 P < 0.0001</p> <p>Void volume per void, mean mL ± SD: 225 ± 162 P < 0.0001</p> <p>Improvement in degree of urgency before voiding was present in 69% of patients</p> <p>Eliminated use of catheterization, %: 58</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: NA</p> <p>Masking: NA</p> <p>Pt selection criteria: -</p> <p>Loss to followup: -</p> <p>Drop-out rates: -</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: ++</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 3. KQ2 Procedural and surgical treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Siegel et al. 2000 (continued)				<p>Adverse events: Pain at neurostimulator site (15.4%), new pain (9.0%), suspected lead migration (8.4%), infection (6.1%), transient electric shock (5.5%), pain at lead site (5.4%), adverse change in bowel function (3.0%), technical problems (1.7%), suspected device problems (1.6%), change in menstrual cycle (1.0%), adverse change in voiding function (0.6%), persistent skin irritation (0.5%), suspected nerve injury (0.5%), device rejection (0.5%), other (9.5%)</p> <p>Surgical revision rate: 33% (73/219)</p>	

Evidence Table 3. KQ2 Procedural and surgical treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Evidence Table 3. KQ2 Procedural and surgical treatments (continued)					
Author: Spinelli et al., 2001	Design: Case series	Inclusion criteria: • placement of a sacral neurostimulator	Incontinence episodes/day, mean ± SD: G2: 5.4 ± 3.9	Mean incontinence episodes/day, UUI patients (including detrusor or urethral instability), %: Completely dry: G1: 39	Quality: Overall quality score: fair
Country and setting: Italy, Registry	sacral neuromodulation with Itriel II or Interstim stimulation system	Exclusion criteria: NR	Detrusor instability QoL index, mean: G2: 33.1	Completely dry, 3 months, %: G1: 23 1-3 episodes: G1: 23 > 3 episodes: G1: 15	INTERNAL VALIDITY: poor Randomization: NA Masking: NA Pt selection criteria: -
Enrollment period: NR			Urgency/frequency QoL index, mean: G2: 40		
Funding: NR	Groups: G1: retrospective cases G2: prospective cases		Detrusor instability, n (%): G1: 44 (47.3) G2: 42 (40.8)	Incontinence episodes/day, 12 mo, mean ± SD: G2: 1.1 ± 1.6 P < 0.001	Loss to followup: + Drop-out rates: NR Power calculation: - Statistical issues: -
Author industry relationship disclosures: NR	N at enrollment: G1: 93 G2: 103 N at follow-up: G1: 65 G2: NR Age, mean (range): G1: 50.9 (17, 79) G2: 50.1 (17, 79) Race/ethnicity: NR Women, %: G1: 80.6 G2: 72.8 Follow-up: G1: median 40 months, range 28-73 months G2: 3, 6, 9, 12, 18 and 24 months		Retention, n (%): G1: 20 (21.5) G2: 25 (24.3) Neurogenic retention, n (%): G1: 0 G2: 10 (9.7) Detrusor hyperreflexia, n (%): G1: 8 (8.6) G2: 15 (14.6) Urgency/frequency, n (%): G1: 5 (5.6) G2: 5 (4.9) Pelvic pain, n (%): G1: 9 (9.7) G2: 4 (3.9) Urethral instability, n (%): G1: 4 (4.3) G2: 2 (1.9) Interstitial cystitis, n (%): G1: 3 (3.2) G2: 0 Detrusor insta-	Completely dry, 6 months, %: G2: 65 P < 0.001 Completely dry, 9 months, %: G2: 55 P < 0.003 Completely dry, 12 months, %: G2: 59 P < 0.001 Completely dry, 18 months, %: G2: 43 P < 0.04 Mean voids/day, %: < 8 voids: G1: 42 8-12 voids: G1: 42 >12 voids: G1: 18	EXTERNAL VALIDITY: good Age: + Baseline OAB status: + Baseline characteristics: + Length of followup: ++ Measurement methods: + Measurement reliability: + Intervention description: +

Evidence Table 3. KQ2 Procedural and surgical treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics bility or urgency/ frequency, n (%): Total: 102 (52)	Outcomes	Quality Rating
-------------------	---	-------------------------------	--	----------	----------------

Evidence Table 3. KQ2 Procedural and surgical treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Spinelli et al., 2001 (continued)				Detrusor instability QoL index, 3 months, mean: G2: 74.7 <i>P</i> < 0.001	
				Detrusor instability QoL index, 6 months, mean: G2: 80.5 <i>P</i> < 0.001	
				Detrusor instability QoL index, 12 months, mean: G2: 69.6 <i>P</i> < 0.001	
				Detrusor instability QoL index, 18 months, mean: G2: 73.7 <i>P</i> = NR	
				Urgency/ frequency QoL index, 3 months, mean: G2: 73 <i>P</i> = NR	
				Urgency/ frequency QoL index, 6 months, mean: G2: 66 <i>P</i> = NS	
				Urgency/ frequency QoL index, 12 months, mean: G2: 69.5 <i>P</i> = NS	
				Not requiring catheterization (but baseline catheterization), 3 months, %: G2: 67 <i>P</i> < 0.001	

Evidence Table 3. KQ2 Procedural and surgical treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Spinelli et al., 2001 (continued)				<p>Not requiring catheterization (but baseline catheterization), 6 months, %: G2: 67 <i>P</i> < 0.001</p> <p>Not requiring catheterization (but baseline catheterization), 9 months, %: G2: 50 <i>P</i> < 0.001</p> <p>Not requiring catheterization (but baseline catheterization), 12 months, %: G2: 50 <i>P</i> < 0.003</p> <p>Any adverse events, %: G1: NR G2: 15.5</p> <p>Pain at implant site or extension cable connector, %: G2: 3.9</p> <p>Hematoma/wound problem, %: G2: 1.9</p> <p>Surgical revision necessary, %: G2: 9.7</p> <p>Lead fracture, %: G2: 3.9</p> <p>Explant of stimulation system, %: G2: 3.9</p> <p>Lead repositioning, %: G2: 1.9</p> <p>Lead displacement, %: G1: 6.5</p>	

Evidence Table 3. KQ2 Procedural and surgical treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Spinelli et al., 2001 (continued)				<p>New lead required due to initial bad position, %: G1: 3.2</p> <p>Explantation of pulse generator due to infection, %: G1: 1</p> <p>Lead breakage, %: G1: 1 System removed due to treatment failure, %: G1: 3.2</p>	

Evidence Table 3. KQ2 Procedural and surgical treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Evidence Table 3. KQ2 Procedural and surgical treatments (continued)					
<p>Author: Sutherland et al. 2007</p> <p>Country and setting: US continence and female urology clinic</p> <p>Enrollment period: December 1993 to December 2004</p> <p>Funding: funded in part by Medtronic, Inc.</p> <p>Author conflict of interest: 3 authors received fees for educational speaking on OAB and incontinence from Pfizer, Indevus, Astellas, and Medtronic, and consulting fees from Medtronic</p>	<p>Design: retrospective case series</p> <p>Intervention: implantable pulse generator for sacral nerve stimulation (device made by Medtronic)</p> <p>N at enrollment: 234 patients underwent the procedure</p> <p>N at follow-up: 104 patients were consented</p> <p>Age, mean yrs ± SD: 50 ± 13.4</p> <p>Women, N: 91</p> <p>Race/ethnicity: NR</p> <p>Gender:</p> <p>Mean follow-up 22 months (range 3-162 months)</p>	<p>Inclusion criteria: • Undergoing SNS</p> <p>Exclusion criteria: NR</p>	<p>Urge incontinence: 96% experienced reduction in pad use after SNS therapy; 50% completely dry; 46% used fewer pads, and 4% had no change</p> <p>Nocturnal voids per 24 hrs, mean ± SD: 2.3 ± 1.8</p> <p>Leaks per 24 hrs, mean ± SD: 5.0 ± 4.7</p> <p>Pads used per 24 hrs, mean ± SD: 2.3 ± 2.6</p> <p>Duration of symptoms, mean mos (range): 116-130 (9-600)</p>	<p>Urge incontinence: 96% experienced reduction in pad use after SNS therapy; 50% completely dry; 46% used fewer pads, and 4% had no change</p> <p>Voids per 24 hrs, mean ± SD: 8.5 ± 5.0 voids/24 hr P < 0.0001</p> <p>Nocturnal voids per 24 hrs, mean ± SD: 1.6 ± 2.2 P = 0.0091</p> <p>Leaks per 24 hrs, mean ± SD: 1.0 ± 1.4 leaks/24 hr P < 0.0001</p> <p>Pads used per 24 hrs, mean ± SD: 0.3 ± 0.7 pads/24 hr P < 0.0001</p> <p>Subjective improvement: >50%: 69% >80%: 50% >90%: 35%</p> <p>Satisfaction: - satisfied: 60.5% - equivocal: 23.3% - dissatisfied: 16.1%</p> <p>Adverse events: - infection: 13 events (4 severe requiring hospitalization for iv antibiotics) - hematoma: 2 events - discomfort at component: 28 events</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: NA</p> <p>Method and blinding: NA</p> <p>Pt selection criteria: +</p> <p>Loss to followup: -</p> <p>Drop-out rates: NA</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: +</p> <p>Length of followup: -</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 3. KQ2 Procedural and surgical treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
--------------------------	--	--------------------------------------	--------------------------------	-----------------	-----------------------

Evidence Table 3. KQ2 Procedural and surgical treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Sutherland et al., 2007 (continued)				battery depletion: 5 events - bowel complaints: 4 events - pelvic pain: 1 event - high impedance: 2 events - malposition of component: 1 event - seizures: 2 events - herpes flare: 3 events	

Evidence Table 3. KQ2 Procedural and surgical treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Evidence Table 3. KQ2 Procedural and surgical treatments (continued)					
<p>Author: van Kerrebroeck et al., 2007</p> <p>Country and setting: US, Canada, and Europe, 17 centers</p> <p>Enrollment period: NR</p> <p>Funding: Medtronic Inc.</p> <p>Author conflict of interest: authors have financial interest and/or other relationship with Medtronic and other pharma and device companies (AMS, Astellas, Pfizer, Advanced Bionics, Uroplasty, Uromedica, J&J, Novartis, UCB, Bayer, Boehringer, Allergan, BioXell SpA, Alza, Solvay Pharma, Lilly Icos, Sanofi-Synthelabo, Janssen-Ortho, Purde Pharma, Negma Lerad, Laserscope, QLT, Lumenis, Schwarz, Glaxo, Eli Lilly, Boston Scientific)</p>	<p>Design: Prospective case series</p> <p>Intervention: implantable Interstim system stimulating the nerve ramus in preoperatively tested sacral nerve foramen</p> <p>N at enrollment: 163; 152 with implants</p> <p>N at follow-up: 1 year 79</p> <p>Age mean ± SD: 44.7 ± 11.2</p> <p>Women, n (%): 141 (87)</p> <p>Race/ethnicity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> patients receiving Interstim for management of voiding dysfunction <p>Exclusion criteria: NR</p>	<p>UUI:</p> <ul style="list-style-type: none"> -average number of leaking episodes per day: 9.6 ±6.0 - number of heavy leaks per day: 2.6± 3.3 - number of pads per day: >5 <p>Frequency:</p> <ul style="list-style-type: none"> - average number of voids per day 19.3±7.0 - average voided volume (mL): 92.3±52.8 -degree of urgency/day: 2.3±0.6 <p>Urinary frequency: 28</p> <p>UUI: 87</p> <p>Urinary retention: 31 *</p>	<p>UUI:</p> <ul style="list-style-type: none"> -average number of leaking episodes per day: 4.7 ±4.9 at one year follow-up; maintained throughout follow-up with mean 3.9± 4.0 at 5 years - number of heavy leaks per day: 1.2± 2.7 at one year follow-up and 0.8± 1.7 at 5 year follow-up - number of pads per day: 1.8 at 5 years follow-up <p>Frequency:</p> <ul style="list-style-type: none"> - average number of voids per day 13.0±7.9 at year 1 and 14.8± 7.6 at year 5 - average voided volume (mL): 169.9±118.2 at year 1 and 165.2± 147.7 at year 5 -degree of urgency/day: 1.9±0.8 at year 1, 2.1±0.7 at year 5 <p>Adverse events: -110 events required surgery in 60 patients - system explanted from 16 due to AE or lack of efficacy - 31 AEs were device related and 240 were therapy related - new pain or undesirable change in stimulation: 43 patients, 60 events Pain at PNE or implant site, INS:</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: NA</p> <p>Masking: NA</p> <p>Pt selection criteria: +</p> <p>Loss to followup: ++</p> <p>Drop-out rates: +</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: ++</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 3. KQ2 Procedural and surgical treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
				30 patients, 40 events	

Evidence Table 3. KQ2 Procedural and surgical treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
van Kerrebroeck et al., 2007 (continued)				<ul style="list-style-type: none"> -infection at PNE or implant site: 12 patients, 14 events - pain at PNE or implant site, lead: 12 patients, 13 events - sensation of electric shock: 12 patients, 14 events - undesirable change in voiding function 11 patients, 12 events - lead migration: 13 patients, 14 events - technical problems during PNE/implant: 8 patients, 8 events - device problem: 16 patients, 19 events - Other: 51 patients, 77 events -surgical intervention required for adverse events in 60 patients (110 events), including device exchange in 36 patients, positioning of pulse generator in 12, reposition of lead and generator in 10, reposition lead in 10, permanent explant in 9, temporary explant in 2, other intervention in 2, bilateral implant in 1, and surgical wound care in 1 	

Evidence Table 3. KQ2 Procedural and surgical treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: van Voskuilen et al., 2006</p> <p>Country and setting: Netherlands, academic medical center</p> <p>Enrollment period: 1990 to 2003</p> <p>Funding: WAMU Foundation</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: retrospective case series</p> <p>Intervention: sacral nerve stimulation with Itrel-I device (1990 – 1994), Itrel-II device (1994 – 1999), or Interstim device (1999-2003)</p> <p>Groups: NA</p> <p>N at enrollment: 190</p> <p>N at follow-up: 149</p> <p>Women, n (%): 122 (81.9)</p> <p>Age, mean yrs ± SD: 46.7 (10.0)</p> <p>Race/ethnicity: NR</p> <p>Follow-up, mean mos ± SD: 64.2 (38.5)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> Receiving a neuromodulation system or revision surgery for urge incontinence or urgency-frequency (with or without pelvic pain complaints), or non-obstructive urinary retention <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Percutaneous tined lead procedure Receiving bilateral stimulation 	<p>Overactive bladder, n (%): 107 (71.8)</p> <p>Urinary retention, n (%): 42 (28.2%)</p> <p>Idiopathic urgency/frequency or retention, n (%): 129 (86.6%)</p> <p>Neurologic urgency/frequency or retention: 42 (28.2%)</p>	<p>Good result, n (%): 89 (59.7)</p> <p>Insufficient results, n (%): 44 (29.5)</p> <p>Reprogramming session at last visit, n (%): 16 (10.7%)</p> <p>Good result after reprogramming, n (%): 11 (68.8)</p> <p>Insufficient results after reprogramming, n (%): 5 (31.3)</p> <p>Adverse events, n: 194 in 106 patients</p> <p>Pain/undesirable change in stimulation, n: 64</p> <p>Undesirable change in voiding function/loss of efficacy, n : 42</p> <p>Pain at IPG implant site, n: 41</p> <p>Adverse change in bowel function, n: 15</p> <p>Suspected lead migration, n : 10</p> <p>Suspected device problem (including lead breakage), n: 6</p> <p>Infection, n: 6</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: NA</p> <p>Masking: NA</p> <p>Pt selection criteria: -</p> <p>Loss to followup: -</p> <p>Drop-out rates: NR</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: poor</p> <p>Age: +</p> <p>Baseline OAB status: NR</p> <p>Baseline characteristics: -</p> <p>Length of followup: ++</p> <p>Measurement methods: -</p> <p>Measurement reliability: -</p> <p>Intervention description: +</p>

Evidence Table 3. KQ2 Procedural and surgical treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
van Voskuilen et al. 2006 (continued)				Technical problem, n: 5 Suspected neuropraxia, n : 2 Other, n: 3 Requiring at least one reoperation due to AE, %: 48.3 Life span of replaced IPGs, mean mos (range): 73.7 (28-127) IPG removed, n (%): 21 (14.1)	

Evidence Table 4. KQ 2 Behavioral Treatment of OAB

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Arruda et al., 2008</p> <p>Country and setting: Brazil, community</p> <p>Enrollment period: August 2001 to September 2005</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: None</p>	<p>Design: RCT</p> <p>Intervention: Oxybutynin vs. functional electrostimulation vs. pelvic floor training</p> <p>Groups: G1: Oxybutynin 5 mg b.i.d. G2: Ambulatory stimulation applied vaginally G3: Pelvic floor exercises with a therapist and at home</p> <p>N Screened: 81</p> <p>N at enrollment: G1: 22 G2: 21 G3: 21</p> <p>N at follow-up: G1: 22 G2: 21 G3: 21</p> <p>Age, range: 35-80</p> <p>Race/ethnicity: NR</p> <p>Women, %: 100</p> <p>Length of follow up: 12 weeks</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> Community-dwelling Dx of OAB and DO Capable of completing a bladder diary and performing a pelvic muscle floor contraction For those with MUI, urge was predominant <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Hx of psychiatric or neurologic illness Persistent UTI Inability to comply with regular follow-up visits Current pregnancy Postvoid residual volume > 100 mL Contraindications to anticholinergic therapy Cardiac pacemaker Type III SUI Uncontrolled metabolic conditions or indwelling catheterization Using medications including anticholinergic drugs, calcium antagonists, beta agonists, dopamine agonists, striated muscle relaxants or estrogens Any uterine prolapse 	<p>UUI episodes/ week, mean ± SD: G1: 13.8 ± 11.6 G2: 13.5 ± 15.6 G3: 16.4 ± 17.2</p> <p>Nocturia episodes/week, mean ± SD: G1: 1.7 ± 1.5 G2: 1.9 ± 1.9 G3: 1.4 ± 1.2</p> <p>Pads/day, mean ± SD: G1: 2.6 ± 2.7 G2: 2.3 ± 2.4 G3: 2.7 ± 1.5</p> <p>Voids/day, mean ± SD: G1: 7.7 ± 2.6 G2: 8.6 ± 3.4 G3: 6.8 ± 2.2</p> <p>Residual volume mean mL ± SD: G1: 3.2 ± 6.3 G2: 1.0 ± 2.6 G3: 1.8 ± 3.3</p> <p>Volume, first desire to void, mean mL ± SD: G1: 117.7 ± 68.9 G2: 102.4 ± 51.1 G3: 86.7 ± 38.9</p> <p>Maximal cystometric capacity, mean mL ± SD: G1: 410.4 ± 194.1 G2: 350.0 ± 212.9 G3: 424.0 ± 149.0</p> <p>Involuntary detrusor contraction volume mean mL ± SD: G1: 189.5 ± 114.1 G2: 220.0 ± 127.2 G3: 239.3 ± 163.0</p>	<p>UUI episodes/ week, mean ± SD: G1: 7.0 ± 10.6 G2: 7.9 ± 13.7 G3: 7.8 ± 15.3 G1/BL: <i>P</i> = 0.007 G2/BL: <i>P</i> = 0.039 G3/BL: <i>P</i> = 0.035</p> <p>Urgency resolved, n (%): G1: 14 (63.6) G2: 11 (52.4) G3: 12 (57.1) <i>P</i> = 0.754</p> <p>Satisfied, n (%): G1: 17 (77.3) G2: 11 (52.4) G3: 16 (76.2) <i>P</i> = 0.142</p> <p>Nocturia episodes/week, mean ± SD: G1: 0.9 ± 0.8 G2: 1.2 ± 1.3 G3: 1.0 ± 1.1 G1/BL: <i>P</i> = 0.003 G2/BL: <i>P</i> = 0.036 G3/BL: <i>P</i> = 0.086</p> <p>Pads/day, mean ± SD: G1: 0.9 ± 1.5 G2: 0.9 ± 1.7 G3: 0.8 ± 1.3 G1/BL: <i>P</i> < 0.001 G2/BL: <i>P</i> = 0.004 G3/BL: <i>P</i> < 0.001</p> <p>Voids/day, mean ± SD: G1: 6.4 ± 1.6 G2: 7.9 ± 2.3 G3: 7.1 ± 2.1 G1/BL: <i>P</i> = 0.014 G2/BL: <i>P</i> = 0.291 G3/BL: <i>P</i> = 0.441</p> <p>Residual volume, mean mL ± SD: G1: 4.7 ± 9.4 G2: 1.1 ± 2.5 G3: 2.1 ± 3.5 G1/BL: <i>P</i> = 0.425 G2/BL: <i>P</i> = 0.760 G3/BL: <i>P</i> = 0.297</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: -</p> <p>Method and blinding: -</p> <p>Pt selection criteria: +</p> <p>Loss to followup: +</p> <p>Drop-out rates: -</p> <p>Power calculation: +</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: +</p> <p>Length of followup: +</p> <p>Measurement methods: -</p> <p>Measurement reliability: -</p> <p>Intervention description: +</p>

Evidence Table 4. KQ 2 Behavioral Treatment of OAB

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Arruda et al., 2008 (continued)			Involuntary detrusor contraction maximal pressure (mm H₂O ± SD): G1: 39.4 ± 26.1 G2: 43.7 ± 22.9 G3: 34.2 ± 19.8	Volume first desire to void, mean mL ± SD: G1: 157.3 ± 63.8 G2: 123.8 ± 59.0 G3: 137.6 ± 76.7 G1/BL: <i>P</i> = 0.019 G2/BL: <i>P</i> = 0.091 G3/BL: <i>P</i> = 0.017 Maximal cystometric capacity mean mL ± SD: G1: 517.3 ± 191.7 G2: 436.6 ± 178.7 G3: 489.0 ± 141.3 G1/BL: <i>P</i> = 0.001 G2/BL: <i>P</i> = 0.017 G3/BL: <i>P</i> = 0.113 Involuntary detrusor contraction volume (mL): G1: 188.6 ± 183.2 G2: 73.3 ± 112.4 G3: 114.3 ± 154.2 G1/BL: <i>P</i> = 0.986 G2/BL: <i>P</i> = 0.001 G3/BL: <i>P</i> = 0.044 Involuntary detrusor contraction maximal pressure, mm H₂O ± SD: G1: 19.6 ± 20.9 G2: 22.4 ± 30.1 G3: 17.2 ± 25.5 G1/BL: <i>P</i> < 0.001 G2/BL: <i>P</i> = 0.002 G3/BL: <i>P</i> = 0.027 Normal urodynamic evaluation, n (%): G1: 8 (36.4) G2: 12 (57.1) G3: 11 (52.4) <i>P</i> = 0.358 Persistent improvement at 1 year: G1: 10/17 G2: 4/11 G3: 9/16	

Evidence Table 4. KQ 2 Behavioral Treatment of OAB

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Arruda et al., 2008 (continued)				<p>Dry mouth, n (%): G1: 16 (72.7) G2: 0 G3: 0</p> <p>Difficulty voiding, n (%): G1: 2 (9.1) G2: 0 G3: 0</p> <p>Dizziness, n (%): G1: 1 (4.5) G2: 0 G3: 0</p> <p>Blurred vision, n (%): G1: 1 (4.5) G2: 0 G3: 0</p> <p>Constipation, n (%): G1: 1 (4.5) G2: 0 G3: 0</p> <p>Dry mouth, n (%): G1: 16 (72.7) G2: 0 G3: 0</p> <p>Difficulty voiding, n (%): G1: 2 (9.1) G2: 0 G3: 0</p> <p>Dizziness, n (%): G1: 1 (4.5) G2: 0 G3: 0</p> <p>Blurred vision, n (%): G1: 1 (4.5) G2: 0 G3: 0</p> <p>Constipation, n (%): G1: 1 (4.5) G2: 0 G3: 0</p>	

Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Bryant et al., 2002</p> <p>Country and setting: Australia, Specialty treatment centers</p> <p>Enrollment period: NR</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: Bladder training plus an educational intervention to reduce caffeine intake to less than 100mg per day vs. bladder training alone</p> <p>Groups: G1: BT + caffeine reduction G2: BT</p> <p>N at enrollment: G1: 48 G2: 47</p> <p>N at follow-up: G1: 36 G2: 39</p> <p>Age, yrs ± SD: G1: 56 ± 18 G2: 58 ± 16</p> <p>Women, %: G1: 94 G2: 87</p> <p>Weight in kg, mean ± SD: G1: 69 ± 17 G2: 68 ± 20</p> <p>Race/ethnicity: NR</p> <p>Parity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Adult • Symptoms of urinary urgency, frequency and/or urge incontinence • Routinely ingested ≥ 100mg caffeine/24 hrs <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Significant cognitive impairment • Pregnant • Symptoms of UTI 	<p>Voids/24 hrs, mean ± SD: G1: 11.4 ± 4.0 G2: 11.2 ± 3.5</p> <p>Urgency episodes/24 hrs, mean ± SD: G1: 4.8 ± 3.5 G2: 4.6 ± 3.0</p> <p>Leakages/24 hrs, mean ± SD: G1: 2.8 ± 3.2 G2: 3.1 ± 3.9</p> <p>Volume/void (mL), mean ± SD: G1: 178 ± 78 G2: 177 ± 101</p> <p>Caffeine intake/day (mg), mean ± SD: G1: 238.7 ± 121 G2: 272 ± 184</p> <p>Duration of urinary symptoms (yrs), mean ± SD: G1: 13.7 ± 14.9 G2: 10.7 ± 15.5</p>	<p>Leakages/24 hrs, mean ± SD: G1: 1.2 ± 1.9 G2: 1.4 ± 1.7</p> <p>Mean reduction in leakages/24 hrs, %: G1: 55 G2: 26 P = 0.219</p> <p>Voids/24 hrs, mean ± SD: G1: 6.8 ± 2.0 G2: 7.9 ± 2.6</p> <p>Urgency episodes per 24 h, mean ± SD: G1: 1.6 ± 1.9 G2: 3.2 ± 2.8</p> <p>Mean reduction in voids/24 hrs, %: G1: 35 G2: 23 P = 0.037</p> <p>Mean reduction in urgency episodes/24 hrs, %: G1: 61 G2: 12 P = 0.002</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Masking: -</p> <p>Pt selection criteria: +</p> <p>Loss to follow up: -</p> <p>Drop-out rates: -</p> <p>Power calculation: +</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of follow up: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 4. KQ 2 Behavioral Treatment of OAB

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Burgio et al., 1998</p> <p>Country and setting: US, academic health center outpatient geriatric medicine clinic</p> <p>Enrollment period: July 1989 to August 1995</p> <p>Funding: National Institutes on Aging</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT, placebo controlled</p> <p>Computer-generated random numbers using a block size of 6, w/ prior stratification by type and severity of incontinence</p> <p>Intervention: Biofeedback-assisted behavioral vs. drug treatment (oxybutynin chloride; possible range of doses 2.5 mg/d-5.0 mg t.i.d.) vs. placebo</p> <p>All patients had 4 visits over an 8-week period. Patients in G1 had biofeedback added to behavioral training in absence of 50% improvement by session 3.</p> <p>Groups: G1: Behavioral ± biofeedback G2: Pharmacologic G3: Placebo</p> <p>N at enrollment: 468 screened 271 not eligible 197 randomized G1: 65 G2: 67 G3: 65</p> <p>N at follow-up: G1: 61 G2: 55 G3: 53</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> Community-dwelling women at least age 55 Ambulatory At least 2 urge accidents per week by baseline bladder diary Urge incontinence as predominant pattern Urodynamic evidence of bladder dysfunction <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Continual leakage Postvoid residual urine volume >200mL Uterine prolapse past the entroitus Narrow-angle glaucoma Unstable angina Decompensated congestive heart failure Hx of malignant arrhythmias MMSE <20 (Dementia) 	<p>Duration of symptoms, mean yrs ± SD: G1: 9.4 ± 10.8 G2: 9.8 ± 11.9 G3: 12.7 ± 15.9</p> <p>Restricted activity, (%): G1: 30.8 G2: 32.8 G3: 38.5</p> <p>Urge UUI only, %: G1: 49.2 G2: 49.3 G3: 47.7</p> <p>Accidents per week, mean ± SD: G1: 15.8 ± 14.5 G2: 15.9 ± 14.1 G3: 15.4 ± 13.4 P = .98</p> <p>Severity classification, %: Mild (<5 accidents/wk) G1: 18.5 G2: 17.9 G3: 18.5 Moderate (5-10 accidents/wk) G1: 29.2 G2: 29.9 G3: 27.7 Severe (>10 accidents/wk) G1: 52.3 G2: 52.2 G3: 53.8</p> <p>Previous treatment with surgery, %: G1: 20.0 G2: 26.9 G3: 29.2</p>	<p>Accidents per week, mean ± SD: G1: 2.8 ± 4.7 G2: 5.7 ± 9.8 G3: 8.2 ± 11.6 P = 0.005</p> <p>Percent reduction, mean ± SD: G1: 80.7 ± 24.8 G2: 68.5 ± 37.2 G3: 39.4 ± 80.0 P < 0.001</p> <p>Percent reduction, range: G1: -0.9 – 100 G2: -85.7 – 100 G3: -400.0 - 100</p> <p>Patient perceptions of progress in treatment Much better G1: 74.1 G2: 50.9 G3: 226.9 Better G1: 25.9 G2: 30.9 G3: 38.5 About the same G1: 0.0 G2: 16.4 G3: 28.8 Worse G1: 0.0 G2: 1.8 G3: 5.8</p> <p>Estimate of % improvement, mean ± SD: G1: 81.6 ± 18.6 G2: 66.4 ± 35.4 G3: 45.1 ± 36.6</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: + Masking: + Pt selection criteria: + Loss to follow up: + Drop-out rates: + Power calculation: - Statistical issues: + EXTERNAL VALIDITY: good</p> <p>Age: + Baseline OAB status: + Baseline characteristics: ++ Length of follow up: + Measurement methods: + Measurement reliability: + Intervention description: +</p>

Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Burgio et al., 1998 (continued)	<p>Women, %: 100</p> <p>Age, yrs ± SD: G1: 67.3 ± 7.6 G2: 68.2 ± 7.5 G3: 67.6 ± 7.6</p> <p>Race/ethnicity: NR</p> <p>Parity mean ± SD: G1: 2.8 ± 2.0 G2: 2.1 ± 1.4 G3: 2.7 ± 1.8 <i>P</i> < 0.05</p>		<p>Previous treatment with medication, %: G1: 27.7 G2: 35.8 G3: 30.8</p> <p>Using estrogen, %: G1: 32.3 G2: 38.8 G3: 35.4</p> <p>Using diuretics, %: G1: 20.0 G2: 14.9 G3: 12.3</p>	<p>Having fewer accidents, %: G1: 100.0 G2: 87.3 G3: 67.3</p> <p>Accidents are smaller, %: G1: 87.3 G2: 78.8 G3: 54.0</p> <p>Able to wear less protection, %: G1: 76.0 G2: 56.0 G3: 334.1</p> <p>Comfortable enough with treatment to continue indefinitely, %: G1: 96.5 G2: 54.7 G3: 43.1</p> <p>Patient satisfaction with progress, %: Completely satisfied G1: 77.6 G2: 54.7 G3: 43.1 Somewhat satisfied G1: 22.4 G2: 40.0 G3: 34.0 Not at all satisfied G1: 0.0 G2: 10.9 G3: 38.0 Wish to receive another form of treatment, %: G1: 14.0 G2: 75.5 G3: 75.5</p> <p><i>P</i> < 0.001 for all comparisons</p>	

Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Burgio et al., 1998 (continued)				<p>Adverse effects, %, p compared to placebo G3:</p> <p>Dry mouth, %: G1: 34.9 G2: 96.9 G3: 54.8 <i>P</i> < 0.001</p> <p>Inability to void, %: G1: 6.3 G2: 21.5 G3: 3.2 <i>P</i> = 0.002</p> <p>Constipation, %: G1: 22.2 G2: 38.5 G3: 37.1 <i>P</i> = 0.10</p> <p>Blurred vision, %: G1: 9.5 G2: 15.4 G3: 9.7 <i>P</i> = 0.50</p> <p>Confusion, %: G1: 6.3 G2: 7.7 G3: 11.3 <i>P</i> = 0.59</p>	

Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Burgio et al.. 2000</p> <p>[See Burgio et al., 1998]</p> <p>Enrollment period: Two weeks after completion of Burgio et al. 1998</p> <p>Funding: National Institute on Aging, grants AG 08010 and K04 00431</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Modified crossover of RCT</p> <p>Intervention: Participants whose treatment was not completely successful were given the opportunity to switch or use combined treatment; further reductions in incontinence were measured.</p> <p>Groups: <u>Treatment</u> <u>Changes:</u> G1: Previous oxybutynin to behavioral modification alone G2: Previous behavior alone to 2.5 mg oxybutynin t.i.d. + behavioral therapy G3: Previous oxybutynin alone to 2.5 mg oxybutynin t.i.d. + behavioral therapy G4: Placebo to behavioral G5: Placebo to oxybutynin</p> <p>N at enrollment G1: 19 G2: 8 G3: 27 G4: 34 G5: 10</p> <p>N at follow-up: G1: 18 G2: 8 G3: 26 G4: NR G5: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> Community-dwelling women at least age 55 Ambulatory At least 2 urge accidents per week by baseline bladder diary Urge incontinence as predominant pattern Urodynamic evidence of bladder dysfunction Not completely dry or satisfied with previous, 8-wk treatment <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Continual leakage Postvoid residual urine volume >200mL Uterine prolapse past the introitus Narrow-angle glaucoma Unstable angina Decompensated congestive heart failure Hx of malignant arrhythmias MMSE <20 (Dementia) 	<p>% Reduction of incontinence after previous study (at baseline), mean: G1: 59.1 G2: 57.5 G3: 72.7 G4: 22.9 G5: 44.8</p>	<p>Final % Reduction of incontinence mean, p: G1: 77.1, 0.109 G2: 88.5, 0.034 G3: 84.3, 0.001 G4: 63.9, .002 G5: 76.5, .012</p> <p>Note: 29.2 of G3 declined to continue with drug therapy once they received behavioral modification.</p> <p>Numbers were too low to compare across groups.</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to follow up: +</p> <p>Drop-out rates: +</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of follow up: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Burgio et al., 2001</p> <p>[See Burgio et al., 1998]</p> <p>Country and setting: US, University based</p> <p>Enrollment period: [See Burgio et al., 1998]</p> <p>Funding: NIH</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: Psychological distress associated with pharmacologic treatment vs. behavioral vs. placebo</p> <p>Groups: G1: Behavioral training with biofeedback, 4 visits G2: Oxybutynin 2.5 mg po t.i.d., up to a max of 5 mg t.i.d. G3: Placebo</p> <p>N at enrollment: 197 women</p> <p>N treated: 169</p> <p>N at follow-up: 155 (completed both pre and post-treatment psychological assessment)</p> <p>Age yrs, mean ± SD: 67.5 ± 7.2</p> <p>Women, %: 100</p> <p>Race/ethnicity, %: White: 97 African American: 3</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • ≥ 55 yrs old • ambulatory • UUI ≥2x/wk (2 wk bladder diary), persisting x 3 mos • Predominant UUI • Urodynamic evidence of bladder dysfunction (DI or maximal capacity ≤350 mL) <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • contraindication to oxybutynin or behavioral treatment • Continual leakage • Post void residual > 200 mL • Uterine prolapse beyond the introitus • Decompensated CHF • Hx of malignant arrhythmias • Impaired mental status (MMSE <20) 	<p>SCL-90-R scores mean (SD):</p> <p>Somatization G1: 56.0 (10.6) G2: 51.4 (10.8) G3: 52.4 (11.1)</p> <p>Obsessive-Compulsive: G1: 56.5 (10.7) G2: 56.6 (11.4) G3: 57.7 (10.0)</p> <p>Interpersonal Sensitivity G1: 53.8 (11.0) G2: 51.4 (11.9) G3: 50.4 (12.0)</p> <p>Depression G1: 54.7 (10.0) G2: 52.5 (9.7) G3: 51.0 (11.9)</p> <p>Anxiety G1: 48.7 (13.9) G2: 46.8 (12.0) G3: 47.2 (12.8)</p> <p>Hostility G1: 49.3 (10.7) G2: 45.9 (10.1) G3: 48.3 (10.4)</p> <p>Phobia G1: 47.5 (10.2) G2: 46.7 (10.3) G3: 45.7 (8.5)</p> <p>Paranoid Ideation G1: 48.6 (12.4) G2: 49.3 (11.1) G3: 46.1 (12.8)</p> <p>Psychoticism G1: 54.1 (10.7) G2: 52.1 (10.3) G3: 50.5 (11.2)</p> <p>Global Severity G1: 54.2 (11.1) G2: 52.5 (10.3) G3: 52.4 (11.2)</p>	<p>Reduced incontinence episodes: G1: 83.3% G2: 74.4% G3: 41.4% <i>P</i> < 0.001</p> <p>SCL-90-R scores (SD):</p> <p>Somatization G1: 51.8 (11.4) G2: 51.2 (9.8) G3: 49.8 (13.0)</p> <p>Obsessive-Compulsive: G1: 53.8 (13.9) G2: 53.9 (10.9) G3: 55.4 (11.0)</p> <p>Interpersonal sensitivity G1: 49.5 (12.0) G2: 48.9 (11.2) G3: 49.2 (11.3)</p> <p>Depression G1: 51.5 (11.5) G2: 50.6 (10.7) G3: 51.4 (11.2)</p> <p>Anxiety G1: 46.1 (14.6) G2: 44.5 (12.3) G3: 45.8 (12.9)</p> <p>Hostility G1: 44.9 (10.8) G2: 44.6 (10.5) G3: 47.3 (11.2)</p> <p>Phobia G1: 47.1 (11.2) G2: 45.0 (8.3) G3: 45.1 (8.5)</p> <p>Paranoid ideation G1: 45.8 (10.9) G2: 47.2 (11.6) G3: 47.2 (12.0)</p> <p>Psychoticism G1: 49.2 (11.7) G2: 50.4 (9.7) G3: 49.6 (10.3)</p> <p>Global Severity G1: 50.8 (12.8) G2: 50.4 (10.0) G3: 51.4 (10.9)</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to follow up: +</p> <p>Drop-out rates: +</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of follow up: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Burgio et al., 2001 (continued)			<p>155 participants were compared to the 42 who did not complete intervention and psychological assessment, higher scores (greater distress) on 6 of 10 SCL-90-R scales (somatization, obsessive/compulsive, depression, hostility, paranoid ideation, global severity index), all p values <0.05</p> <p>Normal range, score 0-63 >75% in normal range (including dropouts) on 9 of 10 scales</p> <p>Highest impairment rate: 33% scored abnormal for obsessive-compulsive</p>	<p>Correlations between reduction of incontinence and changes in psychological symptoms:</p> <p>Somatization G1: 0.28* G2: -0.09 G3: 0.17</p> <p>Obsessive-Compulsive G1: 0.01 G2: -0.14 G3: 0.02</p> <p>Interpersonal Sensitivity G1: -0.09 G2: 0.04 G3: 0.13</p> <p>Depression G1: -0.04 G2: 0.03 G3: 0.07</p> <p>Anxiety G1: -0.10 G2: -0.01 G3: 0.34*</p> <p>Hostility G1: -0.10 G2: 0.09 G3: 0.11</p> <p>Phobic anxiety G1: -0.21 G2: -0.17 G3: 0.02</p> <p>Paranoid Ideation G1: 0.14 G2: -0.04 G3: -0.06</p> <p>Psychoticism G1: -0.01 G2: -0.01 G3: 0.13</p>	

Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Burgio et al., 2001 (continued)				Global Severity Index G1: 0.01 G2: 0.06 G3: 0.45*** * <i>P</i> < 0.05 *** <i>P</i> = 0.001	

Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Burgio et al., 2002</p> <p>Country and setting: US, Academic medical center</p> <p>Enrollment period: April 1995 to March 2001</p> <p>Funding: NIH</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: Behavioral training with or without biofeedback for 8 wks compared to self-training, followed by 2 wks of post-treatment bladder diaries and a patient satisfaction questionnaire</p> <p>Groups: G1: Behavioral training with biofeedback G2: Behavioral training with verbal feedback G3: Self-administered behavioral training N at enrollment: Evaluated: 474 Excluded: 252 Randomized: 222 G1: 73 G2: 74 G3: 75 N at follow-up: G1: 62 G2: 65 G3: 68 Women, %: 100 Age, mean ± SD: G1: 64.8 ± 7.1 G2: 65.8 ± 7.6 G3: 65.8 ± 8.5 Race/ethnicity, n (%): Black: G1: 11 (15.1) G2: 13 (17.6) G3: 11 (14.7)</p>	<p>Inclusion criteria:* Women Age ≥ 55 Community-dwelling Ambulatory Predominant urge incontinence ≥ 2x/wk for at least 3 mos Urodynamic evidence of bladder dysfunction</p> <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Continual leakage • PVR ≥ 150mL • Severe uterine prolapse past uterine introitus • Decompensated congestive heart failure • MMSE score <24 	<p>Incontinence episodes/week, mean ± SD: G1: 15.1 ± 13.5 G2: 17.3 ± 16.3 G3: 15.4 ± 14.2 <i>P</i> = 0.62</p> <p>< 5 incontinence episodes/wk, n (%): G1: 14 (19.2) G2: 14 (18.9) G3: 16 (21.3)</p> <p>5-10 incontinence episodes/wk, n (%): G1: 20 (27.4) G2: 21 (28.4) G3: 21 (28.0)</p> <p>> 10 incontinence episodes/wk, n (%): G1: 39 (53.4) G2: 39 (52.7) G3: 38 (50.7) <i>P</i> > 0.99</p> <p>UUI, n (%): G1: 50 (68.5) G2: 50 (67.6) G3: 50 (66.7)</p> <p>MUI, n (%): G1: 23 (31.5) G2: 24 (32.4) G3: 25 (33.3) <i>P</i> = 0.97</p> <p>Duration of symptoms (yrs), mean ± SD: G1: 7.1 ± 7.8 G2: 6.6 ± 7.7 G3: 6.6 ± 8.7</p> <p>Previous treatment with medication, n (%): G1: 16 (21.9) G2: 18 (24.3) G3: 21 (28.0)</p>	<p>Incontinence episodes/week, mean ± SD: G1: 6.1 ± 10.3 G2: 6.0 ± 10.7 G3: 6.7 ± 11.4 <i>P</i> = 0.78</p> <p>Incontinence episodes/week, % change, mean ± SD: G1: -63.1 ± 42.7 G2: -69.4 ± 32.7 G3: -58.6 ± 38.8 <i>P</i> = 0.23</p> <p>Incontinence episodes/week, % change, median (IQR): G1: -75.0 (-100, 120.0) G2: -82.8 (-100, 0) G3: -70.4 (-100, 29.4)</p> <p>Fewer incontinence episodes, n (%): G1: 51 (96.2) G2: 57 (100) G3: 58 (92.1) <i>P</i> = 0.09</p> <p>Smaller accidents, n (%): G1: 42 (79.2) G2: 49 (89.1) G3: 42 (67.7) G2/G3: <i>P</i> = 0.006 G1/G2/G3: <i>P</i> = 0.02</p> <p>Able to wear less protection, n (%): G1: 33 (71.7) G2: 40 (83.3) G3: 34 (70.8) <i>P</i> = .29</p> <p>Much better progress, n (%): G1: 33 (62.3) G2: 36 (63.2) G3: 20 (30.8)</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Masking: -</p> <p>Pt selection criteria: +</p> <p>Loss to follow up: +</p> <p>Drop-out rates: +</p> <p>Power calculation: +</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of follow up: -</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Burgio et al., 2002 (continued)	<p>Parity, mean \pm SD: G1: 2.5 \pm 1.7 G2: 3.0 \pm 2.0 G3: 2.7 \pm 2.0</p> <p>Using estrogen, n (%): G1: 52 (72.2) G2: 51 (68.9) G3: 44 (59.5)</p>		<p>Previous treatment with surgery, n (%): G1: 16 (21.9) G2: 13 (17.6) G3: 12 (16.0)</p> <p>Bladder capacity (mL), mean \pm SD: G1: 282 \pm 117 G2: 238 \pm 100 G3: 266 \pm 105 <i>P</i> = 0.04</p>	<p>Better progress, n (%): G1: 18 (34.0) G2: 20 (35.1) G3: 36 (55.4)</p> <p>No Progress, n (%): G1: 2 (3.8) G2: 1 (1.8) G3: 8 (12.3)</p> <p>Worse, n (%): G1: 0 G2: 0 G3: 1 (1.5) G2/G3: <i>P</i> < 0.001 G1/G3: <i>P</i> < 0.001 G1/G2/G3: <i>P</i> < 0.001</p> <p>Comfortable enough to continue treatment indefinitely, n (%): G1: 49 (98.0) G2: 54 (100) G3: 54 (88.5) G2/G3: <i>P</i> = 0.01 G1/G2/G3: <i>P</i> = 0.009</p> <p>Completely satisfied with progress, n (%): G1: 39 (75.0) G2: 47 (85.5) G3: 34 (55.7)</p> <p>Somewhat satisfied with progress, n (%): G1: 12 (23.1) G2: 8 (14.5) G3: 24 (39.3)</p> <p>Not at all satisfied with progress, n (%): G1: 1 (1.9) G2: 0 G3: 3 (4.9) G2/G3: <i>P</i> < 0.001 G1/G3: <i>P</i> = 0.03 G1/G2/G3: <i>P</i> < 0.001</p>	

Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Burgio et al., 2002 (continued)				<p>Incontinence restricts activities-not at all, n (%): Not at all: G1: 36 (69.2) G2: 43 (78.2) G3: 31 (50.8)</p> <p>Incontinence restricts activities- some or all, n (%): G1: 16 (30.8) G2: 12 (21.8) G3: 30 (49.2) G2/G3: <i>P</i> = 0.002 G1/G3: <i>P</i> = 0.047 G1/G2/G3: <i>P</i> = 0.007</p> <p>Not at all disturbed about incontinence, n (%) G1: 26 (49.1) G2: 32 (59.3) G3: 23 (39.0)</p> <p>Somewhat disturbed about incontinence, n (%): G1: 26 (49.1) G2: 22 (40.7) G3: 32 (54.2)</p> <p>Extremely disturbed about incontinence, n (%): G1: 1 (1.9) G2: 0 G3: 4 (6.8) <i>P</i> = 0.18</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Burgio et al., 2008</p> <p>Country and setting: US, Academic medical centers</p> <p>Enrollment period: July 2004 to January 2006</p> <p>Funding: NIH Pfizer</p> <p>Author industry relationship disclosures: 20 of 29 Allergan (3) Alza (1) Astellas Pharma (7) Bionovo (1) Bristol-Meyers Squibb (1) Dynogen (1) Elan (1) Ethicon (2) GSK (4) Johnson & Johnson (3) Lilly (7) Medtronic (1) Merck (1) Novartis (6) Ortho-McNeil (3) Pfizer (>10) Procter & Gamble (3) Q-Med (1) Renesa (1) Sanofi (1) Solace (1) Watson (1)</p>	<p>Design: RCT</p> <p>Intervention: Pharmacologic vs. pharmacologic plus behavioral interventions for 10 wks (Stage 1), followed by no drug therapy (Stage 2) with assessments at 10 wks and 8 mos*</p> <p>Groups: G1: Tolterodine tartrate (ER capsules), 4 mg/d plus behavioral training: pelvic floor muscle control and exercises, behavioral strategies to diminish urgency, suppress bladder contractions and prevent both stress and urge incontinence; delayed voiding; fluid management; handout with hints G2: Tolterodine tartrate (ER capsules), 4 mg/d</p> <p>N at enrollment: 4043 screened 2612 not eligible 870 declined 561 consented 254 excluded 307 randomized G1: 153 G2: 154</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Women • Community-dwelling • UUI only, or urge-predominant • ≥ 7 episodes of incontinence in a 7-day bladder diary • Persistent incontinence for at least 3 mos • No current use of antimuscarinics or other medications that could affect UI • No evidence of neurogenic etiology <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Age < 21 • Pregnant, planning a pregnancy in next 8 mos, or not using birth control • < 6 mos postpartum or termination after 20 wks gestation • Unable to contract pelvic floor muscles during evaluation • Behavioral therapy of > 2 mos in past 2 yrs • Continual leakage or always damp • Hypersensitive to study drug • Systemic disease that affects bladder function 	<p>UUI, 7-13 episodes/week, n (%): G1: 2 (1.3) G2: 2 (1.3)</p> <p>UUI, ≥ 14 episodes/week, n (%): G1: 2 (1.3) G2: 4 (2.6)</p> <p>MUI, 7-13 episodes/week, n (%): G1: 46 (29.9) G2: 46 (30.1)</p> <p>MUI, ≥ 14 episodes/week, n (%): G1: 104 (67.5) G2: 101 (66.0)</p> <p>Adjusted incontinence episodes/week, mean: G1: 23.1 G2: 23.2</p> <p>Previous non-surgical treatment for incontinence, n (%): G1: 19 (12) G2: 22 (14)</p>	<p>Success, n (%): G1: 43 (28) G2: 41 (27)</p> <p>Failure, n (%): G1: 75 (49) G2: 78 (51)</p> <p>Success rate, 8 months, lifetable analysis, % (95% CI): G1: 41 (32, 50) G2: 41 (33, 50) G1/G2: 0 (-12, 12)</p> <p>Success rate, 8 months, complete cases, % (95% CI): G1: 36 (27, 45) G2: 34 (25, 43) G1/G2: 2 (-10, 14)</p> <p>Success rate, 8 mos, assuming missings were failures, % (95% CI): G1: 28 (21, 35) G2: 27 (20, 34) G1/G2: 1 (-9, 11)</p> <p>Adjusted incontinence episodes/week, after Stage 1, mean: G1: 2.7 G2: 4.7</p> <p>Adjusted incontinence episodes/week, after Stage 1, mean change: G1: -20.4 G2: -18.5 G1/G2: -1.9 (-5.9, 2.0)</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Masking: -</p> <p>Pt selection criteria: +</p> <p>Loss to followup: -</p> <p>Drop-out rates: ++</p> <p>Power calculation: +</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: ++</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Burgio et al., 2008 (continued)	<p>N at follow-up: G1: 153 Completed treatment: 107 Outcome known at 8 mos: 119 G2: 154 Completed treatment: 101 Outcome known at 8 mos: 118</p> <p>Age, mean ± SD: G1: 55.8 ± 14.2 G2: 58.0 ± 13.5</p> <p>Women, %: 100</p> <p>Race/ethnicity, n (%): Hispanic: G1: 13 (9) G2: 17 (11) NH White: G1: 105 (69) G2: 85 (56) NH Black: G1: 22 (14) G2: 35 (23) Other: G1: 13 (9) G2: 15 (10)</p> <p>BMI, kg/m² ± SD: G1: 33.2 ± 9.5 G2: 32.3 ± 7.6</p>	<ul style="list-style-type: none"> • Currently using catheter to empty bladder • Postvoid residual volume > 150mL • Treatment for prolapse with pessary < 3 mos • Incontinence, vaginal, bladder or prolapse surgery in the past 6 mos • Urethral diverticulum • Previous augmentation cystoplasty or artificial sphincter • Neuromodulation for pelvic indications • Using anticholinergic agents, cholinergic agonists, tricyclic antidepressants, or duloxetine in ≤ 4 wks • Using diuretics with dosage change past 3 mos • Uncontrolled medical problem • History of bladder or pelvic cancer or pelvic radiation therapy • Glaucoma • Gastric retention • Non-ambulatory • Participation in other intervention trial that might influence outcome 		<p>Achieved 70% reduction in incontinence episodes, per bladder diary, 10 weeks (%): G1: 69 G2: 58 G1/G2: 11 (-0.3, 22.1)</p> <p>Totally dry, per bladder diary, 10 weeks (%): G1: 21 G2: 17</p> <p>Voids/day, mean change: G1: 0.5 G2: -0.4 G1/G2: 0.9 (0.3, 1.5)</p> <p>Symptom Distress Scores: G1/G2: <i>P</i> < 0.0001</p> <p>Symptom Bother Scores (OAB-q), Stage 1, mean change: G1: -36.7 G2: -30.4 G1/G2: <i>P</i> < 0.0001</p> <p>Symptom Bother Scores (OAB-q), Stage 2, mean change: G1: -30.9 G2: -20.4 G1/G2: <i>P</i> < 0.0001</p> <p>Patient completely satisfied, Stage 1, %: G1: 53 G2: 40 G1/G2: 13 (1, 25)</p>	

Evidence Table 2. KQ 2 Pharmacologic Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Burgio et al., 2008 (continued)				<p>Patient completely satisfied, 8 months, %: G1: 33 G2: 20 G1/G2: 13 (2, 24)</p> <p>Patient better or much better, Stage 1, %: G1: 90 G2: 77 G1/G2: 13 (4, 22)</p> <p>Patient better or much better, 8 months, %: G1: 69 G2: 43 G1/G2: 26 (14 , 38)</p> <p>Persistence in perceived improvement, 8 mos, women with improvement at Stage 1: G1: 72 G2: 54 G1/G2: 17 (4, 30)</p> <p>Harms: G1: 3 participants 1: blurred vision, syncope, night sweats, stomach cramping and weakness 2: 2 episodes of small-bowel obstruction and an allergic reaction (pruritus and rash) 3: tachycardia during stage 2 G2: 3 participants 1: small bowel obstruction 2: peripheral edema 3: renal cell carcinoma diagnosis during stage 2</p>	

Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Colombo et al., 1995</p> <p>Country and setting: Italy; Setting</p> <p>Enrollment period: May 1990 to March 1993</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: Oxybutynin vs. Bladder training for 6 weeks</p> <p>Groups: G1: Oxybutynin, 3 daily doses of 5 mg each for 6 weeks (dose reduced to half if substantial AEs) G2: Bladder training</p> <p>N at enrollment: G1: 42 G2: 39</p> <p>N at 6 wk follow-up: G1: 38 G2: 37</p> <p>N at 6 mo follow-up: G1: 28 G2: 27</p> <p>Age, yrs ± SD: G1: 48 (31 – 65) G2: 49 (24 – 65)</p> <p>Race/ethnicity, mean ± SD: Black: NR</p> <p>Women, N (%): G1: 42 (100) G2: 39 (100)</p> <p>Postmenopausal n (%): G1: 16 (38) G2: 20 (51)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Socially embarrassing (severe) urinary urge incontinence • On cystometry: detrusor instability, or low-compliance bladder (LCB), or sensory bladder <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Stable bladder at cystometry • Neurologic disease • Detrusor hyperreflexia • Age greater than 65 yr • Coexisting genuine SUI • Genital prolapse • Postvoid residual volume >50mL • Previous gynecologic or urogynecologic surgery • Prior use of any drug to treat UUI • Urethral diverticula • Fistulas • Urinary tract neoplasia • Cystitis • Bladder stones • Previous pelvic radiotherapy 	<p>Detrusor instability, n (%): G1: 14 (37) G2: 13 (35)</p> <p>Low compliance bladder, n (%): G1: 9 (24) G2: 8 (22)</p> <p>Sensory bladder, n (%): G1: 15 (39) G2: 16 (43)</p> <p>Daily UUI episodes, range 9 – 17</p> <p>Diurnal frequency, n (%): G1: 32 (84) G2: 29 (78)</p> <p>Nocturia, n (%): G1: 11 (29) G2: 18 (49)</p> <p>Volume at first desire (mL): G1: 120 ± 59 G2: 134 ± 61</p> <p>Volume at very strong desire (mL): G1: 317 ± 92 G2: 332 ± 87</p>	<p>Clinically cured overall, n (%): G1: 28 (74) G2: 27 (73)</p> <p>Cured among DI, (%): G1: 13 (93) G2: 8 (62) <i>P</i> = 0.07</p> <p>Cured among LCB: G1: 6 (67) G2: 6 (75) <i>P</i> = 0.56</p> <p>Cured among sensory bladder: G1: 9 (60) G2: 13 (81) <i>P</i> = 0.18</p> <p>Cured among 18 G1 patients requiring dosage halving: 12 (67%)</p> <p>Diurnal frequency resolved, n (%): G1: 18 (56) G2: 20 (69)</p> <p>Nocturia resolved n (%): G1: 3 (27) G2: 11 (61)</p> <p>Volume at first desire (mL), mean ± SD: G1: 179 ± 32 <i>P</i> = .0009 G2: 178 ± 49 <i>P</i> = .001</p> <p>Volume at very strong desire (mL) mean ± SD: G1: 408 ± 76 <i>P</i> = .00001 G2: 403 ± 69 <i>P</i> = .0002</p> <p>Patients still cured at 6 mos, n: G1: 16 G2: 26</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Masking: -</p> <p>Pt selection criteria: +</p> <p>Loss to follow up: ++</p> <p>Drop-out rates: +</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of follow up: -</p> <p>Measurement methods: +</p> <p>Measurement reliability: -</p> <p>Intervention description: +</p>

Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Colombo et al., 1995 (continued)				<p>Patients still cured at 6 mos among DI, n: G1: 8 G2: 8</p> <p>Patients still cured at 6 mos among LCB, n: G1: 4 G2: 6</p> <p>Patients still cured at 6 mos among sensory bladder, n: G1: 4 G2: 12</p> <p>Treatment discontinued in 6 cases: G1: 4 (3 cases of severe dry mouth, 1 case of previously unknown glaucoma) G2: 2 (treatment was time consuming)</p> <p>Other adverse effects G1: 18 (47%) with AE requiring halving of dosage: -dry mouth (n=15) -constipation (n=6) -nausea (n=5) -dizziness (n=2) - decrease in visual acuity (n=1) - tachycardia (n=1)</p>	

Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Diokno et al., 1995</p> <p>Country and setting: US; private clinic</p> <p>Enrollment period: January 1992 to December 1992</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Cohorts with comparison; (Series of patients self selected into two groups)</p> <p>Intervention: Bladder training vs. anticholinergic or antispasmodic</p> <p>Groups: G1: Bladder training G2: oxybutynin 2.5-5.0 mg b.i.d. or t.i.d.</p> <p>N at enrollment: G1: 39 G2: 33</p> <p>N at follow-up: G1: 26 G2: 28</p> <p>Age, mean yrs (range): 64 (20-93)</p> <p>Race/ethnicity: NR</p> <p>Women, N (%): 72 (100)</p> <p>Parity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> Incontinent by AHCPR guidelines* Incontinence persisted after treatment for transient incontinence <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Post-void residual of more than 150 mL 	<p>UUI, n (%): G1: 27 (49) G2: 28 (51)</p> <p>MUI with predominant urge, n (%): G1: 12 (71) G2: 5 (29)</p>	<p>Continent, n (%): G1: 1 (4) G2: 1 (4)</p> <p>Improved, n (%): G1: 22 (85) G2: 19 (68)</p> <p>No Change, n (%): G1: 3 (11) G2: 8 (28)</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: NA</p> <p>Masking: NA</p> <p>Pt selection criteria: -</p> <p>Loss to follow up: -</p> <p>Drop-out rates: NR</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: poor</p> <p>Age: -, NR</p> <p>Baseline OAB status: NR</p> <p>Baseline characteristics: -</p> <p>Length of follow up: ++</p> <p>Measurement methods: -</p> <p>Measurement reliability: -</p> <p>Intervention description: +</p>

Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes*	Quality Rating
<p>Author: Dorey et al., 2006</p> <p>Country and setting: UK, Urogynecology clinic</p> <p>Enrollment period: January 2002 to July 2004</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Retrospective review of cases</p> <p>Intervention: Physiotherapeutic treatment, avg. 3 sessions, including: PFME Urge suppression techniques Fluid advice Dietary advice</p> <p>Groups: NA</p> <p>N at enrollment: 87</p> <p>N at follow-up: 66 13 (15%) excluded for SUI 7 (8%) LTFU 1 (1%) self-catherized</p> <p>Women, n (%): 87 (100)</p> <p>Age, mean yrs (range): 56 (29-79)</p> <p>Race/ethnicity: NR</p> <p>BMI >30, %: 20</p> <p>Parous, n (%): 79 (91)</p> <p>Prior pelvic surgery, n (%): 60 (69)</p> <p>Prior hysterectomy, n (%): 30 (35)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> Treated for UUI from Jan 2002 to Jul 2004 <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Severe stress urinary incontinence Self-catherization 	<p>Daily leakage, n (%): 44 (67)</p> <p>Strong pelvic floor muscles, %: 75</p>	<p>Significant improvement in UUI, n (%): 55 (83)</p> <p>Daily leakage, n (%): 8 (12)</p> <p>100 % improvement, n (%): 8 (12)</p> <p>≥80% improvement, n (%): 24 (37)</p> <p>50-75% improvement, n (%): 31(48)</p> <p>Slight improvement, n (%): 6 (9)</p> <p>No improvement, n (%): 4 (6)</p> <p>No UUI at discharge, n (%): 5 (8)</p> <p>Mild UUI at discharge, n (%): 46 (69)</p> <p>Moderate UUI at discharge, n (%): 12 (18)</p> <p>Severe UUI at discharge, n (%): 4 (6)</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: NA</p> <p>Masking: NA</p> <p>Pt selection criteria: -</p> <p>Loss to follow up: NR</p> <p>Drop-out rates: NR</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of follow up: NA</p> <p>Measurement methods: -</p> <p>Measurement reliability: -</p> <p>Intervention description: +</p>

Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes [^]	Quality Rating
<p>Author: Dowd et. al., 2003</p> <p>Country and setting: US, Community</p> <p>Enrollment period: NR</p> <p>Funding: Kidney Foundation of Ohio</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Cohorts with comparison, patients alternately assigned to 3 groups</p> <p>Intervention: G1: Pamphlet plus CS plus coaching G2: Pamphlet plus cognitive strategies (CS) via audiotape daily for 4 wks then 1x wk until 12 wks G3: Pamphlet on bladder health, etc.</p> <p>Groups: G1: Bladder health information + CS + coaching G2: Bladder health information + CS G3: Bladder health information</p> <p>N at enrollment: Total: 58</p> <p>N at follow-up: G1: 18 G2: 18 G3: 14</p> <p>Women, n (%): G1: 17 (94) G2: 18 (100) G3: 13 (93) <i>P</i> = 0.54</p> <p>Age, mean: G1: 68.7 G2: 59.6 G3: 57.1 <i>P</i> = 0.03</p> <p>Race/ethnicity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> Adults Living independently No major hearing problem, Able to read and write English, CUBS for 6 mos or more MMSE score >20 Negative urine screen <p>Exclusion criteria: See inclusion criteria</p>	<p>UUI, %: G1: 26.7 G2: 31.6 G3: 33.3</p> <p>SUI or MUI, %: G1: 73.3 G2: 68.4 G3: 66.7</p>	<p>Comfort (UFIQC): Persons in G1 and G3 saw modest gains over time; G2 did not. Significant group-by-type of UI interaction: G1 and G3 with urge had more improvement than participants with stress or other UI (<i>F</i>=3.61; <i>P</i> = .037)</p> <p>Bladder Function (BFQ): Significant change for all groups (<i>F</i>=13.31; <i>P</i> = 0.0001); No significant interactions by type of incontinence by significant main effect for urge</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: NA</p> <p>Masking: NA</p> <p>Pt selection criteria: +</p> <p>Loss to follow up: -</p> <p>Drop-out rates: ++</p> <p>Power calculation: +</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: +</p> <p>Baseline OAB status: NR</p> <p>Baseline characteristics: -</p> <p>Length of follow up: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Fantl et al., 1981</p> <p>Country and setting: US, University</p> <p>Enrollment period: N/A</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Retrospective chart review</p> <p>Intervention: Bladder retraining drill (BRD) and anticholinergics when BRD not satisfactory</p> <p>Groups: G1: Detrusor muscle contraction approximately 5 seconds after a cough G2: Detrusor muscle contracts spontaneously without prior provocation G3: Both cough contraction and spontaneous contraction</p> <p>N at enrollment: G1: 39 G2: 4 G3: 49</p> <p>N at follow-up: NA</p> <p>Women, %: 100</p> <p>Age, mean ± SD: 42.7 ± 10:3</p> <p>Race/ethnicity: NR</p> <p>Parity, mean: 3.5</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Cystometric tracings showed a rise of 15 cm water or more in intravesical pressure when recorded independently and calculated by subtraction from intra-abdominal transrectal pressures <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Neuropathic conditions • UTI 	<p>Urgency and frequency, n (%): 72 (78.2)</p> <p>Urinary incontinence, n (%): 65 (70.6)</p> <p>Sensation of incomplete voiding, n (%): 59 (64.1)</p> <p>Nocturia, n (%): 66 (71.7)</p> <p>Dysuria, n (%): 3 (3.2)</p> <p>MUI, n (%): 6 (6.5)</p> <p>Enuresis in childhood, n (%): 5 (5.4)</p> <p>Previously operated on for similar urologic symptoms, n (%): 34 (36.9)</p>	<p>Cured, n (%): G1: 32 (82.1) G2: 1 (25) G3: 41 (83.7)</p> <p>Cured by BRD alone, n (%): 44 (78.6)</p> <p>Cured by BRD + anticholinergics, n (%):* 30 (83.5)</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: NA</p> <p>Masking: NA</p> <p>Pt selection criteria: +</p> <p>Loss to follow up: NR</p> <p>Drop-out rates: NR</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of follow up: NA</p> <p>Measurement methods: +</p> <p>Measurement reliability: -</p> <p>Intervention description: -</p>

Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Frewen, 1982</p> <p>Country and setting: UK; Outpatient clinic</p> <p>Enrollment period: NR</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Case series</p> <p>Intervention: Bladder training</p> <p>Groups: NA</p> <p>N at enrollment: 90</p> <p>N at follow-up: 90</p> <p>Age, range: 15-75</p> <p>Race/ethnicity: NR</p> <p>Women, N (%): 90 (100)</p> <p>Parity mean ± SD: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> Consecutive patients presenting with frequency, urgency and/or urge incontinence <p>Exclusion criteria:</p> <ul style="list-style-type: none"> NR 	<p>Urgency incontinence, n (%): 63 (70)</p> <p>Detrusor Instability, n (%): NR (63)</p> <p>Frequency, n: 10</p> <p>Diurnal-nocturnal enuresis, n: 20</p> <p>Duration of symptoms, yrs (range): stable: 4-5 unstable: 12</p> <p>Results of cystometry in 82 patients, n: Unstable: 40</p>	<p>Cure, n (%):* 78 (86.6)</p> <p>Free from incontinence (but residual urgency), n (%): 12 (13.4)</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: NA</p> <p>Masking: NA</p> <p>Pt selection criteria: -</p> <p>Loss to follow up: ++</p> <p>Drop-out rates: NR</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: +</p> <p>Length of follow up: NA</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Ghei et al., 2006</p> <p>Country and setting: UK, Primary care</p> <p>Enrollment period: NR Duration 16 weeks</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Observational cohort</p> <p>Intervention: Bladder retraining vs. Bladder retraining + antimuscarinic (oxybutynin IR/ ER, tolterodine IR/ER, or imipramine combined with either oxybutynin or tolterodine as combination therapy)</p> <p>Groups: G1: Bladder retraining alone G2: Antimuscarinic therapy+ bladder retraining</p> <p>N at enrollment: G1: 52 G2: 656</p> <p>N at follow-up: G1: 46 G2: 501</p> <p>Women, n (%): G1: 45 (86) G2: 618 (94)</p> <p>Age, mean ± SD: G1: 52 ± 14 G2: 54 ± 23</p> <p>Race/ethnicity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Frequency • Urgency with or without UUI <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • SUI • Symptoms of BOO 	<p>Incontinence episodes/day, mean ± SD: G1: 0.47 ± 1.2 G2: 1.12 ± 1.7 <i>P</i> < 0.001</p> <p>Voids/day, mean ± SD: G1: 14 ± 6 G2: 11 ± 6 <i>P</i> = 0.001</p>	<p>Incontinence episodes/day, mean change difference (95% CI): G2/G1: -0.60 (-0.93, -0.27) <i>P</i> = 0.024</p> <p>Voids/day, mean change difference (95% CI): G2/G1: 2.35 (1.4, 3.3) <i>P</i> < 0.001</p> <p>Nocturia episodes/day, mean change difference (95% CI): G2/G1: 0.57 (0.15, 0.99)</p> <p>Attendance visits, n (%) Failed follow-up: G1: 6 (12) G2: 155 (23) 1 follow-up visit: G1: 10 (19) G2: 15 (2) 2 follow-up visits: G1: 18 (35) G2: 86 (13) 3 follow-up visits: G1: 9 (17) G2: 225 (35) 4 follow-up visits: G1: 9 (17) G2: 175 (27)</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: NA</p> <p>Masking: NA</p> <p>Pt selection criteria: +</p> <p>Loss to follow up: -</p> <p>Drop-out rates: NR</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: +</p> <p>Length of follow up: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: -</p> <p>Intervention description: +</p>

Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Goode et al., 2002</p> <p>Country and setting: US; academic health center outpatient geriatric medicine clinic</p> <p>Enrollment period: July 1989 to August 1995</p> <p>Funding: National Institutes on Aging</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT, placebo controlled</p> <p>Computer-generated random numbers using a block size of 6, w/ prior stratification by type and severity of incontinence</p> <p>Intervention: Biofeedback-assisted behavioral vs. drug treatment (oxybutynin chloride; possible range of doses 2.5 mg/d-5.0 mg t.i.d.) vs. placebo</p> <p>All patients had 4 visits over an 8-week period. Patients in G1 had biofeedback added to behavioral training in absence of 50% improvement by session 3.</p> <p>Groups: G1: Behavioral ± biofeedback G2: Pharmacologic G3: Placebo</p> <p>N at enrollment: 468 screened 271 not eligible 197 randomized 105 had pre and post treatment urodynamics G1: 33 G2: 35 G3: 37</p> <p>N at follow-up: NA</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> Community-dwelling women at least age 55 Ambulatory At least 2 urge accidents per week by baseline bladder diary Urge incontinence as predominant pattern Urodynamic evidence of bladder dysfunction <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Continual leakage Postvoid residual urine volume >200mL Uterine prolapse past the introitus Narrow-angle glaucoma Unstable angina Decompensated congestive heart failure Hx of malignant arrhythmias MMSE <20 (Dementia) 	<p>Voids per day, mean ± SD: G1: 10.0 G2: 10.9 G3: 10.0</p> <p>Cystometry at baseline:</p> <p>First desire to void, mL ± SD: G1: 97.1 ± 50.7 G2: 101.1 ± 62.1 G3: 124.6 ± 73.7</p> <p>Strong desire to void, mL ± SD: G1: 188.5 ± 93.1 G2: 212.1 ± 86.7 G3: 222.3 ± 87.0</p> <p>Bladder capacity, mL ± SD: G1: 288.3 ± 117.0 G2: 308.7 ± 93.7 G3: 328.9 ± 107.6</p>	<p>Voids per day, mean ± SD: G1: 8.2 G2: 8.8 G3: 9.7</p> <p>DI on UDS, n (%) +Baseline DI/+DI post-treatment G1: 7 (21.2) G2: 1 (2.9) G3: 5 (13.5)</p> <p>DI on UDS, n (%) +Baseline DI/-DI post-treatment G1: 1 (3.0) G2: 7 (20.0) G3: 7 (18.9)</p> <p>DI on UDS, n (%) -Baseline DI/+DI post-treatment G1: 3 (9.1) G2: 3 (8.6) G3: 3 (8.1)</p> <p>DI on UDS, n (%) -Baseline DI/-DI post-treatment G1: 22 (66.7) G2: 24 (68.6) G3: 22 (59.5)</p> <p>Cystometry post-treatment:</p> <p>First desire to void, mL ± SD: G1: 115.9 ± 64.9 G2: 145.6 ± 74.0 G3: 133.5 ± 59.6</p> <p>Strong desire to void, mL ± SD: G1: 228.9 ± 106.4 G2: 282.0 ± 93.2 G3: 230.1 ± 78.8</p> <p>Bladder capacity, mL ± SD: G1: 305.6 ± 117.9 G2: 377.6 ± 92.1 G3: 323.0 ± 109.0</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to follow up: +</p> <p>Drop-out rates: +</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of follow up: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Goode et al., 2002 (continued)	<p>Age, yrs ± SD: G1: 65.3 ± 4.5 G2: 67.9 ± 7.9 G3: 67.6 ± 7.7</p> <p>Race/ethnicity, %: Black: 2 White: 98</p> <p>Women: 100%</p> <p>Parity mean ± SD: G1: 3.1 ± 1.7 G2: 2.1 ± 1.3 G3: 2.3 ± 1.5</p>			<p>Cystometry change:</p> <p>First desire to void, mL: G1: 18.8 G2: 44.4 G3: 8.9 <i>P</i> = 0.149</p> <p>Strong desire to void, mL: G1: 40.5 G2: 69.9 G3: 7.8 <i>P</i> = 0.018</p> <p>Bladder capacity, mL: G1: 17.3 G2: 68.9 G3: -6.0 <i>P</i> = 0.000</p> <p>Standardized estimates of direct and mediated effects of treatment:</p> <p>G1 v G3: Total effect: 0.28* Direct effect: 0.23 Mediated Effect: 0.05</p> <p>G2 v G3: Total Effect: 0.34* Direct Effect: 0.30* Mediated Effect: 0.04</p> <p>*<i>P</i> < 0.01</p>	

Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Herschorn et al., 2004</p> <p>Country and setting: Canada; Family medicine and urology clinics</p> <p>Enrollment period: June 2000 to December 2001</p> <p>Funding: Pharmacia Pfizer Canada</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: Health education with tolterodine vs. tolterodine alone</p> <p>Groups: G1: Health education with tolterodine G2: Tolterodine alone</p> <p>N at enrollment: G1: 39 G2: 45</p> <p>N at follow-up, 5 weeks: G1: 37 G2: 40</p> <p>N at follow-up, 10 weeks: G1: 35 G2: 32</p> <p>N at follow-up, 16 weeks: G1: 34 G2: 31</p> <p>Age, yrs ± SD: G1: 65.7 ± 14.5 G2: 63.1 ± 15.7</p> <p>Race/ethnicity: NR</p> <p>Women, %: G1: 92.3 G2: 84.4</p> <p>Parity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 50 • Symptoms of OAB • Attend investigators' practice • Normal cognitive function • Able to read English <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Enrollment in another clinical trial • Interstitial cystitis • UTI • Already taking tolterodine 	<p>Duration of OAB, yrs ± SD: G1: 8.7 ± 11.0 G2: 8.7 ± 10.8</p> <p>Mild bladder problems, n (%): G1: 13 (33.3) G2: 13 (28.9)</p> <p>Moderate bladder problems, n (%): G1: 19 (48.7) G2: 28 (62.2)</p> <p>Severe bladder problems, n (%): G1: 7 (18.0) G2: 4 (8.9)</p> <p>Obtained prescription, n (%): G1: 38 (97.4) G2: 37 (82.2) <i>P</i> < 0.05</p> <p>Intends to fill prescription, (%): G1: 0 (0) G2: 6 (7.5)</p>	<p>Change in incontinence episodes/wk, mean ± SD: G1: -7.72 ± 21.16 G2: -10.24 ± 19.56†</p> <p>Change in voids/d, mean ± SD: G1: -1.82 ± 3.41 G2: -2.18 ± 4.89 <i>P</i> = NR</p> <p>Change in nocturnal voids, mean ± SD: G1: -0.44 ± 1.13† G2: -0.07 ± 0.91</p> <p>No change in bladder problem severity, n (%): G1: 14 (42.4) G2: 20 (66.7)</p> <p>Improved bladder problem severity, n (%): G1: 15 (45.4)† G2: 6 (20)</p> <p>Worsened bladder problem severity, n (%): G1: 4 (12.1) G2: 4 (13.3)</p> <p>Compliance, 10 weeks, %: G1: 41 G2: 38 <i>P</i> > 0.05</p> <p>Compliance, 16 weeks, %: G1: 39 G2: 31 <i>P</i> > 0.05</p> <p>Continued or started non-drug OAB treatment, 16 weeks, %: G1: 82 G2: 53 <i>P</i> > 0.05</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: -</p> <p>Masking: -</p> <p>Pt selection criteria: +</p> <p>Loss to followup: -</p> <p>Drop-out rates: NR</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: +</p> <p>Baseline OAB status: NR</p> <p>Baseline characteristics: -</p> <p>Length of followup: ++</p> <p>Measurement methods: +</p> <p>Measurement reliability: -</p> <p>Intervention description: +</p>

Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Herschorn et al., 2004 (continued)				Stopped non-drug OAB treatments, %: G1: 12.8 G2: 28.9	

Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Jarvis et al., 1980</p> <p>Country and setting: UK, Hospital</p> <p>Enrollment period: NR</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: Behavioral: 1) rationale explained 2) pt instructed to pass urine at specific intervals during the day 3) pt encouraged to maintain usual fluid intake and chart fluid balance 4) pt introduced to someone successfully treated by the drill</p> <p>Groups: G1: Bladder drill G2: Control (told they should now be able to hold urine 4 hrs and sent home)</p> <p>N at enrollment: G1: 30 G2: 30</p> <p>N at follow-up: G1: 30 G2: 30</p> <p>Women, %: 100</p> <p>Age, range: 27-79</p> <p>Race/ethnicity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Women • Diagnosis of DI by pressure flow studies • Not taking medications known to affect urinary tract function <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Co-existing genuine SUI 	<p>UUI, n: G1: 30 G2: 30</p> <p>SUI, n: G1: 21 G2: 21</p> <p>Urgency, n: G1: 30 G2: 30</p> <p>Diurnal frequency, n: G1: 30 G2: 30</p> <p>Nocturnal frequency, n: G1: 27 G2: 25</p>	<p>Continent and symptom-free: G1: 27 G2: 7 <i>P</i> < 0.01</p> <p>UUI, n: G1: 3 G2: 23 <i>P</i> < 0.01</p> <p>SUI, n: G1: 3 G2: 16 <i>P</i> < 0.01</p> <p>Urgency, n: G1: 4 G2: 23 <i>P</i> < 0.01</p> <p>Diurnal frequency, n: G1: 5 G2: 23 <i>P</i> < 0.01</p> <p>Nocturnal frequency: G1: 3 G2: 20 <i>P</i> < 0.01</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: -</p> <p>Masking: -</p> <p>Pt selection criteria: -</p> <p>Loss to follow up: ++</p> <p>Drop-out rates: ++</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: poor</p> <p>Age: +</p> <p>Baseline OAB status: NR</p> <p>Baseline characteristics: -</p> <p>Length of follow up: ++</p> <p>Measurement methods: -</p> <p>Measurement reliability: -</p> <p>Intervention description:-</p>

Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Jarvis et al., 1981</p> <p>Country and setting: UK, Academic medical center</p> <p>Enrollment period: NR</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: Inpatient bladder drill vs. outpatient drug therapy</p> <p>Groups: G1: Inpatient bladder drill G2: Flavoxate hydrochloride 200 mg t.i.d. and imipramine 25 mg t.i.d. x 4 wks</p> <p>N at enrollment: G1: 25 G2: 25</p> <p>N at follow-up: G1: 25 G2: 25</p> <p>Women, %: 100</p> <p>Age, mean ± SD (range): G1: 47 ± 15.4 (17-78) G2: 46 ± 12.8 (17-65)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Women • UDS-diagnosed detrusor instability <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • DM • Neurological abnormalities • UTIs • Taking a drug suspected of affecting lower urinary tract function • Genuine stress incontinence 	<p>UUI, n: G1: 25 G2: 25</p> <p>SUI, n: G1: 19 G2: 17</p> <p>Urgency, n: G1: 25 G2: 25</p> <p>Frequency, n: G1: 25 G2: 25</p> <p>Nocturia, n: G1: 21 G2: 19</p> <p>Duration of symptoms (years), mean ± SD (range): G1: 4.3 ± 2.7 (1-15) G2: 5.4 ± 3.2 (1-20)</p> <p>Bladder volume (mL), first sensation, mean: G1: 87 G2: 79</p> <p>Max cystometric capacity (mL), mean: G1: 381 G2: 353</p>	<p>UUI, n: G1: 4 G2: 11</p> <p>SUI, n: G1: 1 G2: 9</p> <p>Urgency, n: G1: 4 G2: 11</p> <p>Frequency, n: G1: 6 G2: 12</p> <p>Nocturia, n: G1: 4 G2: 13</p> <p>Continent, n (%): G1: 21 (84) G2: 14 (56) <i>P</i> < 0.05</p> <p>Symptom-free, n (%): G1: 19 (76) G2: 12 (48) <i>P</i> < 0.05</p> <p>Bladder volume (mL), first sensation, mean: G1: 152 G2: 140</p> <p>Max cystometric capacity (mL), mean: G1: 470 G2: 446</p> <p>Adverse events, drug therapy, n:* Dizziness: 8 Headache: 6 Dry mouth: 6 Nausea: 4 Drowsiness: 2 Vomiting: 1</p> <p>Discontinued due to AEs, n: Dizziness: 1 Headache: 1 Vomiting: 1</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: -</p> <p>Masking: -</p> <p>Pt selection criteria: +</p> <p>Loss to follow up: ++</p> <p>Drop-out rates: -</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Lauti et al., 2008</p> <p>Country and setting: New Zealand, Academic</p> <p>Enrollment period: February 2003 to July 2003</p> <p>Funding: University of Otago</p> <p>Author industry relationship disclosures: None</p>	<p>Design: RCT pilot study, unmasked</p> <p>Intervention: Oxybutynin vs. bladder retraining vs. combination therapy</p> <p>Groups: G1: Oxybutynin 2.5 mg/day (daily dose could be increased by 2.5 mg every 5 days to a maximum of 15 mg/day) G2: Bladder retraining G3: Combination therapy</p> <p>N screened: 120</p> <p>N at enrollment: G1: 21 G2: 16 G3: 19</p> <p>N at 3 month follow-up: G1: 18 G2: 16 G3: 12</p> <p>N at 12 month follow-up: G1: 16 G2: 14 G3: 12</p> <p>Age, mean ± SD: G1: 53.8 ± 14.8 G2: 63.9 ± 17.2 G3: 47.6 ± 16.3</p> <p>Race/ethnicity: NR</p> <p>Women, %: Total: 100</p> <p>Parous, %: G1: 81 G2: 62.5 G3: 73.7</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age > 18 • Predominant UUI <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Predominant SUI • Contraindications to anticholinergic drugs • Current UTI • Neurological disease • Psychiatric disorder • Untreated co-existing pelvic organ prolapse below the hymenal ring • Obstructed voiding • Functional-reversible cause of incontinence • Inability to toilet independently • Limited fluency of written/spoken English • Current or recent use of any of the trial interventions 	<p>Urgency episodes/day, mean ± SD: G1: 3.8 ± 2.7 G2: 3.1 ± 2.2 G3: 3.5 ± 2.0</p> <p>Incontinence episodes/day, mean ± SD: G1: 2.2 ± 1.5 G2: 1.0 ± 1.1 G3: 1.8 ± 1.6</p> <p>Nocturia episodes/day, mean ± SD: G1: 1.1 ± 1.0 G2: 1.4 ± 1.0 G3: 0.8 ± 0.7</p> <p>Voids per day, mean ± SD: G1: 7.8 ± 2.8 G2: 8.0 ± 1.7 G3: 8.4 ± 2.5</p> <p>OAB-q total HRQoL, mean ± SD: G1: 73.1 ± 17.4 G2: 69.5 ± 24.6 G3: 71.6 ± 21.5</p> <p>OAB-q severity, mean ± SD: G1: 47.0 ± 16.2 G2: 42.3 ± 17.7 G3: 45.9 ± 18.7</p> <p>OAB-q coping, mean ± SD: G1: 72.0 ± 21.6 G2: 66.2 ± 31.7 G3: 73.8 ± 26.2</p> <p>OAB-q concern, mean ± SD: G1: 68.2 ± 19.0 G2: 68.8 ± 27.6 G3: 63.8 ± 29.2</p> <p>OAB-q sleep, mean ± SD: G1: 63.1 ± 28.7 G2: 59.8 ± 29.9 G3: 55.1 ± 27.6</p>	<p>Urgency episodes/day, 3 mos, mean ± SD: G1: 2.2 ± 1.8 G2: 1.5 ± 2.1 G3: 1.7 ± 1.8</p> <p>Urgency episodes/day, 12 mos, mean ± SD: G1: 2.3 ± 2.5 G2: 1.9 ± 2.1 G3: 2.0 ± 1.1</p> <p>Incontinence episodes/day, 3 mos, mean ± SD: G1: 0.8 ± 0.8 G2: 0.1 ± 0.3 G3: 0.6 ± 0.8</p> <p>Incontinence episodes/day, 12 mos, mean ± SD: G1: 0.9 ± 0.0 G2: 0.9 ± 1.0 G3: 0.8 ± 0.7</p> <p>Nocturia episodes/day, 3 mos, mean ± SD: G1: 1.0 ± 0.5 G2: 0.8 ± 0.7 G3: 0.6 ± 0.5</p> <p>Nocturia episodes/day, 12 mos, mean ± SD: G1: 1.0 ± 0.9 G2: 1.2 ± 0.6 G3: 0.7 ± 0.7</p> <p>Voids/day, 3 mos, mean ± SD: G1: 6.7 ± 1.8 G2: 6.3 ± 1.6 G3: 6.7 ± 2.2</p> <p>Voids/day, 12 mos, mean ± SD: G1: 7.2 ± 1.1 G2: 6.8 ± 1.4 G3: 7.6 ± 1.5</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Method and blinding: -</p> <p>Pt selection criteria: +</p> <p>Loss to followup: +</p> <p>Drop-out rates: +</p> <p>Power calculation: +</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: ++</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Lauti et al., 2008 (continued)			<p>OAB-q social, mean ± SD: G1: 92.5 ± 14.6 G2: 85.4 ± 19.9 G3: 92.8 ± 18.6</p> <p>SF-12 quality of life, physical, mean ± SD: G1: 49.0 ± 9.6 G2: 41.7 ± 11.5 G3: 46.2 ± 10.6</p> <p>SF-12 quality of life, mental, mean ± SD: G1: 49.1 ± 9.3 G2: 53.1 ± 8.8 G3: 46.3 ± 8.3</p>	<p>OAB-q total HRQoL, 3 mos, mean ± SD: G1: 82.3 ± 16.1 G2: 89.6 ± 9.4 G3: 91.8 ± 7.4</p> <p>OAB-q total HRQoL, 12 mos, mean ± SD: G1: 87.9 ± 11.6 G2: 81.6 ± 19.3 G3: 88.9 ± 9.9</p> <p>OAB-q severity, 3 mos, mean ± SD: G1: 37.2 ± 22.0 G2: 16.8 ± 12.0 G3: 21.6 ± 10.9</p> <p>OAB-q severity, 12 mos, mean ± SD: G1: 24.6 ± 10.6 G2: 33.1 ± 16.6 G3: 21.9 ± 14.8</p> <p>OAB-q coping, 3 mos, mean ± SD: G1: 79.2 ± 22.1 G2: 91.6 ± 9.5 G3: 92.7 ± 9.4</p> <p>OAB-q coping, 12 mos, mean ± SD: G1: 89.2 ± 13.7 G2: 81.5 ± 23.7 G3: 90.5 ± 10.0</p> <p>OAB-q concern, 3 mos, mean ± SD: G1: 78.6 ± 18.0 G2: 87.7 ± 14.5 G3: 90.2 ± 12.4</p> <p>OAB-q concern, 12 mos, mean ± SD: G1: 85.3 ± 15.5 G2: 81.7 ± 19.7 G3: 85.2 ± 13.4</p> <p>OAB-q sleep, 3 mos mean ± SD: G1: 77.7 ± 24.9 G2: 81.3 ± 14.6 G3: 85.0 ± 19.6</p>	

Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Lauti et al., 2008 (continued)				<p>OAB-q sleep, 12 mos, mean ± SD: G1: 79.9 ± 18.3 G2: 72.0 ± 24.5 G3: 83.2 ± 18.4</p> <p>OAB-q social, 3 mos, mean ± SD: G1: 96.4 ± 9.7 G2: 95.6 ± 7.0 G3: 98.9 ± 1.9</p> <p>OAB-q social, 12 mos, mean ± SD: G1: 97.3 ± 7.1 G2: 91.9 ± 14.2 G3: 97.3 ± 6.9</p> <p>SF-12 quality of life, physical G1: 50.6 ± 8.0 G2: 42.1 ± 12.7 G3: 48.4 ± 10.8</p> <p>SF-12 quality of life, physical, 12 mos, mean ± SD: G1: 50.0 ± 7.3 G2: 45.1 ± 13.9 G3: 45.3 ± 13.4</p> <p>SF-12 quality of life, mental, 3 mos, mean ± SD: G1: 50.4 ± 9.6 G2: 51.2 ± 9.5 G3: 46.7 ± 7.6</p> <p>SF-12 quality of life, mental, 12 mos, mean ± SD: G1: 49.6 ± 7.5 G2: 50.1 ± 10.7 G3: 50.6 ± 8.4</p> <p>Dry mouth, n (%): G1: 3 (21) G2: 5 (46) G3: 5 (42)</p> <p>Headaches, n (%): G1: 6 (43) G2: 1 (11) G3: 7 (58)</p> <p>Dizziness, n (%): G1: 4 (29) G2: 2 (20) G3: 3 (25)</p>	

Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Lauti et al., 2008 (continued)				Constipation, n (%): G1: 3 (21) G2: 3 (27) G3: 3 (27) Fatigue, n (%): G1: 9 (64) G2: 5 (46) G3: 7 (64)	

Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Macaulay et al., 1987</p> <p>Country and setting: UK, Specialty treatment center</p> <p>Enrollment period: NR</p> <p>Funding: Wellcome Trust, trustees of St. George's Hospital</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: Brief eclectic psychotherapy, bladder training or medication</p> <p>Groups: G1: Psychotherapy G2: Bladder drill G3: Propantheline</p> <p>N at enrollment: G1: 19 G2: 16 G3: 15</p> <p>N at follow-up: G1: 18 G2: 15 G3: 14</p> <p>Women, %: 100</p> <p>Age: NR</p> <p>Race/ethnicity: NR</p> <p>Follow-up: 3 months</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Previous completion of survey on psych conditions w/ OAB • Detrusor instability on UDS or sensory urgency <p>Exclusion criteria: NR</p>	<p>Detrusor instability, n: G1: 10 G2: 8 G3: 8</p> <p>Sensory urgency, n: G1: 9 G2: 8 G3: 7</p> <p>Voids/day, mean: G1: NR G2: NR G3: 10.8</p> <p>Bladder capacity (mL), mean: G1: 393 G2: NR G3: 323</p> <p>Bladder volume (mL), first sensation, mean: G1: 107 G2: 110 G3: 70</p> <p>Detrusor pressure rise (cm H₂O), mean: G1: NR G2: 45.5 G3: NR</p>	<p>Voids/day, mean: G1: NR G2: NR G3: 8.3 G3/BL: $P < 0.005$</p> <p>Bladder capacity (mL), mean: G1: 414 G2: NR G3: 368 G1/BL: $P = NS$ G3/BL: $P = NS$</p> <p>Bladder volume (mL), first sensation, mean: G1: 142 G2: 150 G3: 137 G1/BL: $P = NS$ G2/BL: $P < 0.05$ G3/BL: $P = 0.06$</p> <p>Detrusor pressure rise (cm H₂O), mean: G1: NR G2: 29.5 G3: NR G2/BL: $P < 0.05$</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: -</p> <p>Masking: -</p> <p>Pt selection criteria: -</p> <p>Loss to follow up: ++</p> <p>Drop-out rates: ++</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: -, NR</p> <p>Baseline OAB status: NR</p> <p>Baseline characteristics: -</p> <p>Length of follow up: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Mattiason et al., 2003</p> <p>Country and setting: Sweden, Denmark, Norway (Tolterodine Scandinavian Study Group)</p> <p>Enrollment period: October 1999 to December 2000</p> <p>Funding: Pharmacia Corp</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT, single-blinded (balanced blocks of 4, computerized randomization list)</p> <p>Intervention: Tolterodine 2mg b.i.d. ± bladder training (BT); Tolterodine dosage could be decreased to 1 mg P.O. b.i.d. during the first 2 wks if intolerable SE; BT taught with a written handout</p> <p>Groups: G1: Tolterodine + BT x 24 wks G2: Tolterodine x 24 wks</p> <p>N at enrollment: Total: 505</p> <p>N at follow-up: G1: 244 G2: 257</p> <p>N Completed treatment: G1: 77% G2: 79% Total: 74%</p> <p>Women, n (%) G1: 177 (73) G2: 201 (78)</p> <p>Age, median (range): G1: 62 (19, 86) G2: 63 (22, 86)</p> <p>Race/ethnicity: NR</p> <p>Follow-up: 24 weeks</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 18 • ≥ 8 voids/day and urinary urgency (± UUI) as determined by 1 wk bladder diary • With or without UUI • Women of reproductive age had to be using reliable birth control <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Contraindication to antimuscarinic therapy • Use of electrostimulation therapy or BT within prior 3 mo • Indwelling catheter or intermittent catheterization • Pregnancy or lactation • Use of anticholinergic agents or concomitant treatment for OAB (estrogen permitted) 	<p>Urgency episodes/day, mean (range): G1: 6.0 (0, 23.0) G2: 6.6 (0, 34.3)</p> <p>Incontinence episodes/day, mean (range): G1: 2 (0.3, 20.3) G2: 2.3 (0.3, 16.3)</p> <p>Voids/day, mean (range): G1: 10.3 (7.3, 27.6) G2: 10.6 (7.7, 24.6)</p> <p>Duration of symptoms > 5 years, n (%): G1: 120 (49) G2: 124 (48)</p> <p>Previous drug therapy for OAB, n (%): G1: 40 (160) G2: 35 (14)</p> <p>Previous surgery affecting lower urinary tract, n (%): G1: 28 (11.5) G2: 30 (12)</p> <p>Symptoms caused at least moderate problems, %: G1: 92 G2: 93</p>	<p>Urgency episodes/day, median % change (IQR): G1: -38 (-76.7, -14.1) G2: -38 (-68.7, -8.0) G1/G2: <i>P</i> = 0.75</p> <p>Incontinence episodes/day, median IQR% change; n=301: G1: -87 (-100, -20) G2: -81 (-100, -41.8) G1/G2: <i>P</i> = 0.28</p> <p>Voids/day, median % change (IQR): G1: -33 (-42.3, 21.3) G2: -25 (-38.8, -13.0) G1/G2: <i>P</i> < 0.001</p> <p>Voided volume (mL), median % change (IQR): G1: 31.5 (13.3, 56.2) G2: 20 (3.1, 45.4) G1/G2: <i>P</i> < 0.001</p> <p>Symptoms are “minor or less”, %: G1: 66.5 G2: 61.5</p> <p>Overall improvement in symptoms, %: G1: 76 G2: 71</p> <p>Worsening of symptoms, %: G1: 3 G2: 5</p> <p>Dry mouth, n (%): G1: 76 (31) G2: 90 (35)</p> <p>Headache, n (%): G1: 15 (6) G2: 21 (8)</p>	<p>Quality: Overall quality score: fair INTERNAL VALIDITY: poor Randomization: + Masking: + Pt selection criteria: + Loss to follow up: - Drop-out rates: - Power calculation: + Statistical issues: + EXTERNAL VALIDITY: good Age: + Baseline OAB status: + Baseline characteristics: ++ Length of follow up: ++ Measurement methods: + Measurement reliability: + Intervention description: +</p>

Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Mattiason et al., 2003 (continued)				<p>Constipation, n (%): G1: 7 (3) G2: 14 (5)</p> <p>≥ 1 SE, n (%): G1: 158 (65) G2: 177 (69) G1/G2: <i>P</i> = NS</p> <p>Withdrawal due to, %: AE: 15</p> <p>Withdrawal due to lack of efficacy: 3</p> <p>Withdrawal of consent : 2</p> <p>Protocol violations: 1</p> <p>Completed treatment: G1: 77% G2: 79%</p>	

Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)					
<p>Author: Millard et al., 1983</p> <p>Country and setting: Australia, Hospital</p> <p>Enrollment period: NR</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Report of hospital-based results</p> <p>Intervention: All patients initially hospitalized for 5-14 days then assigned to a Frewen-type bladder training program or biofeedback, each lasting 3 months. Adjunctive therapy* was used when patients did not show improvement.</p> <p>Groups: NR</p> <p>N at enrollment: 59</p> <p>N at follow-up: 59</p> <p>Women, %: 100</p> <p>N at follow-up: 39</p> <p>Age, mean (range): 49 (14-74)</p> <p>Race/ethnicity: NR</p>	<p>Inclusion criteria*:</p> <ul style="list-style-type: none"> Men, women or children with frequency, urgency, nocturia and urge incontinence <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Neurological lesions 	<p>Frequency, n: 59</p> <p>Nocturia, n: 37</p> <p>UUI, n: 42</p> <p>MUI, n: 24</p> <p>Enuresis, n: 12</p> <p>Giggle, n: 9</p> <p>Urodynamic status, n: Unstable bladder: 38 Unstable bladder & bladder neck incompetence: 6 Sensory urgency: 12 Sensory urgency & bladder neck incompetence: 3</p> <p>Duration of symptoms (years), mean: 11.9</p>	<p>Results, patients with unstable bladder, n: Cure: 18 Significant improvement: 10 Minor improvement: 2 Failure: 8</p> <p>Results, patients with unstable bladder & bladder neck incompetence, n: Cure: 0 Significant improvement: 3 Minor improvement: 1 Failure: 0</p> <p>Results, patients with sensory urgency, n: Cure: 7 Significant improvement: 4 Minor improvement: 1 Failure: 0</p> <p>Results, patients with sensory urgency & bladder neck incompetence, n: Cure: 0 Significant improvement: 0 Minor improvement: 0 Failure: 3</p> <p>Bladder Capacity (mL), by result, patients with unstable bladder, mean change: Cure: 149 Significant improvement: 87 Minor</p>	<p>Quality: Overall quality score: poor INTERNAL VALIDITY: poor Randomization: NA Masking: NA Pt selection criteria: - Loss to follow up: ++ Drop-out rates: NR Power calculation: - Statistical issues: - EXTERNAL VALIDITY: fair Age: + Baseline OAB status: + Baseline characteristics: ++ Length of follow up: + Measurement methods: - Measurement reliability: + Intervention description: +</p>

Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
				improvement: 95 Failure: 73	

Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Millard et al., 1983 (continued)				<p>Bladder Capacity (mL) by result, patients with sensory urgency, mean change: Cure: 83 Significant improvement: 115 Minor improvement: 127 Failure: NA</p> <p>Voiding frequency (hours), by result, patients with unstable bladder, initial/final: Cure: 1.9/4.8 Significant improvement: 1.8/3.2 Minor improvement: 1.0/4.0 Failure: 1.4/3.3</p> <p>Voiding frequency (hours), by result, patients with sensory urgency, initial/final: Cure: 1.7/3.8 Significant improvement: 1.0/3.4 Minor improvement: 3.3/4.0 Fail: NA</p>	

Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)					
<p>Author: Sand et al., 2007</p> <p>Country and setting: US, Multicenter, 327 sites (141 Urology 141, 96 Primary care, 43 Ob-Gyn, 17 Geriatric Med)</p> <p>Enrollment period: NR</p> <p>Funding: Watson Laboratories, Inc</p> <p>Author industry relationship disclosures: 7 of 7 Allergan (1) Astellas (1) Esprit (1) GlaxoSmithKline (2) GSK (1) GTx (1) Indevus-Esprit (1) Lilly (1) Novartis (3) Ortho-McNeil (1) Ortho Urology (1) Pfizer (4) Sanofi (1) USB (1) Watson (6)</p>	<p>Design: RCT, open-label (case series for drug, RCT for educational materials)</p> <p>Intervention: Oxybutynin transdermal system 3.9 mg/day, twice weekly patch for up to 6 months with “standard instruction” or with “educational intervention”.</p> <p>Groups: G1: Standard instructions G2: Educational intervention (educational booklet, OAB newsletters, dosing reminders, calendar reminders, bladder diary)</p> <p>N at enrollment: Total: 2,878 G1: 1,282 G2: 1,596</p> <p>Women, %: 87.2</p> <p>Age, yrs ± SD: 62.5 ± 14.8</p> <p>Race/ethnicity, %: White: 83.6 African American: 9.9 Hispanic: 4.8 Asian: 1.2 Other: 0.5</p> <p>Follow-up: 6 months</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥18 • UUI, urinary urgency or frequency • Willing to discontinue all prescription and OTC medications for OAB • Capable of completing QoL questionnaires without assistance • Negative pregnancy test & medically acceptable contraceptive <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Contraindications to oxybutynin • Reversible etiologies for OAB • Prior treatment with Oxytrol • Long-term care facilities and nursing homes 	<p>KHQ, UUI severity item score, %: A lot: 38.1 Moderately: 31.3 A little: 20.5 Omitted, N/A: 10.1</p> <p>KHQ, urgency severity item score, %: A lot: 49.2 Moderately: 30.3 A little: 14.8 Omitted, N/A: 5.6</p> <p>KHQ, frequency severity item score, %: A lot: 59.3 Moderately: 32.5 A little: 7.2 Omitted, N/A: 1.0</p> <p>KHQ, nocturia severity item score, %: A lot: 47.3 Moderately: 33.5 A little: 16.7 Omitted, N/A: 2.4</p> <p>History of OAB, years, %: <1 year: 12.0 1-2 years: 18.5 2-4 years: 23.1 ≥ 4 years: 46.4</p> <p>OAB Severity: No problem: 1.8 Very minor: 4.6 Minor: 15.5 Moderate: 33.0 Severe: 28.4 Many severe problems: 16.7</p> <p>Previous OAB treatments, n (%): 0: 43.4 1: 38.7 2: 13.1 3: 2.9</p>	<p>KHQ, UUI severity item score, %: A lot: 21.3 Moderately: 29.5 A little: 31.6 Omitted, N/A: 17.7</p> <p>KHQ, urgency severity item score, %: A lot: 27.3 Moderately: 33.8 A little: 27.3 Omitted, N/A: 11.5</p> <p>KHQ, frequency severity item score, %: A lot: 33.5 Moderately: 45.2 A little: 18.7 Omitted, N/A: 2.6</p> <p>KHQ, nocturia severity item score, %: A lot: 27.5 Moderately: 35.4 A little: 31.3 Omitted, N/A: 5.7</p> <p>KHQ, general health perception, mean ± SD (% improvement): -1.2 ± 17.4 (-4.3)</p> <p>KHQ, incontinence impact, mean ± SD (% improvement): -13.5 ± 29.5 (-19.5)</p> <p>KHQ, symptom severity, mean ± SD (% improvement): -12.4 ± 24.8 (-22.2)***</p> <p>KHQ, role limitations, mean ± SD (%)</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: -</p> <p>Masking: NA</p> <p>Pt selection criteria: +</p> <p>Loss to followup: -</p> <p>Drop-out rates: -</p> <p>Power calculation: +</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: ++</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
			4: 1.9	improvement): -13.3 ± 29.2 (-29.5)	

Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Sand et al., 2007 (continued)			<p>KHQ, general health perception, mean ± SD: 28.2 ± 19.8</p> <p>KHQ, incontinence Impact, mean ± SD: 69.3 ± 27.4</p> <p>KHQ, symptom severity, mean ± SD: 55.9 ± 20.5*</p> <p>KHQ, role limitations, mean ± SD: 45.1 ± 31.0</p> <p>KHQ, physical limitations, mean ± SD: 46.7 ± 31.6</p> <p>KHQ, social limitations, mean ± SD: 25.6 ± 28.3</p> <p>KHQ, emotions, mean ± SD: 30.0 ± 29.2</p> <p>KHQ, personal relationships, mean ± SD: 20.6 ± 29.5***</p> <p>KHQ, sleep/energy, mean ± SD: 54.2 ± 27.3</p> <p>KHQ, severity (coping) measures, mean ± SD: 47.9 ± 26.4</p>	<p>KHQ, physical limitations, mean ± SD (% improvement): -11.7 ± 29.9 (-25.1)</p> <p>KHQ, social limitations, mean ± SD (% improvement): -6.7 ± 23.7 (-26.2)</p> <p>KHQ, emotions, mean ± SD (% improvement): -8.8 ± 25.4 (-29.3)</p> <p>KHQ, personal relationships, mean ± SD (% improvement): -6.0 ± 23.5 (-29.1)***</p> <p>KHQ, sleep/energy, mean ± SD (% improvement): -11.2 ± 24.1 (-20.7)</p> <p>KHQ, severity (coping) measures, mean ± SD (% improvement): -8.6 ± 21.3 (-18.0)</p> <p>Side effects, application site, %: Total: 14.0 Pruritis: 4.9 Erythema: 4.6 Dermatitis: 4.4 Irritation: 3.2 Other: 2.0</p> <p>Side effects, %: Rash: 3.0 Dry mouth: 2.6 Pruritis: 2.6 Skin irritation: 2.1</p> <p>Withdrew due to AEs, %: 21.3</p>	

Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Sand et al., 2007 (continued)				<p>Withdrew: 1452 (50.5%)</p> <p>Adverse events, %: 21.3</p> <p>Withdrawn consent, %: 7.5</p> <p>Requirement for alternative therapy, %: 7.4</p> <p>Loss to follow-up, %: 7.2</p> <p>Noncompliance, %: 5.6</p> <p>Administrative decision, %: 0.7</p> <p>Ineligible, %: 0.4</p> <p>Death, %: 0.1</p> <p>No reason given, %: 0.3</p>	

Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)					
<p>Author: Song, et al., 2006</p> <p>Country and setting: Korea, Medical Center</p> <p>Enrollment period: May 2001 to April 2002</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: Bladder training (BT) vs. Tolterodine vs. BT + Tolterodine</p> <p>Groups: G1: BT x 12 wks G2: Tolterodine 2 mg bid x 12 wks G3: Tolterodine 2 mg bid + BT x 12 wks</p> <p>N at enrollment: G1: 46 G2: 47 G3: 46</p> <p>N at follow-up: G1: 26 G2: 32 G3: 31</p> <p>Women, %: 100</p> <p>Age, mean ± SD: G1: 45.73 ± 12.68 G2: 48.41 ± 9.38 G3: 45.42 ± 9.54</p> <p>Race/ethnicity, %: Korean: 100</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 18 • ≥ 8 voids/day • Urge with or without incontinence • Symptom duration ≥ 3 months • No prior history of treatment for OAB <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Active urinary tract infection • Clinically significant SUI • Bladder outlet obstruction • Interstitial cystitis • Glaucoma • Megacolon • Maximal urine flow rate of < 10 mL/sec • Postvoid residual volume > 30% of the total amount voided on uroflowmetry 	<p>Voids/day, mean ± SD: G1: 10.93 ± 2.14 G2: 11.63 ± 2.57 G3: 11.90 ± 1.51</p> <p>Nocturia episodes/day, mean ± SD: G1: 1.45 ± 1.14 G2: 1.72 ± 1.04 G3: 1.96 ± 1.49</p> <p>Urgency, mean score ± SD: G1: 2.58 ± 1.30 G2: 2.81 ± 0.74 G3: 3.00 ± 1.10</p> <p>Maximum flow rate (mL/s), mean ± SD: G1: 20.35 ± 8.44 G2: 22.56 ± 4.94 G3: 21.19 ± 4.96</p> <p>Residual urine (mL), mean ± SD: G1: 9.08 ± 22.56 G2: 7.59 ± 12.39 G3: 6.42 ± 10.16</p> <p>Symptom duration (years), mean ± SD: G1: 6.44 ± 6.84 G2: 4.54 ± 5.15 G3: 4.10 ± 3.99</p>	<p>Voids/day, mean (% decrease) G1: 8.1 (25.9%)* G2: 8.1 (30.2%)* G3: 7.9 (33.5%)* G3/G1: P < 0.05</p> <p>Nocturia episodes/day, mean (% reduction): G1: 0.6 (56.1%)* G2: 0.6 (65.4%)* G3: 0.6 (66.3%)*</p> <p>Urgency, mean score (% reduction): G1: 1.4 (44.8%)* G2: 1.1 (62.2%)* G3: 1.2 (60.2%)* G3/G1: P = 0.021 G2/G1: P = 0.017 G2/G3: P = NS</p> <p>Satisfaction, mean score (% improved): G1: 1.5 (53.9) G2: 1.4 (63.0) G3: 1.3 (71.0)</p> <p>Dry mouth, n (%): G1: 0 (0.0) G2: 7 (21.9) G3: 9 (28.9)</p> <p>Hesitancy, n (%): G1: 0 (0.0) G2: 3 (9.4) G3: 2 (6.5)</p> <p>Decreased appetite/constipation, n (%): G1: 0 (0.0) G2: 2 (6.3) G3: 2 (6.5)</p> <p>Headache, n (%): G1: 0 (0.0) G2: 1 (3.1) G3: 0 (0.0)</p>	<p>Quality: Overall quality score: fair INTERNAL VALIDITY: poor Randomization: - Masking: - Pt selection criteria: + Loss to follow up: - Drop-out rates: + Power calculation: - Statistical issues: - EXTERNAL VALIDITY: good Age: + Baseline OAB status: + Baseline characteristics: ++ Length of follow up: + Measurement methods: + Measurement reliability: + Intervention description: +</p>

Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
--------------------------	--	--------------------------------------	--------------------------------	-----------------	-----------------------

Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics*	Outcomes	Quality Rating
--------------------------	--	--------------------------------------	---------------------------------	-----------------	-----------------------

Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics*	Outcomes	Quality Rating
<p>Author: Szonyi et al., 1995</p> <p>Country and setting: UK</p> <p>Enrollment period: NR</p> <p>Funding: Smith & Nephew Pharmaceuticals Ltd</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: Oxybutynin + bladder training vs. placebo + bladder training</p> <p>Groups: G1: Oxybutynin 2.5 mg bid with dose titration on days 29 and 43 plus bladder training G2: Placebo + bladder training</p> <p>N at enrollment: G1: 30 G2: 30</p> <p>N at follow-up: G1: 16 G2: 23</p> <p>Women, n (%): 56 (93)</p> <p>Age, mean ± SD: 82.2 ± 6.06</p> <p>Race/ethnicity: NR</p> <p>Weight (kg), mean ± SD: 67.4 ± 14.92</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age > 70 • Frequency, urgency and UUI • Mobile • Able to keep diary <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • UTI • Hepatic or renal disease • Glaucoma • Uncontrolled diabetes • Taking imipramine or propantheline 		<p>Voids/2 weeks, median change (95% CI): G1/G2: 577 (-27.0, 6.0) <i>P</i> = 0.0025</p> <p>Nocturia episodes/2 weeks, median change (95% CI): G1/G2: -6 (-5, 7.0)</p> <p>Daytime incontinence episodes/2 weeks, median change (95% CI): G1 vs. G2: -9.5 (-11.0, 3.0)</p> <p>Nocturia episodes/2 weeks, median change (95% CI): G1/G2: -1.0 (-3.0, 2.0)</p> <p>Patient assessment of benefit, %:</p> <p>29 days: G1: 86 G2: 55 <i>P</i> = 0.02</p> <p>43 days: G1: 71 G2: 59 <i>P</i> = 0.41</p> <p>57 days: G1: 79 G2: 55 <i>P</i> = 0.09</p> <p>Patient response, 29 days, n:</p> <p>Cure: G1: 1 G2: 0</p> <p>Significant improvement: G1: 15 G2: 8</p> <p>Marginal improvement: G1: 7 G2: 8</p>	<p>Quality:</p> <p>Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to follow up: -</p> <p>Drop-out rates: -</p> <p>Power calculation: +</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: +</p> <p>Length of follow up: -</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics*	Outcomes	Quality Rating
Szonyi et al., 1995 (continued)				<p>No change: G1: 5 G2: 13</p> <p>Patient response, 57 days, n: Cure: G1: 4 G2: 3 Significant improvement: G1: 14 G2: 8 Marginal improvement: G1: 3 G2: 4 No change: G1: 7 G2: 14</p> <p>Dry mouth, %: G1: 93 G2: 86</p> <p>Blurred vision, %: G1: 50 G2: 59</p> <p>Heartburn, %: G1: 57 G2: 45</p> <p>Constipation, %: G1: 50 G2: 45</p> <p>Dry skin, %: G1: 50 G2: 59</p> <p>Poor compliance (< 75% of tablets), %: G1: 20 G2: 20</p>	

Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics*	Outcomes	Quality Rating
Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)					
<p>Author: Wang et al., 2004</p> <p>Country and setting: Taiwan, NR</p> <p>Enrollment period: July 2001 to December 2002</p> <p>Funding: National Science Council, Taiwan</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: Pelvic floor muscle training (PFMT), biofeedback-assisted PFMT (BAPFMT), electrical stimulation (ES)</p> <p>Groups: G1: PFMT per PERFECT score G2: BAPFMT per PERFECT score G3: ES using biphasic symmetric probe current w/ 10-Hz freq, 400-microsec pulse width, 10/5 duty cycle, varying intensity</p> <p>N at enrollment: G1: 40 G2: 38 G3: 42</p> <p>N at follow-up: G1: 34 G2: 34 G3: 35</p> <p>Women: 100%</p> <p>Age, mean ± SD: G1: 50.09 ± 15.85 G2: 52.32 ± 12.68 G3: 55.74 ± 12.53</p> <p>Race/ethnicity: NR</p> <p>BMI, kg/m² ± SD: G1: 22.69 ± 3.32 G2: 23.70 ± 3.90 G3: 23.93 ± 3.21</p> <p>Parity, mean ± SD: G1: 2.47 ± 1.44</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Female • Age 16-75 • OAB symptoms for > 6 months • Voids/day ≥ 8 • UUI/day ≥ 1 <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Anticholinergic medications • Tricyclic antidepressants • Treatment with pelvic-floor exercises • Treatment with bladder training • Repair of pelvic prolapse • Pregnancy • Deafness • Neurologic disorders • DM • Pacemaker or IUD use • Genital prolapse > stage II by ICS • Residual urine > 100 mL • UTI 	<p>Incontinence episodes/day, mean ± SD: G1: 0.86 ± 1.80 G2: 0.92 ± 1.77 G3: 2.09 ± 2.96</p> <p>Urine leakage during former pregnancy, no (%): G1: 7 (20.59) G2: 3 (8.83) G3: 3 (8.57)</p>	<p>UUI symptoms resolved, n (%): G1: 10 (30.3) G2: 13 (38.24) G3: 14 (40)</p> <p>UUI symptoms modified, n (%): G1: 3 (6.06) G2: 4 (11.76) G3: 4 (11.43)</p> <p>UUI symptoms unchanged, n (%): G1: 21 (63.64) G2: 17 (50) G3: 17 (48.6)</p> <p>Incontinence episodes/day, mean ± SD: G1: 0.73±1.82 G2: 0.70±1.80 G3: 1.95±2.84 G1/G2/G3: P = 0.016</p> <p>OAB symptom improvement/ cure, %: G1: 38.2 G2: 50.0 G3: 51.4 G1/G2/G3: P = 0.567</p> <p>Adherence to treatment, median % (range): G1: 0.833 (0.25-1.00) G2: 0.791 (0.58-1.99) G3: 0.750 (0.54-1.00) G1/G2/G3: P = 0.356</p> <p>Compliance with home program (days), median (range):</p>	<p>Quality: Overall quality score: fair INTERNAL VALIDITY: poor Randomization: + Masking: - Pt selection criteria: + Loss to follow up: + Drop-out rates: - Power calculation: + Statistical issues: - EXTERNAL VALIDITY: good Age: + Baseline OAB status: + Baseline characteristics: ++ Length of follow up: + Measurement methods: + Measurement reliability: + Intervention description: +</p>

Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics*	Outcomes	Quality Rating
	G2: 2.91 ± 1.86 G3: 3.69 ± 1.75			G1: 8.5 (0-44) G2: 14.5 (0-44) G1/G2: <i>P</i> = 0.636	

Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics*	Outcomes	Quality Rating
Wang et al., 2004 (continued)	Menopausal, n (%) G1: 21 (61.76) G2: 17 (50) G3: 8 (22.86)			<p>KHQ, general health perception, mean change ± SD: G1: 14.66 ± 24.57 G2: 12.10 ± 20.28 G3: 16.96 ± 24.58 G1/G2/G3: P = 0.376</p> <p>KHQ, incontinence impact, mean change ± SD: G1: 6.03 ± 120.82 G2: 37.42 ± 37.11 G3: 47.03 ± 35.45 G1/G2/G3: P = 0.067</p> <p>KHQ, role limitation, mean change ± SD: G1: 25.86 ± 26.57 G2: 30.64 ± 30.15 G3: 34.52 ± 31.73 G1/G2/G3: P = 0.376</p> <p>KHQ, physical limitation, mean change ± SD: G1: 25.29 ± 26.58 G2: 33.33 ± 33.88 G3: 28.57 ± 28.99 G1/G2/G3: P = 0.693</p> <p>KHQ, social limitation, mean change ± SD: G1: 17.05 ± 21.20 G2: 22.76 ± 29.20 G3: 20.84 ± 27.45 G1/G2/G3: P = 0.799</p> <p>KHQ, personal relationships, mean change ± SD: G1: 2.30 ± 13.89 G2: 10.75 ± 26.37 G3: 3.57 ± 22.39 G1/G2/G3: P = 0.167</p>	

Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics*	Outcomes	Quality Rating
Wang et al., 2004 (continued)				<p>KHQ, emotions, mean change ± SD: G1: 19.31 ± 29.68 G2: 22.22 ± 27.82 G3: 46.83 ± 37.33 G1/G2/G3: <i>P</i> = 0.005 G2/G3: <i>P</i> = 0.003 G1/G3: <i>P</i> = 0.007 G2/G1: <i>P</i> = 0.751</p> <p>KHQ, sleep/energy, mean change ± SD: G1: 18.83 ± 26.18 G2: 26.88 ± 24.60 G3: 38.10 ± 39.51 G1/G2/G3: <i>P</i> = 0.249</p> <p>KHQ, severity measures, mean change ± SD: G1: 14.71 ± 20.27 G2: 20.65 ± 31.19 G3: 31.23 ± 23.83 G1/G2/G3: <i>P</i> = 0.004 G2/G3: <i>P</i> = 0.029 G1/G3: <i>P</i> = 0.001 G2/G1: <i>P</i> = 0.587</p> <p>KHQ, total score, mean change ± SD: G1: 50.27 ± 171.42 G2: 185.86 ± 176.57 G3: 180.08 ± 176.03 G1/G2/G3: <i>P</i> = 0.003 G2/G3: <i>P</i> = 0.952 G1/G3: <i>P</i> = 0.004 G2/G1: <i>P</i> = 0.003</p> <p>Pelvic muscle strength, scale of power: G2/G3: <i>P</i> < 0.001 G1/G3: <i>P</i> < 0.001</p>	

Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics*	Outcomes	Quality Rating
Wang et al., 2004 (continued)				<p>Times of fast contraction, mean \pm SD: G1: -5.82 \pm 5.05 G2: -6.21 \pm 4.56 G3: -3.03 \pm 4.98 G2/G3: $P = 0.007$ G1/G3: $P = 0.012$</p> <p>Degree of vaginal pressure, mean change \pm SD: G1: -36.03 \pm 21.79 G2: -38.35 \pm 29.62 G3: -8.91 \pm 12.83 G2/G3: $P < 0.001$ G1/G3: $P < 0.01$</p> <p>Vaginal pressure, mean % change (range): G1: -78.95 (-522.22, -10.34) G2: -105.0 (-2450 to 79.55) G3: -12.63 (-138.46 to 50) G1/G2/G3: $P < 0.001$</p> <p>Vaginal pressure, mean change \pm SD: G1: -36.03 \pm 21.79 G2: -38.35 \pm 29.62 G3: -8.91 \pm 12.83 G2/G3: $P < 0.001$ G1/G3: $P < 0.001$ G2/G1: $P = NS$</p>	

Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics*	Outcomes	Quality Rating
<p>Author: Wang et al., 2006</p> <p>Country and setting: Taiwan; Academic medical center</p> <p>Enrollment period: July 2004 to November 2005</p> <p>Funding: National Science Council, Taiwan</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: Electric Stimulation (ES) vs. Oxybutynin vs. Placebo for 12 weeks</p> <p>Groups: G1: ES: intravaginal electrode; biphasic symmetric pulsed current w/ a 10-Hz frequency, 400-millisecond pulse width, 10/5 duty cycle, and varying intensity, 20 min/session, twice weekly G2: Oxybutynin, 2.5 mg t.i.d. G3: Placebo pill t.i.d.</p> <p>N at enrollment: G1: 25 G2: 26 G3: 23</p> <p>N at follow-up: G1: 24 G2: 23 G3: 21</p> <p>Age: NR</p> <p>Race/ethnicity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> Female OAB for ≥ 6 mos Age 16-80 Urgency ≥ 4 times per day <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Use of anticholinergics or TCAs Previous treatment with pelvic-floor muscle training, bladder training Pelvic prolapse repair Pregnancy Neurologic disorders DM Demand cardiac pacemaker IUD use Genital prolapse greater than ICS Stage II PVR>100 mL Overt SI Anti-incontinence surgery UTI 	<p>Warning time, s (range) G1: 41.5 (8-105) G2: 44 (2-215) G3: 65 (26-265)</p> <p>MVV, mL/void (range): G1: 340 (120-450) G2: 310 (130-800) G3: 350 (120-600)</p> <p>Daily voided volume, mL (range) G1: 2160 (1010-2950) G2: 2106 (1560-3153) G3: 2305 (1305-3300)</p> <p>Pads per day (range): G1: 1 (0-4.1) G2: 0 (0-3) G3: 1 (0-4)</p> <p>Urgency (subj): G1: 11.4 (4.0-16.0) G2: 12.1 (5-18) G3: 9.8 (4.2-15.5)</p> <p>Frequency (subj): G1: 12.8 (8.8-16.0) G2: 11.5 (4.3-19.5) G3: 11.5 (6.5-22.8)</p> <p>Nocturia (subj): G1: 1.75 (0-6.5) G2: 0 (0-2.5) G3: 0.65 (0-3.1)</p> <p>UI (subj): G1: 1 (0-2) G2: 0 (0-2) G3: 1 (0-2)</p>	<p>Warning time, s (range) G1: 72 (32-633) G2: 54.5 (138-850) G3: 66.5 (26-219) <i>P</i>(G1-G3) < 0.001</p> <p>MVV, mL/micturition (range): G1: 355 (150-550) G2: 336.5 (138-850) G3: 340 (160-600) <i>P</i>(G1-G3) = 0.035</p> <p>Voided volume/d, mL (range) G1: 2270 (1210-3106) G2: 2100 (1619-3200) G3: 2305 (1351-3221) <i>P</i>(G1-G3) = 0.050</p> <p>Pad count, pad/d (range): G1: 0 (0-2) G2: 0 (0-2.5) G3: 1 (0-3) <i>P</i>(G1-G3) = 0.012</p> <p>Urgency (subj) G1: 1.0 (0.0-12.3) G2: 6 (0.5-13) G3: 7.4 (3.9-13.4) <i>P</i>(G1-G3) < 0.001</p> <p>Frequency (subj) G1: 7.8 (1.8-13.0) G2: 7.4 (2-14) G3: 10 (3.9-13.4) <i>P</i>(G1-G3) = 0.002</p> <p>Nocturia (subj) G1: 0 (0-3.0) G2: 0 (0-2.5) G3: 1 (0-3.6) <i>P</i>(G1-G3) = 0.002</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Masking: -</p> <p>Pt selection criteria: +</p> <p>Loss to followup: ++</p> <p>Drop-out rates: +</p> <p>Power calculation: +</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Wang et al., 2006 (continued)				<p>UI (subj) G1: 0.5 (0-2) G2: 0 (0-2) G3: 1 (0-2) $P(G1-G3) = 0.413$</p> <p>P values before vs. after treatment</p> <p>Warning time, s (range): G1: 0.002 G2: 0.001 G3: 0.532</p> <p>MVV, mL/void (range): G1: 0.018 G2: 0.004 G3: 0.979</p> <p>Daily voided volume, mL (range): G1: 0.024 G2: 0.728 G3: 0.627</p> <p>Pad per day (range): G1: 0.010 G2: 0.662 G3: 0.501</p> <p>Urgency (subj): G1: <0.001 G2: <0.001 G3: 0.003</p> <p>Frequency (subj): G1: <0.001 G2: <0.001 G3: 0.070</p> <p>Nocturia (subj): G1: 0.001 G2: 0.394 G3: 0.176</p> <p>UI (subj): G1: 0.814 G2: 0.083 G3: 0.854</p>	

Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Wyman et al. 1998</p> <p>Country and setting: US, Academic medical center</p> <p>Enrollment period: NR</p> <p>Funding: NIH</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: 12-wk program of patient education, self-monitoring of voiding behaviors, compliance assessment, and positive-reinforcement techniques administered by trained RNs</p> <p>Groups: G1: Bladder Training (BT) G2: Pelvic muscle exercise (PME) G3: Combination therapy (CT)</p> <p>N at enrollment: Total: 204 G1: 68 G2: 69 G3: 67</p> <p>N at follow-up, 3 mos: G1: 68 G2: 64 G3: 61</p> <p>Age, mean ± SD: G1: 60 ± 10 G2: 62 ± 10 G3: 61 ± 9</p> <p>Race/ethnicity, n (%): White: G1: 64 (94) G2: 62 (90) G3: 49 (91)</p> <p>Parity (vaginal births), mean ± SD: G1: 2.4 ± 1.7 G2: 2.6 ± 1.9 G3: 2.5 ± 1.7 <i>P</i> = 0.846</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 45 • Community dwelling • Mentally intact • Able to toilet independently • Urine loss ≥1x/wk • Urodynamic evidence of GUI, DI or both <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Reversible cases of UI • Uncontrolled metabolic conditions • Residual urine volume after voiding > 100mL • UTI • Genitourinary fistula • Indwelling catheterization • Inability to correctly perform a pelvic muscle contraction on digital exam 	<p>Stress urinary incontinence, n (%): G1: 19 (28) G2: 35 (51) G3: 22 (33) <i>P</i> = 0.015</p> <p>Urge urinary incontinence, n (%): G1: 8(12) G2: 6(9) G3: 10(15)</p> <p>Mixed urinary incontinence, n (%): G1: 41 (60) G2: 27 (39) G3: 35 (52) <i>P</i> = 0.044</p> <p>Genuine stress incontinence by urodynamics, n (%): G1: 48 (71) G2: 48 (70) G3: 49 (73)</p> <p>Detrusor Instability by urodynamics, n (%): G1: 9 (13) G2: 10 (14) G3: 10 (15)</p> <p>DI + GSI by urodynamics, n (%): G1: 11 (16) G2: 11 (16) G3: 8 (12)</p> <p>Incontinence episodes/wk, mean ± SD: G1: 14.6 ± 11.2 G2: 16.8 ± 17.1 G3: 14.9 ± 13.8</p>	<p>Incontinence episodes/wk, immediately after treatment, mean ± SD: G1: 10.6 ± 16.3 G2: 9.6 ± 10.8 G3: 6.8 ± 10.7 <i>P</i> = 0.004 G1/G2: <i>P</i> = 0.796 G1/G3: <i>P</i> = 0.006 G2/G3: <i>P</i> = 0.003</p> <p>Incontinence episodes/wk, immediately after treatment, DI ± GSI subgroup, mean ± SD: G1: 6.2 ± 9.1 G2: 11.9 ± 12.7 G3: 5.8 ± 9.5</p> <p>Incontinence episodes, % reduction, immediately after treatment, n (%): 100%: G1: 12 (18) G2: 8 (13) G3: 19 (31) 75-99%: G1: 11 (16) G2: 8 (13) G3: 14 (23) 50-74%: G1: 12 (18) G2: 20 (31) G3: 10 (16) 0-49%: G1: 14 (21) G2: 13 (20) G3: 10 (16) None/worse: G1: 19 (28) G2: 15(23) G3: 8 (13) <i>P</i> = 0.050</p> <p>Incontinence episodes/wk, 3 mos, mean ± SD: G1: 10.0 ± 12.0 G2: 9.4 ± 14.0 G3: 8.1 ± 12.4 <i>P</i> = 0.126</p>	<p>Quality: Overall quality score: fair INTERNAL VALIDITY: poor Randomization: - Masking: - Pt selection criteria: + Loss to follow up: ++ Drop-out rates: - Power calculation: + Statistical issues: + EXTERNAL VALIDITY: good Age: + Baseline OAB status: + Baseline characteristics: ++ Length of follow up: + Measurement methods: + Measurement reliability: + Intervention description: +</p>

Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Wyman et al. 1998 (continued)	BMI, kg/m ² ± SD: G1: 28.4 ± 6.6 G2: 27.6 ± 5.8 G3: 26.8 ± 5.6 P = 0.288		Incontinence episodes/wk, DI ± GSI subgroup, mean ± SD: G1: 14.3 ± 10.3 G2: 20.7 ± 22.4 G3: 16.2 ± 13.7 Previous medical treatment for UI, n (%): G1: 14 (21) G2: 14 (21) G3: 16 (24) Previous surgical treatment for UI, n (%): G1: 15 (22) G2: 14 (21) G3: 22 (33) UDI QoL, mean ± SD: G1: 130.1 ± 48.8 G2: 119.7 ± 49.9 G3: 118.9 ± 46.9 UDI QoL, DI ± GSI subgroup, mean ± SD: G1: 143.2 ± 54.0 G2: 133.2 ± 59.6 G3: 115.4 ± 42.5 IIQ-R incontinence impact, mean ± SD: G1: 93.7 ± 74.2 G2: 75.6 ± 67.1 G3: 84.6 ± 67.8 IIQ-R incontinence impact, DI ± GSI subgroup, mean ± SD: G1: 118.5 ± 84.4 G2: 93.7 ± 90.3 G3: 84.0 ± 66.9	Incontinence episodes, % reduction, 3 mos, n (%): 100%: G1: 10 (16) G2: 13 (20) G3: 16 (27) 75-99%: G1: 9 (15) G2: 12 (22) G3: 13 (22) 50-74%: G1: 9 (15) G2: 9 (14) G3: 6 (10) 0-49%: G1: 15 (24) G2: 12 (18) G3: 15 (25) None/worse: G1: 19 (31) G2: 17 (26) G3: 10 (17) P = 0.587 UDI QoL, immediately after treatment, mean ± SD: G1: 95.5 ± 54.4 G2: 90.8 ± 52.0 G3: 64.4 ± 48.6 UDI QoL, immediately after treatment, DI ± GSI subgroup, mean ± SD: G1: 86.8 ± 54.8 G2: 114.8 ± 70.3 G3: 67.6 ± 48.5 P = 0.054 G1/G2: P = 0.37 G1/G3: P = 0.956 G2/G3: P = 0.038 UDI QoL, 3 months, mean ± SD: G1: 91.7 ± 55.0 G2: 85.0 ± 52.4 G3: 72.8 ± 50.4 P = 0.203	

Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Wyman et al., 1998 (continued)				<p>IIQ-R incontinence impact, immediately after treatment, mean \pm SD: G1: 72.1 \pm 75.2 G2: 56.8 \pm 61.4 G3: 46.6 \pm 65.3</p> <p>IIQ-R incontinence impact, immediately after treatment, DI \pm GSI subgroup, mean \pm SD: G1: 81.2 \pm 88.7 G2: 89.9 \pm 79.4 G3: 31.8 \pm 34.4 <i>P</i> = 0.0301 G1/G2: <i>P</i> = 0.357 G1/G3: <i>P</i> = 0.084 G2/G3: <i>P</i> = 0.009</p> <p>IIQ-R incontinence impact, 3 months, mean \pm SD: G1: 65.7 \pm 80.2 G2: 59.3 \pm 67.7 G3: 59.8 \pm 83.9 <i>P</i> = 0.850</p> <p>Patient perception of improvement, immediately after treatment, n (%): Much: G1: 25 (38) G2: 19 (30) G3: 32 (52) Some: G1: 18 (27) G2: 29 (46) G3: 23 (38) None: G1: 20 (30) G2: 13 (21) G3: 4 (7) Worse: G1: 3 (5) G2: 2 (3) G3: 2 (3) <i>P</i> = 0.011</p>	

Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Wyman et al., 1998 (continued)				<p>Patient perception of improvement, 3 mos, n (%): Much: G1: 21 (35) G2: 24 (37) G3: 31 (53) Some: G1: 16 (27) G2: 21 (33) G3: 13 (22) None: G1: 19 (32) G2: 16 (25) G3: 12 (21) Worse: G1: 4 (7) G2: 3 (5) G3: 2 (4) $P = 0.414$</p> <p>Patient satisfaction, immediately after treatment, n (%): Very satisfied: G1: 42 (64) G2: 46 (73) G3: 50 (82) Slightly satisfied: G1: 6 (9) G2: 10 (16) G3: 7 (11) Neither G1: 12 (21) G2: 6 (10) G3: 3 (5) Dissatisfied: G1: 4 (6) G2: 1 (2) G3: 1 (2)</p> <p>Patient satisfaction, 3 mos, n (%): Very satisfied: G1: 36 (60) G2: 42 (66) G3: 45 (78) Slightly satisfied: G1: 11 (18) G2: 11 (17) G3: 6 (10)</p>	

Evidence Table 4. KQ 2 Behavioral Treatment of OAB (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Wyman et al., 1998 (continued)				Neither: G1: 8 (13) G2: 10 (16) G3: 5 (9)	
				Dissatisfied: G1: 5 (8) G2: 1 (2) G3: 2 (3) P = 0.310	

Evidence Table 5. KQ2 Complementary and Alternative Therapies

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Emmons and Otto, 2005</p> <p>Country and setting: US, Academic medical center</p> <p>Enrollment period: July 2000 to October 2002</p> <p>Funding: Supported in part by grants from the Oregon Health Science Foundation and the Public Health Service</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: Acupuncture</p> <p>Groups: G1: acupuncture with needles placed at sites intended to address overactive bladder with UUI; administered for 20 minutes during 4 weekly sessions G2: acupuncture with needles placed at sites intended to support relaxation; administered for 20 minutes during 4 weekly sessions</p> <p>N at enrollment: G1: 44 G2: 41</p> <p>N at follow-up, 2-4 weeks: G1: 38 G2: 36</p> <p>Women, %: 100</p> <p>Age, mean ± SD: G1: 53 ± 13 G2: 51 ± 12</p> <p>Race/ethnicity: Minority, %</p> <p>Parity, mean ± SD: G1: 1.8 ± 1.6 G2: 1.8 ± 1.6</p> <p>Menopausal, n: G1: 21 G2: 20</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • > 8 voids per day • Subjective urgency to void • Urge-associated incontinence ≥ twice during a 3 day period <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Current treatment with medications for OAB • Current treatment with acupuncture for other indications • Unable to walk • Unable to complete a 3-day voiding diary • Hematuria • Untreated UTI 	<p>UUI episodes/3 days, mean ± SD: G1: 16.2 ± 11.1 G2: 15.4 ± 10.2 P = 0.68</p> <p>Incontinence episodes/3 days, mean ± SD: G1: 6.3 ± 7.3 G2: 8.9 ± 9.2 P = 0.09</p> <p>Voids/3 days, mean ± SD: G1: 30.4 ± 7.8 G2: 32.7 ± 11.5 P = 0.40</p> <p>Urinary distress inventory score, mean ± SD: G1: 8.4 ± 3.6 G2: 8.6 ± 5.5 P = 0.87</p> <p>Incontinence impact questionnaire score, mean ± SD: G1: 8.9 ± 2.8 G2: 9.1 ± 2.6 P = 0.80</p> <p>Functional bladder capacity (mL), mean ± SD: G1: 210 ± 88 G2: 199 ± 84 P = 0.50</p> <p>Cystometric max capacity (mL), mean ± SD: G1: 371 ± 161 G2: 341 ± 126 P = 0.49</p> <p>Cystometric volume (mL), first urge to void, mean ± SD: G1: 61 ± 67 G2: 69 ± 52 P = 0.84</p>	<p>UUI episodes/3 days, mean ± SD (% change): G1: 11.4 ± 8.8 (30) G2: 15.0 ± 9.4 (3) G1/BL: P < 0.003 G1/G2: P = 0.016</p> <p>Incontinence episodes/3 days, mean ± SD (% change): G1: 2.6 ± 3.1 (59) G2: 5.3 ± 5.9 (40) G1/BL: P < 0.003 G2/BL: P < 0.003 G1/G2: P = 0.71</p> <p>Voids/3 days, mean ± SD (% change): G1: 26.2 ± 7.1 (14) G2: 33.1 ± 16.1 (4) G1/G2: P = 0.03</p> <p>Urinary distress inventory score, mean ± SD (% change): G1: 3.6 ± 3.2 (57) G2: 5.8 ± 4.8 (32) G1/BL: P < 0.003 G2/BL: P < 0.003 G1/G2: P = 0.05</p> <p>Incontinence impact questionnaire score, mean ± SD (% change): G1: 4.3 ± 2.7 (52) G2: 7.0 ± 3.5 (23) G1/BL: P < 0.003 G2/BL: P < 0.003 G1/G2: P = 0.004</p> <p>Functional bladder capacity (mL), mean ± SD (% change): G1: 236 ± 99 (12) G2: 196 ± 85 (-2) G1/BL: P < 0.03 G1/G2: P = 0.01</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: fair</p> <p>Randomization: +</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: +</p> <p>Drop-out rates: ++</p> <p>Power calculation: +</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: -</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 5. KQ2 Complementary and Alternative Therapies (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Emmons and Otto, 2005 (continued)			Cystometric volume (mL), strong urge to void, mean ± SD: G1: 274 ± 144 G2: 241 ± 100 <i>P</i> = 0.26	Cystometric max capacity (mL), mean ± SD (% change): G1: 415 ± 205 (12) G2: 356 ± 193 (4) G1/G2: <i>P</i> = 0.049	
			Detrusor contractions during cystometry, n: G1: 7 G2: 11 <i>P</i> = 0.22	Cystometric volume (mL), first urge to void, mean ± SD (% change): G1: 65 ± 80 (6) G2: 57 ± 75 (-19) G1/G2: <i>P</i> = 0.47	
			Incontinence surgery, n: G1: 8 G2: 6	Cystometric volume (mL), strong urge to void, mean ± SD (% change): G1: 297 ± 167 (8) G2: 276 ± 156 (14) G1/BL: <i>P</i> < 0.03 G2/BL: <i>P</i> < 0.03 G1/G2: <i>P</i> = 0.75	
				Detrusor contractions during cystometry, n (% change): G1: 6 (3) G2: 10 (3) G1/G2: <i>P</i> = 0.98	
				Bleeding/bruising from the needles, %: 23	
				Minor discomfort with needle placement, %: 25	

Evidence Table 5. KQ2 Complementary and Alternative Therapies (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Freeman and Baxby, 1982</p> <p>Country and setting: UK, Academic medical center</p> <p>Enrollment period: NR</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Case series</p> <p>Intervention: 12 sessions of hypnotherapy</p> <p>Groups: NA</p> <p>N at enrollment: 63</p> <p>N at follow-up: 44</p> <p>Women, %: 100</p> <p>Age, mean ± SD: 44.0 ± 15.6</p> <p>Race/ethnicity: NR</p> <p>Follow-up: 3-14 months</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Women • Detrusor instability by urodynamics • Normal cystoscopy • Normal neurological findings • No drugs that affect the detrusor <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Genuine stress incontinence • Flow rate < 15 ml/s • Voiding pressure > 70 cm H₂O 	<p>Max cystometric capacity (mL), full study population, mean ± SD (range): 410.0 ± 13.3 (120-650)</p> <p>Max cystometric capacity (mL), urodynamic studies performed, mean ± SD: 378 ± 126</p> <p>Height of unstable contraction (cm H₂O), full study population, mean ± SD (range): 43.6 ± 22.1 (15-100)</p> <p>Height of unstable contraction (cm H₂O), urodynamic studies performed, mean ± SD: 57.5 ± 23.3</p> <p>Bladder volume (mL), first unstable contraction, full study population, mean ± SD (range): 318.0 ± 14.4 (50-650)</p> <p>Bladder volume (mL), first unstable contraction, urodynamic studies performed, mean ± SD: 245 ± 159</p>	<p>Max cystometric capacity (mL), mean ± SD: 487 ± 117</p> <p>Max cystometric capacity (mL), mean change ± SD: 92.3 ± 112.2 P < 0.001</p> <p>Height of unstable contraction (cm H₂O), mean ± SD: 39.0 ± 13.2</p> <p>Height of unstable contraction difference (cm H₂O), mean change ± SD: -18.5 ± 18.9 P < 0.01</p> <p>Bladder volume (mL), first unstable contraction, mean ± SD: 337 ± 123</p> <p>Bladder volume (mL), first unstable contraction, mean change ± SD: 92.2 ± 126 P < 0.02</p> <p>Symptom free, n: 29</p> <p>Considerably improved symptoms, n: 14</p> <p>No change in symptoms, n: 7</p> <p>Relapse, n: 1 year: 3 10 months: 1 8 months: 1 7 months: 1 2 months: 1</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: NA</p> <p>Masking: NA</p> <p>Pt selection criteria: +</p> <p>Loss to followup: -</p> <p>Drop-out rates: -</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: -, NR</p> <p>Baseline OAB status: NR</p> <p>Baseline characteristics: -</p> <p>Length of followup: ++</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 5. KQ2 Complementary and Alternative Therapies (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Mak et al., 2007</p> <p>Country and setting: Hong Kong, China, Gynecological hospital</p> <p>Enrollment period: January 2003 to March 2003</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: Foot reflexology vs nonspecific foot massage</p> <p>Groups: G1: foot reflexology for 45 min G2: nonspecific foot massage for 45 min</p> <p>N at enrollment: G1: 60 G2: 60</p> <p>N at follow-up: G1: 54 G2: 43</p> <p>Women, %: 100</p> <p>Age, mean ± SD: G1: 56.0 ± 13.4 G2: 56.4 ± 9.1</p> <p>Race/ethnicity: NR</p> <p>Menopausal, n (%): G1: 33 (57.8) G2: 29 (67.4) <i>P</i> = 0.533</p> <p>BMI, kg/m², median (IQR): G1: 24.2 (21.6, 27.4) G2: 24.7 (21.8, 26.7)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> Majority of leakage related to UI Taking antimuscarinics with required 3 week washout <p>Exclusion criteria:</p> <ul style="list-style-type: none"> SUI UTI Interstitial cystitis Urinary tract obstruction Organic disease of bladder or urethra Pregnant Breast-feeding Pelvic, vaginal, or bladder surgery within 6 mos Heart disease Abnormal EKG 	<p>UUI episodes, median (IQR): G1: 0.5 (0, 2) G2: 1.0 (0, 2) <i>P</i> = 0.86</p> <p>Urgency episodes, mean ± SD: G1: 2.22 ± 1.42 G2: 2.14 ± 1.12 <i>P</i> = 0.75</p> <p>Voids/day, mean ± SD: G1: 11.3 ± 4.68 G2: 10.23 ± 3.53 <i>P</i> = 0.22</p> <p>Daytime voids/day, mean ± SD: G1: 9.07 ± 4.59 G2: 8.09 ± 3.83 <i>P</i> = 0.26</p> <p>Nocturia episodes, median (IQR): G1: 2 (1, 3) G2: 2 (1, 3) <i>P</i> = 0.94</p> <p>Perceived severity, median (IQR): G1: 5.6 (3.9, 7.1) G2: 5.3 (3.1, 7.0) <i>P</i> = 0.18</p> <p>KHQ, general health, mean score ± SD: G1: 56.0 ± 18.1 G2: 55.8 ± 17.1 <i>P</i> = 0.955</p> <p>KHQ, incontinence impact, mean score ± SD: G1: 68.5 ± 30.6 G2: 69.7 ± 27.9 <i>P</i> = 0.836</p>	<p>UUI episodes, median change (IQR): G1: 0 (-1, 0) G2: 0 (-1, 0)</p> <p>Urgency episodes, mean change ± SD: G1: 0.27 ± 0.94 G2: 0.48 ± 1.12 <i>P</i> = 0.32</p> <p>Voids/day, mean change ± SD: G1: 2.18 ± 2.80 G2: 1.04 ± 2.92 <i>P</i> = 0.055</p> <p>Daytime voids/day, mean change ± SD: G1: 1.90 ± 2.76 G2: 0.55 ± 3.21 <i>P</i> = 0.029</p> <p>Nocturia episodes, median change (IQR): G1: -1 (-1, 0) G2: -1 (-1, 0)* <i>P</i> = 0.46</p> <p>Frequency, change from > 8 to ≤ 8 after treatment, n (%): G1: 11 (20.3) G2: 7 (16.2) <i>P</i> = 0.482</p> <p>Perceived severity, median change (IQR): G1: 1.5 (-0.65, 3.4) G2: 1.2 (-0.8, 3) <i>P</i> = 0.7</p> <p>Perceived change of bladder symptoms, median (IQR): G1: 1.5 (0.5, 3) G2: 1.1 (0, 2) <i>P</i> = 0.172</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Masking: -</p> <p>Pt selection criteria: +</p> <p>Loss to followup: -</p> <p>Drop-out rates: +</p> <p>Power calculation: +</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: -</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 5. KQ2 Complementary and Alternative Therapies (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Mak et al., 2007 (continued)			<p>KHQ, role limitations, mean score ± SD: G1: 39.8 ± 33.7 G2: 41.4 ± 33.8 <i>P</i> = 0.811</p> <p>KHQ, physical limitations, mean score ± SD: G1: 37.3 ± 35.0 G2: 54.2 ± 78.6 <i>P</i> = 0.160</p> <p>KHQ, social limitations, mean score ± SD: G1: 28.6 ± 30.2 G2: 35.1 ± 30.5 <i>P</i> = 0.294</p> <p>KHQ, personal limitations, mean score ± SD: G1: 17.2 ± 29.9 G2: 24.4 ± 31.5 <i>P</i> = 0.358</p> <p>KHQ, emotion, mean score ± SD: G1: 36.2 ± 33.5 G2: 44.1 ± 35.9 <i>P</i> = 0.262</p> <p>KHQ, sleep/energy, mean score ± SD: G1: 36.1 ± 33.9 G2: 33.3 ± 29.9 <i>P</i> = 0.674</p> <p>KHQ, severity measures, mean score ± SD: G1: 25.8 ± 21.1 G2: 25.5 ± 20.8 <i>P</i> = 0.959</p> <p>Peak flow rate, ml/s ± SD: G1: 16.7 ± 9.75 G2: 18.5 ± 7.69</p>	<p>KHQ, general health, mean change ± SD: G1: -5.55 ± 19.8 G2: -5.23 ± 19.3 <i>P</i> = 0.936</p> <p>KHQ, incontinence impact, mean change ± SD: G1: -9.87 ± 32.7 G2: -14.72 ± 33.5 <i>P</i> = 0.477</p> <p>KHQ, role limitations, mean change ± SD: G1: -3.08 ± 31.0 G2: -2.32 ± 28.5 <i>P</i> = 0.902</p> <p>KHQ, physical limitations, mean change ± SD: G1: -1.54 ± 27.9 G2: -12.8 ± 71.9 <i>P</i> = 0.337</p> <p>KHQ, social limitations, mean change ± SD: G1: -0.51 ± 29.1 G2: -6.97 ± 27.6 <i>P</i> = 0.267</p> <p>KHQ, personal limitations, mean change ± SD: G1: 2.15 ± 29.1 G2: -10.1 ± 29.8 <i>P</i> = 0.116</p> <p>KHQ, emotion, mean change ± SD: G1: 1.44 ± 30.4 G2: -10.0 ± 31.6 <i>P</i> = 0.074</p> <p>KHQ, sleep/energy, mean change ± SD: G1: -5.8 ± 36.9 G2: -6.2 ± 36.2 <i>P</i> = 0.964</p>	

Evidence Table 5. KQ2 Complementary and Alternative Therapies (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Mak et al., 2007 (continued)			Max cystometric capacity, ml ± SD: G1: 378.1 ± 89.0 G2: 347.8 ± 94.0 <i>P</i> = 0.11	KHQ, severity measures, mean change ± SD: G1: -0.37 ± 8.8 G2: -1.86 ± 21.3 <i>P</i> = 0.720	
			Duration of symptoms, median (IQR): G1: 5 (3, 9) G2: 6 (3, 10)	Believed to have received intervention, %: G1: 88.9 G2: 67.4 <i>P</i> = 0.012	
			Previous treatment for idiopathic detrusor overactivity, n (%): G1: 27 (47.3) G2: 22 (44.9)		
			Previous continence surgery, n (%): G1: 5 (8.8) G2: 7 (16.3)		

Evidence Table 6. KQ 3 Comparison of Treatments

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Abrams et al., 1998</p> <p>Country and setting: UK, Ireland, Sweden, Academic medical center</p> <p>Enrollment period: July 1995 to July 1996</p> <p>Funding: Pharmacia Upjohn</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: Tolterodine vs. oxybutynin vs. placebo x 12 wks</p> <p>Groups: G1: Tolterodine 2mg bid (dose could be reduced to 1.5 mg b.i.d.) G2: Oxybutynin 5mg t.i.d. (dose could be reduced to 2.5 mg t.i.d.) G3: Placebo</p> <p>N at enrollment: G1: 118 G2: 118 G3: 57</p> <p>N at follow-up: G1: 118 G2: 117 G3: 56</p> <p>Women, n (%): G1: 91 (77.1) G2: 88 (74.5) G3: 43 (75.4)</p> <p>Age, mean (range): G1: 55 (19-80) G2: 58 (21-80) G3: 58 (26-78)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age \geq 18 • UDS confirmed bladder overactivity • \geq 8 voids/day • \geq 1 episode UUI/ day <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • SUI • Detrusor hyperreflexia • Hepatic, renal, hematological disorders • Symptomatic or recurrent UTI • BOO • Bladder retraining • Electrical stimulation therapy • Indwelling catheter • Intermittent catheterization • Pregnant/nursing • Women without reliable BC 	<p>Incontinence episodes, n (%): G1: 93 (79) G2: 88 (75) G3: 40 (70)</p> <p>Incontinence episodes/day, mean (range): G1: 2.9 (0.1-24.0) G2: 2.6 (0.1-24.0) G3: 3.3 (0.1-23.5)</p> <p>Voids/day, mean (range): G1: 11.5 (6.3-37.0) G2: 10.7 (5.3-27.7) G3: 11.7 (6.9-34.4)</p> <p>\geq 8 voids/day, n (%): G1: 112 (95) G2: 109 (92) G3: 53 (93)</p> <p>Voided volume (mL), mean (range): G1: 166 (55-426) G2: 176 (57-423) G3: 157 (59-276)</p> <p>Previous OAB therapy, n (%): G1: 61 (52) G2: 71 (60) G3: 43 (75)</p>	<p>Incontinence episodes/day, mean change \pm SD: G1: -1.3 \pm 3.2 G2: -1.7 \pm 3.1 G3: -0.9 \pm 1.5 G1/G3: P = 0.22 G2/G3: P = 0.023</p> <p>Voids/day, mean change \pm SD: G1: -2.7 \pm 3.8 G2: -2.3 \pm 2.7 G3: -1.6 \pm 3.6 G1/G3: P = 0.002 G2/G3: P = 0.068</p> <p>Voided volume (mL), mean change \pm SD: G1: 38 \pm 54 G2: 47 \pm 58 G3: 6 \pm 42 G1/G3: P < 0.001 G2/G3: P < 0.001</p> <p>Subjective improvement in bladder symptoms, %: G1: 50 G2: 49 G3: 47</p> <p>Dose reductions, n (%): G1: 9 (8) G2: 38 (32) G3: 1 (2) G2/G1: P < 0.001 G2/G3: P < 0.001</p> <p>Dry mouth, n (%): G1: 59 (50) G2: 102 (86) G3: 12 (21) G1/G3: P < 0.001 G2/G1: P < 0.001 G2/G3: P < 0.001</p> <p>Dyspepsia, n (%): G1: 11 (9) G2: 27 (23) G3: 3 (5)</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: fair</p> <p>Randomization: +</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: ++</p> <p>Drop-out rates: -</p> <p>Power calculation: +</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Abrams et al., 1998 (continued)				<p>Nausea, n (%): G1: 4 (3) G2: 7 (6) G3: 6 (11)</p> <p>Upper respiratory infection, n (%): G1: 12 (10) G2: 3 (3) G3: 8 (14)</p> <p>Adverse events reported, N: G1: 302 G2: 412 G3: 117</p> <p>Patients with ≥ 1 AE, n (%): G1: 105 (89) G2: 114 (97) G3: 46 (81) G2/G1: P = 0.023 G2/G3: P < 0.001</p> <p>Discontinued due to AEs, n (%): G1: 10 (8) G2: 20 (17) G3: 1 (2)</p>	

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Arruda et al., 2008</p> <p>Country and setting: Brazil, community</p> <p>Enrollment period: August 2001 to September 2005</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: None</p>	<p>Design: RCT</p> <p>Intervention: Oxybutynin vs. functional electrostimulation vs. pelvic floor training</p> <p>Groups: G1: Oxybutynin 5 mg b.i.d. G2: Ambulatory stimulation applied vaginally G3: Pelvic floor exercises with a therapist and at home</p> <p>N Screened: 81</p> <p>N at enrollment: G1: 22 G2: 21 G3: 21</p> <p>N at follow-up: G1: 22 G2: 21 G3: 21</p> <p>Age, range: 35-80</p> <p>Race/ethnicity: NR</p> <p>Women, %: 100</p> <p>Length of follow up: 12 weeks</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> Community-dwelling Dx of OAB and DO Capable of completing a bladder diary and performing a pelvic muscle floor contraction For those with MUI, urge was predominant <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Hx of psychiatric or neurologic illness Persistent UTI Inability to comply with regular follow-up visits Current pregnancy Postvoid residual volume > 100 ml Contraindications to anticholinergic therapy Cardiac pacemaker Type III SUI Uncontrolled metabolic conditions or indwelling catheterization Using medications including anticholinergic drugs, calcium antagonists, beta agonists, dopamine agonists, striated muscle relaxants or estrogens Any uterine prolapse 	<p>UII episodes/week, mean ± SD: G1: 13.8 ± 11.6 G2: 13.5 ± 15.6 G3: 16.4 ± 17.2</p> <p>Nocturia episodes/week, mean ± SD: G1: 1.7 ± 1.5 G2: 1.9 ± 1.9 G3: 1.4 ± 1.2</p> <p>Pads/day, mean ± SD: G1: 2.6 ± 2.7 G2: 2.3 ± 2.4 G3: 2.7 ± 1.5</p> <p>Voids/day, mean ± SD: G1: 7.7 ± 2.6 G2: 8.6 ± 3.4 G3: 6.8 ± 2.2</p> <p>Residual volume mean mL ± SD: G1: 3.2 ± 6.3 G2: 1.0 ± 2.6 G3: 1.8 ± 3.3</p> <p>Volume, first desire to void, mean mL ± SD: G1: 117.7 ± 68.9 G2: 102.4 ± 51.1 G3: 86.7 ± 38.9</p> <p>Maximal cystometric capacity, mean mL ± SD: G1: 410.4 ± 194.1 G2: 350.0 ± 212.9 G3: 424.0 ± 149.0</p>	<p>UII episodes/week, mean ± SD: G1: 7.0 ± 10.6 G2: 7.9 ± 13.7 G3: 7.8 ± 15.3 G1/BL: <i>P</i> = 0.007 G2/BL: <i>P</i> = 0.039 G3/BL: <i>P</i> = 0.035</p> <p>Urgency resolved, n (%): G1: 14 (63.6) G2: 11 (52.4) G3: 12 (57.1) <i>P</i> = 0.754</p> <p>Satisfied, n (%): G1: 17 (77.3) G2: 11 (52.4) G3: 16 (76.2) <i>P</i> = 0.142</p> <p>Nocturia episodes/week, mean ± SD: G1: 0.9 ± 0.8 G2: 1.2 ± 1.3 G3: 1.0 ± 1.1 G1/BL: <i>P</i> = 0.003 G2/BL: <i>P</i> = 0.036 G3/BL: <i>P</i> = 0.086</p> <p>Pads/day, mean ± SD: G1: 0.9 ± 1.5 G2: 0.9 ± 1.7 G3: 0.8 ± 1.3 G1/BL: <i>P</i> < 0.001 G2/BL: <i>P</i> = 0.004 G3/BL: <i>P</i> < 0.001</p> <p>Voids/day, mean ± SD: G1: 6.4 ± 1.6 G2: 7.9 ± 2.3 G3: 7.1 ± 2.1 G1/BL: <i>P</i> = 0.014 G2/BL: <i>P</i> = 0.291 G3/BL: <i>P</i> = 0.441</p> <p>Residual volume, mean mL ± SD: G1: 4.7 ± 9.4 G2: 1.1 ± 2.5 G3: 2.1 ± 3.5 G1/BL: <i>P</i> = 0.425 G2/BL: <i>P</i> = 0.760 G3/BL: <i>P</i> = 0.297</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: -</p> <p>Method and blinding: -</p> <p>Pt selection criteria: +</p> <p>Loss to followup: +</p> <p>Drop-out rates: -</p> <p>Power calculation: +</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: +</p> <p>Length of followup: +</p> <p>Measurement methods: -</p> <p>Measurement reliability: -</p> <p>Intervention description: +</p>

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Arruda et al., 2008 (continued)			Involuntary detrusor contraction volume mean mL ± SD: G1: 189.5 ± 114.1 G2: 220.0 ± 127.2 G3: 239.3 ± 163.0	Volume first desire to void, mean mL ± SD: G1: 157.3 ± 63.8 G2: 123.8 ± 59.0 G3: 137.6 ± 76.7 G1/BL: <i>P</i> = 0.019 G2/BL: <i>P</i> = 0.091 G3/BL: <i>P</i> = 0.017	
			Involuntary detrusor contraction maximal pressure (mm H₂O): G1: 39.4 ± 26.1 G2: 43.7 ± 22.9 G3: 34.2 ± 19.8	Maximal cystometric capacity mean mL ± SD: G1: 517.3 ± 191.7 G2: 436.6 ± 178.7 G3: 489.0 ± 141.3 G1/BL: <i>P</i> = 0.001 G2/BL: <i>P</i> = 0.017 G3/BL: <i>P</i> = 0.113	
				Involuntary detrusor contraction volume (mL): G1: 188.6 ± 183.2 G2: 73.3 ± 112.4 G3: 114.3 ± 154.2 G1/BL: <i>P</i> = 0.986 G2/BL: <i>P</i> = 0.001 G3/BL: <i>P</i> = 0.044	
				Involuntary detrusor contraction maximal pressure, mm H₂O ± SD: G1: 19.6 ± 20.9 G2: 22.4 ± 30.1 G3: 17.2 ± 25.5 G1/BL: <i>P</i> < 0.001 G2/BL: <i>P</i> = 0.002 G3/BL: <i>P</i> = 0.027	
				Normal urodynamic evaluation, n (%): G1: 8 (36.4) G2: 12 (57.1) G3: 11 (52.4) <i>P</i> = 0.358	
				Persistent improvement at 1 year: G1: 10/17 G2: 4/11 G3: 9/16	

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Arruda et al., 2008 (continued)				<p>Dry mouth, n (%): G1: 16 (72.7) G2: 0 G3: 0</p> <p>Difficulty voiding, n (%): G1: 2 (9.1) G2: 0 G3: 0</p> <p>Dizziness, n (%): G1: 1 (4.5) G2: 0 G3: 0</p> <p>Blurred vision, n (%): G1: 1 (4.5) G2: 0 G3: 0</p> <p>Constipation, n (%): G1: 1 (4.5) G2: 0 G3: 0</p> <p>Dry mouth, n (%): G1: 16 (72.7) G2: 0 G3: 0</p> <p>Difficulty voiding, n (%): G1: 2 (9.1) G2: 0 G3: 0</p> <p>Dizziness, n (%): G1: 1 (4.5) G2: 0 G3: 0</p> <p>Blurred vision, n (%): G1: 1 (4.5) G2: 0 G3: 0</p> <p>Constipation, n (%): G1: 1 (4.5) G2: 0 G3: 0</p>	

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Burgio et al. 1998</p> <p>Country and setting: US; academic health center outpatient geriatric medicine clinic</p> <p>Enrollment period: July 1989 to August 1995</p> <p>Funding: National Institutes on Aging</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT, placebo controlled</p> <p>Computer-generated random numbers using a block size of 6, w/ prior stratification by type and severity of incontinence</p> <p>Intervention: Biofeedback-assisted behavioral vs. drug treatment (oxybutynin chloride; possible range of doses 2.5 mg/d-5.0 mg t.i.d.) vs. placebo</p> <p>All patients had 4 visits over an 8-week period. Patients in G1 had biofeedback added to behavioral training in absence of 50% improvement by session 3.</p> <p>Groups: G1: Behavioral ± biofeedback G2: Pharmacologic G3: Placebo</p> <p>N at enrollment: 468 screened 271 not eligible 197 randomized G1: 65 G2: 67 G3: 65</p> <p>N at follow-up: G1: 61 G2: 55 G3: 53</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> Community-dwelling women ≥ 55 years old Ambulatory ≥ 2 urge accidents per week by baseline bladder diary Urge incontinence as predominant pattern Urodynamic evidence of bladder dysfunction <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Continual leakage Postvoid residual urine volume >200mL Uterine prolapse past the introitus Narrow-angle glaucoma Unstable angina Decompensated congestive heart failure Hx of malignant arrhythmias MMSE <20 (Dementia) 	<p>Duration of symptoms, mean yrs ± SD: G1: 9.4 ± 10.8 G2: 9.8 ± 11.9 G3: 12.7 ± 15.9</p> <p>Restricted activity, (%): G1: 30.8 G2: 32.8 G3: 38.5</p> <p>Urge UUI only, %: G1: 49.2 G2: 49.3 G3: 47.7</p> <p>Accidents per week, mean n ± SD: G1: 15.8 ± 14.5 G2: 15.9 ± 14.1 G3: 15.4 ± 13.4 P=.98</p> <p>Severity classification, %: Mild (<5 accidents/wk) G1: 18.5 G2: 17.9 G3: 18.5</p> <p>Moderate (5-10 accidents/wk) G1: 29.2 G2: 29.9 G3: 27.7</p> <p>Severe (>10 accidents/wk) G1: 52.3 G2: 52.2 G3: 53.8</p> <p>Previous treatment with surgery, %: G1: 20.0 G2: 26.9 G3: 29.2</p>	<p>Accidents per week, mean ± SD: G1: 2.8 ± 4.7 G2: 5.7 ± 9.8 G3: 8.2 ± 11.6 P=.005</p> <p>Percent reduction, mean ± SD: G1: 80.7 ± 24.8 G2: 68.5 ± 37.2 G3: 39.4 ± 80.0 P<0.001</p> <p>Percent reduction, range: G1: -0.9 – 100 G2: -85.7 – 100 G3: -400.0 - 100</p> <p>Patient perceptions of progress in treatment Progress, %: Much better G1: 74.1 G2: 50.9 G3: 226.9</p> <p>Better G1: 25.9 G2: 30.9 G3: 38.5</p> <p>About the same G1: 0.0 G2: 16.4 G3: 28.8</p> <p>Worse G1: 0.0 G2: 1.8 G3: 5.8</p> <p>Estimate of % improvement, mean ± SD: G1: 81.6 ± 18.6 G2: 66.4 ± 35.4 G3: 45.1 ± 36.6</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: +</p> <p>Drop-out rates: +</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Burgio et al. 1998 (continued)	<p>Women, %: 100</p> <p>Age, yrs ± SD: G1: 67.3 ± 7.6 G2: 68.2 ± 7.5 G3: 67.6 ± 7.6</p> <p>Race/ethnicity,: NR</p> <p>Parity mean ± SD: G1: 2.8 ± 2.0 G2: 2.1 ± 1.4 G3: 2.7 ± 1.8</p>		<p>Previous treatment with medication, %: G1: 27.7 G2: 35.8 G3: 30.8</p> <p>Using estrogen, %: G1: 32.3 G2: 38.8 G3: 35.4</p> <p>Using diuretics, %: G1: 20.0 G2: 14.9 G3: 12.3</p>	<p>Having fewer accidents, %: G1: 100.0 G2: 87.3 G3: 67.3</p> <p>Accidents are smaller, %: G1: 87.3 G2: 78.8 G3: 54.0</p> <p>Able to wear less protection, %: G1: 76.0 G2: 56.0 G3: 334.1</p> <p>Comfortable enough with treatment to continue indefinitely, %: G1: 96.5 G2: 54.7 G3: 43.1</p> <p>Patient satisfaction with progress, %: Completely satisfied G1: 77.6 G2: 54.7 G3: 43.1</p> <p>Somewhat satisfied G1: 22.4 G2: 40.0 G3: 34.0</p> <p>Not at all satisfied G1: 0.0 G2: 10.9 G3: 38.0</p> <p>Wish to receive another form of treatment, %: G1: 14.0 G2: 75.5 G3: 75.5</p> <p>p<0.001 for all comparisons</p>	

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Burgio et al. 1998 (continued)				<p>Adverse effects, % compared to placebo G3:</p> <p>Dry mouth G1: 34.9 G2: 96.9 G3: 54.8 p<0.001</p> <p>Inability to void G1: 6.3 G2: 21.5 G3: 3.2 p=0.002</p> <p>Constipation G1: 22.2 G2: 38.5 G3: 37.1 p=0.10</p> <p>Blurred vision G1: 9.5 G2: 15.4 G3: 9.7 p=0.50</p> <p>Confusion G1: 6.3 G2: 7.7 G3: 11.3 p=0.59</p>	

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Burgio et al. 2000 [See Burgio et al., 1998]</p> <p>Enrollment period: two weeks after completion of Burgio et al. 1998, #2440</p> <p>Funding: National Institute on Aging, grants AG 08010 and K04 00431</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Modified crossover of RCT</p> <p>Intervention: Participants whose treatment was not completely successful were given the opportunity to switch or use combined treatment; further reductions in incontinence were measured.</p> <p>Groups: <u>Treatment</u> <u>Changes:</u> G1: Previous oxybutynin to behavioral modification alone G2: Previous behavior alone to 2.5 mg oxybutynin t.i.d. + behavioral therapy G3: Previous oxybutynin alone to 2.5 mg oxybutynin t.i.d. + behavioral therapy G4: Placebo to behavioral G5: Placebo to oxybutynin</p> <p>N at enrollment G1: 19 G2: 8 G3: 27 G4: 34 G5: 10</p> <p>N at follow-up: G1: 18 G2: 8 G3: 26 G4: NR G5: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> Community-dwelling women at least age 55 Ambulatory At least 2 urge accidents per week by baseline bladder diary Urge incontinence as predominant pattern Urodynamic evidence of bladder dysfunction not completely dry or satisfied with previous, 8-wk treatment <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Continual leakage Postvoid residual urine volume >200mL Uterine prolapse past the introitus Narrow-angle glaucoma Unstable angina Decompensated congestive heart failure Hx of malignant arrhythmias MMSE <20 (Dementia) 	<p>% Reduction of incontinence after previous study (at baseline), mean: G1: 59.1 G2: 57.5 G3: 72.7 G4: 22.9 G5: 44.8</p>	<p>Final % Reduction of incontinence mean, p: G1: 77.1, 0.109 G2: 88.5, 0.034 G3: 84.3, 0.001 G4: 63.9, .002 G5: 76.5, .012</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: +</p> <p>Drop-out rates: +</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Note: 29.2 of G3 declined to continue with drug therapy once they received behavioral modification.

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Burgio et al., 2001</p> <p>[See Burgio et al., 1998]</p> <p>Country and setting: US, University based</p> <p>Enrollment period: [See Burgio et al., 1998]</p> <p>Funding: NIH</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: Psychological distress associated with pharmacologic treatment vs. behavioral vs. placebo</p> <p>Groups: G1: Behavioral training with biofeedback, 4 visits G2: Oxybutynin 2.5 mg P.O. t.i.d., up to a max of 5 mg t.i.d. G3: Placebo</p> <p>N at enrollment: 197 women</p> <p>N treated: 169</p> <p>N at follow-up: 155 (completed both pre and post-treatment psychological assessment)</p> <p>Age yrs, mean ± SD: 67.5 ± 7.2</p> <p>Women, %: 100</p> <p>Race/ethnicity, %: White: 97 African American: 3</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • ≥ 55 yrs old • Ambulatory • UUI ≥2x/wk (2 wk bladder diary), persisting x 3 mo • Predominant UUI • UDS evidence of bladder dysfunction (DI or maximal capacity ≤350 mL) <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Contraindication to oxybutynin or behavioral treatment • Continual leakage • PVR > 200 mL • Uterine prolapse beyond the introitus • Decompensated CHF • Hx of malignant arrhythmias • Impaired mental status (MMSE <20) 	<p>SCL-90-R scores ± SD:</p> <p>Somatization G1: 56.0 (10.6) G2: 51.4 (10.8) G3: 52.4 (11.1)</p> <p>Obsessive-Compulsive: G1: 56.5 (10.7) G2: 56.6 (11.4) G3: 57.7 (10.0)</p> <p>Interpersonal Sensitivity G1: 53.8 (11.0) G2: 51.4 (11.9) G3: 50.4 (12.0)</p> <p>Depression G1: 54.7 (10.0) G2: 52.5 (9.7) G3: 51.0 (11.9)</p> <p>Anxiety G1: 48.7 (13.9) G2: 46.8 (12.0) G3: 47.2 (12.8)</p> <p>Hostility G1: 49.3 (10.7) G2: 45.9 (10.1) G3: 48.3 (10.4)</p> <p>Phobia G1: 47.5 (10.2) G2: 46.7 (10.3) G3: 45.7 (8.5)</p> <p>Paranoid Ideation G1: 48.6 (12.4) G2: 49.3 (11.1) G3: 46.1 (12.8)</p> <p>Psychoticism G1: 54.1 (10.7) G2: 52.1 (10.3) G3: 50.5 (11.2)</p> <p>Global Severity G1: 54.2 (11.1) G2: 52.5 (10.3) G3: 52.4 (11.2)</p>	<p>Reduction in incontinence:</p> <p>G1: 83.3% G2: 74.4% G3: 41.4% P<0.001</p> <p>SCL-90-R scores ± SD:</p> <p>Somatization G1: 51.8 (11.4) G2: 51.2 (9.8) G3: 49.8 (13.0)</p> <p>Obsessive-Compulsive: G1: 53.8 (13.9) G2: 53.9 (10.9) G3: 55.4 (11.0)</p> <p>Interpersonal Sensitivity G1: 49.5 (12.0) G2: 48.9 (11.2) G3: 49.2 (11.3)</p> <p>Depression G1: 51.5 (11.5) G2: 50.6 (10.7) G3: 51.4 (11.2)</p> <p>Anxiety G1: 46.1 (14.6) G2: 44.5 (12.3) G3: 45.8 (12.9)</p> <p>Hostility G1: 44.9 (10.8) G2: 44.6 (10.5) G3: 47.3 (11.2)</p> <p>Phobia G1: 47.1 (11.2) G2: 45.0 (8.3) G3: 45.1 (8.5)</p> <p>Paranoid Ideation G1: 45.8 (10.9) G2: 47.2 (11.6) G3: 47.2 (12.0)</p> <p>Psychoticism G1: 49.2 (11.7) G2: 50.4 (9.7) G3: 49.6 (10.3)</p> <p>Global Severity G1: 50.8 (12.8) G2: 50.4 (10.0) G3: 51.4 (10.9)</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: +</p> <p>Drop-out rates: +</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Burgio et al., 2001 (continued)			155 participants were compared to the 42 who did not complete intervention and psychological assessment, higher scores (greater distress) on 6 of 10 SCL-90-R scales (somatization, obsessive/compulsive, depression, hostility, paranoid ideation, global severity index), all p values <0.05 Normal range, score 0-63 >75% in normal range (including dropouts) on 9 of 10 scales Highest impairment rate: 33% scored abnormal for obsessive-compulsive	Correlations between Reduction of Incontinence and Changes in Psychological Symptoms Somatization G1: 0.28* G2: -0.09 G3: 0.17 Obsessive-Compulsive G1: 0.01 G2: -0.14 G3: 0.02 Interpersonal Sensitivity G1: -0.09 G2: 0.04 G3: 0.13 Depression G1:-0.04 G2: 0.03 G3: 0.07 Anxiety G1: -0.10 G2: -0.01 G3: 0.34* Hostility G1: -0.10 G2: 0.09 G3: 0.11 Phobic anxiety G1:-0.21 G2: -0.17 G3: 0.02 Paranoid Ideation G1: 0.14 G2: -0.04 G3: -0.06 Psychoticism G1: -0.01 G2: -0.01 G3: 0.13 Global Severity Index G1: 0.01 G2: 0.06 G3: 0.45*** *p<0.05 ***P=0.001	

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Burgio et al., 2008</p> <p>Country and setting: US, Academic medical centers</p> <p>Enrollment period: July 2004 to January 2006</p> <p>Funding: NIH Pfizer</p> <p>Author industry relationship disclosures: 20 of 29 Allergan (3) Alza (1) Astellas Pharma (7) Bionovo (1) Bristol-Meyers Squibb (1) Dynogen (1) Elan (1) Ethicon (2) GSK (4) Johnson & Johnson (3) Lilly (7) Medtronic (1) Merck (1) Novartis (6) Ortho-McNeil (3) Pfizer (>10) Procter & Gamble (3) Q-Med (1) Renesa (1) Sanofi (1) Solace (1) Watson (1)</p>	<p>Design: RCT</p> <p>Intervention: Pharmacologic vs. pharmacologic plus behavioral interventions for 10 wks (Stage 1), followed by no drug therapy (Stage 2) with assessments at 10 wks and 8 mos*</p> <p>Groups: G1: tolterodine tartrate (ER capsules), 4 mg/d plus behavioral training: pelvic floor muscle control and exercises, behavioral strategies to diminish urgency, suppress bladder contractions and prevent both stress and urge incontinence; delayed voiding; fluid management; handout with hints G2: tolterodine tartrate (ER capsules), 4 mg/d</p> <p>N at enrollment: 4043 screened 2612 not eligible 870 declined 561 consented 254 excluded 307 randomized G1: 153 G2: 154</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Women • Community-dwelling • UUI only, or urge-predominant incontinence • ≥ 7 episodes of incontinence in a 7-day bladder diary • Persistent incontinence for at least 3 mos • No antimuscarinics or other medications that could affect UI • No evidence of neurogenic etiology <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Age < 21 • Pregnant, or planning a pregnancy in next 8 mos, or not using birth control • < 6 mos postpartum or termination after 20 wks gestation • Unable to contract pelvic floor muscles • Behavioral therapy of > 2 mos in past 2 yrs • Continual leakage or always damp • Hypersensitive to study drug • Systemic disease that affects bladder function 	<p>UUI, 7-13 episodes/week, n (%): G1: 2 (1.3) G2: 2 (1.3)</p> <p>UUI, ≥ 14 episodes/week, n (%): G1: 2 (1.3) G2: 4 (2.6)</p> <p>MUI, 7-13 episodes/week, n (%): G1: 46 (29.9) G2: 46 (30.1)</p> <p>MUI, ≥ 14 episodes/week, n (%): G1: 104 (67.5) G2: 101 (66.0)</p> <p>Adjusted incontinence episodes/week, mean: G1: 23.1 G2: 23.2</p> <p>Previous non-surgical treatment for incontinence, n (%): G1: 19 (12) G2: 22 (14)</p>	<p>Success, n (%): G1: 43 (28) G2: 41 (27)</p> <p>Failure, n (%): G1: 75 (49) G2: 78 (51)</p> <p>8 month success rate, lifetable analysis, % (95% CI): G1: 41 (32, 50) G2: 41 (33, 50) G1/G2: 0 (-12, 12)</p> <p>8 mo success rate, complete cases, % (95% CI): G1: 36 (27, 45) G2: 34 (25, 43) G1/G2: 2 (-10, 14)</p> <p>8 mo success rate, assuming missings were failures, % (95% CI): G1: 28 (21, 35) G2: 27 (20, 34) G1/G2: 1 (-9, 11)</p> <p>Adjusted incontinence episodes/week, after Stage 1, mean: G1: 2.7 G2: 4.7</p> <p>Adjusted incontinence episodes/week, after Stage 1, mean change: G1: -20.4 G2: -18.5 G1/G2: -1.9 (-5.9, 2.0)</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Masking: -</p> <p>Pt selection criteria: +</p> <p>Loss to followup: -</p> <p>Drop-out rates: ++</p> <p>Power calculation: +</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: ++</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Burgio et al., 2008 (continued)	<p>N at follow-up: G1: 153 Completed treatment: 107 Outcome known at 8 mos: 119 G2: 154 Completed treatment: 101 Outcome known at 8 mos: 118</p> <p>Age, mean ± SD: G1: 55.8 ± 14.2 G2: 58.0 ± 13.5</p> <p>Women, %: 100</p> <p>Race/ethnicity, n (%): Hispanic: G1: 13 (9) G2: 17 (11) NH White: G1: 105 (69) G2: 85 (56) NH Black: G1: 22 (14) G2: 35 (23) Other: G1: 13 (9) G2: 15 (10)</p> <p>BMI, kg/m² ± SD: G1: 33.2 ± 9.5 G2: 32.3 ± 7.6</p>	<ul style="list-style-type: none"> • Currently using catheter to empty bladder • Postvoid residual volume > 150mL • Treatment for prolapse with pessary < 3 mos • Incontinence, vaginal, bladder or prolapse surgery in the past 6 mos • Urethral diverticulum • Previous augmentation cystoplasty or artificial sphincter • Neuromodulation for pelvic indications • Using anticholinergic agents, cholinergic agonists, tricyclic antidepressants, or duloxetine in ≤ 4 wks • Using diuretics with dosage change past 3 mos • Uncontrolled medical problem • History of bladder or pelvic cancer or pelvic radiation therapy • Glaucoma • Gastric retention • Non-ambulatory • Participation in other intervention trial that might influence outcome 		<p>Achieved 70% reduction in incontinence episodes, 10 wks (%): G1: 69 G2: 58 G1/G2: 11 (-0.3, 22.1)</p> <p>Totally dry, per bladder diary, 10 wks (%): G1: 21 G2: 17</p> <p>Voids/day, mean change: G1: 0.5 G2: -0.4 G1/G2: 0.9 (0.3, 1.5)</p> <p>Symptom Distress Scores: G1/G2: P < 0.0001</p> <p>Symptom Bother Scores (OAB-q), Stage 1, mean change: G1: -36.7 G2: -30.4 G1/G2: P < 0.0001</p> <p>Symptom Bother Scores (OAB-q), Stage 2, mean change: G1: -30.9 G2: -20.4 G1/G2: P < 0.0001</p> <p>Patient completely satisfied, Stage 1, %: G1: 53 G2: 40 G1/G2: 13 (1, 25)</p>	

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Burgio et al., 2008 (continued)				<p>Patient completely satisfied, 8 months, %: G1: 33 G2: 20 G1/G2: 13 (2, 24)</p> <p>Patient better or much better, Stage 1, %: G1: 90 G2: 77 G1/G2: 13 (4, 22)</p> <p>Patient better or much better, 8 months, %: G1: 69 G2: 43 G1/G2: 26 (14 , 38)</p> <p>Persistence in perceived improvement, 8 months, women with improvement at Stage 1: G1: 72 G2: 54 G1/G2: 17 (4, 30)</p> <p>Harms: G1: 3 participants 1: blurred vision, syncope, night sweats, stomach cramping and weakness 2: 2 episodes of small-bowel obstruction and an allergic reaction (pruritus and rash) 3: tachycardia during stage 2 G2: 3 participants 1: small bowel obstruction 2: peripheral edema 3: renal cell carcinoma diagnosis during stage 2</p>	

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Chancellor, Kianifard et al., 2008</p> <p>Country and setting: US</p> <p>Enrollment period: May 2005 to February 2006</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: 6 of 7 Genaera (1) Novartis (6)</p>	<p>Design: Randomized open-label</p> <p>Intervention: Behavioral modification in addition to darifenacin (allowed to increase dose from darifenacin 7.5 mg to 15 mg daily after 2 weeks)</p> <p>Groups:* G1: Darifenacin G2: Darifenacin and behavioral modification</p> <p>N at enrollment: G1: 190 G2: 205</p> <p>N at follow-up, 12 weeks: G1: 173 G2: 175</p> <p>Age, mean ± SD: G1: 57.4 ± 13.1 G2: 58.4 ± 14.6</p> <p>Race/ethnicity: White: G1: 90 G2: 88.3 Other (non-white): G1: 10 G2: 11.7</p> <p>Women, %: G1: 90 G2: 88.3</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • ≥ 8 voids/ day • ≥ 2 UUI episodes/ day • ≥ 2 episodes of urgency/day <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Any drug with bladder effects for 2 weeks prior to study participation • Participation in any formal bladder-training program within 30 days of screening • Predominant SUI • Bladder or neurologic condition that could affect bladder function or in which use of anticholinergics was contra-indicated 	<p>UUI episodes/day, mean ± SD: G1: 2.78 ± 2.57 G2: 3.00 ± 2.56</p> <p>UUI episodes/day, median: G1: 2.33 G2: 2.58</p> <p>Mean UUI episodes/day, total population, n (%): 0 episodes: G1: 36 (19.1) G2: 29 (14.2) 1-6 episodes: G1: 138 (73.0) G2: 155 (76.0) 7-13 episodes: G1: 15 (7.9) G2: 20 (9.8) ≥14 episodes: G1: 0 (0.0) G2: 0 (0.0)</p> <p>Mean UUI episodes/day, age ≥ 65, n (%): 0 episodes G1: 9 (17.0) G2: 8 (9.9) 1-6 episodes G1: 40 (75.5) G2: 63 (77.8) 7-13 episodes G1: 4 (7.5) G2: 10 (12.3)</p> <p>Urgency episodes/day, mean ± SD: G1: 10.88 ± 3.80 G2: 10.58 ± 4.00</p> <p>Urgency episodes/day, median: G1: 10.67 G2: 10.33</p> <p>Voids/day, mean ± SD: G1: 11.92 ± 3.03 G2: 11.75 ± 3.37</p>	<p>UUI episodes/day mean change ± SD: G1: -1.89 ± 2.29 G2: -2.10 ± 2.32 G1/G2: P = 0.268</p> <p>UUI episodes/day median change (95% CI): G1: -1.33 (-2.00, -1.00) G2: -2.00 (-2.00, -1.33)</p> <p>Urgency episodes/day mean change ± SD: G1: -2.87 ± 3.59 G2: -2.68 ± 3.54 G1/G2: P = 0.882</p> <p>Urgency episodes/day median change (95% CI): G1: -2.33 (-3.00, -1.67) G2: -2.67 (-3.00, -2.00)</p> <p>Voids/day, mean change ± SD: G1: -2.96 ± 2.91 G2: -2.82 ± 2.87 G1/G2: P = 0.681</p> <p>Voids/day, median change (95% CI): G1: -2.67 (-3.33, -2.00) G2: -2.67 (-3.00, -2.33)</p> <p>Nocturia episodes/day, mean change ± SD: G1: -0.65 ± 1.26 G2: -0.67 ± 1.21 G1/G2: P = 0.315</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Masking: -</p> <p>Pt selection criteria: +</p> <p>Loss to followup: +</p> <p>Drop-out rates: -</p> <p>Power calculation: +</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Chancellor, Kianifard et al., 2008 (continued)			<p>Voids/day, median: G1: 11.33 G2: 11.33</p> <p>Nocturia episodes/day, mean ± SD: G1: 1.77 ± 1.43 G2: 1.87 ± 1.35</p> <p>Nocturia episodes/day, median: G1: 1.67 G2: 1.67</p> <p>Pads used/day, mean ± SD: G1: 1.12 ± 1.93 G2: 0.99 ± 1.67</p> <p>Pads used/day, median: G1: 0 G2: 0</p>	<p>Nocturia episodes/day, median change (95% CI): G1: -0.67 (-0.67, -0.33) G2: -0.67 (-0.67, -0.33)</p> <p>Pads used/day mean change ± SD: G1: -0.72 ± 1.54 G2: -0.61 ± 1.28 G1/G2: P = 0.978</p> <p>Pads used/day median change (95% CI): G1: 0 (0,0) G2: 0 (0,0)</p> <p>Side effects, %:** Constipation: 18.5 Dry mouth 25 UTI: 4.8 Headache: 3.8</p> <p>Discontinued due to adverse event(s), n (%) G1: 6 (3.2) G2: 21 (10.25)</p> <p>Discontinued due to constipation, %: 2</p> <p>Discontinued due to dry mouth, %: 1.8</p>	

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Chancellor, Zinner et al., 2008</p> <p>Country and setting: USA, multicenter</p> <p>Enrollment period: NR</p> <p>Funding: Astellas Pharma US, Glaxo-SmithKline</p> <p>Author industry relationship disclosures: 12 of 12 Abbot (1) AEterna Zentaris (1) Akros (1) Allergan (3) American Medical Systems (2) Amgen (1) Astellas (10) AstraZeneca (1) Bard (1) Bayer (1) Boehringer Ingelheim (3) Cephalon (1) Coloplast (1) Cook (1) CooperSurgical (1) Eli Lilly (2) Ferring GSK (6) Gynecare (1) Indevus (2) J&J (1) Mankind (1) Medtronic (1) Merck (1) Novartis (3) Novo Nordisk (1) Ortho-McNeil (1)</p>	<p>Design: Prospective cohort, open label, flexible-dose, 14 day washout</p> <p>Intervention: Solifenacin 5-10 mg daily</p> <p>Groups: G1: Solifenacin 5-10 mg daily</p> <p>N at enrollment: G1: 440</p> <p>N at follow-up (%): G1: 390 (88.4)</p> <p>Age, mean ± SD: G1: 61.4 ± 13.8</p> <p>Race/ethnicity, n (%): White: G1: 392 (88.9)</p> <p>Women, N (%): G1: 389 (88.2)</p> <p>BMI, kg/m² ± SD: G1: 30.1 ± 7.1</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 18 • OAB symptoms > 3 months • Treatment with tolterodine ER 4 mg, solifenacin, darifenacin, or trospium x ≥ 4 weeks • Desired change in therapy • ≥ 3 mean UUI episodes/day <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Treatment < 4 weeks with tolterodine, solifenacin, darifenacin, or trospium • SUI • MUI with primary stress • UTI • Chronic bladder inflammation • Bladder cancer • Severe constipation • Elevated PVR • Neurological deficit • Renal/ hepatic disease • Narrow angle glaucoma • Urinary retention • Gastric retention • Hypersensitivity to drugs • BOO • Women of childbearing age/ lactation 	<p>Urgency episodes/day, mean change: G1: 6.9 ± 4.4</p> <p>Incontinence episodes/day, mean ± SD: G1: 3.8 ± 3.6</p> <p>Voids/day, mean ± SD: G1: 11.3 ± 3.8</p> <p>Nocturia episodes/day, mean ± SD: G1: 1.9 ± 1.2</p> <p>PPBC score, mean: G1: 4.2</p> <p>OAB-q score, mean: Symptom bother: 57.3 Coping: 52.2 Concern: 51.3 Sleep: 50.6 Social Interaction: 77.7 Total score: 56.7</p>	<p>Urgency episodes/day, mean change ± SD (95% CI): G1: -4.2 ± 4.2 (-4.6, -3.8) G1/BL: P < 0.001</p> <p>Incontinence episodes/day, mean change ± SD (95% CI): G1: -2.6 ± 3.2 (-3.0, -2.3) G1/BL: P < 0.001</p> <p>Voids/day, mean change ± SD (95% CI): G1: -2.3 ± 3.2 (-2.6, -2.0) G1/BL: P < 0.001</p> <p>Nocturia episodes/day, mean change ± SD (95% CI): G1: -0.8 ± 1.0 (-0.9, -0.6) G1/BL: P < 0.001</p> <p>PPBC score, mean change (95% CI): G1: -1.2 (-1.3, 1.0) G1/BL: P < 0.001</p> <p>OAB-q score, mean: Symptom bother: 27.8 Coping: 80.1 Concern: 81.1 Sleep: 75.1 Social Interaction: 92.7 Total score: 81.9 G1/BL: P < 0.001 all domains and total score</p> <p>Dry Mouth, n (%): G1: 77 (17.5)</p> <p>Constipation, n (%): G1: 51 (11.6)</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: fair</p> <p>Randomization: -</p> <p>Method and blinding: NA</p> <p>Pt selection criteria: +</p> <p>Loss to followup: +</p> <p>Drop-out rates: +</p> <p>Power calculation: +</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Chancellor, Zinner, et al., 2008 (continued)				Blurred vision, n (%): G1: 10 (2.3)	
Author industry relationship disclosures:				UTI, n (%): G1: 19 (4.3)	
Pfizer (7)				Headache, n (%): G1: 13 (2.9)	
Pharmacia (1)					
Pri Med (1)					
Reliant (1)					
Solvay (1)					
Sanofi-Aventis (1)				URI, n (%): G1: 11 (2.5)	
Schering-Plough (1)					
Takeda (1)					
TAP (1)					
Watson (1)					

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Chapple et al. 2003</p> <p>Country and setting: 98 centers – United States, United Kingdom, Poland, Russia, New Zealand, Belgium, Netherlands</p> <p>Enrollment period: NR</p> <p>Funding: Yamanouchi Pharmaceutical Co., Ltd, Tokyo, Japan</p> <p>Author industry relationship disclosures: Not reported</p>	<p>Design: RCT, double-blind placebo controlled, with 2-wk placebo run-in period</p> <p>Intervention: tolterodine vs. solifenacin vs. placebo</p> <p>Groups: G1: tolterodine 2 mg bid G2: solifenacin 5 mg qd G3: solifenacin 10 mg qd G4: placebo qd</p> <p>N at enrollment: G1: 279 G2: 269 G3: 266 G4: 267</p> <p>N at follow-up: G1: 250 G2: 266 G3: 264 G4: 253</p> <p>Female, n (%) G1: 200 (80.0) G2: 188 (71.2) G3: 194 (72.9) G4: 193 (76.3)</p> <p>Age, yrs ± SD: G1: 56.9 +/-12.8 G2: 58.1± 13.4 G3: 57.2 ±13.4 G4: 57.8 ± 13.7</p> <p><65 years of age, n (%) G1: 169 (63.5) G2: 172 (65.2) G3: 172 (68.8) G4: 168 (66.4)</p> <p>≥65 years of age, n (%) G1: 97 (36.5) G2: 92 (34.8) G3: 78 (31.2) G4: 85 (33.6)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • ≥ 18 years of age • symptoms of OAB (including urinary frequency with urgency and/or urge incontinence) for ≥3 months • ≥8 voids per 24 hr • ≥3 episodes urinary incontinence during 3-day diary period <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • BOO • PVR >200 mL • stress predominant factor • neurological cause for detrusor overactivity • UTI or bladder stones • previous pelvic irradiation • malignant disease of pelvic organs • contraindication to antimuscarinic medication (including narrow-angle glaucoma, urinary or gastric retention) • non pharmacologic treatment for OAB 2 wks before study • diabetic neuropathy 	<p>UI only, n (%) G1: 142 (56.8) G2: 162 (61.4) G3: 172 (64.7) G4: 177(70.0)</p> <p>Mixed SI/UI, n (%) G1: 90 (36.0) G2: 81 (30.7) G3: 79 (29.7) G4: 59(23.3)</p> <p>No incontinence, n (%) G1: 18 (7.2) G2: 20 (7.6) G3: 15 (5.6) G4: 17 (6.7)</p> <p>Urgency episodes/24 h, mean ± SD G1: 5.77 ± 4.89 G2: 5.82 ± 4.45 G3: 5.45 ± 3.87 G4: 5.30 ± 3.92</p> <p>UI/24 h, mean ± SD G1: 2.33 ± 2.50 G2: 2.14 ± 2.44 G3: 1.86 ± 1.54 G4: 2.02 ± 2.50</p> <p>Incontinence episodes/24 h, mean ± SD: G1: 2.64 ± 2.55 G2: 2.59 ± 2.88 G3: 2.32 ± 1.94 G4: 2.71 ± 2.83</p> <p># Voids/24 h, mean ± SD: G1: 12.08± 3.86 G2: 12.32 ±3.95 G3: 12.08 ± 3.43 G4: 12.20 ± 4.11</p> <p>Volume voided/void, mL, mean ± SD: G1: 149.6 ± 54.6 G2: 147.2 ±51.2 G3: 147.0 ± 50.3 G4: 143.8 ± 53.6</p>	<p>Urgency episodes/24 h mean Δ from baseline: G1: -2.05 (3.58) G2: -2.85 (3.74) G3: -3.07 (3.90) G4: -1.41 (3.67)</p> <p>Urgency episodes/24 h Δ from baseline, %: G1: -38 G2: -55 G3: -52 G4: -33</p> <p>Urgency episodes estimated difference vs. tolterodine (95%CI) G1: -0.791 (-14.34, -0.148) p<0.001 G2: -1.015 (-1.659, -0.370) p<0.001 G3: p=0.0511</p> <p>UI/24 h mean Δ from baseline: G1: -0.91 (2.01) G2: -1.41 (1.74) G3: -1.36 (2.13) G4: -0.62 (1.96)</p> <p>UI/24 h Δ from baseline, %: G1: -58 G2: -65 G3: -63 G4: -40</p> <p>UI estimated difference vs. tolterodine (95%CI) G1: -0.487 (-0.988, 0.014) p=0.002 G2: -0.436 (-0.921, 0.048) p=0.0028 G3: p=0.2390</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: -</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: +</p> <p>Drop-out rates: ++</p> <p>Power calculation: +</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Chapple et al. 2003 (continued)	<p><75 years of age, n (%) G1: 236 (88.7) G2: 241 (91.3) G3: 233 (93.2) G4: 225 (88.9)</p> <p>≥75 years of age, n (%) G1: 30 (11.3) G2: 23 (8.7) G3: 17 (16.8) G4: 28 (11.1)</p> <p>Weight, mean kg ± SD G1: 74.8 +/-14.8 G2: 74.6±14.3 G3: 75.5±14.2 G4: 72.6±14.4</p> <p>Race, n (%) White G1: 261(98.1) G2: 260 (98.5) G3: 247(98.8) G4: 248 (98.0) Black G1: 2 (0.8) G2: 0 G3: 1 (0.4) G4: 1 (0.4) Asian G1: 2 (0.8) G2: 0 G3: 2(0.8) G4: 1 (0.4) Other G1: 1 (0.4) G2: 4 (1.5) G3: 0 G4: 3 (1.2)</p>	<ul style="list-style-type: none"> • use of drugs intended to treat incontinence • use of drugs with cholinergic or anticholinergic side-effects • participation in a clinical trial within 30 days • childbearing potential, pregnant or nursing or not using reliable contraceptive methods 	<p>Time from start of symptoms, mean months (%) G1: 57.4 (60.5) G2: 72.6 (105.4) G3: 62.9 (82.5) G4: 61.0 (83.9)</p> <p>Prior drug therapy, n (%) G1: 93 (34.9) G2: 106 (40.1) G3: 77 (30.8) G4: 83 (32.8)</p> <p>Any non drug therapy, n (%) G1: 92 (34.6) G2: 92 (34.8) G3: 88 (35.2) G4: 76 (30.0)</p>	<p>Incontinence episodes/24 h mean Δ from baseline: G1: -1.42 (1.82) G2: -1.45 (2.24) G3: -1.14 (2.15) G4: -0.76 (2.26)</p> <p>Incontinence episodes/24 h Δ from baseline, %: G1: -59 G2: -47 G3: -59 G4: -29</p> <p>Incontinence episodes estimated difference vs. tolterodine (95%CI) G1: -0.276 (-0.761, 0.208) p=0.008 G2: -0.316 (-0.786, 0.154) p=0.0038 G3: p=0.1122</p> <p>Voids/24 h mean Δ from baseline: G1: -1.88 (3.00) G2: -2.19 (2.87) G3: -2.61 (3.24) G4: -1.20 (3.26)</p> <p>Voids/24 h Δ from baseline, %: G1: -15 G2: -20 G3: -17 G4: -18</p> <p>Voids estimated difference vs. tolterodine (95%CI) G1: -0.312 (-0.844, 0.219) p=0.003 G2: -0.737 (-1.269, -0.204) p<0.001 G3: p=0.0145</p>	

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Chapple et al. 2003 (continued)				<p>Volume voided, mL mean Δ from baseline: G1: 24.4 (49.2) G2: 32.9 (47.7) G3: 39.2 (50.4) G4: 07.4 (36.3)</p> <p>Volume voided, mL Δ from baseline, %: G1: 20 G2: 25 G3: 29 G4: 9</p> <p>Volume voided, mL estimated difference vs. tolterodine (95%CI) G1: 8.4 (0.496, 16.34) p,0.001 G2: 14.8 (6.855, 22.72) p<0.001 G3: p<0.001</p> <p>Discontinued due to AEs, n (%) G1: 5 (1.9) G2: 9 (3.2) G3: 7 (2.6) G4: 10 (3.7)</p> <p>Dry mouth, n (%): G1: 49 (18.6) G2: 39 (14.0) G3: 57 (21.3) G4: 13 (4.9)</p> <p>Constipation, n (%) G1: 7 (2.6) G2: 20 (7.2) G2: 21 (7.8) G4: 5 (1.9)</p> <p>Blurred Vision, n (%) G1: 4 (1.5) G2: 10 (3.6) G3: 15 (5.6) G4: 7 (2.6)</p>	

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Chapple et al., 2005</p> <p>Chapple et al., 2007†</p> <p>Country and setting: European, Multicenter</p> <p>Enrollment period: May 2003 to October 2004</p> <p>Funding: Yamanouchi (now Astellas)</p> <p>Author industry relationship disclosures: NR* 1 of 9† Astellas (1) Novartis (1) Pfizer (1) Schwartz (1) UCB (1)</p>	<p>Design: RCT</p> <p>Intervention: Solifenacin 5mg vs. tolterodine ER 4mg</p> <p>Groups: G1: solifenacin 5mg G2: tolterodine ER 4mg</p> <p>G1a: G1 who elected not to increase dose at 4 weeks G2a: G2 who elected not to increase dose at 4 weeks</p> <p>N at enrollment: G1: 578 G2: 599</p> <p>N at follow-up: G1a: 297 G2a: 267</p> <p>Age, mean: G1: 56.5 G2: 56.4 G1a: 56.5 G2a: 56.9</p> <p>Race/ethnicity, %: White: G1: 99.3 G2: 99.5 G1a: 99.3 G2a: 99.3</p> <p>Women, %: G1: 85.3 G2: 88.3 G1a: 87.5 G2a: 87.1</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 18 • OAB symptoms for ≥ 3 months • outpatient treatment • ≥ 8 voids/day • ≥ 1 incontinence episode/day or ≥ 1 urgency episode/day <p>Exclusion criteria: NR</p>	<p>UII episodes/day, mean ± SD:* G1: 2.31 ± 2.35 G2: 2.12 ± 2.14</p> <p>UII episodes/day, mean:† G1a: 1.97 G2a: 1.66</p> <p>Urgency episodes/day, mean ± SD:* G1: 6.01 ± 4.66 G2: 5.84 ± 4.12</p> <p>Urgency episodes/day, mean:† G1a: 5.67 G2a: 5.30</p> <p>Incontinence episodes/day, mean ± SD:* G1: 2.77 ± 2.65 G2: 2.55 ± 2.37</p> <p>Incontinence episodes/day, mean:† G1a: 2.32 G2a: 2.01</p> <p>Voids/day, mean ± SD:* G1: 11.78 ± 3.58 G2: 11.66 ± 3.25</p> <p>Voids/day, mean:† G1: 11.10 G2: 11.36</p> <p>Nocturia, episodes/day, mean ± SD:* G1: 2.02 ± 1.33 G2: 1.92 ± 1.22</p> <p>Nocturia, episodes/day, mean:† G1a: 1.83 G2a: 1.84</p>	<p>UII episodes/day, 4 wks, mean change:† G1: -1.22 G2: -0.91 P = NS</p> <p>UII episodes/day, 12 wks, mean change:* G1: -1.42 G2: -0.83 P = 0.001</p> <p>UII episodes/day, 12 wks, mean change:† G1a: -1.46 G2a: -1.03</p> <p>Urgency episodes/day, 4 wks, mean change:† G1: -1.98 G2: -1.67 P = NS</p> <p>Urgency episodes/day, 12 wks, mean change:* G1: -2.85 G2: -2.42 P = 0.035</p> <p>Urgency episodes/day, 12 wks, mean change:† G1a: -3.08 G2a: -2.62</p> <p>Incontinence episodes/day, 4 wks, mean change:† G1: -1.30 G2: -0.90 P = 0.0181</p>	<p>Quality: Overall quality score: good</p> <p>INTERNAL VALIDITY: good</p> <p>Randomization: +</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: ++</p> <p>Drop-out rates: ++</p> <p>Power calculation: +</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Chapple et al., 2005*				Incontinence episodes/day, 12 wks, mean change:* G1: -1.60 G2: -1.11 P = 0.006	
Chapple et al., 2007† (continued)				Incontinence episodes/day, 12 wks, mean change:† G1a: -1.56 G2a: -1.23	
				Voids/day, 4 wks, mean change:† G1: -1.71 G2: -1.47 P = NS	
				Voids/day, 12 wks, mean change:* G1: -2.45 G2: -2.24 P = 0.004 for non-inferiority	
				Voids/day, 12 wks, mean change:† G1a: -2.47 G2a: -2.49	
				Nocturia, episodes/day, 4 wks, mean change:† G1: -0.51 G2: -0.44 P = NS	
				Nocturia, episodes/day, 12 wks, mean change:* G1: -0.71 G2: -0.63 P = NS	
				Nocturia episodes/day, 12 wks, mean change:† G1a: -0.72 G2a: -0.69	

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Chapple et al., 2005* Chapple et al., 2007† (continued)				<p>Perception of bladder condition, 4 wks, mean change:† G1: -0.96 G2: -0.88 P = NS</p>	
				<p>Perception of bladder condition, 12 wks, mean change:* G1: -1.51 G2: -1.33 P < 0.0061</p>	
				<p>Perception of bladder condition, 12 wks, mean change:† G1a: -1.72 G2a: -1.62</p>	
				<p>Pad use, 4 wks, mean change:† G1: -1.21 G2: -0.80 P = 0.0089</p>	
				<p>Pad use, 12 wks, mean change:* G1: -1.72 G2: -1.19 P < 0.0023</p>	
				<p>Pad use, 12 wks, mean change:† G1a: -1.55 G2a: -1.40</p>	
				<p>Dry rate, 4 wks, mean % change:† G1: 39 G2: 34 P = NS</p>	
				<p>Dry rate, 12 wks, mean % change:† G1a: 65.4 G2a: 58.3</p>	

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Chapple et al., 2005* Chapple et al., 2007† (continued)				<p>Voided volume (mL), 4 wks, mean change:† G1: 28.51 G2: 24.29 P = NS</p> <p>Voided volume (mL), 12 wks, mean change:* G1: 38 G2: 31 P = 0.010</p> <p>Voided volume (mL), 12 wks, mean change:† G1a: 39.95 G2a: 37.84</p> <p>Dry mouth, mild, %:*† G1: 17.5 G2: 14.8 G1a: 6.5 G2a: 5.0</p> <p>Dry mouth, moderate, %:*† G1: 10.8 G2: 7.7 G1a: 10.4 G2a: 7.0</p> <p>Dry mouth, severe, %:*† G1: 1.7 G2: 1.5 G1a: 0.7 G2a: 2.1</p> <p>Constipation, mild, %:*† G1: 3.2 G2: 1.3 G1a: 2.0 G2a: 1.0</p> <p>Constipation, moderate, %:*† G1: 2.7 G2: 1.0 G1a: 1.7 G2a: 1.4</p>	

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Chapple et al., 2005*				Constipation, severe, %:*†	
Chapple et al., 2007† (continued)				G1: 0.5	
				G2: 0.2	
				G1a: 0.3	
				G2a: 0.0	
				Blurred vision, mild, %:*†	
				G1: 0.7	
				G2: 0.7	
				G1a: 0.3	
				G2a: 0.7	
				Blurred vision, moderate, %:*†	
				G1: 0.0	
				G2: 1.0	
				G1a: 0.0	
				G2a: 1.7	
				Blurred vision, severe, %:*†	
				G1: 0.0	
				G2: 0.0	
				G1a: 0.0	
				G2a: 0.0	

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Chapple et al., 2008</p> <p>Country and setting: European, multicenter (academic and private)</p> <p>Enrollment period: NR</p> <p>Funding: Pfizer, Schwarz BioSciences</p> <p>Author industry relationship disclosures: 4 of 5 Pfizer (4)</p>	<p>Design: Multicenter randomized placebo-controlled</p> <p>Intervention: Tolterodine 4 mg ER vs. fesoterodine 8 mg</p> <p>Groups: G1: Tolterodine ER 4mg G2: Fesoterodine 8mg G3: Placebo</p> <p>N at enrollment: G1: 290 G2: 287 G3: 283</p> <p>Age, mean ± SD: Total: 57 ± 14</p> <p>Race/ethnicity, %: White: Total: >95</p> <p>Women, %: Total: 80</p> <p>Follow-up: 12 weeks</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 18 • OAB sx w/ urinary urgency for ≥6 mos • ≥ 8 voids/day • ≥ 6 urgency episodes or ≥ 3 UUI episodes/day • At least moderate problems recorded via a Likert scale • Negative pregnancy test • Adequate contraception throughout trial <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • LUT pathology: SI, bladder stones, interstitial cystitis, urothelial tumors • Grade III or higher pelvic prolapsed • Bladder-outlet obstruction • Polyuria (>3L/d) • Symptomatic or recurrent UTIs • PVR urine volume >100 mL • Antimuscarinic agent w/in 2 wks • Electrostimulation for bladder training in prior 4 wks • Active UTI • Underlying neurological disease • Clinically relevant cardiac arrhythmia • Unstable angina • QTcB interval > 500ms 	<p>Time since first diagnosis or onset of OAB (years), mean: Total: 8-9</p> <p>Incontinent at BL, n: G1: 213 G2: 217 G3: 203</p>	<p>KHQ severity score, mean change: G1: -12.6 G2: -14.0 G3: -9.0 G1/G3: $P < 0.05$ G2/G3: $P < 0.05$</p> <p>KHQ severity score, patients incontinent at BL, mean change: G1: -14.9 G2: -15.8 G3: -10.8 G1/G3: $P < 0.05$ G2/G3: $P < 0.05$</p> <p>KHQ emotions score, mean change: G1: -16.3 G2: -17.4 G3: -10.1 G1/G3: $P < 0.05$ G2/G3: $P < 0.05$</p> <p>KHQ emotions score, patients incontinent at BL, mean change: G1: -17.3 G2: -18.6 G3: -11.3 G1/G3: $P < 0.05$ G2/G3: $P < 0.05$</p> <p>KHQ role limitations score, mean change: G1: -22.1 G2: -21.7 G3: -11.8 G1/G3: $P < 0.05$ G2/G3: $P < 0.05$</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Method and blinding: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: NR</p> <p>Drop-out rates: NR</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Chapple et al., 2008 (continued)				<p>KHQ role limitations score, patients incontinent at BL, mean change: G1: -23.2 G2: -23.7 G3: -12.4 G1/G3: <i>P</i> < 0.05 G2/G3: <i>P</i> < 0.05</p> <p>KHQ physical limitations score, mean change: G1: -19.7 G2: -21.7 G3: -11.4 G1/G3: <i>P</i> < 0.05 G2/G3: <i>P</i> < 0.05</p> <p>KHQ Physical limitations score, patients incontinent at BL, mean change: G1: -20.5 G2: -23.3 G3: -11.1 G1/G3: <i>P</i> < 0.05 G2/G3: <i>P</i> < 0.05</p> <p>KHQ social limitations score, mean change: G1: -14.1 G2: -15.4 G3: -8.7 G1/G3: <i>P</i> < 0.05 G2/G3: <i>P</i> < 0.05</p> <p>KHQ social limitations score, patients incontinent at BL, mean change: G1: -15.7 G2: -16.2 G3: -9.5 G1/G3: <i>P</i> < 0.05 G2/G3: <i>P</i> < 0.05</p>	

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Chapple et al., 2008 (continued)				<p>KHQ sleep/energy score, mean change: G1: -11.7 G2: -13.6 G3: -9.6 G1/G3: P = NS G2/G3: P < 0.05</p> <p>KHQ sleep/energy score, patients incontinent at BL, mean change: G1: -12.5 G2: -15.3 G3: -10.4 G1/G3: P = NS G2/G3: P < 0.05</p> <p>KHQ personal relationship score, mean change: G1: -10.4 G2: -11.9 G3: -6.2 G1/G3: P = NS G2/G3: P < 0.05</p> <p>KHQ personal relationship score, patients incontinent at BL, mean change: G1: -12.7 G2: -12.3 G3: -6.8 G1/G3: P < 0.05 G2/G3: P = NS</p> <p>KHQ incontinence impact score, mean change: G1: -23.3 G2: -24.6 G3: -16.1 G1/G3: P < 0.05 G2/G3: P < 0.05</p>	

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Chaple et al., 2008 (continued)				<p>KHQ incontinence impact score, patients incontinent at BL, mean change: G1: -23.8 G2: -26.5 G3: -17.7 G1/G3: $P < 0.05$ G2/G3: $P < 0.05$</p> <p>KHQ general health score, mean change: G1: -4.3 G2: -4.0 G3: -3.8 G1/G3: $P = NS$ G2/G3: $P = NS$</p> <p>KHQ general health score, patients incontinent at BL, mean change: G1: -4.3 G2: -4.5 G3: -5.5 G1/G3: $P = NS$ G2/G3: $P = NS$</p> <p>ICIQ-SF total score, mean: G1: -3.95 G2: -4.41 G3: -2.55 G1/G3: $P < 0.05$ G2/G3: $P < 0.05$</p> <p>ICIQ-SF total score, patients incontinent at BL, mean: G1: -4.56 G2: -5.29 G3: -3.12 G1/G3: $P < 0.05$ G2/G3: $P < 0.05$</p>	

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Chapple et al., 2008 (continued)				<p>Major improvement in severity of bladder-related problems, %: G1: 34 G2: 39 G3: 25 G1/G3: $P = 0.01$ G2/G3: $P = 0.01$</p> <p>Dry mouth, n (%): G1: 49 (17) G2: 97 (34) G3: 20 (7)</p> <p>Constipation, n (%): G1: 8 (3) G2: 13 (5) G3: 4 (1)</p> <p>Nasopharyngitis, n (%): G1: 10 (3) G2: 5 (2) G3: 7 (3)</p> <p>Dry eye, n (%): G1: 1 (1) G2: 12 (4) G3: 0 (0)</p> <p>Nausea, n (%): G1: 6 (2) G2: 4 (1) G3: 1 (1)</p> <p>Fatigue, n (%): G1: 10 (3) G2: 1 (1) G3: 1 (1)</p> <p>Dry throat, n (%): G1: 3 (1) G2: 8 (3) G3: 0 (0)</p>	

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Colombo et al., 1995</p> <p>Country and setting: Italy; Setting</p> <p>Enrollment period: May 1990 to March 1993</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT, Computer generated random assignment</p> <p>Intervention: Oxybutynin vs. Bladder training for 6 weeks</p> <p>Groups: G1: Oxybutynin, 3 daily doses of 5 mg each for 6 weeks (dose reduced to half if substantial AEs) G2: Bladder training</p> <p>N at enrollment: G1: 42 G2: 39</p> <p>N at 6 wk follow-up: G1: 38 G2: 37</p> <p>N at 6 mo follow-up: G1: 28 G2: 27</p> <p>Age, yrs ± SD: G1: 48 (31 – 65) G2: 49 (24 – 65)</p> <p>Race/ethnicity, mean ± SD: Black: NR</p> <p>Women, N (%): G1: 42 (100) G2: 39 (100)</p> <p>Postmenopausal n (%): G1: 16 (38) G2: 20 (51)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Socially embarrassing (severe) urinary urge incontinence • On cystometry: detrusor instability, or LCB, or sensory bladder <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Stable bladder at cystometry • Neurologic disease • Detrusor hyperreflexia • Age greater than 65 y • Coexisting genuine SUI • Genital prolapse • Postvoid residual volume >50mL • Previous gynecologic or urogynecologic surgery • Prior use of any drug to treat UUI • Urethral diverticula • Fistulas • Urinary tract neoplasia • Cystitis • Bladder stones • Previous pelvic radiotherapy 	<p>Detrusor instability, n (%): G1: 14 (37) G2: 13 (35)</p> <p>Low compliance bladder, n (%): G1: 9 (24) G2: 8 (22)</p> <p>Sensory bladder, n (%): G1: 15 (39) G2: 16 (43)</p> <p>Daily UUI episodes, range 9 – 17</p> <p>Diurnal frequency, n (%): G1: 32 (84) G2: 29 (78)</p> <p>Nocturia, n (%): G1: 11 (29) G2: 18 (49)</p> <p>Volume at first desire (mL): G1: 120 ± 59 G2: 134 ± 61</p> <p>Volume at very strong desire (mL): G1: 317 ± 92 G2: 332 ± 87</p>	<p>Clinically cured overall, n (%): G1: 28 (74) G2: 27 (73)</p> <p>Cured among DI, (%): G1: 13 (93) G2: 8 (62) P=0.07</p> <p>Cured among LCB: G1: 6 (67) G2: 6 (75) P=0.56</p> <p>Cured among sensory bladder: G1: 9 (60) G2: 13 (81) P=0.18</p> <p>Cured among 18 G1 patients requiring dosage halving: 12 (67%)</p> <p>Diurnal frequency resolved, n (%): G1: 18 (56) G2: 20 (69)</p> <p>Nocturia resolved n (%): G1: 3 (27) G2: 11 (61)</p> <p>Volume at first desire (mL), mean ± SD: G1: 179 ± 32 P=.0009 G2: 178 ± 49 P=.001</p> <p>Volume at very strong desire (mL) mean ± SD: G1: 408 ± 76 P=.00001 G2: 403 ± 69 P=.0002</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Masking: -</p> <p>Pt selection criteria: +</p> <p>Loss to followup: ++</p> <p>Drop-out rates: +</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: -</p> <p>Measurement methods: +</p> <p>Measurement reliability: -</p> <p>Intervention description: +</p>

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Colombo et al. 1995 (continued)				<p>Patients still cured at 6 mos, n: G1: 16 G2: 26</p> <p>Patients still cured at 6 mos among DI, n: G1: 8 G2: 8</p> <p>Patients still cured at 6 mos among LCB, n: G1: 4 G2: 6</p> <p>Patients still cured at 6 mos among sensory bladder, n: G1: 4 G2: 12</p> <p>Treatment discontinued in 6 cases: G1: 4 (3 cases of severe dry mouth, 1 case of previously unknown glaucoma) G2: 2 (treatment was time consuming)</p> <p>Other adverse effects G1: 18 (47%) with AE requiring halving of dosage: -dry mouth (n=15) -constipation (n=6) -nausea (n=5) -dizziness (n=2) - decrease in visual acuity (n=1) - tachycardia (n=1)</p>	

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Diokno et al. 1995</p> <p>Country and setting: US; private clinic</p> <p>Enrollment period: January 1992 to December 1992</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Cohorts with comparison; (Series of patients self selected into two groups)</p> <p>Intervention: Bladder training vs. anticholinergic or antispasmodic</p> <p>Groups: G1: Bladder training G2: oxybutynin 2.5-5.0 mg bid or t.i.d.</p> <p>N at enrollment: G1: 39 G2: 33</p> <p>N at follow-up: G1: 26 G2: 28</p> <p>Age, mean yrs (range): 64 (20-93)</p> <p>Race/ethnicity: NR</p> <p>Women, N (%): 72 (100)</p> <p>Parity mean ± SD: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> Incontinent by AHCPR guidelines* Incontinence persisted after treatment for transient incontinence <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Post-void residual of more than 150 mL 	<p>UII, n (%): G1: 27 (49) G2: 28 (51)</p> <p>MUI with predominant urge, n (%): G1: 12 (71) G2: 5 (29)</p>	<p>Continent, n (%): G1: 1 (4) G2: 1 (4)</p> <p>Improved, n (%): G1: 22 (85) G2: 19 (68)</p> <p>No Change, n (%): G1: 3 (11) G2: 8 (28)</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: NA</p> <p>Masking: NA</p> <p>Pt selection criteria: -</p> <p>Loss to followup: -</p> <p>Drop-out rates: NR</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: poor</p> <p>Age: -, NR</p> <p>Baseline OAB status: NR</p> <p>Baseline characteristics: -</p> <p>Length of followup: ++</p> <p>Measurement methods: -</p> <p>Measurement reliability: -</p> <p>Intervention description: +</p>

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Diokno et al., 2003*</p> <p>Chu et al., 2005†</p> <p>Country and setting: US, Academic medical center</p> <p>Enrollment period: November 2000 to October 2001</p> <p>Funding: ALZA Corporation, Ortho-McNeil Pharmaceuticals</p> <p>Author industry relationship disclosures: 5 of 7* ALZA (1) Indevus (1) Ortho-McNeil (4) Pharmacia (1) Watson (1) NR†</p>	<p>Design: RCT</p> <p>Intervention: Oxybutynin ER vs. Tolterodine ER 12 week treatment period</p> <p>Groups: G1: Oxybutynin ER 10 mg daily G2: Tolterodine ER 4mg daily</p> <p>N at enrollment: G1: 391 G2: 399</p> <p>N at follow-up: G1: 339 G2: 357</p> <p>Age, mean (range) : G1: 60 (23-92) G2: 60 (18-85)</p> <p>Race/ethnicity, N (%): White: G1: 329 (84) G2: 342 (86) Black: G1: 32 (8) G2: 35 (9) Hispanic: G1: 27 (7) G2: 19 (5) Other: G1: 3 (1) G2: 3 (1)</p> <p>Women, N (%): G1: 391 (100) G2: 399 (100)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 18 • ≥ 21 episodes UUI/ week • ≥ 10 voids/ day • Urge > non-urge incontinence <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Treatable GU conditions causing incontinence • PVR > 150 mL by ultrasound x 2 • Risk of developing complete urinary retention • Medical problems worsened by anticholinergic effects • Hematuria • Uncontrolled narrow angle glaucoma • Obstructive uropathy • Reduced GI motility • Hypersensitivity to medications 	<p>UUI episodes/wk, mean ± SD: G1: 37.2 ± 15.2 G2: 36.9 ± 14.1</p> <p>Total incontinence episodes/wk, mean ± SD: G1: 43.3 ± 19.3 G2: 42.6 ± 18.2</p> <p>Voids/wk, mean ± SD: G1: 94.8 ± 25.4 G2: 96.2 ± 24.5</p>	<p>UUI episodes/wk, mean: G1: 10.8 G2: 11.2 P = 0.28</p> <p>Incontinence episodes/wk, mean: G1: 12.3 G2: 13.8 P = 0.08</p> <p>Voids/wk, mean: G1: 66.4 G2: 71.1 P = 0.03</p> <p>UUI episodes/wk, % reduction ± SD: G1: 72.0 ± 34.0 G2: 70.2 ± 33.2 P = 0.13</p> <p>Incontinent episodes/wk, % reduction ± SD: G1: 72.8 ± 31.0 G2: 69.1 ± 34.9 P = 0.08</p> <p>Voids/wk, % reduction ± SD: G1: 27.7 ± 19.7 G2: 24.9 ± 18.9 P = 0.05</p> <p>Total dryness, %: G1: 23 G2: 16.8 P = 0.03</p> <p>No UUI episodes, %: G1: 26.7 G2: 20.9 P = 0.06</p> <p>Dry mouth, any degree, n (%): G1: 116 (29.7) G2: 89 (22.3) P = 0.02</p> <p>Dry Mouth, mild, n (%): G1: 87 (22.3) G2: 69 (17.3)</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: fair</p> <p>Randomization: -</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: ++</p> <p>Drop-out rates: ++</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Diokno et al., 2003*				Dry mouth, moderate-severe, n (%): G1: 29 (7.4) G2: 20 (5.0)	
Chu et al., 2005† (continued)				Constipation, n (%): G1: 25 (6.4) G2: 31 (7.8)	
				Diarrhea, n (%): G1: 31 (7.9) G2: 25 (6.3)	
				Headache, n (%): G1: 22 (5.6) G2: 24 (6.0)	
				UTI, n (%): G1: 20 (5.1) G2: 13 (3.3)	
				CNS AE, n (%):† G1: 35 (9.0) G2: 33 (8.3)	
				Dizziness, n (%):† G1: 15 (3.8) G2: 10 (2.5)	
				Dizziness, mild,** %:† G1: 1.8 G2: 1.5	
				Dizziness, moderate,** %:† G1: 0.8 G2: 0.3	
				Somnolence, n (%):† G1: 4 (1.0) G2: 9 (2.3)	
				Somnolence, mild,** %:† G1: 0.5 G2: 1.5	
				Somnolence, moderate,** %:† G1: 0.3 G2: 0.5	
				Insomnia, n (%):† G1: 7 (1.8) G2: 3 (0.8)	

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Diokno et al., 2003*				Insomnia, mild**	
				%:†	
				G1: 0.8	
				G2: 0	
Chu et al., 2005† (continued)				Insomnia, moderate,** %:†	
				G1: 0.5	
				G2: 0	
				Depression, n	
				(%):†	
				G1: 5 (1.3)	
				G2: 3 (0.8)	
				Hypertonia, n	
				(%):†	
				G1: 2 (0.5)	
				G2: 4 (1.0)	
				Hypertonia, mild,** %:†	
				G1: 0	
				G2: 0	
				Hypertonia, moderate,** %:†	
				G1: 0.3	
				G2: 0	
				Anxiety, mild,** %:†	
				G1: 0.5	
				G2: 0	
				Anxiety, moderate,** %:†	
				G1: 0.3	
				G2: 0	
				Nervousness, mild,** %:†	
				G1: 0	
				G2: 0	
				Nervousness, moderate,** %:†	
				G1: 0.3	
				G2: 0	
				Tremor, mild,** %:†	
				G1: 0.3	
				G2: 0.3	
				Tremor, moderate,** %:†	
				G1: 0	
				G2: 0	

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Diokno et al., 2003*				Confusion, mild,** %:†	
Chu et al., 2005† (continued)				G1: 0.3 G2: 0.3	
				Confusion, moderate,** %:†	
				G1: 0 G2: 0	

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Giannitsas et al., 2004</p> <p>Country and setting: Greece, Specialty treatment center</p> <p>Enrollment period: NR</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Randomized for which drug to receive first two-way crossover, table of random numbers</p> <p>Intervention: Oxybutynin 15 mg t.i.d. vs. Tolterodine 4mg bid; 6 weeks treatment with 3-4 weeks washout</p> <p>Groups: G1: Oxybutynin 15 mg t.id. G2: Tolterodine 4mg bid</p> <p>Stratified by UDS findings: a: high volume (> 250mL); low pressure (< 25cmH₂O) b: high volume (≥ 250mL); high pressure (> 25cmH₂O) c: low volume (< 250mL); low pressure (< 25cmH₂O) d: low volume (< 250mL); high pressure (> 25cmH₂O)</p> <p>N at enrollment: Total: 128</p> <p>N at follow-up: Total: 107 Ga: 6 Gb: 25 Gc: 36 Gd: 40</p> <p>Women, %: 100</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 18 • DO on urodynamics <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Symptomatic or recurrent UTI • BOO • Neurologic disease • History of previous pelvic surgery • Narrow angle glaucoma • SUI • History of anticholinergic side effects • Interstitial cystitis • Child-bearing age without BC 	<p>Urinary voids/day, mean ± SD: Total: 8.5 ± 2.63 Ga: 7.2 ± NR Gb: 8.0 ± 2.40 Gc: 8.3 ± 2.31 Gd: 9.3 ± 2.91</p> <p>Volume (mL)/day, mean ± SD: Total: 1568.5 ± 398.64 Ga: 1756 ± NR Gb: 1594.6 ± 326.12 Gc: 1678.8 ± 402.15 Gd: 1420.5 ± 384.96</p> <p>Voided volume (mL), mean ± SD: Total: 196.1 ± 60.19 Ga: 253 ± NR Gb: 213.5 ± 53.57 Gc: 209.8 ± 57.43 Gd: 163.4 ± 51.97</p> <p>Bladder volume (mL), first void, mean ± SD: Total: 105.6 ± 39.38 Ga: 109 ± NR Gb: 128.0 ± 41.78 Gc: 94.3 ± 35.40 Gd: 101.1 ± 38.14</p> <p>Bladder volume (mL), first contraction, mean ± SD: Total: 172 ± 98.4 Ga: 258 ± NR Gb: 303.3 ± 60.51 Gc: 111.9 ± 48.11 Gd: 124.4 ± 56.66</p>	<p>Urinary voids/day, mean ± SD: G1: 7.7 ± 1.87 G2: 7.6 ± 2.18 G1a: 6.1 ± NR G2a: 6.3 ± NR G1b: 7.3 ± 1.69 G2b: 7.0 ± 1.87 G1c: 7.5 ± 1.41 G2c: 7.2 ± 1.58 G1d: 8.3 ± 2.23 G2d: 8.4 ± 2.53</p> <p>Volume (mL)/day, mean ± SD: G1: 1764.4 ± 333.03 G2: 1670.7 ± 338.6 G1a: 1862 ± NR G2a: 1720 ± NR G1b: 1715.6 ± 292.54 G2b: 1665.8 ± 251.19 G1c: 1847.9 ± 333.81 G2c: 1808.1 ± 317.59 G1d: 1694.7 ± 331.33 G2d: 1550.3 ± 373.65</p> <p>Voided volume (mL), mean ± SD: G1: 239.9 ± 64.98 G2: 236.7 ± 63.03 G1a: 321 ± NR G2a: 286 ± NR G1b: 243.3 ± 59.56 G2b: 248.3 ± 53.91 G1c: 252.9 ± 55.75 G2c: 258.6 ± 60.63 G1d: 217.6 ± 67.07 G2d: 203.1 ± 55.43</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Masking: NA</p> <p>Pt selection criteria: +</p> <p>Loss to followup: ++</p> <p>Drop-out rates: -</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: -</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating	
Giannitsas et al., 2004 (continued)	<p>Age, mean ± SD: Total: 56 ± 16.3 Ga: 53 ± 17.2 Gb: 57 ± 16.2 Gc: 57 ± 16.3 Gd: 54 ± 16.6</p> <p>Weight (kg), mean ± SD: Total: 63 ± 5.6 Ga: 63 ± 5.6 Gb: 70 ± 9.1 Gc: 67 ± 8.8 Gd: 69 ± 7.5</p>		<p>Pressure (cmH₂O), first contraction, mean ± SD: Total: 34.8 ± 21.97 Ga: 17.4 ± NR Gb: 37.7 ± 14.03 Gc: 18.5 ± 4.60 Gd: 50.3 ± 25.14</p> <p>Overactivity index, mean ± SD: Total: 36.8 ± 31.36 Ga: 15.3 ± NR Gb: 24.8 ± 19.66 Gc: 26.3 ± 16.14 Gd: 57.0 ± 38.85</p> <p>Cystometric capacity (mL), mean ± SD: Total: 362.8 ± 119.10 Ga: 403 ± NR Gb: 410.0 ± 97.78 Gc: 357.6 ± 127.52 Gd: 331.4 ± 114.17</p>	<p>Bladder volume (mL), first desire void, mean ± SD: G1: 129.0 ± 30.14 G2: 117.9 ± 27.62 G1a: 144 ± NR G2a: 140 ± NR G1b: 153.5 ± 25.72 G2b: 132.0 ± 31.01 G1c: 120.8 ± 25.07 G2c: 113.2 ± 23.16 G1d: 119.4 ± 29.43 G2d: 110.1 ± 26.15 G1/BL: P < 0.05 G2/BL: P < 0.05</p> <p>Bladder volume (mL), first contraction, mean ± SD: G1: 212.9 ± 106.10 G2: 206.9 ± 103.56 G1a: 382 ± NR G2a: 364 ± NR G1b: 355.28 ± 74.79 G1c: 142.4 ± 43.51 G2c: 144.6 ± 48.43 G1d: 173.3 ± 57.37 G2d: 162.7 ± 53.87 G1d/BL: P < 0.01 G2d/BL: P < 0.01</p>		

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Giannitsas et al., 2004 (continued)				<p>Pressure (cmH₂O), first contraction, mean ± SD: G1: 30.9 ± 22.63 G2: 30.9 ± 19.01 G1a: 14.0 ± NR G2a: 17.4 ± NR G1b: 35.6 ± 12.86 G2b: 29.6 ± 13.56 G1c: 17.2 ± 6.80 G2c: 17.9 ± 6.69 G1d: 42.9 ± 28.91 G2d: 44.1 ± 20.75</p> <p>Overactivity index, mean ± SD: G1: 24.4 ± 22.61 G2: 24.7 ± 23.46 G1a: 7.0 ± NR G2a: 9.5 ± NR G1b: 14.1 ± 12.09 G2b: 14.1 ± 12.71 G1c: 18.3 ± 15.89 G2c: 16.9 ± 16.84 G1d: 38.9 ± 26.50 G2d: 40.7 ± 26.58</p> <p>Cystometric capacity (mL), mean ± SD: G1: 419.3 ± 120.86 G2: 415.63 ± 114.06 G1a: 465 ± NR G2a: 453 ± NR G1b: 449.6 ± 106.23 G2b: 459.4 ± 101.17 G1c: 409.9 ± 130.22 G2c: 411.05 ± 132.49 G1d: 401.8 ± 118.34 G2d: 386.7 ± 96.53</p> <p>Dry mouth, n (%): G1: 52 (40.6) G2: 20 (15.6)</p>	

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Giannitsas et al., 2004 (continued)				Constipation, n (%): G1: 11 (10.3) G2: 3 (2.8) Discontinued due to AEs, n: Dry mouth: 12 Palpitations: 1	

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Goode et al., 2002</p> <p>Country and setting: US; academic health center outpatient geriatric medicine clinic</p> <p>Enrollment period: July 1989 to August 1995</p> <p>Funding: National Institutes on Aging</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT, placebo controlled</p> <p>Computer-generated random numbers using a block size of 6, w/ prior stratification by type and severity of incontinence</p> <p>Intervention: Biofeedback-assisted behavioral vs. drug treatment (oxybutynin chloride; possible range of doses 2.5 mg/d-5.0 mg t.i.d.) vs. placebo</p> <p>All patients had 4 visits over an 8-week period. Patients in G1 had biofeedback added to behavioral training in absence of 50% improvement by session 3.</p> <p>Groups: G1: Behavioral ± biofeedback G2: Pharmacologic G3: Placebo</p> <p>N at enrollment: 468 screened 271 not eligible 197 randomized 105 had pre and post treatment urodynamics</p> <p>G1: 33 G2: 35 G3: 37</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> Community-dwelling women at least age 55 Ambulatory At least 2 urge accidents per week by baseline bladder diary Urge incontinence as predominant pattern Urodynamic evidence of bladder dysfunction <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Continual leakage Postvoid residual urine volume >200mL Uterine prolapse past the introitus Narrow-angle glaucoma Unstable angina Decompensated congestive heart failure Hx of malignant arrhythmias MMSE <20 (Dementia) 	<p>Voids per day, mean n ± SD: G1: 10.0 G2: 10.9 G3: 10.0</p> <p>Cystometry at baseline, mL</p> <p>First desire to void G1: 97.1 ± 50.7 G2: 101.1 ± 62.1 G3: 124.6 ± 73.7</p> <p>Strong desire to void G1: 188.5 ± 93.1 G2: 212.1 ± 86.7 G3: 222.3 ± 87.0</p> <p>Bladder capacity G1: 288.3 ± 117.0 G2: 308.7 ± 93.7 G3: 328.9 ± 107.6</p>	<p>Voids per day, mean n ± SD: G1: 8.2 G2: 8.8 G3: 9.7</p> <p>DI on UDS, n (%) +Baseline DI/+DI post-treatment G1: 7 (21.2) G2: 1 (2.9) G3: 5 (13.5)</p> <p>DI on UDS, n (%) +Baseline DI/-DI post-treatment G1: 1 (3.0) G2: 7 (20.0) G3: 7 (18.9)</p> <p>DI on UDS, n (%) -Baseline DI/+DI post-treatment G1: 3 (9.1) G2: 3 (8.6) G3: 3 (8.1)</p> <p>DI on UDS, n (%) -Baseline DI/-DI post-treatment G1: 22 (66.7) G2: 24 (68.6) G3: 22 (59.5)</p> <p>Cystometry post-treatment, mL First desire to void G1: 115.9 ± 64.9 G2: 145.6 ± 74.0 G3: 133.5 ± 59.6</p> <p>Strong desire to void G1: 228.9 ± 106.4 G2: 282.0 ± 93.2 G3: 230.1 ± 78.8</p> <p>Bladder capacity G1: 305.6 ± 117.9 G2: 377.6 ± 92.1 G3: 323.0 ± 109.0</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: +</p> <p>Drop-out rates: +</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Goode et al., 2002 (continued)	<p>Age, yrs ± SD: G1: 65.3 ± 4.5 G2: 67.9 ± 7.9 G3: 67.6 ± 7.7</p> <p>Race/ethnicity, %: Black: 2 White: 98</p> <p>Women: 100%</p> <p>Parity mean ± SD: G1: 3.1 ± 1.7 G2: 2.1 ± 1.3 G3: 2.3 ± 1.5</p>			<p>Cystometry, change in mean volume by group</p> <p>First desire to void G1: 18.8 G2: 44.4 G3: 8.9 P= 0.149</p> <p>Strong desire to void G1: 40.5 G2: 69.9 G3: 7.8 P= 0.018</p> <p>Bladder capacity G1: 17.3 G2: 68.9 G3: -6.0 P= 0.000</p> <p><u>Standardized Estimates of Direct and Mediated Effects of Treatment</u></p> <p>G1 v G3 Total effect: 0.28* Direct effect: 0.23 Mediated Effect: 0.05</p> <p>G2vG3 Total Effect: 0.34* Direct Effect: 0.30* Mediated Effect: 0.04</p> <p>*P<0.01</p>	

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Halaska et al., 2003</p> <p>Country and setting: Europe and Asia, Academic medical center</p> <p>Enrollment period: May 1996 to May 1999</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Randomized controlled double blind crossover</p> <p>Intervention: Tropium vs. Oxybutynin</p> <p>Groups: G1: Trospium 20 mg bid G2: Oxybutynin 5mg bid</p> <p>N at enrollment: G1: 268 G2: 90</p> <p>N at follow-up: G1: 200 G2: 66</p> <p>Women, n (%): G1: 228 (85) G2: 78 (87)</p> <p>Age, mean (range): G1: 54.2 (19, 89) G2: 52.2 (19, 85)</p> <p>Weight (kg), mean (range): G1: 72.3 (50-120) G2: 70.4 (50-90)</p> <p>Height (cm), mean (range): G1: 164.8 (144-185) G2: 165.5 (145-183)</p> <p>Smokers, n (%): G1: 38 (14) G2: 10 (11)</p> <p>Previous illness, n (%): G1: 184 (69) G2: 66 (73)</p> <p>Previous medication, n (%): G1: 101 (38) G2: 46 (51)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age \geq 18 yrs • Urge syndrome • Urge incontinence • UUI as one component of MUI • UUI due to neurologic condition (detrusor hyperreflexia) <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Absolute tachycardia • Closed angle glaucoma • Myasthenia gravis • Arteriosclerosis of cerebral vessels • SUI • Heart or renal failure • Frequency from diuretics • BOO • Acute UTI • Hiatus hernia with reflux esophagitis • Stenosis of GI tract • Megacolon • Colonic ulceration • Allergy to study medications • Anticholinergics, TCAs, alpha blockers, beta sympathomimetics \leq 7 days • Urological or gynecologic surgery \leq 3 mos • Pregnant or lactating • In another study 	<p>Urgency episodes/day, mean: G1: 10.2 G2: 11.0</p> <p>Incontinence episodes/ day, mean: G1: 1.5 G2: 2.1</p> <p>Voids/day, mean: G1: 11.4 G2: 12.5</p> <p>Max cystometric capacity (mL), mean: G1: 205 G2: 205</p>	<p>Urgency episodes/day, mean: G1: 6.7 G2: 7.4</p> <p>Incontinence episodes/ day, mean: G1: 0.5 G2: 1.1</p> <p>Voids/day, mean: G1: 7.9 G2: 8.3</p> <p>Max cystometric capacity (mL), 26 wks, mean change: G1: 92.0 G2: 117.0 G1/BL: $P \leq 0.001$ G2/BL: $P \leq 0.001$</p> <p>Max cystometric capacity (mL), 52 wks, mean change: G1: 115.0 G2: 119.4 G1/BL: $P \leq 0.001$ G2/BL: $P \leq 0.001$</p> <p>Bladder volume (mL), first contraction, 26 wks, mean change: G1: 63.5 G2: 61.2</p> <p>Bladder volume (mL), first contraction, 52 wks, mean change: G1: 46.1 G2: 36.7</p> <p>Bladder volume (mL), first sensation, 26 wks, mean change: G1: 73.6 G2: 76.93</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: -</p> <p>Masking: -</p> <p>Pt selection criteria: +</p> <p>Loss to followup: ++</p> <p>Drop-out rates: ++</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: ++</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Halaska et al., 2003 (continued)				<p>Bladder volume (mL), first sensation, 26 wks, mean change: G1: 78.6 G2: 70.2</p> <p>Abnormal EKG, n (%): G1: 4 (0.1) G2: 2 (0.2)</p> <p>Any adverse event, n (%): G1: 10 (3.7) G2: 6 (6.7)</p> <p>Poor efficacy, n (%): G1: 8 (3) G2: 2 (2.2)</p> <p>Poor compliance, n (%): G1: 15 (15.6) G2: 6 (6.7)</p> <p>Abdominal pain, n (%): G1: 5 (2) G2: 0 (0)</p> <p>Constipation, n (%): G1: 18 (7) G2: 4 (4)</p> <p>Diarrhea, n (%): G1: 2 (1) G2: 2 (2)</p> <p>Dyspepsia, n (%): G1: 13 (5) G2: 3 (3)</p> <p>Dysphagia, n (%): G1: 9 (3) G2: 3 (3)</p> <p>Dry mouth, n (%): G1: 87 (33) G2: 45 (50) P < 0.001</p> <p>Nausea, n (%): G1: 6 (2) G2: 2 (2)</p>	

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Halaska et al., 2003 (continued)				UTI, n (%): G1: 33 (12) G2: 10 (11) Headache, n (%): G1: 11 (4) G2: 8 (9) Visual disturbances, n (%): G1: 9 (3) G2: 5 (6) Virus infection, n (%): G1: 9 (3) G2: 4 (4) Abnormal EKG, n (%): G1: 4 (0.1) G2: 2 (0.2) Sleeplessness, n (%): G1: 10 (4) G2: 2 (2)	

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Herschorn et al., 2004</p> <p>Country and setting: Canada; Family medicine and urology clinics</p> <p>Enrollment period: June 2000 to December 2001</p> <p>Funding: Pharmacia Pfizer Canada</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: Health education with tolterodine vs. tolterodine alone</p> <p>Groups: G1: Health education with tolterodine G2: tolterodine alone</p> <p>N at enrollment: G1: 39 G2: 45</p> <p>N at follow-up, 5 weeks: G1: 37 G2: 40</p> <p>N at follow-up, 10 weeks: G1: 35 G2: 32</p> <p>N at follow-up, 16 weeks: G1: 34 G2: 31</p> <p>Age, yrs ± SD: G1: 65.7 ± 14.5 G2: 63.1 ± 15.7</p> <p>Race/ethnicity: NR</p> <p>Women, %: G1: 92.3 G2: 84.4</p> <p>Parity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 50 • Symptoms of OAB[^] • Attend investigators' practice • Normal cognitive function • Able to read English <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Enrollment in another clinical trial • Interstitial cystitis • UTI • Already taking tolterodine 	<p>Duration of OAB, yrs ± SD: G1: 8.7 ± 11.0 G2: 8.7 ± 10.8</p> <p>Mild bladder problems, n (%): G1: 13 (33.3) G2: 13 (28.9)</p> <p>Moderate bladder problems, n (%): G1: 19 (48.7) G2: 28 (62.2)</p> <p>Severe bladder problems, n (%): G1: 7 (18.0) G2: 4 (8.9)</p> <p>Obtained prescription, n (%): G1: 38 (97.4) G2: 37 (82.2) P < 0.05</p> <p>Intends to fill prescription, (%): G1: 0 (0) G2: 6 (7.5)</p>	<p>Change in incontinence episodes/wk, mean ± SD: G1: -7.72 ± 21.16 G2: -10.24 ± 19.56†</p> <p>Change in voids/d, mean ± SD: G1: -1.82 ± 3.41 G2: -2.18 ± 4.89 P = NR</p> <p>Change in nocturnal voids, mean ± SD: G1: -0.44 ± 1.13† G2: -0.07 ± 0.91</p> <p>No change in bladder problem severity, n (%): G1: 14 (42.4) G2: 20 (66.7)</p> <p>Improved bladder problem severity, n (%): G1: 15 (45.4)† G2: 6 (20)</p> <p>Worsened bladder problem severity, n (%): G1: 4 (12.1) G2: 4 (13.3)</p> <p>Compliance, 10 weeks, %: G1: 41 G2: 38 P > 0.05</p> <p>Compliance, 16 weeks, %: G1: 39 G2: 31 P > 0.05</p> <p>Continued or started non-drug OAB treatment, 16 weeks, %: G1: 82 G2: 53 P > 0.05</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: -</p> <p>Masking: -</p> <p>Pt selection criteria: +</p> <p>Loss to followup: -</p> <p>Drop-out rates: NR</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: +</p> <p>Baseline OAB status: NR</p> <p>Baseline characteristics: -</p> <p>Length of followup: ++</p> <p>Measurement methods: +</p> <p>Measurement reliability: -</p> <p>Intervention description: +</p>

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Herschorn et al., 2004 (continued)				Stopped non-drug OAB treatments, %: G1: 12.8 G2: 28.9	

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Jarvis et al., 1981</p> <p>Country and setting: UK, Academic medical center</p> <p>Enrollment period: NR</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: Inpatient bladder drill vs. outpatient drug therapy</p> <p>Groups: G1: inpatient bladder drill G2: flavoxate hydrochloride 200 mg tds and imipramine 25 mg tds x 4 wks</p> <p>N at enrollment: G1: 25 G2: 25</p> <p>N at follow-up: G1: 25 G2: 25</p> <p>Women, %: 100</p> <p>Age, mean ± SD (range): G1: 47 ± 15.4 (17-78) G2: 46 ± 12.8 (17-65)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Women • UDS-diagnosed detrusor instability <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • DM • Neurological abnormalities • UTIs • Taking a drug suspected of affecting lower urinary tract function • Genuine stress incontinence 	<p>UII, n: G1: 25 G2: 25</p> <p>SUI, n: G1: 19 G2: 17</p> <p>Urgency, n: G1: 25 G2: 25</p> <p>Frequency, n: G1: 25 G2: 25</p> <p>Nocturia, n: G1: 21 G2: 19</p> <p>Duration of symptoms (years), mean ± SD (range): G1: 4.3 ± 2.7 (1-15) G2: 5.4 ± 3.2 (1-20)</p> <p>Bladder volume (mL), first sensation, mean: G1: 87 G2: 79</p> <p>Max cystometric capacity (mL), mean: G1: 381 G2: 353</p>	<p>UII, n: G1: 4 G2: 11</p> <p>SUI, n: G1: 1 G2: 9</p> <p>Urgency, n: G1: 4 G2: 11</p> <p>Frequency, n: G1: 6 G2: 12</p> <p>Nocturia, n: G1: 4 G2: 13</p> <p>Continent, n (%): G1: 21 (84) G2: 14 (56) <i>P</i> < 0.05</p> <p>Symptom-free, n (%): G1: 19 (76) G2: 12 (48) <i>P</i> < 0.05</p> <p>Bladder volume (mL), first sensation, mean: G1: 152 G2: 140</p> <p>Max cystometric capacity (mL), mean: G1: 470 G2: 446</p> <p>Adverse events, drug therapy, n:* Dizziness: 8 Headache: 6 Dry mouth: 6 Nausea: 4 Drowsiness: 2 Vomiting: 1</p> <p>Discontinued due to AEs, n: Dizziness: 1 Headache: 1 Vomiting: 1</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: -</p> <p>Masking: -</p> <p>Pt selection criteria: +</p> <p>Loss to follow up: ++</p> <p>Drop-out rates: -</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of follow up: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Lauti et al., 2008</p> <p>Country and setting: New Zealand, Academic</p> <p>Enrollment period: February 2003 to July 2003</p> <p>Funding: University of Otago</p> <p>Author industry relationship disclosures: None</p>	<p>Design: RCT pilot study, unmasked</p> <p>Intervention: Oxybutynin vs. bladder retraining vs. combination therapy</p> <p>Groups: G1: Oxybutynin 2.5 mg/day (daily dose could be increased by 2.5 mg every 5 days to a maximum of 15 mg/day) G2: Bladder retraining G3: Combination therapy</p> <p>N screened: 120</p> <p>N at enrollment: G1: 21 G2: 16 G3: 19</p> <p>N at 3 month follow-up: G1: 18 G2: 16 G3: 12</p> <p>N at 12 month follow-up: G1: 16 G2: 14 G3: 12</p> <p>Age, mean ± SD: G1: 53.8 ± 14.8 G2: 63.9 ± 17.2 G3: 47.6 ± 16.3</p> <p>Race/ethnicity: NR</p> <p>Women, %: Total: 100</p> <p>Parous, %: G1: 81 G2: 62.5 G3: 73.7</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age > 18 • Predominant UI <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Predominant SUI • Contraindications to anticholinergic drugs • Current UTI • Neurological disease • Psychiatric disorder • Untreated co-existing pelvic organ prolapse below the hymenal ring • Obstructed voiding • Functional-reversible cause of incontinence • Inability to toilet independently • Limited fluency of written/spoken English • Current or recent use of any of the trial interventions 	<p>Urgency episodes/day, mean ± SD: G1: 3.8 ± 2.7 G2: 3.1 ± 2.2 G3: 3.5 ± 2.0</p> <p>Incontinence episodes/day, mean ± SD: G1: 2.2 ± 1.5 G2: 1.0 ± 1.1 G3: 1.8 ± 1.6</p> <p>Nocturia episodes/day, mean ± SD: G1: 1.1 ± 1.0 G2: 1.4 ± 1.0 G3: 0.8 ± 0.7</p> <p>Voids per day, mean ± SD: G1: 7.8 ± 2.8 G2: 8.0 ± 1.7 G3: 8.4 ± 2.5</p> <p>OAB-q total HRQL, mean ± SD: G1: 73.1 ± 17.4 G2: 69.5 ± 24.6 G3: 71.6 ± 21.5</p> <p>OAB-q severity, mean ± SD: G1: 47.0 ± 16.2 G2: 42.3 ± 17.7 G3: 45.9 ± 18.7</p> <p>OAB-q coping, mean ± SD: G1: 72.0 ± 21.6 G2: 66.2 ± 31.7 G3: 73.8 ± 26.2</p> <p>OAB-q concern, mean ± SD: G1: 68.2 ± 19.0 G2: 68.8 ± 27.6 G3: 63.8 ± 29.2</p> <p>OAB-q sleep, mean ± SD: G1: 63.1 ± 28.7 G2: 59.8 ± 29.9 G3: 55.1 ± 27.6</p>	<p>Urgency episodes/day, 3 mos, mean ± SD: G1: 2.2 ± 1.8 G2: 1.5 ± 2.1 G3: 1.7 ± 1.8</p> <p>Urgency episodes/day, 12 mos, mean ± SD: G1: 2.3 ± 2.5 G2: 1.9 ± 2.1 G3: 2.0 ± 1.1</p> <p>Incontinence episodes/day, 3 mos, mean ± SD: G1: 0.8 ± 0.8 G2: 0.1 ± 0.3 G3: 0.6 ± 0.8</p> <p>Incontinence episodes/day, 12 mos, mean ± SD: G1: 0.9 ± 0.0 G2: 0.9 ± 1.0 G3: 0.8 ± 0.7</p> <p>Nocturia episodes/day, 3 mos, mean ± SD: G1: 1.0 ± 0.5 G2: 0.8 ± 0.7 G3: 0.6 ± 0.5</p> <p>Nocturia episodes/day, 12 mos, mean ± SD: G1: 1.0 ± 0.9 G2: 1.2 ± 0.6 G3: 0.7 ± 0.7</p> <p>Voids/day, 3 mos, mean ± SD: G1: 6.7 ± 1.8 G2: 6.3 ± 1.6 G3: 6.7 ± 2.2</p> <p>Voids/day, 12 mos, mean ± SD: G1: 7.2 ± 1.1 G2: 6.8 ± 1.4 G3: 7.6 ± 1.5</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Method and blinding: -</p> <p>Pt selection criteria: +</p> <p>Loss to followup: +</p> <p>Drop-out rates: +</p> <p>Power calculation: +</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: ++</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Lauti et al., 2008 (continued)			<p>OAB-q social, mean ± SD: G1: 92.5 ± 14.6 G2: 85.4 ± 19.9 G3: 92.8 ± 18.6</p> <p>SF-12 quality of life, physical, mean ± SD: G1: 49.0 ± 9.6 G2: 41.7 ± 11.5 G3: 46.2 ± 10.6</p> <p>SF-12 quality of life, mental, mean ± SD: G1: 49.1 ± 9.3 G2: 53.1 ± 8.8 G3: 46.3 ± 8.3</p>	<p>OAB-q total HRQL, 3 mos, mean ± SD: G1: 82.3 ± 16.1 G2: 89.6 ± 9.4 G3: 91.8 ± 7.4</p> <p>OAB-q total HRQL, 12 mos, mean ± SD: G1: 87.9 ± 11.6 G2: 81.6 ± 19.3 G3: 88.9 ± 9.9</p> <p>OAB-q severity, 3 mos, mean ± SD: G1: 37.2 ± 22.0 G2: 16.8 ± 12.0 G3: 21.6 ± 10.9</p> <p>OAB-q severity, 12 mos, mean ± SD: G1: 24.6 ± 10.6 G2: 33.1 ± 16.6 G3: 21.9 ± 14.8</p> <p>OAB-q coping, 3 mos, mean ± SD: G1: 79.2 ± 22.1 G2: 91.6 ± 9.5 G3: 92.7 ± 9.4</p> <p>OAB-q coping, 12 mos, mean ± SD: G1: 89.2 ± 13.7 G2: 81.5 ± 23.7 G3: 90.5 ± 10.0</p> <p>OAB-q concern, 3 mos, mean ± SD: G1: 78.6 ± 18.0 G2: 87.7 ± 14.5 G3: 90.2 ± 12.4</p> <p>OAB-q concern, 12 mos, mean ± SD: G1: 85.3 ± 15.5 G2: 81.7 ± 19.7 G3: 85.2 ± 13.4</p> <p>OAB-q sleep, 3 mos mean ± SD: G1: 77.7 ± 24.9 G2: 81.3 ± 14.6 G3: 85.0 ± 19.6</p>	

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Lauti et al., 2008 (continued)				<p>OAB-q sleep, 12 mos, mean ± SD: G1: 79.9 ± 18.3 G2: 72.0 ± 24.5 G3: 83.2 ± 18.4</p> <p>OAB-q social, 3 mos, mean ± SD: G1: 96.4 ± 9.7 G2: 95.6 ± 7.0 G3: 98.9 ± 1.9</p> <p>OAB-q social, 12 mos, mean ± SD: G1: 97.3 ± 7.1 G2: 91.9 ± 14.2 G3: 97.3 ± 6.9</p> <p>SF-12 quality of life, physical G1: 50.6 ± 8.0 G2: 42.1 ± 12.7 G3: 48.4 ± 10.8</p> <p>SF-12 quality of life, physical, 12 mos, mean ± SD: G1: 50.0 ± 7.3 G2: 45.1 ± 13.9 G3: 45.3 ± 13.4</p> <p>SF-12 quality of life, mental, 3 mos, mean ± SD: G1: 50.4 ± 9.6 G2: 51.2 ± 9.5 G3: 46.7 ± 7.6</p> <p>SF-12 quality of life, mental, 12 mos, mean ± SD: G1: 49.6 ± 7.5 G2: 50.1 ± 10.7 G3: 50.6 ± 8.4</p> <p>Dry mouth, n (%): G1: 3 (21) G2: 5 (46) G3: 5 (42)</p> <p>Headaches, n (%): G1: 6 (43) G2: 1 (11) G3: 7 (58)</p> <p>Dizziness, n (%): G1: 4 (29) G2: 2 (20) G3: 3 (25)</p>	

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Lauti et al., 2008 (continued)				Constipation, n (%): G1: 3 (21) G2: 3 (27) G3: 3 (27) Fatigue, n (%): G1: 9 (64) G2: 5 (46) G3: 7 (64)	

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Lee et al., 2002</p> <p>Country and setting: South Korea, University</p> <p>Enrollment period: NR</p> <p>Funding: Pharmacia Corp</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: Tolterodine 2mg bid vs. Oxybutynin 5mg bid</p> <p>Groups: G1: Tolterodine 2mg bid G2: Oxybutynin 5mg bid</p> <p>N at enrollment: G1: 112 G2: 116</p> <p>N at follow-up: G1: 97 G2: 90</p> <p>Women, n (%): G1: 84 (74) G2: 92 (79)</p> <p>Age, mean (range): G1: 52 (27, 82) G2: 52 (20, 86)</p> <p>Race/ethnicity (%): Asian: G1: 100 G2: 100</p> <p>BMI, kg/m² (range): G1: 23 (17, 32.5) G2: 23.5 (16, 38)</p> <p>Previous drug therapy: N (%) G1: 36 (32) G2: 26 (22)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 18 • OAB symptoms > 6 mos • ≥ 8 voids/day, with or without incontinence (measured by diary) <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • SUI • Women not using reliable contraception • Pregnant or nursing • Prior treatment with anticholinergic < 2 wks • Renal or hepatic disease • Narrow angle glaucoma • Urinary retention • Gastric retention • Hypersensitivity to drugs • UTI • IC • Hematuria • BOO • Concomitant bladder training, e-stim treatment • Indwelling catheter • Intermittent catheterization • Concomitant treatment for OAB ≤ 2 mos 	<p>Incontinence episodes/day, mean (range): G1: 2.6 (0.3, 9.3) G2: 2.4 (3.0, 14.7)</p> <p>Patients with incontinence episodes, n (%): G1: 46 (41) G2: 42 (36)</p> <p>Voids/day, mean (range): G1: 12.2 (8.0, 23.7) G2: 12.4 (7.7, 29.7)</p>	<p>Incontinence episodes/day, mean change ± SD (% change): G1: -2.2 ± 2.3 (-85) G2: -1.4 ± 1.8 (-58)</p> <p>G1/BL: P = 0.0001 G1/BL: P = 0.0001 G1/G2: P = 0.10</p> <p>Voids/ day, mean change ± SD (% change): G1: -2.6 ± 2.9 (-21) G2: -1.8 ± 4.2 (-15)</p> <p>G1/BL: P = 0.0001 G1/BL: P = 0.0001 G1/G2: P = 0.14</p> <p>Benefit of Tx, %: G1: 45 G2: 46 G1/G2: P = NS</p> <p>Patients reporting adverse events, n (%): G1: 62 (55) G2: 94 (82)</p> <p>Discontinued due to AEs, n (%): G1: 11 G2: 18</p> <p>Dry mouth, n (%): G1: 39 (35) G2: 72 (63) G1/G2: P = 0.001</p> <p>Dry mouth, mild, n (%): G1: 29 (26) G2: 40 (35)</p> <p>Dry mouth, moderate, n (%): G1: 9 (8) G2: 26 (23)</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: -</p> <p>Drop-out rates: -</p> <p>Power calculation: +</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: -</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Lee et al., 2002 (continued)				<p>Dry mouth, severe, n (%): G1: 1 (1) G2: 6 (5)</p> <p>Voiding disorder, n (%): G1: 10 (9) G2: 16 (14)</p> <p>Dyspepsia, n (%): G1: 8 (7) G2: 6 (5)</p> <p>Abdominal pain, n (%): G1: 6 (5) G2: 6 (5)</p> <p>Headache, n (%): G1: 4 (4) G2: 6 (5)</p>	

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Macaulay et al., 1987</p> <p>Country and setting: UK, Specialty treatment center</p> <p>Enrollment period: NR</p> <p>Funding: Wellcome Trust, trustees of St. George's Hospital</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT (randomization not specified)</p> <p>Intervention: Brief eclectic psychotherapy, bladder training or medication</p> <p>Groups: G1: psychotherapy G2: bladder drill G3: propantheline</p> <p>N at enrollment: G1: 19 G2: 16 G3: 15</p> <p>N at follow-up: G1: 18 G2: 15 G3: 14</p> <p>Women, %: 100</p> <p>Age: NR</p> <p>Race/ethnicity: NR</p> <p>Follow-up: 3 months</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Previous completion of survey on psych conditions w/ OAB • Detrusor instability on UDS or sensory urgency <p>Exclusion criteria: NR</p>	<p>Detrusor instability, n: G1: 10 G2: 8 G3: 8</p> <p>Sensory urgency, n: G1: 9 G2: 8 G3: 7</p> <p>Voids/day, mean: G1: NR G2: NR G3: 10.8</p> <p>Bladder capacity (mL), mean: G1: 393 G2: NR G3: 323</p> <p>Bladder volume (mL), first sensation, mean: G1: 107 G2: 110 G3: 70</p> <p>Detrusor pressure rise (cm H₂O), mean: G1: NR G2: 45.5 G3: NR</p>	<p>Voids/day, mean: G1: NR G2: NR G3: 8.3 G3/BL: $P < 0.005$</p> <p>Bladder capacity (mL), mean: G1: 414 G2: NR G3: 368 G1/BL: $P = NS$ G3/BL: $P = NS$</p> <p>Bladder volume (mL), first sensation, mean: G1: 142 G2: 150 G3: 137 G1/BL: $P = NS$ G2/BL: $P < 0.05$ G3/BL: $P = 0.06$</p> <p>Detrusor pressure rise (cm H₂O), mean: G1: NR G2: 29.5 G3: NR G2/BL: $P < 0.05$</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: -</p> <p>Masking: -</p> <p>Pt selection criteria: -</p> <p>Loss to followup: ++</p> <p>Drop-out rates: ++</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: -, NR</p> <p>Baseline OAB status: NR</p> <p>Baseline characteristics: -</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Mattiasson et al., 2003</p> <p>Country and setting: Sweden, Denmark, Norway (Tolterodine Scandinavian Study Group)</p> <p>Enrollment period: October 1999 to December 2000</p> <p>Funding: Pharmacia Corp</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT, single-blinded (balanced blocks of 4, computerized randomization list)</p> <p>Intervention: Pharmacologic ± behavioral therapy; Tolterodine 2mg b.i.d. ± BT; Tolterodine dosage could be decreased to 1mg P.O. b.i.d. during the first 2 wks if intolerable SE; BT taught with a written handout</p> <p>Groups: G1: Tolterodine + BT x 24 wks G2: Tolterodine x 24 wks</p> <p>N at enrollment: Total: 505</p> <p>N at follow-up: G1: 244 G2: 257</p> <p>N Completed treatment: G1: 77% G2: 79% Total: 74%</p> <p>Women, n (%) G1: 177 (73) G2: 201 (78)</p> <p>Age, median (range): G1: 62 (19, 86) G2: 63 (22, 86)</p> <p>Race/ethnicity: NR</p> <p>Follow-up: 24 weeks</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 18 • ≥ 8 voids/day and urinary urgency (± UUI) as determined by 1 wk bladder diary • With or without UUI • Women of reproductive age had to be using reliable birth control <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Contraindication to antimuscarinic therapy • Use of electrostimulation therapy or BT within prior 3 mo • Indwelling catheter or intermittent catheterization • Pregnancy or lactation • Use of anticholinergic agents or concomitant treatment for OAB (estrogen permitted) 	<p>Urgency episodes/day, mean (range): G1: 6.0 (0, 23.0) G2: 6.6 (0, 34.3)</p> <p>Incontinence episodes/day, mean (range): G1: 2 (0.3, 20.3) G2: 2.3 (0.3, 16.3)</p> <p>voids/day, mean (range): G1: 10.3 (7.3, 27.6) G2: 10.6 (7.7, 24.6)</p> <p>Duration of symptoms > 5 years, n (%): G1: 120 (49) G2: 124 (48)</p> <p>Previous drug therapy for OAB, n (%): G1: 40 (160) G2: 35 (14)</p> <p>Previous surgery affecting lower urinary tract, n (%): G1: 28 (11.5) G2: 30 (12)</p> <p>Symptoms caused at least moderate problems, %: G1: 92 G2: 93</p>	<p>Urgency episodes/day, median % change (IQR): G1: -38 (-76.7, -14.1) G2: -38 (-68.7, -8.0) G1/G2: <i>P</i> = 0.75</p> <p>Incontinence episodes/day, median IQR% change; n=301: G1: -87 (-100, -20) G2: -81 (-100, -41.8) G1/G2: <i>P</i> = 0.28</p> <p>voids/day, median % change (IQR): G1: -33 (-42.3, 21.3) G2: -25 (-38.8, -13.0) G1/G2: <i>P</i> < 0.001</p> <p>Voided volume (mL), median % change (IQR): G1: 31.5 (13.3, 56.2) G2: 20 (3.1, 45.4) G1/G2: <i>P</i> < 0.001</p> <p>Symptoms are "minor or less", %: G1: 66.5 G2: 61.5</p> <p>Overall improvement in symptoms, %: G1: 76 G2: 71</p> <p>Worsening of symptoms, %: G1: 3 G2: 5</p> <p>Dry mouth, n (%): G1: 76 (31) G2: 90 (35)</p> <p>Headache, n (%): G1: 15 (6) G2: 21 (8)</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: -</p> <p>Drop-out rates: -</p> <p>Power calculation: +</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: ++</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Mattiasson et al., 2003 (continued)				<p>Constipation, n (%): G1: 7 (3) G2: 14 (5)</p> <p>≥ 1 SE, n (%): G1: 158 (65) G2: 177 (69) G1/G2: <i>P</i> = NS</p> <p>Withdrawal due to, %: AE: 15</p> <p>Withdrawal due to lack of efficacy: 3</p> <p>Withdrawal of consent : 2</p> <p>Protocol violations: 1</p> <p>Completed treatment: G1: 77% G2: 79%</p>	

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Millard, 2004</p> <p>Country and setting: International, Multicenter study, 54 sites</p> <p>Enrollment period: NR</p> <p>Funding: Pharmacia Corp</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: Tolterodine ± simplified pelvic-floor exercise regimen for 24 weeks</p> <p>Groups: G1: tolterodine 2 mg b.i.d. and simple PFME program G2: tolterodine 2 mg b.i.d. daily</p> <p>N at enrollment: G1: 227 G2: 253</p> <p>N at follow-up, 12 weeks, n (%): G1: 181 (79.7) G2: 205 (81.0)</p> <p>N at follow-up, 24 weeks, n (%): G1: 164 (72.2) G2: 190 (75.1)</p> <p>Women, n (%): G1: 169 (75.4%) G2: 190 (75.4%)</p> <p>Age, mean ± SD: G1: 53.6 ± 16.9 G2: 53.2 ± 17.4</p> <p>Race/ethnicity: Asian: G1: 176 (78.6%) G2: 203 (80.6%) White/mixed: G1: 48 (21.4%) G2: 49 (19.4%)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age 18-90 • ≥ 8 voids/day • Urgency • ≥ 1 UI episode/d • Sx for ≥ 6 months <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Symptomatic SI • Significant PVR volume • Neuropathy • Glaucoma • UTI • + urine cytology • Use of concomitant anticholinergic therapy w/in 14 days of randomization 	<p>Urgency episodes/day, mean ± SD: G1: 4.2 ± 3.6 G2: 4.1 ± 4.0</p> <p>Urgency episodes/day, median (IQR): G1: 3.6 (1.3, 6.0) G2: 3.0 (1.3, 6.0)</p> <p>Incontinence episodes/day, mean ± SD: G1: 3.44 ± (3.4) G2: 3.21 ± (3.4)</p> <p>Incontinence episodes/day, median (IQR): G1: 2.3 (1.3, 4.0) G2: 2.9 (1.3, 3.7)</p> <p>Voids/day, mean ± SD: G1: 11.87 ± 4.3 G2: 12.78 ± 5.6</p> <p>Voids/day, median (IQR): G1: 10.7 (9.0, 13.7) G2: 11.3 (9.0, 15.0)</p> <p>Voided volume (mL), mean ± SD: G1: 146.1 ± 67.7 G2: 146.0 ± 83.3</p> <p>Voided volume (mL), median (IQR): G1: 137 (98, 186) G2: 132 (99, 189)</p> <p>Duration of symptoms, n (%): ≤ 5 years: G1: 166 (74.1) G2: 173 (68.7) > 5 years: G1: 56 (25) G2: 78 (31)</p>	<p>Urgency episodes/day, wk 12, mean change ± SD (% change): G1: -1.9 ± 4.0 (-64.5) G2: -2.2 ± 3.6 (-69.8) G1/BL: P = 0.001 G1/BL: P = 0.001 G1/G2: P = NS</p> <p>Urgency episodes/day, wk 12, median change: G1: -1.6 G2: -1.3 G1/G2: P = 0.7658</p> <p>Urgency episodes/day, wk 24, mean change ± SD (% change): G1: -2.2 ± 3.6 (-78.7) G2: -2.7 ± 3.5 (-83) G1/BL: P = 0.001 G1/BL: P = 0.001 G1/G2: P = NS</p> <p>Urgency episodes/day, wk 24, median change: G1: -1.9 G2: -2.0 G1/G2: P = 0.3029</p> <p>Incontinence episodes/day, wk 12, mean change ± SD: G1: -2.15 ± 3.0 G2: -2.15 ± 2.7 G1/BL: P = 0.001 G1/BL: P = 0.001 G1/G2: P = 0.2215</p>	<p>Quality: Overall quality score: fair INTERNAL VALIDITY: poor Randomization: - Masking: - Pt selection criteria: + Loss to followup: ++ Drop-out rates: - Power calculation: + Statistical issues: - EXTERNAL VALIDITY: good Age: + Baseline OAB status: + Baseline characteristics: ++ Length of followup: ++ Measurement methods: + Measurement reliability: + Intervention description: +</p>

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Millard, 2004 (continued)			<p>Previous UT surgery, n (%): G1: 44 (19.6) G2: 43 (17.1)</p> <p>Prev anticholinergics, n (%): G1: 19 (8.5) G2: 14 (5.6)</p> <p>Prev UUI drug, n (%): G1: 101 (45.1) G2: 91 (36.1)</p> <p>Patient subjective rating of UI symptoms as “severe” or “many severe”, %: G1: 54.0 G2: 56.8</p>	<p>Incontinence episodes/day, wk 12, median change: G1: -1.6 G2: -1.6 G1/G2: P = 0.8251</p> <p>Incontinence episodes/day, wk 24, mean change ± SD (% change): G1: -2.23 ± 3.0 (-64) G2: -2.26 ± 3.0 (-70) G1/BL: P = 0.001 G1/BL: P = 0.001 G1/G2: P = NS</p> <p>Incontinence episodes/day, wk 24, median change: G1: -1.6 G2: -1.6 G1/G2: P = 0.8341</p> <p>Voids/day, wk 12, mean change ± SD (% change): G1: -2.68 ± 3.8 (-22) G2: -3.42 ± 4.6 (-26) G1/BL: P = 0.001 G1/BL: P = 0.001 G1/G2: P = 0.9478</p> <p>Voids/day, wk 24, mean change ± SD (% change): G1: -2.58 ± 5.0 (-22) G2: -3.58 ± 5.2 (-26) G1/BL: P = 0.001 G1/BL: P = 0.001 G1/G2: P = 0.3549</p>	

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Millard, 2004 (continued)				<p>Voided volume (mL), wk 12, median change (% change): G1: 20.4 (17.2) G2: 17.5 (15.8)</p> <p>Voided volume (mL), wk 12, median change (% change): G1: 21.1 (18.1) G2: 19.1 (15.4)</p> <p>Patient subjective report of improvement in bladder condition, wk 12, %: G1: 82.6 G2: 83.9</p> <p>Patient subjective report of improvement in bladder condition, wk 24, %: G1: 81.7 G2: 85.9</p> <p>Adverse events, n (%): G1: 22 (9.7) G2: 23 (9.1)</p> <p>Mild dry mouth, %: G1: 18.1 G2: 21.3</p> <p>Moderate dry mouth, %: G1: 7.5 G2: 5.1</p> <p>Severe dry mouth, %: G1: 4.0 G2: 3.2</p> <p>Headache, %: 6</p> <p>Constipation, %: 4.8</p> <p>Nausea, %: 2.7</p>	

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Millard, 2004 (continued)				Dry eyes, %: 2.5 Dizziness, %: 2.4	

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Sand et al., 2004</p> <p>Country and setting: US, Specialty clinic</p> <p>Enrollment period: Subanalysis of OBJECT trial, women only</p> <p>Funding: ALZA Corp</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: 10mg ER Oxybutynin daily vs. 4mg Tolterodine (2mg bid)</p> <p>Groups: G1: 10mg ER Oxybutynin daily x 12 wks G2: 4mg Tolterodine (2mg bid) x 12 wks</p> <p>Stratified by age: a: Age ≤ 64 b: Age 65-74 c: Age ≥ 75</p> <p>N at enrollment: G1: 152 G2: 163</p> <p>N at follow-up: G1: 132 G2: 146</p> <p>Women, %: 100</p> <p>Age, yrs ± SD: G1: 58.4 G2: 58.8</p> <p>Race/ethnicity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • ≥ 7 and ≤ 50 UUI episodes/week • ≥ 10 voids/day • MUI included if predominant UUI <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • UTI • IC • Urethral diverticulum • Bladder tumor • Bladder stone • Delivery within 6 mos • Pelvic, bladder, vaginal surgery in ≤ 6 mos • PVR ≥ 150 mL • Cardiovascular, renal, pulmonary, gastrointestinal, endocrine, neurologic, autoimmune, hematological, urological, psychiatric, or hepatic disease • Hematuria • Positive urine culture • Narrow angle glaucoma • Obstructive uropathy • Myasthenia gravis • POP to hymenal ring • GI obstruction • Decreased GI motility • GI narrowing • GI retention • Investigational drugs within 1 month of screening • Hypersensitivity to drugs 	<p>UUI episodes/week, mean: G1: 25.2 G2: 25.1</p> <p>Incontinence episodes/week, mean: G1: 28.1 G2: 28.9</p> <p>voids/week, mean: G1: 91.7 G2: 91.6</p> <p>Naïve to anti-cholinergics, %: G1: 60.5 G2: 60.7</p>	<p>UUI episodes/week, %: G1: 6.2 G2: 8.5 G1a: 5.0 G2a: 8.4 G1b: 5.5 G2b: 7.5 G1c: 8.5 G2c: 11.1 G1/G2: P = 0.038 G1a/G2a: P = 0.005 G1b/G2b: P = 0.337 G1c/G2c: P = 0.568</p> <p>Incontinence episodes/week, mean: G1: 7.3 G2: 10.1 G1a: 5.8 G2a: 10.0 G1b: 6.1 G2b: 9.2 G1c: 10.5 G2c: 12.5 G1/G2: P = 0.030 G1a/G2a: P = 0.005 G1b/G2b: P = 0.164 G1c/G2c: P = 0.714</p> <p>voids/week, mean: G1: 68.0 G2: 71.2 G1a: 63.7 G2a: 71.2 G1b: 73.8 G2b: 71.9 G1c: 66.8 G2c: 65.5 G1/G2: P = 0.272 G1a/G2a: P = 0.024 G1b/G2b: P = 0.706 G1c/G2c: P = 0.838</p>	<p>Quality: Overall quality score: fair INTERNAL VALIDITY: fair Randomization: + Masking: + Pt selection criteria: + Loss to followup: + Drop-out rates: + Power calculation: - Statistical issues: + EXTERNAL VALIDITY: good Age: + Baseline OAB status: + Baseline characteristics: ++ Length of followup: + Measurement methods: + Measurement reliability: + Intervention description: +</p>

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Sand et al., 2004 (continued)		<ul style="list-style-type: none"> • Current drug/ EtOH abuse • Pregnant • Breastfeeding • Inability to follow protocol 		<p>Dry mouth, %: G1: 28.3 G2: 33.7</p> <p>Constipation, %: G1: 8.6 G2: 6.7</p> <p>Retention, %: G1: 4.0 G2: 1.2</p> <p>Blurred vision, %: G1: 2.6 G2: 0.6</p> <p>Dizziness, %: G1: 3.9 G2: 4.3</p> <p>Insomnia, %: G1: 0.7 G2: 1.8</p> <p>Somnolence, %: G1: 3.3 G2: 1.8</p> <p>Nervousness, %: G1: 0 G2: 1.2</p> <p>Headache, %: G1: 9.2 G2: 10.4</p> <p>Dyspepsia, %: G1: 5.3 G2: 6.1</p> <p>Nausea, %: G1: 3.3 G2: 1.8</p> <p>Vomiting, %: G1: 2.0 G2: 1.8</p>	

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Schmidt et al., 1999</p> <p>Country and setting: 16 centers in US, Canada, and Europe</p> <p>Enrollment period: 1993- April 1997</p> <p>Funding: Medtronic Inc.</p> <p>Author conflict of interest: Schmidt, Jonas, Oleson, Janknegt, Siegel: financial interest and/or other relationship with Medtronic Inc.</p>	<p>Design: randomized controlled trial</p> <p>Intervention: implantable Interstim system stimulating the nerve ramus in preoperatively tested sacral nerve foramen</p> <p>Groups:</p> <p>G1: implantation with immediate stimulation (n=34 for efficacy analysis)</p> <p>G2: control group receiving standard medical therapy for 6 months; allowed to cross over at 6 months if remained eligible for implantation (n=42 for efficacy analysis)</p> <p>N at enrollment: 155</p> <p>N randomized: 98 (the remaining 57 did not respond to test stimulation so were not randomized)</p> <p>N at follow-up: 76 at 6 months</p> <p>Age: mean 46.6 ±13.0 (range 20.2 – 78.9)</p> <p>Gender: 125 F (80.6%), 30 M (19.4%)</p> <p>Race/ethnicity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • age > 16 years • refractory to standard medical therapy • ≥ 100 mL bladder capacity with normal upper urinary tract • good surgical candidate • able to complete study documentation and return for follow-up evaluation <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Neurological conditions • Stress urinary incontinence • Primary pelvic pain 	<p>Duration of urinary symptoms before enrollment: mean 9.0±7.4 (range 0.6-35.4)</p> <p>Mean number of daily incontinence episodes: 8.9± 5.9</p> <p>G1: 9.7 ± 6.3</p> <p>G2: 9.3 ± 4.8</p> <p>Mean leak severity rating: 1.9± 0.6</p> <p>G1: 2.0 ± 0.7</p> <p>G2: 1.8 ± 0.6</p> <p>Mean absorbent pads/diapers replaced daily 4.8± 4.8</p> <p>G1: 6.2 ± 5.0</p> <p>G2: 5.0 ± 3.7</p> <p>Mean heavy incontinence episodes per day:</p> <p>G1: 3.4 ±3.8</p> <p>G2: 2.6 ±3.5</p> <p>3-day diary data: 90.3% of patients had at least 1 moderate/heavy and 74.8% had at least 1 heavy incontinence episode</p> <p>Previous treatment for urinary problems: 153 (89.7%), including pharmacological in 144 (92.5%), nonsurgical in 55 (35.5%), and surgical in 88 (56.8%)</p>	<p>At 6 months:</p> <p>Mean number of daily incontinence episodes:</p> <p>G1: 2.6± 5.1 (p<0.0001)</p> <p>G2: 11.3 ± 5.9 (p=0.002)</p> <p>Mean leak severity rating:</p> <p>G1: 0.8 ± 0.9 (p<0.0001)</p> <p>G2: 2.0± 0.6 (p=0.006)</p> <p>Mean absorbent pads/diapers replaced daily</p> <p>G1: 1.1 ± 2.0 (p<0.0001)</p> <p>G2: 6.3 ± 3.6 (p=0.003)</p> <p>Mean heavy incontinence episodes per day:</p> <p>G1: 0.3 ±0.9 (P<0.0001)</p> <p>G2: 3.9 ±3.8</p> <p>G1: 47% completely dry at 6 months; 3 patients had no reduction in incontinence after 6 months (one underwent device explantation and 2 had increased frequency of urination and incontinence)</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: -</p> <p>Masking: -</p> <p>Pt selection criteria: +</p> <p>Loss to followup: -</p> <p>Drop-out rates: NR</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: ++</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Schmidt et al., 1999 (continued)				<p>Safety data (pooled for 157 patients): adverse events requiring surgical repositioning or replacement of implant devices document in 51 (32.5%)</p> <p>168 post-implant events reported by 83 patients, including pain at the neurostimulator site in 15.9%, pain at implant site in 19.1, and lead migration in 7.0%</p> <p>Infection or skin irritation led to device explantation in 2 patients and temporary explant in 2 patients</p> <p>No permanent injuries or nerve damage reported.</p>	

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Song, et al., 2006</p> <p>Country and setting: Korea, Medical Center</p> <p>Enrollment period: May 2001 to April 2002</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: bladder training (BT) vs. Tolterodine vs. BT + Tolterodine</p> <p>Groups: G1: BT x 12 wks G2: Tolterodine 2 mg bid x 12 wks G3: Tolterodine 2 mg bid + BT x 12 wks</p> <p>N at enrollment: G1: 46 G2: 47 G3: 46</p> <p>N at follow-up: G1: 26 G2: 32 G3: 31</p> <p>Women, %: 100</p> <p>Age, mean ± SD: G1: 45.73 ± 12.68 G2: 48.41 ± 9.38 G3: 45.42 ± 9.54</p> <p>Race/ethnicity, %: Korean: 100</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 18 • ≥ 8 voids/day • Urge with or without incontinence • Symptom duration ≥ 3 months • No prior history of treatment for OAB <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Active urinary tract infection • Clinically significant SUI • Bladder outlet obstruction • Interstitial cystitis • Glaucoma • Megacolon • Maximal urine flow rate of < 10 mL/sec • Postvoid residual volume > 30% of the total amount voided on uroflowmetry 	<p>voids/day, mean ± SD: G1: 10.93 ± 2.14 G2: 11.63 ± 2.57 G3: 11.90 ± 1.51</p> <p>Nocturia episodes/day, mean ± SD: G1: 1.45 ± 1.14 G2: 1.72 ± 1.04 G3: 1.96 ± 1.49</p> <p>Urgency, mean score ± SD: G1: 2.58 ± 1.30 G2: 2.81 ± 0.74 G3: 3.00 ± 1.10</p> <p>Maximum flow rate (mL/s), mean ± SD: G1: 20.35 ± 8.44 G2: 22.56 ± 4.94 G3: 21.19 ± 4.96</p> <p>Residual urine (mL), mean ± SD: G1: 9.08 ± 22.56 G2: 7.59 ± 12.39 G3: 6.42 ± 10.16</p> <p>Symptom duration (years), mean ± SD: G1: 6.44 ± 6.84 G2: 4.54 ± 5.15 G3: 4.10 ± 3.99</p>	<p>voids/day, mean (% decrease) G1: 8.1 (25.9%)* G2: 8.1 (30.2%)* G3: 7.9 (33.5%)* G3/G1: P < 0.05</p> <p>Nocturia episodes/day, mean (% reduction): G1: 0.6 (56.1%)* G2: 0.6 (65.4%)* G3: 0.6 (66.3%)*</p> <p>Urgency, mean score (% reduction): G1: 1.4 (44.8%)* G2: 1.1 (62.2%)* G3: 1.2 (60.2%)* G3/G1: P = 0.021 G2/G1: P = 0.017 G2/G3: P = NS</p> <p>Satisfaction, mean score (% improved): G1: 1.5 (53.9) G2: 1.4 (63.0) G3: 1.3 (71.0)</p> <p>Dry mouth, n (%): G1: 0 (0.0) G2: 7 (21.9) G3: 9 (28.9)</p> <p>Hesitancy, n (%): G1: 0 (0.0) G2: 3 (9.4) G3: 2 (6.5)</p> <p>Decreased appetite/constipation, n (%): G1: 0 (0.0) G2: 2 (6.3) G3: 2 (6.5)</p> <p>Headache, n (%): G1: 0 (0.0) G2: 1 (3.1) G3: 0 (0.0)</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: -</p> <p>Masking: -</p> <p>Pt selection criteria: +</p> <p>Loss to followup: -</p> <p>Drop-out rates: +</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Swift et al., 2003</p> <p>Country and setting: Europe (167 centers), North America (74 centers), Australia and New Zealand (4 centers), University</p> <p>Enrollment period: February 1999 to October 1999</p> <p>Funding: Pharmacia Corp</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT double blind placebo-controlled double dummy, random permuted blocks of 6</p> <p>Intervention: Tolterodine ER vs. Tolterodine IR</p> <p>Groups: G1: Tolterodine ER 4mg daily G2: Tolterodine IR 2 mg BID G3: placebo</p> <p>N at enrollment: G1: 417 G2: 408 G3: 410</p> <p>N at follow-up: Total: 1092</p> <p>Women, %: 100</p> <p>Age, yrs ± SD: G1: 59 ± 14 G2: 59 ± 14 G3: 60 ± 14</p> <p>Race/ethnicity, n (%): White: G1: 396 (95) G2: 389 (95) G3: 383 (93) Black: G1: 15 (4) G2: 12 (3) G3: 20 (5) Asian/Pacific: G1: 5 (1) G2: 4 (1) G3: 2 (1) Mixed: G1: 0 G2: 3 (1) G3: 5 (1) Unknown: G1: 1 (<1) G2: 0 G3: 0</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Women • Age ≥ 18 • ≥ 8 voids/day • > 5 UUI/ week • Symptoms x ≥ 6 months (per voiding diary) <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • SUJ • Total daily urine volume > 3 liters • Hepatic/ renal disease • UTI • IC • Hematuria • BOO • Current e-stim • Current bladder training • Indwelling catheter • Intermittent catheterization • Pregnant/ nursing • Childbearing age without BC • Anticholinergic meds • Meds inhibit cytochrome P450 	<p>Incontinence episodes/week, mean ± SD: G1: 22.1 ± 22.5 G2: 22.9 ± 21.9 G3: 23.9 ± 21.2</p> <p>Pads/day, mean ± SD: G1: 1.6 ± 2.1 G2: 1.5 ± 2.0 G3: 1.7 ± 2.4</p> <p>voids/day, mean ± SD: G1: 10.8 ± 4.2 G2: 11.1 ± 3.7 G3: 11.2 ± 3.9</p> <p>Voided volume (mL), mean ± SD: G1: 141.2 ± 43.1 G2: 137.2 ± 41.2 G3: 135.7 ± 43.2</p> <p>Previous drug therapy: n (%): G1: 235 (56) G2: 222 (54) G3: 225 (55)</p>	<p>Incontinence episodes/week, mean ± SD: G1: 10.3 ± 17.2 G2: 12.8 ± 19.8 G3: 16.7 ± 19.7 G1/G3: P = 0.001 G2/G3: P = 0.001</p> <p>Pads/day, mean ± SD: G1: 1.0 ± 1.8 G2: 1.0 ± 1.5 G3: 1.5 ± 2.2 G1/G3: P = 0.001 G2/G3: P = 0.001</p> <p>voids/day, mean ± SD: G1: 9.0 ± 3.2 G2: 9.3 ± 4.0 G3: 9.9 ± 3.8 G1/G3: P = 0.001 G2/G3: P = 0.005</p> <p>Voided volume (mL), mean ± SD: G1: 179.1 ± 66.6 G2: 169.7 ± 65.6 G3: 149.0 ± 56.3 G1/G3: P = 0.001 G2/G3: P = 0.001</p> <p>Clinical effectiveness*, dry mouth: G1: 0.53 G2: 0.39 G3: 0.30</p> <p>Dry mouth, n (%): G1: 105 (25.3) G2: 127 (31.2) G3: 33 (8.0) G1/G3: P < 0.01 G2/G3: P < 0.01</p> <p>Abdominal pain, n (%): G1: 18 (4.3) G2: 12 (2.9) G3: 7 (1.7) G1/G3: P = 0.03</p> <p>Flatulence, n (%): G1: 8 (1.9) G2: 11 (2.7) G3: 6 (1.5)</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: fair</p> <p>Randomization: +</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: ++</p> <p>Drop-out rates: ++</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Swift et al., 2003 (continued)	BMI, kg/m² ± SD: G1: 28.8 ± 13.8 G2: 29.0 ± 11.0 G3: 28.8 ± 6.7			Constipation, n (%): G1: 27 (6.5) G2: 27 (6.6) G3: 14 (3.4) Dyspepsia, n (%): G1: 11 (2.7) G2: 14 (3.4) G3: 6 (1.5) Nausea, n (%): G1: 7 (1.7) G2: 9 (2.2) G3: 9 (2.2) Diarrhea, n (%): G1: 10 (2.4) G2: 14 (3.4) G3: 9 (2.2) Xerophthalmia, n (%): G1: 16 (3.9) G2: 8 (2.0) G3: 8 (2.0) Abnormal vision, n (%): G1: 5 (1.2) G2: 4 (1.0) G3: 2 (0.5) Headache, n (%): G1: 29 (7.0) G2: 14 (3.4) G3: 19 (4.6) UTI, n (%): G1: 15 (3.6) G2: 11 (2.7) G3: 19 (4.6) Insomnia, n (%): G1: 7 (1.7) G2: 2 (0.5) G3: 9 (2.2) Somnolence, n (%): G1: 12 (2.9) G2: 11 (2.7) G3: 8 (2.0) Dizziness, n (%): G1: 7 (1.7) G2: 7 (1.7) G3: 4 (1.0)	

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Swift et al., 2003 (continued)				<p>Hypertension, n (%): G1: 6 (1.4) G2: 4 (1.0) G3: 4 (1.0)</p> <p>Sinusitis, n (%): G1: 8 (1.9) G2: 2 (0.5) G3: 3 (0.7)</p> <p>Arthritis, n (%): G1: 1 (0.2) G2: 5 (1.2) G3: 1 (0.2)</p> <p>Dry skin, n (%): G1: 2 (0.5) G2: 5 (1.2) G3: 1 (0.2)</p>	

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Szonyi et al., 1995</p> <p>Country and setting: UK</p> <p>Enrollment period: NR</p> <p>Funding: Smith & Nephew Pharmaceuticals Ltd</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: Oxybutynin plus bladder training vs. placebo plus bladder training</p> <p>Groups: G1: Oxybutynin 2.5 mg bid with dose titration on days 29 and 43 plus bladder training G2: placebo + bladder training</p> <p>N at enrollment: G1: 30 G2: 30</p> <p>N at follow-up: G1: 16 G2: 23</p> <p>Women, n (%): 56 (93)</p> <p>Age, mean ± SD: 82.2 ± 6.06</p> <p>Race/ethnicity: NR</p> <p>Weight (kg), mean ± SD: 67.4 ± 14.92</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age > 70 • Frequency, urgency and UUI • Mobile • Able to keep diary <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • UTI • Hepatic or renal disease • Glaucoma • Uncontrolled diabetes • Taking imipramine or propantheline 		<p>Voids/2 weeks, median change (95% CI): G1/G2: 577 (-27.0, 6.0) P = 0.0025</p> <p>Nocturia episodes/2 weeks, median change (95% CI): G1/G2: -6 (-5, 7.0)</p> <p>Daytime incontinence episodes/2 weeks, median change (95% CI): G1 vs. G2: -9.5 (-11.0, 3.0)</p> <p>Nocturia episodes/2 weeks, median change (95% CI): G1/G2: -1.0 (-3.0, 2.0)</p> <p>Patient assessment of benefit, %:</p> <p>29 days: G1: 86 G2: 55 P = 0.02</p> <p>43 days: G1: 71 G2: 59 P = 0.41</p> <p>57 days: G1: 79 G2: 55 P = 0.09</p> <p>Patient response, 29 days, n: Cure: G1: 1 G2: 0 Significant improvement: G1: 15 G2: 8 Marginal improvement: G1: 7 G2: 8</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: -</p> <p>Drop-out rates: -</p> <p>Power calculation: +</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: +</p> <p>Length of followup: -</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Szonyi et al., 1995 (continued)				<p>No change: G1: 5 G2: 13</p> <p>Patient response, 57 days, n: Cure: G1: 4 G2: 3 Significant improvement: G1: 14 G2: 8 Marginal improvement: G1: 3 G2: 4 No change: G1: 7 G2: 14</p> <p>Dry mouth, %: G1: 93 G2: 86</p> <p>Blurred vision, %: G1: 50 G2: 59</p> <p>Heartburn, %: G1: 57 G2: 45</p> <p>Constipation, %: G1: 50 G2: 45</p> <p>Dry skin, %: G1: 50 G2: 59</p> <p>Poor compliance (< 75% of tablets), %: G1: 20 G2: 20</p>	

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Tseng et al., 2009</p> <p>Country and setting: Taiwan, University</p> <p>Enrollment period: January 2005 to November 2005</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Prospective cohort randomized</p> <p>Intervention: Tolterodine 2 mg b.i.d. vs. tolterodine 2 mg b.i.d. + conjugated equine estrogen 0.625 mg twice per week</p> <p>Groups: G1: tolterodine 2 mg b.i.d. G2: tolterodine 2 mg b.i.d. + conjugated equine estrogen 0.625 mg twice per week</p> <p>N at enrollment: G1: 40 G2: 40</p> <p>N at follow-up: G1: 40 G2: 40</p> <p>Age, mean ± SD: G1: 64.5 ± 7.4 G2: 66.2 ± 6.8</p> <p>Race/ethnicity: NR</p> <p>Women, N (%): G1: 40 (100) G2: 40 (100)</p> <p>BMI, kg/m² ± SD: G1: 24.5 ± 3.9 G2: 25.3 ± 3.8</p> <p>Previous antimuscarinic Rx, n (%): G1: 5 (12.5) G2: 6 (15)</p> <p>Prior hysterectomy, n (%): G1: 12 (30) G2: 14 (35)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 18 • OAB symptoms • Amenorrheic <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Advanced POP > Stage 2 • Women with storage and voiding dysfunction • undiagnosed • Severe constipation • Elevated PVR • Neurological deficit • Renal/ hepatic disease • Narrow angle glaucoma • Urinary retention • Gastric retention • Hypersensitivity to drugs • BOO • Cardiac conduction disorders • Myasthenia gravis • History of CVA • History of VTE • Gallbladder disease • Known or suspected breast carcinoma • Undiagnosed genital bleeding • HRT within 3 months of study enrollment 	<p>UII episodes/ day, mean ± SD: G1: 1.8 ± 0.7 G2: 2.1 ± 1.1</p> <p>Urgency episodes/ day, mean ± SD: G1: 4.5 ± 0.8 G2: 4.3 ± 0.7</p> <p>Nocturia episodes/ day, mean ± SD: G1: 3.5 ± 0.8 G2: 3.3 ± 0.8</p> <p>Voids/day, mean ± SD: G1: 14.1 ± 1.3 G2: 14.8 ± 1.5</p> <p>UDI-6 score, mean ± SD: G1: 9.5 ± 3.9 G2: 8.6 ± 3.8</p> <p>IIQ-7 score, mean ± SD: G1: 10.2 ± 4.5 G2: 9.4 ± 3.6</p> <p>Voided volume (mL), mean ± SD: G1: 108.5 ± 14.0 G2: 115.8 ± 15.1</p>	<p>UII episodes/day, mean ± SD: G1: 1.5 ± 0.5 G2: 1.5 ± 0.5 P = NS</p> <p>Urgency episodes/ day, mean ± SD: G1: 3.5 ± 0.5 G2: 3.3 ± 0.6</p> <p>Nocturia episodes/ day, mean ± SD: G1: 2.9 ± 0.6 G2: 2.6 ± 0.7</p> <p>Voids/day, mean ± SD: G1: 6.4 ± 1.9 G2: 5.8 ± 0.9 P = 0.001</p> <p>UDI-6 score, mean ± SD: G1: 7.2 ± 2.9 G2: 6.9 ± 2.7 P < 0.001</p> <p>IIQ-7 score, mean ± SD: G1: 6.5 ± 2.7 G2: 6.1 ± 2.5 P < 0.001</p> <p>Voided volume (mL), mean ± SD: G1: 134.5 ± 15.8 G2: 141.9 ± 16.1 P = 0.007</p> <p>Adverse events: None</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: fair</p> <p>Randomization: +</p> <p>Method and blinding: -</p> <p>Pt selection criteria: +</p> <p>Loss to followup: ++</p> <p>Drop-out rates: ++</p> <p>Power calculation: +</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Van Kerrebroeck et al., 2001</p> <p>Freeman et al., 2003*</p> <p>Country and setting: North America (74 centers), Australasia (4 centers), Europe (89 centers)</p> <p>Enrollment period: NR</p> <p>Funding: Pharmacia Corporation</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: Tolterodine ER vs. Tolterodine IR vs. placebo</p> <p>Groups: G1: tolterodine ER 4 mg t.i.d. G2: tolterodine IR 2 mg b.i.d. G3: placebo</p> <p>N at enrollment: G1: 507 G2: 514 G3: 508</p> <p>N at follow-up: Total: 1442 G1: 398 G3: 374</p> <p>Women, n (%): G1: 417 (82) G2: 408 (79) G3: 410 (81)</p> <p>Age, mean (range): G1: 60 (20, 89) G2: 60 (22, 92) G3: 61 (22, 93)</p> <p>Race/ethnicity, %: White: G1: 95.7 G3: 94.7 Black: G1: 3.0 G3: 3.5 Asian/Pacific Islander: G1: 1.0 G3: 0.8 Other: G1: 0.3 G3: 1.1</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 18 • Urinary frequency (≥ 8 voids/day) <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • SUI • Total daily urine volume > 3 L • Contra-indications to antimuscarinic treatment • Hepatic or renal disease • UTIs • Interstitial cystitis • Hematuria • BOO • Current electrostimulation or bladder training therapy • Indwelling catheter or intermittent self-catheterization • Pregnant or nursing • Women not using reliable contraception • Being treated for OAB with other anticholinergic drugs or drugs that inhibit cytochrome P450 3A4 isoenzymes • Estrogen therapy < 2 months • Treatment w/ investigational drug < 2 months 	<p>Incontinence episodes/week, mean (range): G1: 22.1 (0, 168.0) G2: 23.2 (0, 168.0) G3: 23.3 (0, 168.0)</p> <p>≥ 5 incontinence episodes/week, n (%): G1: 492 (97) G2: 498 (97) G3: 494 (97)</p> <p>Pads/day, mean (range): G1: 1.4 (0-18) G2: 1.4 (0-25) G3: 1.5 (0-22)</p> <p>Voids/day, mean (range): G1: 10.9 (2.3, 51.3) G2: 11.1 (2.0, 48.6) G3: 11.3 (2.0, 37.4)</p> <p>≥ 8 voids/day, n (%): G1: 458 (90) G2: 469 (91) G3: 467 (92)</p> <p>Previous drug therapy, n (%): G1: 270 (53) G2: 276 (54) G3: 263 (52)</p> <p>Poor efficacy, %: G1: 43 G2: 38.4 G3: 40.7</p> <p>Able to finish tasks before visiting a toilet, %: G1: 5 G3: 6</p>	<p>Urinary Urgency, subjective assessment, 12 wks, n (%):* Improvement: G1: 173 (44) G3: 118 (32) G1/G3: P < 0.001 No change: G1: 201 (51) G3: 212 (57) Deterioration: G1: 22 (6) G3: 44 (12) G1/G3: P < 0.002</p> <p>Urinary urgency, improvement, 12 wks, women only, %: G1: 46.6 G3: 26.6 G1/G3: P = 0.001 OR 1.81 (95% CI: 1.31, 2.49)</p> <p>Not able to hold urine, 12 wks, %: G1: 58 G3: 32 G1/G3: P < 0.001</p> <p>Incontinence episodes/week, mean change ± SD (%) G1: -11.8 ± 17.8 G2: -10.6 ± 16.9 G3: -6.9 ± 15.4 G1/G3: P = 0.00001 G2/G3: P = 0.0005 G1/G2: P < 0.05</p> <p>Incontinence episodes/week, median % change: G1: -71 G2: -60 G3: -30 G1/G2: P < 0.05</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: + Masking: + Pt selection criteria: + Loss to followup: + Drop-out rates: - Power calculation: - Statistical issues: + EXTERNAL VALIDITY: good</p> <p>Age: + Baseline OAB status: + Baseline characteristics: ++ Length of followup: + Measurement methods: + Measurement reliability: + Intervention description: +</p>

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Van Kerrebroeck et al., 2001 Freeman et al., 2003* (continued)			Voided volume (mL), mean (range): G1: 141 (36, 338) G2: 137 (38, 283) G3: 136 (31, 374)	Pads/day, mean ± SD: G1: -0.5 ± 1.4 G2: -0.5 ± 18 G3: -0.2 ± 1.4 G1/G3: P = 0.0145 G2/G3: P = 0.0035 Voids/day, mean ± SD: G1: -3.5 ± 4.9 G2: -3.3 ± 4.4 G3: -2.2 ± 4.0 G1/G3: P = 0.00001 G2/G3: P = 0.0002 Voluntary voids/day, mean ± SD: G1: -1.8 ± 3.4 G2: -1.7 ± 3.3 G3: -1.2 ± 2.9 G1 vs. G3 G1/G3: P = 0.00047 G2/G3: P = 0.0079 Bladder symptoms, improvement, 12 wks, women only, %:* G1: 62.8 G3: 48.4 G1/G3: P = 0.001 OR 1.78 (95% CI: 0.34, 2.37) Treatment benefit, 12 wks, n (%):* Much benefit: G1: 172 (43.2) G3: 88 (23.5) G1/G3: P < 0.001 Little benefit G1: 138 (34.7) G3: 118 (31.6) No benefit G1: 88 (22.1) G3: 168 (44.9)	

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Van Kerrebroeck et al., 2001 Freeman et al., 2003* (continued)				<p>Able to finish tasks before visiting a toilet, 12 wks, %: G1: 33 G3: 18 G1/G3: P < 0.001</p> <p>Voided volume (mL), mean change ± SD: G1: +34 ± 51 G2: +29 ± 47 G3: +14 ± 41 G1/G3: P = 0.00001 G2/G3: P = 0.0001</p> <p>Discontinued due to AEs, n (%): G1: 27 (5) G2: 28 (5) G3: 33 (6)</p> <p>Reported serious adverse events, n: G1: 7 G2: 12 G3: 18</p> <p>Parasympathetic Dry mouth, n (%): G1: 118 (23) G2: 156 (30) G3: 39 (8)</p> <p>Xerophthalmia, n (%): G1: 17 (3) G2: 12 (2) G3: 10 (2)</p> <p>Abnormal vision, n (%): G1: 6 (1) G2: 4 (1) G3: 2 (0.5)</p> <p>Dry skin, n (%): G1: 2 (0.5) G2: 6 (1) G3: 1 (0.5)</p>	

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Van Kerrebroeck et al., 2001 Freeman et al., 2003* (continued)				Gastrointestinal Constipation, n (%): G1: 30 (6) G2: 35 (7) G3: 22 (4) Dyspepsia, n (%): G1: 15 (3) G2: 16 (3) G3: 7 (1) Abdominal pain, n (%): G1: 19 (4) G2: 13 (3) G3: 8 (2) Diarrhea, n (%): G1: 10 (2) G2: 16 (3) G3: 11 (2) Flatulence, n (%): G1: 10 (2) G2: 14 (3) G3: 9 (2) Nausea, n (%): G1: 7 (1) G2: 10 (2) G3: 10 (2) Headache, n (%): G1: 32 (6) G2: 19 (4) G3: 23 (5) Somnolence, n (%): G1: 14 (3) G2: 13 (3) G3: 9 (2) Dizziness, n (%): G1: 11 (2) G2: 9 (2) G3: 5 (1) Fatigue, n (%): G1: 11 (2) G2: 6 (1) G3: 4 (1) Insomnia, n (%): G1: 7 (1) G2: 2 (0.5) G3: 9 (2)	

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Van Kerrebroeck et al., 2001				Urinary tract infection, n (%): G1: 16 (3) G2: 13 (3) G3: 20 (4)	
Freeman et al., 2003* (continued)				Dysuria, n (%): G1: 5 (1) G2: 8 (2) G3: 1 (0.5)	
				Peripheral edema, n (%): G1: 7 (1) G2: 7 (1) G3: 4 (1)	

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Wang et al., 2006</p> <p>Country and setting: Taiwan; Academic medical center</p> <p>Enrollment period: July 2004 to November 2005</p> <p>Funding: National Science Council, Taiwan</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: 12-wk RCT</p> <p>Intervention: Electric Stimulation (ES) vs. Oxybutynin vs. Placebo for 12 weeks</p> <p>Groups: G1: ES: intravaginal electrode; biphasic symmetric pulsed current w/ a 10-Hz frequency, 400-millisecond pulse width, 10/5 duty cycle, and varying intensity, 20 min/session, twice weekly G2: Oxybutynin, 2.5 mg 3 times daily G3: Placebo pill 3 times daily</p> <p>N at enrollment: G1: 25 G2: 26 G3: 23</p> <p>N at follow-up: G1: 24 G2: 23 G3: 21</p> <p>Age: NR</p> <p>Race/ethnicity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Female • OAB for ≥ 6 mos • Age 16-80 • Urgency ≥ 4 times per day <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Use of anticholinergics or TCAs • Previous treatment with pelvic-floor muscle training, bladder training • Pelvic prolapse repair • Pregnancy • Neurologic disorders • DM • Demand cardiac pacemaker • IUD use • Genital prolapse greater than ICS Stage II • PVR>100 mL • Overt SI • Anti-incontinence surgery • UTI 	<p>Warning time, s (range) G1: 41.5 (8-105) G2: 44 (2-215) G3: 65 (26-265)</p> <p>MVV, mL/void (range): G1: 340 (120-450) G2: 310 (130-800) G3: 350 (120-600)</p> <p>Daily voided volume, mL (range) G1: 2160 (1010-2950) G2: 2106 (1560-3153) G3: 2305 (1305-3300)</p> <p>Pads per day (range): G1: 1 (0-4.1) G2: 0 (0-3) G3: 1 (0-4)</p> <p>Urgency (subj): G1: 11.4 (4.0-16.0) G2: 12.1 (5-18) G3: 9.8 (4.2-15.5)</p> <p>Frequency (subj): G1: 12.8 (8.8-16.0) G2: 11.5 (4.3-19.5) G3: 11.5 (6.5-22.8)</p> <p>Nocturia (subj): G1: 1.75 (0-6.5) G2: 0 (0-2.5) G3: 0.65 (0-3.1)</p> <p>UI (subj): G1: 1 (0-2) G2: 0 (0-2) G3: 1 (0-2)</p>	<p>Warning time, s (range) G1: 72 (32-633) G2: 54.5 (138-850) G3: 66.5 (26-219) p-value (among 3 groups): <0.001</p> <p>MVV, mL/micturition (range): G1: 355 (150-550) G2: 336.5 (138-850) G3: 340 (160-600) p-value (among 3 groups): 0.035</p> <p>Voided volume/d, mL (range) G1: 2270 (1210-3106) G2: 2100 (1619-3200) G3: 2305 (1351-3221) p-value (among 3 groups): 0.050</p> <p>Pad count, pad/d (range): G1: 0 (0-2) G2: 0 (0-2.5) G3: 1 (0-3) p-value (among 3 groups): 0.012</p> <p>Urgency (subj) G1: 1.0 (0.0-12.3) G2: 6 (0.5-13) G3: 7.4 (3.9-13.4) p-value (among 3 groups): <0.001</p> <p>Frequency (subj) G1: 7.8 (1.8-13.0) G2: 7.4 (2-14) G3: 10 (3.9-13.4) p-value (among 3 groups): 0.002</p> <p>Nocturia (subj) G1: 0 (0-3.0) G2: 0 (0-2.5) G3: 1 (0-3.6) p-value (among 3 groups): 0.002</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Masking: -</p> <p>Pt selection criteria: +</p> <p>Loss to followup: ++</p> <p>Drop-out rates: +</p> <p>Power calculation: +</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Wang et al., 2006 (continued)				<p>UI (subj) G1: 0.5 (0-2) G2: 0 (0-2) G3: 1 (0-2) p-value (among 3 groups): 0.413</p> <p>P values before vs. after treatment</p> <p>Warning time, s (range): G1: 0.002 G2: 0.001 G3: 0.532</p> <p>MVV, mL/void (range): G1: 0.018 G2: 0.004 G3: 0.979</p> <p>Daily voided volume, mL (range): G1: 0.024 G2: 0.728 G3: 0.627</p> <p>Pad per day (range): G1: 0.010 G2: 0.662 G3: 0.501</p> <p>Urgency (subj): G1: <0.001 G2: <0.001 G3: 0.003</p> <p>Frequency (subj): G1: <0.001 G2: <0.001 G3: 0.070</p> <p>Nocturia (subj): G1: 0.001 G2: 0.394 G3: 0.176</p> <p>UI (subj): G1: 0.814 G2: 0.083 G3: 0.854</p>	

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Zinner et al., 2005</p> <p>Country and setting: US, Specialty treatment center</p> <p>Enrollment period: NR</p> <p>Funding: Pfizer Novartis Pharma Thomson ACUMED</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Randomized double-blind placebo-controlled four way crossover</p> <p>Intervention: Darifenacin 15 mg qd vs. darifenacin 30 mg qd vs. Oxybutynin vs. placebo</p> <p>Groups:* G1: Darifenacin 15mg daily G2: Darifenacin 30mg daily G3: Oxybutynin 5 mg t.i.d. G4: placebo</p> <p>N at enrollment: G1-G4: 76</p> <p>N included, tolerability: G1-G4: 61</p> <p>N included, efficacy: G1-G4: 58</p> <p>Women, n (%): G1-G4: 71 (93.4)</p> <p>Age, mean (range): G1-G4: 59.9 (33-84)</p> <p>Race/ethnicity: NR</p> <p>Weight (kg), mean (range): G1-G4: 75.7 (42-157)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age 18-85 • ≥ 4 UUI episodes/week • ≥ 8 voids/day (from 14 day run in placebo voiding diary) <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Neurogenic bladder • SUJ • Contraindication to antimuscarinic treatment • UTI • Bladder symptoms • Bladder stones • BOO • Concomitant disease • Concurrent bladder training • Concurrent thyroid or HRT • Other bladder meds • Pregnant or breastfeeding 	<p>Urgency episodes/day, mean \pm SD: G1-G 4: 9.3 \pm 3.4</p> <p>Urgency severity, mean \pm SD: G1-G4: 2.0 \pm 0.4</p> <p>Incontinence episodes/week, mean \pm SD: G1-G4: 20.4 \pm 17.7</p> <p>Voids/ day, mean \pm SD: G1-G4: 10.4 \pm 3.0</p>	<p>Urgency episodes/day, mean: G1: 7.95 G2: 7.59 G3: 8.12 G4: 8.71 G1-G3/G4: P < 0.05</p> <p>Urgency severity, mean: G1: 1.93 G2: 1.84 G3: 1.89 G4: 2.03 G1-G3/G4: P < 0.05</p> <p>Incontinence episodes/week, mean: G1: 10.93 G2: 8.82 G3: 9.45 G4: 14.64 G1-G3/G4: P < 0.05</p> <p>Voids/day, mean \pm SD: G1: 9.33 G2: 8.85 G3: 9.24 G4: 9.62 G2/G4: P < 0.05</p> <p>Dry mouth, (%): G1: 13.1 G2: 34.4 G3: 36.1 G4: 4.9 G1/G3: P < 0.05 G2/G3: P < 0.05 G2/G4: P < 0.05 G3/G4: P < 0.05</p> <p>Constipation, (%): G1: 9.8 G2: 21.3 G3: 8.2 G4: 3.3 G2/G3: P < 0.05 G2/G4: P < 0.05</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: + Masking: + Pt selection criteria: + Loss to followup: - Drop-out rates: NR Power calculation: + Statistical issues: + EXTERNAL VALIDITY: fair Age: -, NR Baseline OAB status: NR Baseline characteristics: - Length of followup: - Measurement methods: + Measurement reliability: + Intervention description: +</p>

Evidence Table 6. KQ 3 Comparison of Treatments (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Zinner et al., 2005 (continued)				Blurred vision, (%): G1: 0 G2: 0 G3: 3.3 G4: 0 Dizziness, (%): G1: 0 G2: 0 G3: 1.6 G4: 0	

Evidence Table 7. KQ4 Modifiers of outcomes

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Anderson et al., 2006</p> <p>[See evidence table for Diokno et al. 2003]</p> <p>Country and setting: US, Multicenter</p> <p>Enrollment period: NR</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: 1 of 6 ALZA Corp (1)</p>	<p>Design: RCT</p> <p>Intervention: different extended release anticholinergic medications in women with or without prior anticholinergic treatment for OAB</p> <p>Groups: G1: Prior anticholinergic treatment oxybutynin 10 mg po daily G1b: ER tolterodine, 4mg po daily G2: No prior anticholinergic treatment G2a: ER oxybutynin 10 mg po daily G2b: ER tolterodine, 4mg po daily Ga: ER oxybutynin 10 mg po daily Gb: ER tolterodine, 4mg po daily</p> <p>N at enrollment: Total: 790 G1: 373 G1a: 180 G1b: 193 G2: 417 G2a: 211 G2b: 206</p> <p>N at follow-up, 12 weeks: G1: 341 G1a: 165 G1b: 176 G2: 355 G2a: 174 G2b: 181</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Women • Age ≥ 18 • Mean of 21-60 UUI episodes per week and mean of ≥ 10 voids per 24 hr <p>Exclusion criteria: NR</p>	<p>UUI episodes/wk, mean ± SD: G1a: 36.8 ± 16.4 G1b: 37.4 ± 14.0 G2a: 37.5 ± 14.0 G2b: 36.2 ± 13.9</p> <p>Incontinence episodes/wk, mean ± SD: G1a: 41.5 ± 19.0 G1b: 43.0 ± 18.0 G2a: 45.0 ± 19.4 G2b: 41.9 ± 17.9</p> <p>Voids/week, mean ± SD: G1a: 92.8 ± 23.7 G1b: 94.6 ± 25.2 G2a: 96.5 ± 27.1 G2b: 97.9 ± 24.2</p>	<p>UUI episodes/wk, mean ± SD: G1a: 11.4 ± 17.9 G1b: 13.3 ± 15.1 G2a: 10.2 ± 13.7 G2b: 9.3 ± 13.3 G1a/G1b: P = 0.3</p> <p>UUI episodes/wk, completed wk 12, mean ± SD: G1a: 9.9 ± 14.1 G1b: 12.9 ± 14.9 P = 0.049</p> <p>No UUI, 12 wk, %: G1a: 25.2 G1b: 16.4 G2a: 29.4 G2b: 26.4 G1a/G1b: P = 0.046 G2a/G2b: P = 0.495</p> <p>Incontinence episodes/wk, mean ± SD: G1a: 12.7 ± 18.7 G1b: 16.5 ± 19.8 G2a: 11.9 ± 15.1 G2b: 11.3 ± 16.0 G1a/G1b: P = 0.09 (P = 0.012 if completed wk 12) G2a/G2b: P = 0.886</p> <p>Voids/week, mean ± SD: G1a: 68.4 ± 17.2 G1b: 72.8 ± 25.4 G2a: 64.8 ± 22.0 G2b: 69.4 ± 21.3 G1a/G1b: P = 0.05 (P = 0.026 iw 12) G2a/G2b: P = 0.035 (P = 0.026 wk 12)</p> <p>Dry mouth, % G1a: 32 G1b: 19 G2a: 27.5 G2b: 25.2 G1a/G1b: P = 0.004 G2a/G2b: P = 0.6</p>	<p>Quality: Overall quality score: fair INTERNAL VALIDITY: poor Randomization: - Masking: + Pt selection criteria: + Loss to followup: ++ Drop-out rates: ++ Power calculation: - Statistical issues: + EXTERNAL VALIDITY: good Age: + Baseline OAB status: + Baseline characteristics: ++ Length of followup: + Measurement methods: + Measurement reliability: + Intervention description: +</p>

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Anderson et al., 2006 (continued)	<p>Women, %: 100</p> <p>Age, mean ± SD: G1a: 62.6 ± 12.9 G1b: 62 ± 12.6 G2a: 57.5 ± 13.4 G2b: 58.8 ± 12.4</p> <p>Race/ethnicity, %: White: G1a: 87 G1b: 88 G2a: 82 G2b: 84 Black: G1a: 7 G1b: 9 G2a: 9 G2b: 8.7 Asian: G1a: 0.6 G1b: 0 G2a: 0 G2b: 1 Hispanic: G1a: 5.6 G1b: 3.1 G2a: 8.1 G2b: 6.3 Other G1a: 0 G1b: 0.5 G2a: 0.9 G2b: 0</p>			<p>Constipation, %: G1a: 7.8 G1b: 5.2 G2a: 5.2 G2b: 10.2</p> <p>Diarrhea, %: G1a: 7.8 G1b: 5.7 G2a: 8.1 G2b: 6.8</p> <p>Headache, %: G1a: 4.4 G1b: 5.2 G2a: 6.6 G2b: 6.8</p> <p>Discontinued d/t AE, n (%): G1a: 7 (3.9) G1b: 6 (3.1) G2a: 13 (6.2) G2b: 13 (6.3)</p> <p>Withdrew, %: G1: 3.5 G2: 6.2</p>	

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Appell, Abrams et al., 2001</p> <p>Country and setting: Multinational, Multicenter</p> <p>Enrollment period: NR</p> <p>Funding: Pharmacia</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Open-label extension study (after 12-wk RCT and 1 wk washout)</p> <p>Intervention: Tolterodine 2 mg BID with option for patients to self-lower their dosage to 1mg bid</p> <p>Groups: NA</p> <p>N at enrollment: 854</p> <p>N at follow-up, n (%): 594 (70)</p> <p>Age, yrs, mean (range): 59.5 (19-89)</p> <p>Race/ethnicity: NR</p> <p>Women, %: 76</p> <p>Parity mean ± SD: NR</p> <p>Previous drug therapy for OAB, n (%) 472 (55)</p> <p>Previous lower urinary tract surgery, n (%) 291 (34)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • 18+ years old • Cystometric evidence of detrusor overactivity (phasic detrusor contraction with amplitude 10+ cm H2O • ≥ 8 voids/day • ≥ 1+ urinary incontinence episode/day <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • SUI • Hepatic or renal disease • UTI • IC • Hematuria • Contraindication to antimuscarinic therapy • Previous serious AE on oxybutynin • Voiding difficulty w/ treatment of urinary retention • Treatment w/in 14 days prior to baseline visit • Initiation of antimuscarinic or any drug for UI during study • Electro-stimulation or bladder training • Indwelling or intermittent catheter • Total voided volume >3L/day 	<p>Detrusor instability, %: 93.7</p> <p>Urge incontinence episodes, n (%): 724 (85)</p> <p>Symptom duration > 5 yrs, n (%): 412 (48)</p> <p>Urgency, n (%): 841 (98)</p> <p>Severe/very severe problems, n (%): 384 (45)</p> <p>voids/day, mean (range): 11.4 (5.3-37.0)</p> <p>Urge incontinence episodes/day, mean (range): 3.5 (0.1-24.0)</p> <p>Volume voided mean mL (range): 159 (25-423)</p> <p>Adverse events at end of 12-wk RCT, n (%): Any: 358 (76) ANS: 203 (43) CNS: 59 (12) GI: 125 (26) Respiratory: 68 (14) Urinary: 50 (11) Dry mouth: 187 (39) UTI: 26 (5) Headache: 49 (10) Constipation: 31 (7) Abdominal pain: 36 (8) URI: 28 (6)</p>	<p>Voids/day, 3 mos, mean (range) 8.8 (2.0-23.4)</p> <p>Voids/day, 3 months, mean change (95% CI): -2.6 (-2.8 to -2.3) P = 0.0001</p> <p>Voids/day, 9 mos, mean (range): 8.9 (1.9-31.6)</p> <p>Voids/day, 9 months, mean change (95% CI): -2.5 (-2.9 to -2.4) P = 0.0001</p> <p>Void frequency, 9 mos, median change: -22%</p> <p>Urge incontinence episodes/day, 3 mos, mean (range): 1.3 (0.0-24.0)</p> <p>Urge incontinence episodes/day, mean change (95% CI): -2.1 (-2.4 to -1.9) P = 0.0001</p> <p>Urge incontinence episodes/day, 3 mos, mean (range): 1.5 (0.0-24.0)</p> <p>Urge incontinence episodes/day, mean change (95% CI): -2.0 (-2.2 to -1.7) P = 0.0001</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: -</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: +</p> <p>Drop-out rates: +</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Appell, Abrams et al., 2001 (continued)				<p>Urge Incontinence episodes/day, median change, %: -76%</p> <p>Volume voided, 3 mos, mean mL (range): 201 (33-444)</p> <p>Volume voided, 3 mos, mean change mL (95% CI): +41 (36-45) P = 0.0001</p> <p>Volume voided, 9 mos, mean change mL (range): 199 (34-514)</p> <p>Volume voided mean change mL (95% CI): +40 (35-45) P = 0.0001</p> <p>Volume voided, 9 mos, median change, %: +22%</p> <p>Improvement, 9 mos, %: 65</p> <p>Any adverse event, n (%): 652 (76)</p> <p>ANS, n (%): 268 (31)</p> <p>General, n (%): 219 (26)</p> <p>CNS/PNS, n (%): 82 (10)</p> <p>GI, n (%): 201 (24)</p> <p>Respiratory, n (%): 139 (16)</p> <p>Urinary, n (%): 165 (19)</p>	

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Appell, Abrams et al., 2001 (continued)				<p>Dry mouth, n (%): 236 (28)</p> <p>Mild dry mouth, %: 19%</p> <p>Moderate dry mouth, %: 7%</p> <p>Severe dry mouth, %: 2%</p> <p>UTI, n (%): 106 (12)</p> <p>Headache, n (%): 57 (7)</p> <p>Constipation, n (%): 57 (7)</p> <p>Abdominal pain, n (%): 50 (6)</p> <p>Upper respiratory tract infection, n (%): 45 (5)</p> <p>Serious adverse events, n: 72</p> <p>Discontinued due to AE, n (%): 73 (9)</p> <p>Reduced dosage to 1 mg bid, n (%): 108 (13%)</p>	

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Capo et al. 2008</p> <p>Country and setting: US, 207 centers</p> <p>Enrollment period: NR</p> <p>Funding: Astellas Pharma US, Inc., and GlaxoSmithKline</p> <p>Author industry relationship disclosures: 5/5 Astellas (4) Boehringer Mannheim Pharmaceuticals (1) Cephalon Inc. (1) GlaxoSmithKline (3) Reliant Pharmaceuticals (1) Schering Plough Corporation (1) Abbott (1) Akros (1) AstraZenica (1) Johnson & Johnson (1) Lilly (1) Manking (1) Merck (1) Novartis (1) NovoNordisk (1) Pfizer Inc. (1) Solvay (1) Takeda & TAP (1) Pri Med (1) Lucente (1)</p>	<p>Design: RCT</p> <p>Intervention: 12-wk treatment with 5 mg solifenacin succinate. Dosage could be increased at wk 4 and wk 8 to 10 mg, maintained or decreased in response to pt perceived efficacy and tolerability</p> <p>Groups: G1: Hispanic subgroup of G2 G2: full study population including G1</p> <p>N at enrollment: G1: 94 G2: 2205</p> <p>N at follow-up, range: G1: 47-80 G2: 1254-2005</p> <p>Age, yrs ± SD: G1: 51.8 ± 12.8 G2: 59.7 ± 14.4</p> <p>Weight in lbs, mean ± SD: G1: 171.1±36.4 G2: 182.2±47.0</p> <p>Race/ethnicity, n (%): G1: Hispanic 94 (100) G2: White 1761 (79.9) Hispanic 94 (4.3) Black 274 (12.4) Asian 59 (2.7) Other 17 (0.8)</p> <p>Women, N (%): G1: 74 (78.7) G2: 1813 (82.2)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • ≥ 18 yrs of age • OAB symptoms ≥3 mos • Toilet without difficulty • Washout of other OAB meds ≥ 7 days • Non-drug treatment of OAB if established ≥ 4 wks prior to study and continued <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • SUI • Stress predominant MUI • UTI or chronic inflammation clinically significant outflow • Obstruction due to BPH • Narrow-angle glaucoma • Urinary or gastric retention • Severe renal or hepatic impairment • Chronic severe constipation or gastrointestinal obstructive disease • Bladder cancer • Women not using a reliable contraceptive • Hypersensitivity to the study medication, any of its components, or to anticholinergic medication 	<p>Urge incontinence, n (%): G1: 63 (67.0) G2: 1586 (71.9)</p> <p>Urinary urgency, n %: G1: 88 (93.6) G2: 2007 (91.0)</p> <p>Frequency, n (%): G1: 86 (91.5) G2: 1969 (89.3)</p> <p>Nocturia, n (%): G1: 79 (84.0) G2: 1792 (81.3)</p> <p>Patient perception of bladder condition, mean score: G1: 4.0 G2: 4.0</p> <p>Most bothersome OAB symptom-frequency, n %: G1: 38 (40.4) G2: 619 (28.1)</p> <p>Most bothersome OAB symptom-urinary urgency, n %: G1: 22 (23.4) G2: 508 (23.0)</p> <p>Most bothersome OAB symptom-urge incontinence, n %: G1: 17 (18.1) G2: 602 (27.3)</p>	<p>Increased dosage 5 mg/day, n (%)-Wk 4: G1: 46 (48.9)</p> <p>Increased dosage 10 mg/day, n (%)-Wk 4: G1: 37 (39.4)</p> <p>Discontinued, n Wk 4: G1: 1</p> <p>Increased dosage 5 mg/day, n (%)-Wk 8: G1: 31 (33.0)</p> <p>Increased dosage 10 mg/day, n (%)-Wk 8: G1: 39 (41.5)</p> <p>Discontinued-Wk 8, n: 1</p> <p>Returned to 5 mg/day-Wk 8, n: 5</p> <p>Increased dosage to 10 mg/day-Wk 8, n: 12</p> <p>Patient Perception of Bladder, mean Wk 4: G1: 3.3* G2: 3.3*</p> <p>Patient Perception of Bladder, mean Wk 8: G1: 2.8* G2: 2.8*</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: NA</p> <p>Masking: NA</p> <p>Pt selection criteria: +</p> <p>Loss to followup: -</p> <p>Drop-out rates: -</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: +</p> <p>Baseline OAB status: NR</p> <p>Baseline characteristics: -</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Capo et al. 2008 (continued)	Parity : NR		<p>Most bothersome OAB symptom-nocturia, n %: G1: 12 (12.8) G2: 337 (15.3)</p> <p>Most bothersome OAB symptom-none specified, n %: G1: 5 (5.3) G2: 139 (6.3)</p>	<p>Patient Perception of Bladder, mean Wk 12: G1: 3.0* G2: 2.9* *P<0.001 vs BL</p> <p>Symptom bother from VAS, change from Baseline, mean scores (mm), p-value, CI: Urinary urgency: G1: -39.2 P<0.001, -45.9, -32.5 G2: -39.5 P<0.001, -41.0, -38.1</p> <p>Urge Incontinence: G1: -37.7 P<0.001, -46.0, -29.4 G2: -40.1, P<0.001, -41.8, -38.4</p> <p>Frequency: G1: -40.0 P<0.001, -46.9, -33.2 G2: -41.8 P<0.0001, -43.3, -40.3</p> <p>Nocturia: G1: -43.2 P<0.001, -50.4, -36.1 G2: -36.9, -38.4, -35.4</p> <p>OAB-q subscales at end-point, change from baseline, mean, p-value, CI:</p> <p>Symptom bother: G1: -29.1 P<0.001, -34.8, -23.4 G2: -29.6 P<0.001, -30.7, -28.6</p>	

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Capo et al. 2008 (continued)				<p>Coping: G1: 24.0 P<0.001, 18.0, 30.1 G2: 27.4 P< 0.001 26.2, 28.5</p> <p>Concern: G1: 25.8 P<0.001, 19.5, 32.2 G2: 29.6 P< 0.001, 28.4, 30.8</p> <p>Sleep: G1: 25.7 P<0.001, 19.4, 32.0 G2: 27.3 P< 0.001, 26.1, 28.5</p> <p>Social Interaction G1: 15.9 P<0.001, 11.5, 20.3 G2: 14.7 P< 0.001, 13.7, 15.6</p> <p>Overall Health-related QoL: G1: 23.2 P<0.001, 18.0, 28.5 G2: 25.4 P< 0.001, 24.4, 26.4</p> <p>Any Adverse Event, n (%): G1: 48 (51.4) G2: 1321 (59.4)</p> <p>Dry mouth, n (%): G1: 17 (18.1) G2: 477 (21.4)</p> <p>Constipation, n (%): G1: 12 (12.8) G2: 295 (13.3)</p> <p>Headache, n (%): G1: 3 (3.2) G2: 76 (3.4)</p>	

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Capo et al. 2008 (continued)				<p>Dizziness, n (%): G1: 2 (2.1) G2: 27 (1.2)</p> <p>Blurred vision, n (%): G1: 4 (4.3) G2: 57 (2.6)</p> <p>Palpitations, n (%): G1: 2 (2.1) G2: 6 (0.3)</p> <p>Upper respiratory tract infection, n (%): G1: 3 (3.2) G2: 68 (3.1)</p> <p>UTI, n (%): G1: 3 (3.2) G2: 76 (3.4)</p> <p>Bronchitis, n (%): G1: 3 (3.2) G2: 31 (1.4)</p> <p>Nasopharyngitis, n (%): G1: 2 (2.1) G2: 50 (2.3)</p> <p>Influenza, n (%): G1: 2 (2.1) G2: 11 (0.5)</p> <p>Insomnia, n (%): G1: 2 (2.1) G2: 17 (0.8)</p> <p>Depression, n (%): G1: 2 (2.1) G2: 19 (0.9)</p> <p>Hypertension, n (%): G1: 2 (2.1) G2: 16 (0.7)</p>	

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Dmochowski et al., 2002</p> <p>Country and setting: US, Community</p> <p>Enrollment period: NR 2 week washout 12 week treatment period</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: 7 of 7 Abbot (1) Alza (3) Amgen (1) AstraZeneca (1) Bayer (1) Bioform (1) Genyx (1) Glaxo (1) Interneuron (2) Lilly (2) Merck (1) Otsulta (1) Pharmacia (3) Pfizer (1) Praecis (2) Roche (1) Seprecor (1) Surx (1) Synthelabo (1) Vivus (1) Watson (7) Yamanouchi (1)</p>	<p>Design: RCT Followed by: 12 week open label dose titration</p> <p>Intervention: Oxybutynin TDS vs placebo</p> <p>Groups: G1: OXY TDS 1.3 mg G2: OXY TDS 2.6 mg G3: OXY TDS 3.9 mg G4: placebo</p> <p>N at enrollment: G1: 130 G2: 133 G3: 125 G4: 132</p> <p>N at follow-up: G1: 128 G2: 131 G3: 123 G4: 130</p> <p>Women, N (%): G1: 120 (92.3) G2: 123 (92.5) G3: 114 (91.2) G4: 121 (91.7)</p> <p>Age, mean ± SD: G1: 61.5 ± 11.8 G2: 61.9 ± 13.5 G3: 59.4 ± 14.5 G4: 62.7 ± 13.1</p> <p>Race/ethnicity, n (%): White: G1: 119 (91.5) G2: 118 (88.7) G3: 118 (94.4) G4: 118 (89.4) Black: G1: 7 (5.4) G2: 10 (7.5) G3: 3 (2.4) G4: 11 (8.3)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 18 • History of OAB • ≥ 10 episodes UUI either pure urge or predominant urge on 7 day voiding diary • ≥ 56 voids/ 7 day diary • Average recorded urinary volume of ≤ 350 mL <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Incontinence related to chronic illness • Concomitant medications • History of lower urinary tract surgery in ≤ 6 mos • IC • Urethral syndrome • Painful bladder syndrome • Overflow urinary incontinence • Alcohol/drug abuse • Known hypersensitivity to oxybutynin, similar compounds or transdermal systems • Active skin disorder • Narrow angle glaucoma • Excessive caffeine intake > 5 cups/day 	<p>Incontinence episodes/week, median: G1: 31.0 G2: 30.0 G3: 31.0 G4: 30.0</p> <p>Voids/day, mean ± SD: G1: 12.4 ± 2.9 G2: 12.1 ± 3.3 G3: 12.3 ± 3.3</p> <p>Voided volume (mL), mean: G1: 175 G2: 165 G3: 156 G4: 170</p> <p>Open Label: Incontinence episodes/week, open label, median: G1: 30.0 G2: 29.0 G3: 37.0 G4: NA</p> <p>Prior years incontinent, years, mean ± SD: G1: 9.1 ± 10.3 G2: 8.9 ± 8.8 G3: 9.9 ± 9.8 G4: 9.1 ± 9.1</p> <p>IIQ (QoL), total score, mean: G1: 167 G2: 161 G3: 144 G4: 160</p> <p>Prior anticholinergic treatment, n, (%): G1: 30 (23.1) G2: 28 (21.1) G3: 28 (22.4) G4: 27 (20.5)</p>	<p>Incontinence episodes/ week, median change: G1: -16.0 G2: -14.0 G3: -19.0 G4: -14.5 G3/G4: P < 0.0165</p> <p>Voids/day, mean change ± SD: G1: -1.8 ± 2.6 G2: -1.8 ± 2.4 G3: -2.3 ± 2.5 G4: -1.7 ± 3.0 G3/G4: P = 0.0457</p> <p>Voided volume, mL, mean increase ± SD: G1: -2 G2: 19 G3: 24 G4: 6 G3/G4: P = 0.0063 G2/G4: P = 0.0157</p> <p>Open Label: Incontinence episodes/week, median change: G1: -18.0 G2: -17.0 G3: -19.0 G4: NA G3/G4: P = 0.0327</p> <p>IIQ (QoL), total score, mean ± SD: G1: NR G2: NR G3: 78.8 ± 51.9 G4: 94.7 ± 50.0 G3/G4: P = 0.0266</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: - Masking: + Pt selection criteria: + Loss to followup: + Drop-out rates: - Power calculation: + Statistical issues: + EXTERNAL VALIDITY: good</p> <p>Age: + Baseline OAB status: + Baseline characteristics: ++ Length of followup: + Measurement methods: + Measurement reliability: + Intervention description: +</p>

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Dmochowski et al., 2002 (continued)	Other: G1: 4 (3.1) G2: 5 (3.8) G3: 4 (3.2) G4: 3 (2.3)			Dry mouth, n (%): G1: 6 (4.6) G2: 9 (6.8) G3: 12 (9.6) G4: 11 (8.3) Dizziness, n (%): G1: 2 (1.5) G2: 4 (3.0) G3: 5 (4.0) G4: 5 (3.8) Dysuria, n (%): G1: 1 (0.8) G2: 3 (2.3) G3: 3 (2.4) G4: 0 (0) Somnolence, n (%): G1: 1 (0.8) G2: 0 (0) G3: 2 (1.6) G4: 1 (0.8) Nausea, n (%): G1: 6 (4.6) G2: 5 (3.8) G3: 2 (1.6) G4: 7 (5.3) Constipation, (%): G1: 7 (5.4) G2: 3 (2.3) G3: 1 (0.8) G4: 4 (3.0) Palpitations, n (%): G1: 1 (0.8) G2: 0 (0) G3: 1 (0.8) G4: 0 (0) Vision abnormal, n (%): G1: 3 (2.3) G2: 2 (1.5) G3: 0 (0) G4: 2 (1.5) Localized application site reactions, n (%): G1: 32 (26.4) G2: 7 (5.7) G3: 8 (6.9)	

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Dmochowski et al., 2002 (continued)				Application site erythema, n (%): G1: 23 (19) G2: 29 (23.6) G3: 14 (12.0)	
				Application site erythema, mild, double blind period, n (%): G1: 79 (31.5) G2: 29 (36.2) G3: NA	
				Application site erythema, moderate, double blind period, n (%): G1: 46 (18.3) G2: 46 (18.1) G3: NA	
				Application site erythema, severe, double blind period, n (%): G1: 6 (2.4) G2: 8 (3.1) G3: NA	
				Application site erythema, mild, open label, n (%): G1: 18 (34.6) G2: 58 (38.4) G3: 87 (43.9)	
				Application site erythema, moderate, open label, n (%): G1: 7(13.5) G2: 23 (15.2) G3: 24 (12.1)	
				Application site erythema, severe, open label, n (%): G1: 0 (0) G2: 2 (1.3) G3: 0 (0)	

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Dmochowski et al. 2003</p> <p>Country and setting: US; Community</p> <p>Enrollment period: NR</p> <p>2 week washout 12 week treatment period</p> <p>Funding:</p> <p>Watson Pharma</p> <p>Author industry relationship disclosures: 6 of 6</p> <p>Watson Pharma</p>	<p>Design: RCT</p> <p>Intervention: Oxybutynin TDS vs Tolterodine ER vs placebo, 2 week washout plus 12 weeks treatment period; drug/placebo applied transdermally twice weekly to the abdomen and oral capsule ingested once daily</p> <p>Groups: G1: OXY TDS 3.9 mg/day G2: Tolterodine ER 4 mg daily G3: placebo</p> <p>N at enrollment: G1: 121 G2: 123 G3: 117 Total: 361</p> <p>N at follow-up: G1: NR G2: NR G3: NR Total: 320 (89%)</p> <p>Age, mean yrs ± SD: G1: 63.1 ± 12.0 G2: 62.9 ± 13.5 G3: 64.5 ± 12.3</p> <p>Race/ethnicity, mean (%): White: G1: 111 (91.7) G2: 120 (97.6) G3: 110 (94.0)</p> <p>Race/ethnicity, mean (%): Black: G1: 8 (6.6) G2: 1(0.8) G3: 4 (3.4)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • ≥ 18yo men and women • Current pharmacological treatment for OAB with beneficial response • ≥4 episodes UUI episodes either pure urge or predominant urge on 3 day voiding diary • ≥ 24 voids/ 3 day diary • Average recorded urinary volume of ≤350 ml <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • History of lower urinary tract surgery in previous 6 months • IC • Urethral syndrome • Painful bladder syndrome • Overflow urinary incontinence 	<p>Incontinence episodes/ day, mean ± SD: G1: 4.7 ± 2.9 G2: 5.0 ± 2.9 G3: 5.0± 3.2</p> <p>Incontinence episodes/day, median: G1: 4 G2: 4 G3: 4</p> <p>Void frequency/ day, mean ± SD: G1: 12.4 ± 2.9 G2: 12.1 ± 3.3 G3: 12.3 ± 3.3</p> <p>Void frequency/day, median: G1: 12 G2: 12 G3: 12</p> <p>Voided volume, mL, mean ± SD: G1: 165 ± 62 G2: 165 ± 61 G3: 175 ± 68</p> <p>Voided volume, mL, median: G1: 160 G2: 150 G3: 171</p> <p>Global assessment of disease, QOL, baseline, (SD): G1: 65 (19) G2: 63 (20) G3: 65 (20)</p> <p>IIQ (QOL)- travel domain, baseline, (SD): G1: 49 (28) G2: 47 (29) G3: 43 (29)</p>	<p>Incontinence episodes/ day, mean ± SD: G1: 1.9 ± 2.7 G2: 1.9 ± 3.0 G3: 2.9 ± 3.8</p> <p>P=0.0137 G1 v G3 P=0.0011 G2 v G3 P=0.5878 G1 v G2</p> <p>Incontinence episodes/day, mean change±SD G1: 2.9±3.0 G2: 3.2±2.8 G3: 2.1±3.0</p> <p>Incontinence episodes/day, median G1: 1 G2: 1 G3: 2</p> <p>Incontinence episodes/day, median change G1: 3 G2: 3 G3: 2</p> <p>Void frequency/ day, mean ± SD: G1: 10.4 ± 3.2 G2: 9.9 ± 3.1 G3: 10.9 ± 3.8</p> <p>P=0.101 G1 v G3 P= 0.0025 G2 v G3 P=0.276 G1 v G2</p> <p>Void frequency/day, mean change±SD: G1: -1.9±2.7 G2: -2.2±2.6 G3: -1±1.4</p> <p>Void frequency/day, median: G1: 10 G2: 10 G3: 10</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: - Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: +</p> <p>Drop-out rates: +</p> <p>Power calculation: +</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Dmochowski et al. 2003 (continued)	<p>Race/ethnicity, mean (%): Other: G1: 2 (1.6) G2: 2 (1.6) G3: 3 (2.6)</p> <p>Women, N (%): G1: 109 (90.1) G2: 117 (95.1) G3: 109 (93.2)</p> <p>Parity : NR</p> <p>Prior antimuscarinic treatment, Tolterodine, n, (%): G1: 57 (47) G2: 60 (49) G3: 54 (46)</p> <p>Prior antimuscarinic treatment, Oxybutynin, n, (%): G1: 61 (51) G2: 59 (48) G3: 59 (50)</p> <p>Prior antimuscarinic treatment, Other, n, (%): G1: 7 (6) G2: 6 (5) G3: 6 (5)</p>		<p>UDI (QOL), irritative symptoms, baseline, (SD): G1: 62 (20) G2: 66 (18) G3: 63 (20)</p>	<p>Void frequency/day, median change: G1: 2 G2: 2 G3: 1</p> <p>Voided volume, mL, mean ± SD: G1: 198 ± 84 G2: 193 ± 75 G3: 182 ± 84 P=0.0010 G1 v G3 P=0.0017 G2 v G3 P=0.7690 G1 v G2</p> <p>Voided volume, mL, mean change±SD: G1: 32±55 G2: 29±57 G3: 9±63</p> <p>Voided volume, mL, median: G1: 188 G2: 189 G3: 165</p> <p>Voided volume, mL, median change: G1: 24 G2: 29 G3: 5.5</p> <p>Global assessment of disease, QOL, change, (SD): G1: 30 (30) G2: 33 (28) G3: 21 (31) P=0.0106 G1 v G3 P=0.001 G2 v G3 P=0.1861 G1 v G3</p> <p>IIQ (QOL)- travel domain, change, (%): G1: 23 (25) G2: 22 (29) G3: 11 (30) P=0.0018 G1 v G3 P=0.0045 G2 v G3</p>	

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Dmochowski et al. 2003 (continued)				<p>UDI (QOL), irritative symptoms, change, (SD): G1: 25 (26) G2: 28 (26) G3: 18 (24) P=0.0156 G1 v G3 P= 0.0010 G2 v G3</p> <p>Treatment compliance with assigned dosage regimen: 92%:</p> <p>Frequency decreased to a greater extent for patients with > 14 micturations per day at baseline G1: -2.9/day, p=0.0036 (data for other groups not reported)</p> <p>Adverse effects:</p> <p>Dry mouth, %: G1: 4.1 G2: 7.3 G3: 1.7 P=0.2678 G1 v G3 P=0.0379 G2 v G3</p> <p>Constipation, (%): G1: 3.3 G2: 5.7 G3: NR</p> <p>Mild systemic adverse effects, n (%): G1: 15 (12.4) G2: 13 (10.6) G3: 6 (5.1)</p> <p>Moderate systemic adverse effects, n (%): G1: 7 (5.8) G2: 13 (10.6) G3: 7 (6.0)</p>	

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Dmochowski et al. 2003 (continued)				<p>Severe systemic adverse effects, n (%) G1: 1 (0.8) G2: 3 (2.4) G3: 1 (0.9)</p> <p>Mild localized application site reactions, n (%): G1: 9 (7.4) G2: 2 (1.6) G3: 5 (4.3)</p> <p>Moderate localized application site reactions, n (%): G1: 17 (14.0) G2: 4 (3.3) G3: 2 (1.7)</p> <p>Severe localized application site reactions, n (%): G1: 6 (5.0) G2: 1 (0.8) G3: 1 (0.9)</p> <p>Treatment discontinuation due to adverse effects, n (%): G1: 13 (10.7) (12 due to application site reactions, 1 due to hot flushes) G2: 2 (1.6) (1 due to fatigue and 1 due to dizziness) G3: NR</p> <p>Postvoid residual >150 mL at end of treatment: G1: 4 G2: 4 G3: 3 No reports of symptomatic urinary retention</p> <p>Withdraw due to AEs: 23</p>	

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Elinoff et al. 2006 Roberts et al. 2006</p> <p>Country and setting: US; 82 Primary care & Ob-Gyn offices</p> <p>Enrollment period: NR</p> <p>Funding: Pfizer</p> <p>Author industry relationship disclosures: 6 of 6 Pfizer (6) Astellas (1) Novartis (1)</p>	<p>Design: Open-label, single-arm cohort</p> <p>Intervention: Tolterodine ER 4 mg qd for 12 wks</p> <p>Groups: NA</p> <p>N at enrollment: 896</p> <p>N at follow-up: 758</p> <p>Age, yrs ± SD: 56 ± 15</p> <p>Race/ethnicity, N (%): White: 691 (80) Black: 94 (11) Asian: 22 (3) Other: 53 (6)</p> <p>Women, N (%): 708 (82)</p> <p>Parity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • 18+ years old • OAB symptoms for ≥3 mos • ≥ 8 voids/day • 2+ episodes of urgency or UUI in a 3-d period • 3+ on the OAB Bother Rating Scale <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Stress, functional, or overflow incontinence • Acute UTI • Clinically significant LUT pathology • Indwelling catheter • Intermittent self-cath • Use of anticholinergics, antimuscarinics for OAB, drugs to trt UUI, potent cytochrome P450 3A4 inhibitors, or investigational drugs 	<p>Duration of OAB, yrs ± SD: 6 (8)</p> <p>Trt duration w/ study drug, days (SD): 80 (19)</p> <p>Daytime frequency episodes, #/d, mean ± SD: 9.3 ± 2.8</p> <p>Nocturnal frequency episodes, #/d, mean ± SD: 3.0 ± 1.7</p> <p>UUI episodes, #/d, mean ± SD: 2.7 ± 3.2</p> <p>Urgency episodes, #/d, mean ± SD: 5.0 ± 3.8</p> <p>OAB-q scores Symptom bother: 60.0 Coping: 50.0 Concern: 51.4 Sleep: 44.4 Social Interaction: 84.0 Total HRQL: 55.2</p> <p>AUA-SI Total: 18 Irritative: 10 Obstructive: 8</p> <p>Most bothersome symptom, n (%) Daytime frequency: 256 (30) Nocturnal frequency: 241 (28) UUI: 206 (24) Urgency: 154 (18)</p>	<p>Urgency, mean change% (95% CI), wk 12:[^] -81.3 (-85.7, -73.3)</p> <p>UUI episodes, mean change % (95% CI), wk 12:[^] -80.0 (-85.7, -69.7)</p> <p>Nighttime frequency, mean change % (95% CI), wk 12:[^] -40.0 (-44.4, -33.3)</p> <p>Daytime frequency, mean change % (95% CI), wk 12:[^] -31.6 (-34.6, -28.1)</p> <p>OAB-q scores, Median change from baseline to 12 wks (95% CI): Symptom bother: -37.5 (-37.5, -35.0) Coping: 32.5 (30.0, 35.0) Concern: 34.3 (31.4, 37.1) Sleep: 28.0 (28.0, 32.0) Social Interaction: 12.0 (12.0 to 16.0) Total HRQL: 28.9 (27.2 to 31.2)</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: NA</p> <p>Masking: NA</p> <p>Pt selection criteria: +</p> <p>Loss to followup: +</p> <p>Drop-out rates: +</p> <p>Power calculation: +</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Elinoff et al. 2006 Roberts et al. 2006 (continued)				<p><u>AUA-SI, Median change from baseline to 12 wks (95% CI)</u> *Total: -9 (-9 to -8) *Irritative: -5 (-5 to -5) *Obstructive: -4 (-4 to -3) *: p<0.001</p> <p>Urgency, mean change, wk 12 % (95% CI): -75.0 (-80.0, -71.4)</p> <p>Change from baseline to wk 12, UUI episodes, mean % (95% CI): -86.1(-80.0, -71.4)</p> <p>Nighttime frequency, mean change % (95% CI): -37.5 (-40.0, -33.3)</p> <p>Daytime frequency, mean change, wk 12 % (95% CI): -29.0 (-31.0, -27.3)</p> <p>All-cause AE: 51%</p> <p>Discontinued treatment, n %: 15 (7)</p> <p>Trt-related AE: 23%</p> <p>Dry mouth, %: 10.0</p> <p>Constipation, %: 3.7%</p> <p>Headache, %: 3.0%</p> <p>UTI, %: 2.7%</p>	

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Garely et al., 2007*</p> <p>Mallett et al., 2007^</p> <p>Country and setting: US, multicenter (207 sites)</p> <p>Enrollment period: NR</p> <p>Funding: Astellas Pharma US Glaxo-SmithKline</p> <p>Author industry relationship disclosures: 4 of 4* Astellas (4) Pfizer (1) Novartis (1)</p> <p>4 of 4^ Astellas (3) GlaxoSmithKline (2) Novartis (1) Watson (1) Yamanouchi (1)</p>	<p>Design: Prospective case series</p> <p>Intervention: Solifenacin qd (started on 5 mg/d; option of increasing to 10 mg/d at week 4; option to maintain 10 mg dose or decrease to 5 mg at wk 8); for 12 weeks of treatment</p> <p>10 mg dose, n (%): 4 wks: G1: 55%</p> <p>8 wks: G1: 59% (8% went back to 5mg; 19% increased to 10 mg)</p> <p>Groups: G1: Urge incontinence as most bothersome symptom G2: Black participants G3: Full study population</p> <p>N at enrollment: G1: 582 G2: 274 G3: 2205</p> <p>Age, mean yrs ± SD: G1: 60.9 ± 13.1 G2: 54.7 ± 13.6 G3: 59.7 ± 14.4</p> <p>Race/ethnicity, n (%): White: G1: 493 (84.7) G3: 1761 (79.9) African American: G1: 62 (10.7%) G3: 274 (12.4%)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> -Age ≥ 18 Symptoms of OAB for ≥ 3 mos Ambulatory Able to use the toilet without difficulty Urgency, UUI, frequency or nocturia ≥ 3 mos <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Previous use of solifenacin 	<p>UUI, n (%): G1: 582 (100%) G2: NR G3: 1596 (71.9%)</p> <p>Urgency, n (%): G1: 534 (91.8) G2: NR G3: 2007 (91.0)</p> <p>Frequency, n (%): G1: 488 (83.8) G2: NR G3: 1969 (89.3)</p> <p>Nocturia, n (%): G1: 438 (75.3) G2: NR G3: 1792 (81.3)</p> <p>Most bothersome OAB symptom</p> <p>Urgency, n (%): G2: 48 (17.5) G3: 508 (23.0)</p> <p>UUI, n (%): G1: 582 (100) G2: 63 (23.0) G3: 602 (27.3)</p> <p>Frequency, n (%): G2: 106 (38.7) G3: 619 (28.1)</p> <p>Nocturia, n (%): G2: 34 (12.4) G3: 337 (15.3)</p> <p>None specified, n (%): G2: 23 (8.4) G3: 139 (6.3)</p> <p>OAB for 3 mos -1 yr, n (%): G1: 57 (9.8) G3: 349 (15.8%)</p> <p>OAB for 1-5 yrs, n (%): G1: 294 (50.5) G3: 1124 (51.0)</p> <p>OAB for >5 yrs: G1: 231 (39.7) G3: 732 (33.2)</p>	<p>Patient perception of bladder condition scale, 4 wks, mean: G1: 3.3* G2: 3.1* G3: 3.3*</p> <p>Patient Perception of Bladder Condition Scale, 8 wks, mean: G1: 3.0* G2: 2.7* G3: 2.9*</p> <p>Patient Perception of Bladder Condition Scale, 12 wks or early termination, mean: G1: 2.9* G2: 2.6* G3: 2.9*</p> <p>Symptom severity, mean VAS, wk 4: Urinary urgency: G1: 40.9 G2: 39.7 G3: 40.6 UUI: G1: 36.9 G2: 29.9 G3: 32.8 Frequency: G1: 34.5 G2: 39.3 G3: 39.9 Nocturia: G1: 30.4 G2: 35.4 G3: 38.2</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: NA</p> <p>Masking: NA</p> <p>Pt selection criteria: -</p> <p>Loss to followup: NR</p> <p>Drop-out rates: +</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Garely et al., 2007*	Other: G1: 27 (4.6%) G3: 170 (7.7%)		Perception of bladder condition scale (mean): G1: 4.6 G2: 4.2 G3: 4.4	Symptom severity, mean VAS, wk 8: Urinary urgency: G1: 31.1 G2: 29.4 G3: 30.0	
Mallett et al., 2007^ (continued)	Women, n (%): G1: 536 (92.1) G2: 213 (77.7) G3: 1813 (82.2) Weight (lbs), mean ± SD (range): G2: 198 ± 51.6 (100-400) G3: 182.2 ± 47.0 (80-413)		Symptom severity, mean VAS: Urinary urgency: G1: 72.3 G2: 70.8 G3: 68.7 UUI: G1: 78.5 G2: 70.8 G3: 64.1 Frequency: G1: 65.7 G2: 70.8 G3: 70.6 Nocturia: G1: 57.9 G2: 66.5 G3: 65.2 OAB-q, symptom severity, mean ± SE: G1: 63.1 ± 0.84 G2: 59.4 (NR) G3: 56.9 (NR) OAB-q, coping, mean ± SE: G1: 48.9 ± 1.18 G2: 46.5 (NR) G3: 53.1 (NR) OAB-q, concern, mean ± SE : G1: 43.1 ± 1.12 G2: 46.6 (NR) G3: 50.8 (NR) OAB-q, sleep, mean ± SE: G1: 55.2 ± 1.22 G2: 43.8 (NR) G3: 49.2 (NR) OAB-q, social, mean ± SE: G1: 73.6 ± 1.14 G2: 68.9 (NR) G3: 76.0 (NR)	UUI: G1: 28.9 G2: 22.1 G3: 24.5 Frequency: G1: 24.9 G2: 27.0 G3: 28.6 Nocturia: G1: 25.4 G2: 36.4 G3: 29.0 Symptom severity, mean VAS, wk 12 (or early termination): Urinary urgency: G1: 27.5 G2: 25.3 G3: 28.0 UUI: G1: 24.3 G2: 20.0 G3: 22.7 Frequency: G1: 21.6 G2: 24.5 G3: 27.4 Nocturia: G1: 22.0 G2: 21.4 G3: 27.2 Symptom severity, VAS, mean change from baseline (95% CI): Urinary urgency*: G1: -43.1 (-45.8, -40.4) G2: -43.0 (-47.6, -38.5) G3: -39.5 (-41.0, -38.1)	

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Garely et al., 2007*			OAB-q, HRQoL, mean ± SE:	UUI*: G1: -51.7 (-54.5, -49.0)	
Mallett et al., 2007^ (continued)			G1: 53.4 ± 1.01 G2: 50.5 (NR) G3: 56.3 (NR)	G2: -42.3 (-47.8, -36.8) G3: -40.1 (-41.8, -38.4) Frequency*: G1: -42.0 (-45.0, -39.0) G2: -44.2 (-48.8, -39.6) G3: -41.8 (-43.3, -40.3) Nocturia*: G1: -34.4 (-37.3, -31.5) G2: -42.6 (-47.5, -37.8) G3: -36.9 (-38.4, -35.4) OAB-q, symptom severity, mean change (± SE or 95% CI)*: G1: -35.9 ± 1.08 G2: -33.6 (-37.0, -30.3) G3: -29.6 (-30.7, -28.6) OAB-q, coping, mean change (± SE or 95% CI)*: G1: 32.5 ± 1.16 G2: 32.9 (29.3, 36.6) G3: 27.4 (26.2, 28.5) OAB-q, concern, mean change (± SE or 95% CI)*: G1: 37.0 ± 1.21 G2: 34.0 (30.4, 37.6) G3: 29.6 (28.4, 30.8) OAB-q, sleep, mean change (± SE or 95% CI)*: G1: 26.0 ± 1.19 G2: 33.8 (29.7, 37.9) G3: 27.3 (26.1, 28.5)	

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Garely et al., 2007* Mallett et al., 2007^ (continued)				<p>OAB-q, social, mean change \pm SE (95% CI)*: G1: 17.7 \pm 1.03 G2: 20.1 (16.7, 23.5) G3: 14.7 (13.7, 15.6)</p> <p>OAB-q, HRQoL, mean change (\pm SE or 95% CI)*: G1: 29.6 \pm 1.02 G2: 30.8 (27.5, 34.1) G3: 25.4 (24.4, 26.4)</p> <p>Side effects, n (%): G1: 357 (61.3) G2: 128 (46.4) G3: 1321 (59.4)</p> <p>Dry mouth, n (%): G1: 104 (17.9) G2: 36 (13.0) G3: 477 (21.4)</p> <p>Constipation, n (%): G1: 85 (14.6) G2: 19 (6.9) G3: 295 (13.3)</p> <p>Nausea, n (%): G1: NR G2: 7 (2.5) G3: 39 (1.8)</p> <p>Headache, n (%): G1: 21 (3.6) G2: 9 (3.3) G3: 76 (3.4)</p> <p>Blurred vision, n (%): G1: 20 (3.4) G2: 7 (2.5) G3: 57 (2.6)</p>	

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Garely et al., 2007* Mallett et al., 2007^ (continued)				<p>Upper respiratory tract infection, n (%): G1: 27 (4.6) G2: 7 (2.5) G3: 69 (3.10)</p> <p>UTI, n (%) G1: 21 (3.6%) G2: NR G3: NR</p> <p>Rash, n (%): G1: NR G2: 6 (2.2) G3: 22 (0.99)</p> <p>Nasopharyngitis, n (%) G1: 13 (2.2%) G2: NR G3: NR</p> <p>Cough, n (%) G1: 12 (2.1%) G2: NR G3: NR</p> <p>Withdrew, n (%): G1: NR G2: 21 (7.6) G3: 216 (9.7)</p>	

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Giannitsas et al., 2004</p> <p>Country and setting: Greece, Specialty treatment center</p> <p>Enrollment period: NR</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Randomized for which drug to receive first two-way crossover, table of random numbers</p> <p>Intervention: Oxybutynin 15 mg tid vs Tolterodine 4mg bid; 6 weeks treatment with 3-4 weeks washout</p> <p>Groups: G1: Oxybutynin 15 mg tid G2: Tolterodine 4mg bid</p> <p>Stratified by UDS findings: a: high volume (> 250mL); low pressure (< 25cmH₂O) b: high volume (≥ 250mL); high pressure (> 25cmH₂O) C: low volume (< 250mL); low pressure (< 25cmH₂O) d: low volume (< 250mL); high pressure (> 25cmH₂O)</p> <p>N at enrollment: Total: 128</p> <p>N at follow-up: Total: 107 Ga: 6 Gb: 25 Gc: 36 Gd: 40</p> <p>Women, %: 100</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 18 • DO on urodynamics <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Symptomatic or recurrent UTI • BOO • Neurologic disease • History of previous pelvic surgery • Narrow angle glaucoma • SUI • History of anticholinergic side effects • Interstitial cystitis • Child-bearing age without BC 	<p>Voids/day, mean ± SD: Total: 8.5 ± 2.63 Ga: 7.2 ± NR Gb: 8.0 ± 2.40 Gc: 8.3 ± 2.31 Gd: 9.3 ± 2.91</p> <p>Volume (mL)/day, mean ± SD: Total: 1568.5 ± 398.64 Ga: 1756 ± NR Gb: 1594.6 ± 326.12 Gc: 1678.8 ± 402.15 Gd: 1420.5 ± 384.96</p> <p>Voided volume (mL), mean ± SD: Total: 196.1 ± 60.19 Ga: 253 ± NR Gb: 213.5 ± 53.57 Gc: 209.8 ± 57.43 Gd: 163.4 ± 51.97</p> <p>Bladder volume (mL), first void, mean ± SD: Total: 105.6 ± 39.38 Ga: 109 ± NR Gb: 128.0 ± 41.78 Gc: 94.3 ± 35.40 Gd: 101.1 ± 38.14</p> <p>Bladder volume (mL), first contraction, mean ± SD: Total: 172 ± 98.4 Ga: 258 ± NR Gb: 303.3 ± 60.51 Gc: 111.9 ± 48.11 Gd: 124.4 ± 56.66</p>	<p>Voids/day, mean ± SD: G1: 7.7 ± 1.87 G2: 7.6 ± 2.18 G1a: 6.1 ± NR G2a: 6.3 ± NR G1b: 7.3 ± 1.69 G2b: 7.0 ± 1.87 G1c: 7.5 ± 1.41 G2c: 7.2 ± 1.58 G1d: 8.3 ± 2.23 G2d: 8.4 ± 2.53</p> <p>Volume (mL)/day, mean ± SD: G1: 1764.4 ± 333.03 G2: 1670.7 ± 338.6 G1a: 1862 ± NR G2a: 1720 ± NR G1b: 1715.6 ± 292.54 G2b: 1665.8 ± 251.19 G1c: 1847.9 ± 333.81 G2c: 1808.1 ± 317.59 G1d: 1694.7 ± 331.33 G2d: 1550.3 ± 373.65</p> <p>Voided volume (mL), mean ± SD: G1: 239.9 ± 64.98 G2: 236.7 ± 63.03 G1a: 321 ± NR G2a: 286 ± NR G1b: 243.3 ± 59.56 G2b: 248.3 ± 53.91 G1c: 252.9 ± 55.75 G2c: 258.6 ± 60.63 G1d: 217.6 ± 67.07 G2d: 203.1 ± 55.43</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Masking: NA</p> <p>Pt selection criteria: +</p> <p>Loss to followup: ++</p> <p>Drop-out rates: -</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: -</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating	
Giannitsas et al., 2004 (continued)	<p>Age, mean ± SD: Total: 56 ± 16.3 Ga: 53 ± 17.2 Gb: 57 ± 16.2 Gc: 57 ± 16.3 Gd: 54 ± 16.6</p> <p>Weight (kg), mean ± SD: Total: 63 ± 5.6 Ga: 63 ± 5.6 Gb: 70 ± 9.1 Gc: 67 ± 8.8 Gd: 69 ± 7.5</p>		<p>Pressure (cmH₂O), first contraction, mean ± SD: Total: 34.8 ± 21.97 Ga: 17.4 ± NR Gb: 37.7 ± 14.03 Gc: 18.5 ± 4.60 Gd: 50.3 ± 25.14</p> <p>Overactivity index, mean ± SD: Total: 36.8 ± 31.36 Ga: 15.3 ± NR Gb: 24.8 ± 19.66 Gc: 26.3 ± 16.14 Gd: 57.0 ± 38.85</p> <p>Cystometric capacity (mL), mean ± SD: Total: 362.8 ± 119.10 Ga: 403 ± NR Gb: 410.0 ± 97.78 Gc: 357.6 ± 127.52 Gd: 331.4 ± 114.17</p>	<p>Bladder volume (mL), first desire void, mean ± SD: G1: 129.0 ± 30.14 G2: 117.9 ± 27.62 G1a: 144 ± NR G2a: 140 ± NR G1b: 153.5 ± 25.72 G2b: 132.0 ± 31.01 G1c: 120.8 ± 25.07 G2c: 113.2 ± 23.16 G1d: 119.4 ± 29.43 G2d: 110.1 ± 26.15 G1/BL: P < 0.05 G2/BL: P < 0.05</p> <p>Bladder volume (mL), first contraction, mean ± SD: G1: 212.9 ± 106.10 G2: 206.9 ± 103.56 G1a: 382 ± NR G2a: 364 ± NR G1b: 355.28 ± 74.79 G1c: 142.4 ± 43.51 G2c: 144.6 ± 48.43 G1d: 173.3 ± 57.37 G2d: 162.7 ± 53.87 G1d/BL: P < 0.01 G2d/BL: P < 0.01</p> <p>Pressure (cmH₂O), first contraction, mean ± SD: G1: 30.9 ± 22.63 G2: 30.9 ± 19.01 G1a: 14.0 ± NR G2a: 17.4 ± NR G1b: 35.6 ± 12.86 G2b: 29.6 ± 13.56</p>		

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Giannitsas et al., 2004 (continued)				<p>Pressure (cmH₂O), first contraction, mean ± SD: G1c: 17.2 ± 6.80 G2c: 17.9 ± 6.69 G1d: 42.9 ± 28.91 G2d: 44.1 ± 20.75</p> <p>Overactivity index, mean ± SD: G1: 24.4 ± 22.61 G2: 24.7 ± 23.46 G1a: 7.0 ± NR G2a: 9.5 ± NR G1b: 14.1 ± 12.09 G2b: 14.1 ± 12.71 G1c: 18.3 ± 15.89 G2c: 16.9 ± 16.84 G1d: 38.9 ± 26.50 G2d: 40.7 ± 26.58</p> <p>Cystometric capacity (mL), mean ± SD: G1: 419.3 ± 120.86 G2: 415.63 ± 114.06 G1a: 465 ± NR G2a: 453 ± NR G1b: 449.6 ± 106.23 G2b: 459.4 ± 101.17 G1c: 409.9 ± 130.22 G2c: 411.05 ± 132.49 G1d: 401.8 ± 118.34 G2d: 386.7 ± 96.53</p> <p>Dry mouth, n (%): G1: 52 (40.6) G2: 20 (15.6)</p> <p>Constipation, n (%): G1: 11 (10.3) G2: 3 (2.8)</p> <p>Discontinued due to AEs, n: Dry mouth: 12 Palpitations: 1</p>	

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Gleason et al. 1999</p> <p>Country and setting: US, Community</p> <p>Enrollment period: NR 12 week follow up</p> <p>Funding: Alza Corporation</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Multicenter Open label Single treatment</p> <p>Intervention: Oxybutynin ER</p> <p>Groups: G1: Oxybutynin ER 5- 30 mg/ day (dose adjustment period)</p> <p>N at enrollment: G1: 256</p> <p>N at follow-up: G1: 219 37 missing data 20 from AEs 4 dc'd lack of effectiveness, included in analysis</p> <p>Age, yrs ± SD: NR 38.8% were 65+</p> <p>Race/ethnicity, mean (%): white: G1: 235 (91.8)</p> <p>Women, N (%): G1: 234 (91.4)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • ≥ adult men and women • Idiopathic urge incontinence • Mixed incontinence with clinically significant urge component <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Uncontrolled medical condition • PVR > 100 ml • Significant bacteriuria • Significant pyuria 	<p>UII episodes/ week, mean ± SD: G1: 18.8 ± 1.2</p> <p>Total incontinent episodes/ week, mean ± SD: G1: 22.2 ± 1.3</p> <p>Voids/ week, mean ± SD: G1: 81.1 ± 1.8</p> <p>PVR, ml, mean ± SD: G1: 17.4 ± 1.3</p> <p>Voided volume, ml, ± SD: G1: 121 ± 79.1</p>	<p>UII episodes/ week, mean ± SD: G1: 2.8 ± 0.6 P<0.001</p> <p>Total incontinent episodes/ week, mean ± SD: G1: 4.0 ± 0.7 P<0.001</p> <p>Voids/ week, mean ± SD: G1: 66.8 ± 1.4</p> <p>Reduction Voids/week, mean ± SD: G1: 14.3 ± 1.3 95%CI: 11.7-16.8</p> <p>PVR, ml, mean ± SD: G1: 21.0 ± 2.2 P = NR</p> <p>Voided volume, ml, ± SD: G1: 113.4 ± 84.6 P = NR</p> <p>Participants free UII episodes, %: 55.7</p> <p>Dry Mouth, n (%): G1: 128 (58.6)</p> <p>Discontinuation d/t AE, n (%): 20 (7.8)</p> <p>Nausea (%): 2.3</p> <p>Dry Mouth (%): 1.6</p> <p>Somnolence (%): 1.2</p> <p>Urinary retention, n (%): 2 (0.8)</p> <p>Increased PVR, n, (%): 1 (0.4)</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: NA</p> <p>Masking: NA</p> <p>Pt selection criteria: +</p> <p>Loss to followup: +</p> <p>Drop-out rates: ++</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Hill et al., 2007 [See evidence table for Haab et al., 2006]</p> <p>Country and setting: France, Denmark, Sweden, Australia, Canada, US, Multicenter</p> <p>Enrollment period: NR</p> <p>Funding: Novartis Pharma AG (educational grant)</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: 2-yr, non-comparative, open-label extension study</p> <p>Intervention: Darifenacin following 2 feeder studies of darifenacin 3.75/7.5/15 mg qd or placebo x 12 wks</p> <p>Groups: NA</p> <p>N at enrollment: 214</p> <p>N at follow-up (%): 137 (64.0)</p> <p>Patient years of exposure: 308</p> <p>Women, n (%): 172 (80.4)</p> <p>Age, n (%): 65-74 years: 166 (77.6) 75-89 years: 48 (22.4)</p> <p>Race/ethnicity: NR</p> <p>Menopausal, n (%): 13 (6.1)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 65 • Successful completion of one of 2 previous, 12-wk feeder studies with no major protocol violation <p>Exclusion criteria: NR</p>	<p>Urgency episodes/day, median: 8.1</p> <p>Urgency severity, median VAS score: 51.8</p> <p>Incontinence episodes/week, median: 18.7</p> <p>Voids/day, median: 10.1</p> <p>Voided volume (mL), median: 156</p> <p>Concomitant drug treatment, n (%): 210 (98.1)</p>	<p>Urgency episodes/day, median change (median % change): -3.7 (-52.0) P < 0.05</p> <p>Urgency severity, VAS score, median change (median % change): -12.6 (-23.3) P < 0.05</p> <p>Incontinence episodes/week, median change (median % change): -11.0 (-83.7) P < 0.05</p> <p>Significant leaks/week, median change (median % change): -4.9 (-100.0) P < 0.05</p> <p>Incontinence episodes, responders, % reduction, %: ≥ 50%: 74.1 ≥ 70%: 60.0 ≥ 90%: 44.4</p> <p>Voids/day, median change (median % change): -1.2 (-12.4) P < 0.05</p> <p>Normalization of voiding frequency (< 8 voids/day) after darifenacin treatment, %: 3 months: 34 2 years: 33.8</p>	<p>Quality: Overall quality score: fair INTERNAL VALIDITY: poor</p> <p>Randomization: NA</p> <p>Masking: NA</p> <p>Pt selection criteria: +</p> <p>Loss to followup: -</p> <p>Drop-out rates: -</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: +</p> <p>Length of followup: ++</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Hill et al., 2007 (continued)				<p>Nocturia episodes/week, median change (median % change): -1.4 (-10.9) G1/BL: P < 0.05</p> <p>Increased dose to 15 mg at 2 wks and maintained, n (%): 110 (51.4)</p> <p>Remained on 7.5 mg dose, n (%): 55 (25.7)</p> <p>Had dose adjusted at other times, n (%): 49 (22.9)</p> <p>Compliance, ≥ 80% of doses, n (%): 214 (84)</p> <p>Voided volume (mL), median change (median % change): 11.1 (6.3)</p> <p>Dry mouth, n (%): 50 (23.4)</p> <p>Constipation, n (%): 48 (22.4)</p> <p>CVD, %: 1.4</p> <p>Peripheral/CNS, %: 3.3</p>	

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Jacquetin and Wyndaele, 2001</p> <p>Country and setting: France (22 centers) and Belgium (10 centers)</p> <p>Enrollment period: NR</p> <p>Funding: Pharmacia</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Double-blind, placebo-controlled, parallel-group, multicenter phase III study w/ 2-wk washout period</p> <p>Intervention: Tolterodine vs placebo for 4 weeks</p> <p>Groups: G1: Tolterodine 2mg bid G2: Tolterodine 1 mg bid G3: placebo (frequency NR)</p> <p>N at enrollment: NR</p> <p>N at follow-up: G1: 103 G2: 97 G3: 51</p> <p>Women, n (%): G1: 84 (82) G2: 74 (76) G3: 41 (80)</p> <p>Age, mean (range): G1: 58 (21-88) G2: 53 (18-85) G3: 56 (19-89)</p> <p>Race/ethnicity: NR</p> <p>BMI, kg/m² (range): G1: 26.4 (17.7-39.7) G2: 25.5 (16.7-46.3) G3: 24.8 (17.6-36.9)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 18 • UDS-verified detrusor overactivity • ≥ 1 UUI episode/ day • ≥ 8 voids/day <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • SI • Hepatic or renal dz • +UTI or recurrent UTIs • Interstitial cystitis • Hematuria • Clinically significant voiding difficulty • Pts receiving bladder training, electrostimulation therapy • Indwelling catheter • Intermittent cath • Pregnant or nursing • Women of childbearing age w/o reliable contraception 	<p>UUI, n (%) G1: 75 (73) G2: 75 (77) G3: 39 (76)</p> <p>UUI episodes/day, mean (range): G1: 3.2 (0.1-24.0) G2: 2.7 (0.1-24.0) G3: 2.4 (0.1-8.4)</p> <p>Voids/day, mean (range): G1: 10.8 (6.2-34.7) G2: 10.7 (4.9-26.4) G3: 11.7 (6.3-26.3)</p> <p>≥ 8 voids/day, n (%): G1: 96 (93) G2: 89 (92) G3: 49 (96)</p> <p>Urinary symptoms > 5 years, n (%): G1: 46 (45) G2: 42 (43) G3: 17 (33)</p> <p>Previous lower urinary tract surgery, n (%): G1: 36 (35) G2: 29 (30) G3: 16 (31)</p> <p>Previous drug therapy for OAB, n (%): G1: 66 (64) G2: 59 (61) G3: 36 (71)</p> <p>Previous poor efficacy response, n (%): G1: 50 (76) G2: 45 (76) G3: 26 (72)</p>	<p>UUI episodes/day, mean change ± SD: G1: -1.3 ± 1.8 G2: -1.1 ± 2.2 G3: -0.4 ± 1.9 G1/G3: P = 0.0089 G2/G3: P = 0.045</p> <p>Voids/day, mean change ± SD: G1: -1.4 ± 4.3 G2: -1.4 ± 2.8 G3: -1.2 ± 2.7 G1/G3: P = NS G2/G3: P = NS</p> <p>Good efficacy response in current study, previous poor efficacy response, n/N (%) G1: 20/39 (51) G2: 18/37 (49) G3: 7/19 (37) G1/G3: P = NS G2/G3: P = NS</p> <p>Voided volume (mL), mean change ± SD: G1: 19 ± 46 G2: 20 ± 42 G3: 7 ± 40 G1/G3: P = 0.056 G2/G3: P = 0.055</p> <p>Dry mouth, n (%): G1: 35 (34) G2: 20 (21) G3: 3 (6) G1/G2: P < 0.05 G1/G3: P < 0.05 G2/G3: P < 0.05</p> <p>Abdominal pain, n (%): G1: 4 (4) G2: 6 (6) G3: 2 (4)</p> <p>Constipation, n (%): G1: 2 (2) G2: 4 (4) G3: 2 (4)</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: fair</p> <p>Randomization: -</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: NR</p> <p>Drop-out rates: ++</p> <p>Power calculation: +</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: -</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Jacquetin and Wyndaele, 2001 (continued)			Voided volume (mL), mean (range): G1: 158 (43-382) G2: 150 (46-320) G3: 148 (23-284)	Headache, n (%): G1: 3 (3) G2: 3 (3) G3: 2 (4) Total AEs reported, n: G1: 84 G2: 78 G3: 26 Any AE, n (%): G1: 55 (53) G2: 39 (40) G3: 16 (31) G1/G3: P < 0.05 Discontinued due to AEs, n (%):* G1: 2 (2) G2: 3 (3) G3: 1 (2)	

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Koonings et al., 1991</p> <p>Country and setting: US</p> <p>Enrollment period: January 1986 to October 1987</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Cohort</p> <p>Intervention: Oxybutynin chloride 5 mg tid x 4 wks</p> <p>Groups: NA</p> <p>N at enrollment: 126</p> <p>N at follow-up: 114</p> <p>Women, %: 100</p> <p>Age, mean (range): 39 (21-74)</p> <p>Race/ethnicity: NR</p> <p>Menopausal, n: 44</p> <p>Parity, mean (range): 2 (0-12)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> Confirmed diagnosis of detrusor instability Uninhibited detrusor contraction > 15 cm H₂O on standing provocative urethra cystometry <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Urethritis (on urethroscopy) Cystitis (on cystoscopy) Neurologic findings on screening test of S2-S4 lower voiding center MUI Glaucoma 	<p>Uninhibited detrusor contraction starting prior to urethral pressure change, n (%): 73 (64%)</p> <p>Urethral pressure drop (≥ 20 cmH₂O) prior to detrusor contraction, n (%): 41 (6)</p>	<p>Good response, n (%): 66 (58)</p> <p>Poor response, n (%): 48 (42)</p> <p>Response, women with uninhibited detrusor contraction starting prior to urethral pressure, n (%): Good: 61 (81) Poor: 5 (12) P < 0.01</p> <p>Response, women with urethral pressure drop (≥ 20 cmH₂O) prior to detrusor contraction, n (%): Good: 12 (16) Poor: 36 (88) P < 0.01</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: NA</p> <p>Masking: NA</p> <p>Pt selection criteria: +</p> <p>Loss to followup: ++</p> <p>Drop-out rates: NR</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: -</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Kreder et al., 2003</p> <p>Country and setting: US and Canada, Community</p> <p>Enrollment period: NR 16 week follow up</p> <p>Funding: Pharmacia Corporation</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Subgroup analysis of multicenter, single blind cohort</p> <p>Intervention: Tolterodine 1-2 mg bid</p> <p>Groups: G1: MUI with primary urge G2: UUI</p> <p>N at enrollment: G1: 171 G2: 552</p> <p>N at follow-up: G1: 170 G2: 551</p> <p>Women, n (%): G1: 165 (96.5) G2: 464 (84)</p> <p>Age, median (range): G1: 62 (21, 88) G2: 65 (20, 88)</p> <p>Race/ethnicity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 65 • History, PE, UDS consistent with urge incontinence • ≥ 4 episodes UI on 5 day voiding diary • ≥ 8 voids/day • Either urgency or ≥ 1 UI episodes / day <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • MUI with predominate stress component • Contraindications to antimuscarinic therapy • Hepatic/renal disease • Symptomatic or recurrent UTI • Hematuria • IC • Voiding difficulty with risk of retention • Concurrent bladder training • Electrostimulation therapy • Indwelling catheter • CIC • Women with reproduction potential • Pregnant/nursing • Concomitant anticholinergic meds/ treatment 	<p>Incontinence episodes/day, median (range): G1: 3 (1, 19) G2: 3 (1, 15)</p> <p>≥ 2 nocturia episodes/day, n (%): G1: 100 (58.5) G2: 334 (60.5)</p> <p>Using pads, n (%): G1: 122 (71) G2: 342 (62)</p> <p>Voided volume (mL), median (range): G1: 169 (62, 506) G2: 164 (31, 524)</p> <p>Duration of symptoms > 5 years, n (%): G1: 91 (53) G2: 259 (47)</p> <p>Previous drug therapy for OAB, n (%): G1: 85 (50) G2: 275 (50)</p>	<p>Incontinence episodes/day, median % change: G1: -67 G2: -75 P = NS</p> <p>Cure rate, dryness, n (%): G1: 66 (39) G2: 243 (44) P = NS</p> <p>voids/day, median % change: G1: -15 G2: -17 P = NS</p> <p>≤ 8 voids/day dryness, n (%): G1: 40 (23.5) G2: 130 (24) P = NS</p> <p>Nocturia episodes/day, median % change: G1: -50 G2: -33 P = NS</p> <p>≤ 2 nocturnal voiding episodes/day, n (%): G1: 83 (83) G2: 254 (76) P = NS</p> <p>Achieved no pad usage, n (%): G1: 26 (21) G2: 93 (27) P = NS</p> <p>Voided volume (mL), median change (range): G1: 26.5 (-261, 195) G2: 27 (-345, 389) G1/BL: P < 0.001 G2/BL: P < 0.001</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: NA</p> <p>Masking: NA</p> <p>Pt selection criteria: +</p> <p>Loss to followup: -</p> <p>Drop-out rates: NR</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Landis et al., 2004</p> <p>[See evidence table for van Kerrebroeck et al., 2001]</p> <p>Country and setting: Multinational, Community</p> <p>Enrollment period: NR 12 weeks</p> <p>Funding: Pharmacia</p> <p>Author industry relationship disclosures: 2 of 4 ALZA (1) Bristol-Myers Squibb (1) Pharmacia (2)</p>	<p>Design: RCT</p> <p>Intervention: Tolterodine ER vs placebo 1 week run-in period with 12 weeks treatment</p> <p>Groups: G1: Tolterodine ER 4 mg daily G2: placebo</p> <p>Stratified: a: 5-20 UUI episodes/week b: ≥ 21 UUI episodes/week</p> <p>N at enrollment: G1: 492 G1a: 321 G1b: 171 G2: 494 G2a: 284 G2b: 210</p> <p>N at follow-up: G1a: 321 G1b: 171 G2a: 284 G2b: 210</p> <p>Women, N (%): G1a: 262 (81.6) G1b: 143 (83.6) G2a: 223 (78.5) G2b: 178 (84.8)</p> <p>Age, mean ± SD: G1a: 60.86 ± 14.45 G1b: 60.04 ± 14.10 G2a: 60.61 ± 13.59 G2b: 61.84 ± 13.85</p> <p>Race/ethnicity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 18 • ≥ 5 UUI episodes/ week • ≥ 8 voids/day • Symptoms of OAB x 5 months <p>Exclusion criteria: NR</p>	<p>UUI episodes/ week, median (range): G1a: 11 (5-20) G1b: 34 (21-168) G2a: 11 (5-20) G2b: 31.5 (21-168)</p> <p>Voids/day, median (range): G1a: 9.57 (2.57-34.85) G1b: 10.71 (2.28-51.28) G2a: 10.28 (2.28-28.42) G2b: 11.14 (2-37.42)</p> <p>Voided volume (mL), median (range): G1a: 142.72 (36.19-338) G1b: 135.12 (51.45-270.52) G2a: 136.25 (32.51-237.5) G2b: 126.94 (21.16-373.46)</p> <p>Number prior Tx for OAB, n (%): G1a: 162 (50.6) G1b: 100 (58.5) G2a: 139 (49.1) G2b: 117 (55.7)</p> <p>Number prior Tx for OAB w/ good efficacy, n (%): G1a: 85 (52.8) G1b: 63 (63.0) G2a: 84 (60.4) G2b: 68 (58.6)</p>	<p>UUI episodes/ week, median % change: G1a: -71.42 G1b: -67.56 G2a: -38.46 G2b: -29.81 G1a/G2a: P = 0.0264 G1b/G2b: P = 0.0221</p> <p>Voids/day, median change: G1a: -1.22 G1b: -1.9 G2a: -0.85 G2b: -0.4 G1a/G2a: P < 0.04 G1b/G2b: P < 0.02</p> <p>Voided volume (mL), median change: G1a: 24.0 G1b: 27.0 G2a: 4.0 G2b: 2.9 G1a/G2a: P < 0.0001 G1a/G2a: P < 0.0001</p>	<p>Quality: Overall quality score: fair INTERNAL VALIDITY: poor Randomization: + Masking: + Pt selection criteria: + Loss to followup: + Drop-out rates: - Power calculation: - Statistical issues: + EXTERNAL VALIDITY: good Age: + Baseline OAB status: + Baseline characteristics: ++ Length of followup: + Measurement methods: + Measurement reliability: + Intervention description: +</p>

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Layton et al. 2001</p> <p>Country and setting: England; Community</p> <p>Enrollment period: November 1998 to May 1999</p> <p>Funding: Pharmacia Upjohn</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Prospective observational cohort</p> <p>Intervention: Tolterodine prescription event monitoring in UK</p> <p>Groups: NA</p> <p>N at enrollment: 35,295 had commenced treatment</p> <p>26,991 green forms mailed out</p> <p>14,526 returned forms</p> <p>Response rate: 53.8%</p> <p>Age, yrs ± SD: 62.7 ± 16.4</p> <p>Women, N (%): 9965 (68.6)</p> <p>Parity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> Patients prescribed tolterodine in general practice in UK <p>Exclusion criteria:</p> <ul style="list-style-type: none"> None 		<p>Dry mouth, n: 250</p> <p>Unspecified adverse effects, n: 168</p> <p>Headache, n: 123</p> <p>Constipation, n: 78</p> <p>General malaise, n: 78</p> <p>Hallucinations, n: 23</p> <p>Palpitations/tachycardia, n: 42</p> <p>Other cardiac arrhythmias, n: 29</p> <p>Chest pains, n: 87</p>	<p>Quality: Overall quality score: good</p> <p>INTERNAL VALIDITY: good</p> <p>Randomization: NA</p> <p>Masking: NA</p> <p>Pt selection criteria: +</p> <p>Loss to followup: NA</p> <p>Drop-out rates: NA</p> <p>Power calculation: +</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: NR</p> <p>Baseline characteristics: +</p> <p>Length of followup: ++</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: -</p>

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Lee et al., 2002</p> <p>Country and setting: South Korea, University</p> <p>Enrollment period: NR</p> <p>Funding: Pharmacia Corp</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: Tolterodine 2mg bid vs Oxybutynin 5mg bid</p> <p>Groups: G1: Tolterodine 2mg bid G2: Oxybutynin 5mg bid</p> <p>N at enrollment: G1: 112 G2: 116</p> <p>N at follow-up: G1: 97 G2: 90</p> <p>Women, n (%): G1: 84 (74) G2: 92 (79)</p> <p>Age, mean (range): G1: 52 (27, 82) G2: 52 (20, 86)</p> <p>Race/ethnicity (%): Asian: G1: 100 G2: 100</p> <p>BMI, kg/m² (range): G1: 23 (17, 32.5) G2: 23.5 (16, 38)</p> <p>Previous drug therapy: N (%) G1: 36 (32) G2: 26 (22)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 18 • OAB symptoms > 6 mos • ≥ 8 voids/day, with or without incontinence (measured by diary) <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • SUI • Women not using reliable contraception • Pregnant or nursing • Prior treatment with anticholinergic < 2 wks • Renal or hepatic disease • Narrow angle glaucoma • Urinary retention • Gastric retention • Hypersensitivity to drugs • UTI • IC • Hematuria • BOO • Concomitant bladder training, e-stim treatment • Indwelling catheter • CIC • Concomitant treatment for OAB ≤ 2 mos 	<p>Incontinence episodes/day, mean (range): G1: 2.6 (0.3, 9.3) G2: 2.4 (3.0, 14.7)</p> <p>Patients with incontinence episodes, n (%): G1: 46 (41) G2: 42 (36)</p> <p>Voids/day, mean (range): G1: 12.2 (8.0, 23.7) G2: 12.4 (7.7, 29.7)</p>	<p>Incontinence episodes/day, mean change ± SD (% change): G1: -2.2 ± 2.3 (-85) G2: -1.4 ± 1.8 (-58)</p> <p>G1/BL: P = 0.0001 G1/BL: P = 0.0001 G1/G2: P = 0.10</p> <p>Voids/ day, mean change ± SD (% change): G1: -2.6 ± 2.9 (-21) G2: -1.8 ± 4.2 (-15)</p> <p>G1/BL: P = 0.0001 G1/BL: P = 0.0001 G1/G2: P = 0.14</p> <p>Benefit of Tx, %: G1: 45 G2: 46 G1/G2: P = NS</p> <p>Patients reporting adverse events, n (%): G1: 62 (55) G2: 94 (82)</p> <p>Discontinued due to AEs, n (%): G1: 11 G2: 18</p> <p>Dry mouth, n (%): G1: 39 (35) G2: 72 (63) G1/G2: P = 0.001</p> <p>Dry mouth, mild, n (%): G1: 29 (26) G2: 40 (35)</p> <p>Dry mouth, moderate, n (%): G1: 9 (8) G2: 26 (23)</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: -</p> <p>Drop-out rates: -</p> <p>Power calculation: +</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: -</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Lee et al., 2002 (continued)				Dry mouth, severe, n (%): G1: 1 (1) G2: 6 (5) Voiding disorder, n (%): G1: 10 (9) G2: 16 (14)	

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Malone-Lee et al., 2003</p> <p>Country and setting: UK, Hospital</p> <p>Enrollment period: 1993-1999</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Prospective observational</p> <p>Intervention: Oxybutynin 2.5mg bid and bladder retraining</p> <p>Groups: G1: urinary frequency and urgency w/ detrusor instability G2: urinary frequency and urgency w/o detrusor instability</p> <p>N at enrollment: G1: 266 G2: 86</p> <p>N at follow-up: Total: 347 G1: NR G2: NR</p> <p>Age, mean (range): G1: 54.8 (20, 90) G2: 51.8 (21, 88)</p> <p>Race/ethnicity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Women • Age ≥ 18 • ≥ 8 voids/day <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Neurological disease • Significant stress incontinence • Symptomatic UTI • Interstitial cystitis 	<p>Incontinence episodes/day, median (95% CI): G1: 2 (0, 7) G2: 2 (0, 6.3)</p> <p>Voids/day median (95% CI): G1: 14 (8, 24) G2: 12 (8, 22)</p>	<p>Incontinence episodes/day, median change (95% CI): G1: 0 (2, 6) G2: 0 (2, 6) G1/G2: P = 0.73</p> <p>Voids/day, median change (95% CI): G1: -5 (1.9-14) G2: -5 (1-12.3) G1/G2: P = 0.61</p> <p>Dry mouth, n (%): G1: 219 (84) G2: 69 (70)</p> <p>Constipation, n (%): G1: 84 (32) G2: 19 (22)</p> <p>Heartburn, n (%): G1: 71 (27) G2: 20 (23)</p> <p>Dry skin, n (%): G1: 46 (18) G2: 4 (5) G1/G2: P = 0.01</p> <p>Headache, n (%): G1: 25 (10) G2: 3 (3.5)</p> <p>Dry eyes, n (%): G1: 12 (5) G2: 1 (1)</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: fair</p> <p>Randomization: NA</p> <p>Masking: NA</p> <p>Pt selection criteria: +</p> <p>Loss to followup: ++</p> <p>Drop-out rates: NR</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: +</p> <p>Length of followup: -</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Michel et al., 2002</p> <p>Country and setting: Germany, Community</p> <p>Enrollment period: NR 12 week follow up</p> <p>Funding: Pharmacia GmbH</p> <p>Author industry relationship disclosures: 1 of 4 Pharmacia(1) Sanofi-Synthelabo(1)</p>	<p>Design: Open-label observational post-marketing surveillance</p> <p>Intervention: Tolterodine</p> <p>Groups: Tolterodine Median dose: 2 mg daily</p> <p>Dose (mg), mean ± SD: 3.81 ± 1.16</p> <p>N at enrollment: 2,250</p> <p>N at follow-up: 1,979</p> <p>Women, n (%): 1,730 (76.9)</p> <p>Age, mean ± SD: 61.1 ± 13.8</p> <p>Race/ethnicity: NR</p> <p>BMI, kg/m² ± SD: 73.2 ± 11.3</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> Physician medical judgement <p>Exclusion criteria: NR</p>	<p>Urgency episodes/day, mean ± SD: 8.4 ± 5.1</p> <p>Incontinence episodes/day, mean ± SD: 3.4 ± 4.2</p> <p>Voids/day, mean ± SD: 12.4 ± 4.3</p>	<p>Urgency episodes/day, mean ± SD: 2.0 ± 3.0 P = NR</p> <p>Incontinence episodes/day, mean ± SD: 0.8 ± 2.0 P = NR</p> <p>Voids/day, mean ± SD: 7.7 ± 2.7 P = NR</p> <p>Urgency, successful treatment, OR (95% CI): Gender, M/F: 0.764 (0.583, 1.001) P = 0.0508</p> <p>Age, years: 0.981 (0.973, 0.990) P < 0.001</p> <p>Frequency, BL episodes/day: 1.038 (0.997- 1.080) P = 0.0724</p> <p>Urgency, BL episodes/day: 0.851 (0.823, 0.880) P < 0.001</p> <p>Incontinence, BL episodes/day: 0.979 (0.950, 1.000) P = 0.1735</p> <p>Tolterodine dose, mg/day: 0.913 (0.830, 1.005) P = 0.0623</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: NA</p> <p>Masking: NA</p> <p>Pt selection criteria: -</p> <p>Loss to followup: +</p> <p>Drop-out rates: -</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Michel et al., 2002 (continued)				<p>Incontinence, successful treatment, OR (95% CI): Gender, M/F: 1.453 (1.062, 1.990) P = 0.0196 Age, years: 0.978 (0.968, 0.987) P < 0.001 Frequency, BL episodes/day: 1.034 (0.993, 1.076) P = 0.1036 Urgency, BL episodes/day: 1.053 (1.018, 1.088) P = 0.0027 Incontinence, BL baseline episodes/day: 0.744 (0.716, 0.774) P < 0.001 Tolterodine dose, mg/day: 0.866 (0.784, 0.956) P = 0.0043</p> <p>Frequency, successful treatment, OR (95% CI): Gender, M/F: 0.745 (0.552, 1.004) P = 0.0532 Age, years: 0.981 (0.971, 0.991) P = 0.001 Frequency, BL episodes/day: 0.735 (0.701, 0.771) P < 0.001 Urgency, BL episodes/day: 1.008 (0.975, 1.041) P = 0.6526</p>	

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Michel et al., 2002 (continued)				<p>Incontinence, BL episodes/day: 0.969 (0.937, 1.002) P = 0.0644</p> <p>Tolterodine dose, mg/day: 1.070 (0.957, 1.196) P = 0.2335</p> <p>Effect on Global efficacy, OR (95% CI):</p> <p>Gender, M/F: 0.656 (0.526, 0.818) P = 0.0002</p> <p>Age, years: 0.986 (0.980, 0.993) P < 0.001</p> <p>Frequency, BL epsidoes/day: 1.002 (0.972, 1.033) P = 0.8896</p> <p>Urgency, BL episodes/day: 1.009 (0.984, 1.033) P = 0.4906</p> <p>Incontinence, BL episodes/day: 0.963 (0.939, 0.987) P < 0.0026</p> <p>Tolterodine dose, mg/day: 1.000 (0.926, 1.080) P = 0.9971</p> <p>Effect on global tolerability of Tolterodine, OR (95% CI):</p> <p>Gender, M/F: 0.993 (0.790, 1.249) P = 0.9519</p> <p>Age, years: 0.995 (0.988, 1.002) P = 0.1915</p> <p>Frequency, BL episodes/day: 1.002 (0.971, 1.034) P = 0.8995</p>	

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Michel et al., 2002 (continued)				Urgency, BL episodes/day: 1.022 (0.997, 1.048) P = 0.0889	
				Incontinence, BL episodes/day: 0.990 (0.965, 1.016) P < 0.459	
				Tolterodine dose, mg/day: 1.114 (1.028, 1.206) P = 0.0085	

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Michel et al., 2007*</p> <p>Michel et al., 2005</p> <p>Country and setting: Germany, NR</p> <p>Enrollment period: November 2001 to June 2003</p> <p>Funding: Pharmacia (became Pfizer before publication)</p> <p>Author industry relationship disclosures: 5/5* 4SC (1) Astellas (1) Boehringer Ingelheim (2) Eli Lilly (2) Pfizer (4) Theravance (1) Anformed (1)</p>	<p>Design: Retrospective cohort, open label</p> <p>Intervention: Tolterodine ER</p> <p>Groups: G1: Incontinent G2: Continent</p> <p>N at enrollment: Total: 3,824 G1: 2,571 G2: 1,253</p> <p>N at follow-up: Total: 3,416</p> <p>Women, %: Total: 75.8 G1: 81.7 G2: 62.6 P < 0.001</p> <p>Age, mean ± SD: Total: 64.8 ± 13.3 G1: 66.3 ± 12.6 G2: 61.4 ± 14.1 P < 0.001</p> <p>Race/ethnicity: NR</p> <p>Follow-up: 9 months</p>	<p>Inclusion criteria: NR</p> <p>Exclusion criteria: NR</p>	<p>Urgency episodes/day, mean ± SD: Total: 7.9 ± 5.2 G1: 8.0 ± 5.2 G2: 7.7 ± 5.2 P = 0.1937</p> <p>Incontinence episodes/day, mean ± SD: G1: 4.8 ± 3.7 G2: NA</p> <p>Voids/day, mean ± SD: Total: 14.0 ± 4.6 G1: 14.1 ± 4.6 G2: 13.5 ± 4.4 P < 0.001</p> <p>Daytime voids/day, mean ± SD: Total: 10.7 ± 3.7 G1: 10.8 ± 3.7 G2: 10.4 ± 3.4 P = 0.0128</p> <p>Frequency ≥ 8 voids/day, %: G1: 95.7 G2: 95.2</p> <p>Nocturia episodes/day, mean ± SD: Total: 3.4 ± 1.7 G1: 3.5 ± 1.8 G2: 3.2 ± 1.6 P < 0.001</p> <p>Pad use/day, mean ± SD: G1: 3.4 ± 2.8 G2: 0.1 ± 0.6 P < 0.001</p> <p>Duration of symptoms (mo), mean ± SD: G1: 50 ± 53 G2: 40 ± 46 P < 0.001</p> <p>Previous treatment, %: G1: 50.8 G2: 44.8 P = 0.0007</p>	<p>Incontinence episodes/day, mean change ± SD: G1: -3.8 ± 3.5 G2: NA</p> <p>Urgency episodes/day, mean ± SD: Total: 1.6 ± 2.8</p> <p>No urgency, %: G1: 53.4 G2: 63.2</p> <p>Voids/day, mean ± SD: Total: 7.5 ± 3.0</p> <p>Daytime voids/day, mean ± SD: Total: 6.2 ± 2.3</p> <p>Frequency ≥ 8 voids/day, %: G1: 43.7 G2: 39.1</p> <p>Nocturia episodes/day, mean ± SD: Total: 1.4 ± 1.1</p> <p>Pad use/day, mean change ± SD: G1: -2.4 ± 2.5</p> <p>Limitation of daily activities, score change ± SD: G1: 4.49 ± 2.65 G2: 4.10 ± 2.51</p> <p>Limitations in daily life caused by bladder problems, score ± SD: G1: 7.59 ± 1.65 G2: 6.66 ± 1.69 P < 0.001</p> <p>Total adverse events, n (%): Total: 496 (13.0)</p> <p>Dry mouth, n (%): 299 (7.8)</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: NA</p> <p>Masking: NA</p> <p>Pt selection criteria: -</p> <p>Loss to followup: NR</p> <p>Drop-out rates: NA</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: +</p> <p>Length of followup: ++</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Michel et al., 2007*				Withdrew: 11%	
Michel et al., 2005 (continued)				Unable to tolerate medication: 2.8%	
				Administrative reasons: 2.6%	
				Lack of efficacy : 2.4%	
				Patient request: 1.2%	

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Moore et al., 1990</p> <p>Moore and Sutherst 1990</p> <p>Country and setting: UK, academic health center</p> <p>Enrollment period: NR</p> <p>Funding: Tillots Laboratories</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT with crossover</p> <p>Intervention: oxybutynin hydrochloride vs placebo</p> <p>Groups: G1:oxybutynin hydrochloride 3 mg tid G2:placebo</p> <p>N at enrollment: 53</p> <p>N at follow-up: G1: 27 G2: 22</p> <p>Age, mean ± SD: G1: 46.0 ± 11.7 G2: 46.4 ± 12.4</p> <p>Race/ethnicity: NR</p> <p>Parity, mean: G1: 2.3 G2: 2.2</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • UDS-defined idiopathic detrusor instability • Involuntary detrusor contractions > 30 cm H2O during filling phase of cystometry <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Neurological disorder • Urologic disorder • Age > 75 years • Genuine SI • Low-compliance bladder • Bacterial or interstitial cystitis • Previous treatment with oxybutynin 	<p>Voids/day, mean ± SD: G1: 11.1 ± 7.9 G2: 12.1 ± 5.7</p> <p>Volume at first desire to void mean mL ± SD: G1: 94.2 ± 66 G2: 107.4 ± 101</p> <p>Max detrusor filling pressure mean cm H2O, ±SD: G1: 55.3 ± 23 G2: 59.2 ± 26</p> <p>Max cystometric capacity (mL) G1: 275 ± 164 G2: 290 ± 168</p> <p>Residual urine volume mean mL ± SD: G1: 23.1 ± 34.2 G2: 22.7 ± 34.0</p> <p>Duration of OAB, mean yrs ± SD): G1: 7.5 ± 7.9 G2: 8.9 ± 9.0</p>	<p>Change in volume at first desire to void, period 1, mean mL ± SD (95% CI) G1: 70.0 ± 103 G2: 7.7 ± 76 G1/G2: P = 0.02 95% CI: (10, 113)</p> <p>Change in maximum detrusor filling pressure, mean ± SD: G1:-16.9 ± 19.5 G2:1.0 ± 28.7 G1/G2: P = 0.02 95% CI: (-32.5, -3.2)</p> <p>Change in maximum cystometric capacity, mean ± SD: G1:104.0 ± 131 G2:7.0 ± 103 G1/G2:p=0.006 95% CI: (29, 165)</p> <p>Change in residual urine Volume, period 1: G1: 22.4 (79.3) G2: 7.7 (45.7) P=0.42</p> <p>Change in residual urine Volume, period 2: G1: 10.0 ± 27.1 G2: 25.7 ± 55.6 P = 0.33</p> <p>Dry mouth on oxybutynin, %: 88</p> <p>Dry mouth on placebo, %: 33</p> <p>Mouth ulcers on oxybutynin, %: 16</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: -</p> <p>Drop-out rates: -</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: ++</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Moore et al., 1990 Moore and Sutherst 1990 (continued)				<p>Mouth ulcers on placebo, %: 0</p> <p>Constipation on oxybutynin, % 12.5</p> <p>Constipation on placebo, % 0</p> <p>Drowsiness on oxybutynin, %: 12.5</p> <p>Drowsiness on placebo, %: 7</p> <p>Nausea on oxybutynin, %: 8.3</p> <p>Nausea on placebo, %: 2.3</p> <p>Initial hesitancy on oxybutynin, %: 4.2</p> <p>Initial hesitancy on placebo, %: 2.3</p> <p>Dizziness on oxybutynin, %: 4.2</p> <p>Dizziness on placebo, %: 7.0</p> <p>Metallic taste on oxybutynin, %: 2.4</p> <p>Metallic taste on placebo, %: 2.3</p> <p>Crown Crisp Experimental Index, >50% improvement:*</p> <p>Free floating anxiety: 6.4</p>	

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Moore et al., 1990 Moore and Sutherst 1990 (continued)				Phobic anxiety: 5.3 Obsessionalism: 6.2 Somatic complaints: 6.2 Depression: 4.0 Hysteria: 2.6 Total: 30.7 Crown Crisp Experimental Index, <49% improvement:* Free floating anxiety: 8.8 Phobic anxiety: 6.8 Obsessionalism: 8.1 Somatic complaints: 8.9	

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Rudy, et al. 2006 Rudy, et al. 2006 Staskin and Harnett, 2004</p> <p>Country and setting: US; 52 sites</p> <p>Enrollment period: NR</p> <p>Funding: Indevus</p> <p>Author industry relationship disclosures: 5 of 5 Indevus (5)</p>	<p>Design: RCT</p> <p>Intervention: Trospium chloride vs placebo x 12 wks</p> <p>Groups: G1: trospium chloride 20 mg BID G2: matching placebo</p> <p>N at enrollment: G1: 329 G2: 329</p> <p>N at follow-up: G1: 323 G2: 325</p> <p>Age, mean yrs ± SE: G1: 61.1 ± 0.69 G2: 61.0 ± 0.70</p> <p>Women, N (%): G1: 269 (81.8) G2: 267 (81.2)</p> <p>Race/ethnicity, n (%): Black: G1: 26 (7.9) G2: 21 (6.4) White: G1: 284 (86.3) G2: 300 (91.2) Hispanic: G1: 13 (4.0) G2: 5 (1.5) Asian: G1: 5 (1.5) G2: 3 (0.9)</p> <p>Parity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • 18+ years old • 6+ mos of OAB sx • 10+ voids/d • Sx of urgency • 7+ urge UIE/wk <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Predominantly SUI, insensate UI, or overflow UI • Neurogenic bladder d/o's • Significant renal dz • Uninvestigated hematuria • UTI at washout or more than 2x during the prior 12 mos • PVR>100mL • Use of any anticholinergic drug or other drug therapy for OAB w/in 21 days before randomization • Bladder surgery w/in 6 mos before randomization • Cancer • Interstitial cystitis • PSA>10ng/mL • Diuretic use • Estrogen therapy • Nonmedical bladder therapy not part of a stable, long-term program • Pregnancy • Contraindication to antimuscarinic therapy 	<p>Daily voids, mean n: G1: 12.94 G2: 13.17 P=0.3169</p> <p>Urgency severity score associated with toilet voids, median: G1: 1.79 G2: 1.75 P=0.4100</p> <p>Volume voided in mL per void/24h, mean: G1: 154.80 ± NR G2: 154.64 ± NR P=0.9667</p> <p>Median daily urge incontinence episodes: G1: 2.86 G2: 2.86 P=0.9849</p> <p>Nocturnal voids/24 hr, median: G1: 2.00 G2: 2.00 P=0.9048</p> <p>Nocturnal urgency severity score associated with toilet voids, mean: G1: 2.03 G2: 2.01 P=0.6863</p> <p>OAB-SCS, median: G1: 36.56 G2: 36.88 P=0.7176</p> <p>Stanford Sleepiness Scale, mean: G1: 1.98 G2: 1.94</p>	<p>Change from baseline to day 1: Daily voids, mean: G1: -0.66 G2: -0.35 P=0.14</p> <p>Urgency severity score associated with toilet voids, mean: G1: 0.00 G2: 0.02 P=0.47</p> <p>Median daily urge incontinence episodes: G1: -1.00 G2: -0.57 P=0.042</p> <p>OAB-SCS, median: G1: -2.50 G2: -0.14 P=0.014</p> <p>Change from baseline to day 2: Daily voids, mean: G1: -1.11 G2: -0.75 P=0.09</p> <p>Urgency severity score associated with toilet voids, mean: G1: -0.08 G2: -0.03 P=0.13</p> <p>Median daily UI episodes: G1: -1.29 G2: -0.86 P=0.0022</p> <p>OAB-SCS, median: G1: -3.86 G2: -2.14 P=0.015</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: NA</p> <p>Masking: NA</p> <p>Pt selection criteria: +</p> <p>Loss to followup: -</p> <p>Drop-out rates: NR</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: +</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: -</p> <p>Intervention description: +</p>

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Rudy et al., 2006			Stanford Sleepiness Scale, age group <65 years:	Change from baseline to <u>day 3</u>:	
Rudy et al., 2006			G1: 1.99	Daily voids, mean:	
Staskin and Harnett, 2004			G2: 2.09	G1: -1.30	
(continued)				G2: -0.77	
			Stanford Sleepiness Scale, age group >65 years:	P=0.012	
			G1: 1.96	Urgency severity score associated with toilet voids, mean:	
			G2: 1.71	G1: -0.11	
			Stanford Sleepiness Scale, age group <75 years:	G2: -0.04	
			G1: 2	P=0.015	
			G2: 1.97	Median daily UI episodes:	
			Stanford Sleepiness Scale, age group >75 years:	G1: -1.57	
			G1: 1.86	G2: -0.86	
			G2: 1.79	P<0.0001	
			Prior OAB medications, n (%):	OAB-SCS, median:	
			G1: 162 (49.2)	G1: -4.71	
			G2: 169 (51.4)	G2: -2.71	
			Prior hx of pelvic-floor training, n (%):	P=0.0019	
			G1: 62 (18.8)	Change from baseline to <u>day 4</u>:	
			G2: 76 (23.1)	Daily voids, mean:	
			Currently practice undergarment change d/t incontinence, n (%):	G1: -1.33	
			G1: 227 (69.0)	G2: -0.95	
			G2: 239 (72.6)	P=0.086	
				Urgency severity score associated with toilet voids, mean:	
				G1: -0.12	
				G2: -0.05	
				P=0.031	
				Median daily UI episodes:	
				G1: -1.57	
				G2: -1.00	
				P<0.0001	
				OAB-SCS, median:	
				G1: -4.86	
				G2: -3.36	
				P=0.011	

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Rudy et al., 2006 Rudy et al., 2006 Staskin and Harnett, 2004 (continued)				<p>Change from baseline to <u>day 5</u>:</p> <p>Daily voids, mean: G1: -1.73 G2: -1.12 P=0.0037</p> <p>Urgency severity score associated with toilet voids, mean: G1: -0.13 G2: -0.05 P=0.021</p> <p>Median daily UI episodes: G1: -1.57 G2: -1.00 P<0.0001</p> <p>OAB-SCS, median: G1: -6.43 G2: -3.36 P<0.0001</p> <p>Change from baseline to <u>day 6</u>:</p> <p>Daily voids, mean: G1: -1.80 G2: -1.28 P=0.017</p> <p>Urgency severity score associated with toilet voids, mean: G1: -0.08 G2: 0.04 P<0.0001</p> <p>Median daily UI episodes: G1: -1.71 G2: -1.00 P<0.0001</p> <p>OAB-SCS, median: G1: -5.43 G2: -2.93 P<0.0001</p>	

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Rudy et al., 2006 Rudy et al., 2006 Staskin and Harnett, 2004 (continued)				<p>Change from baseline to <u>week 1</u>:</p> <p>Daily voids, mean: G1: -1.42 G2: -0.96 P=0.0039</p> <p>Urgency severity score associated with toilet voids, mean: G1: -0.09 G2: -0.01 P=0.0023</p> <p>Volume voided in mL per void/24h: G1: 29.23 G2: 6.05 P<0.0001</p> <p>Median daily urge incontinence episodes: G1: -1.43 G2: -0.86 P<0.0001</p> <p>Nocturnal voids/24 hr, median: G1: -0.29 G2: -0.29 P=0.8454</p> <p>Nocturnal urgency severity score associated with toilet voids, mean: G1: -0.05 G2: 0.03 P=0.0442</p> <p>OAB-SCS, median: G1: -4.71 G2: -2.29 P<0.0001</p>	

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Rudy et al., 2006 Rudy et al., 2006 Staskin and Harnett, 2004 (continued)				<p>Change from baseline to <u>4 weeks</u>:</p> <p>Daily voids, mean: G1: -2.34 G2: -1.55 P<0.0001</p> <p>Urgency severity score associated with toilet voids, mean: G1: -0.19 G2: -0.04 P<0.0001</p> <p>Volume voided in mL per void/24h: G1: 39.50 G2: 9.45 P<0.0001</p> <p>Median daily urge incontinence episodes: G1: -1.71 G2: -1.14 P<0.0001</p> <p>Nocturnal voids/24 hr, median: G1: -0.43 G2: -0.29 P=0.0299</p> <p>Nocturnal urgency severity score associated with toilet voids, mean: G1: -0.13 G2: =0.01 P=0.0062</p> <p>OAB-SCS, median: G1: -8.14 G2: -3.86 P<0.0001</p>	

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Rudy et al., 2006 Rudy et al., 2006 Staskin and Harnett, 2004 (continued)				<p>Change from baseline to <u>12 weeks</u>:</p> <p>Daily voids, mean: G1: -2.67 G2: -1.76 P<0.0001</p> <p>Urgency severity score associated with toilet voids: G1: -0.21 G2: -0.02 P<0.0001</p> <p>Volume voided in mL per void/24h: G1: 35.59 G2: 9.44 P<0.0001</p> <p>Daily UI episodes, median: G1: -1.86 G2: -1.29 P<0.0001</p> <p>Nocturnal voids/d, median: G1: -0.57 G2: -0.29 P=0.0026</p> <p>Nocturnal urgency severity score, mean: G1: -0.17 G2: 0.01 P=0.0005</p> <p>OAB-SCS, median: G1: -8.43 G2: -4.62 P<0.0001</p> <p>Adverse events Any, n (%) G1: 196 (59.6) G2: 153 (46.5)</p> <p>Dry mouth: G1: 65 (19.8) G2: 17 (5.2)</p> <p>Constipation: G1: 36 (10.9) G2: 19 (5.8)</p>	

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Rudy et al., 2006				Headache: G1: 18 (5.5) G2: 15 (4.6)	
Rudy et al., 2006				UTI not otherwise specified: G1: 16 (4.9) G2: 8 (2.4)	
Staskin and Harnett, 2004				Nasopharyngitis: G1: 13 (4.0) G2: 12 (3.6)	
(continued)				Cough: G1: 8 (2.4) G2: 1 (0.3)	
				Diarrhea: G1: 7 (2.1) G2: 13 (4.0)	
				AEs leading to treatment discontinuation, %: G1: 7.3 G2: 4.6	
				Most common AEs leading to discontinuation:	
				Constipation: G1: 1.8% G2: 0.6%	
				Dry mouth: G1: 1.5% G2: 0.0%	
				Stanford Sleepiness Scale, change, wk 1: G1: -0.2 G2: -0.12	
				Stanford Sleepiness Scale, change wk 4: G1: -0.17 G2: -0.14	
				Stanford Sleepiness Scale, change, wk 12: G1: -0.16 G2: -0.11	

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Rudy et al., 2006				Stanford Sleepiness Scale including T-max time point values only (G1=93, G2=97), change BL to week 1: G1: -0.42 G2: -0.2	
Rudy et al., 2006				Stanford Sleepiness Scale including T-max time point values only (G1=144, G2=148), change BL to week 4: G1: -0.2 G2: -0.19	
Staskin and Harnett, 2004				Stanford Sleepiness Scale including T-max time point values only (G1=182, G2=179), change BL to week 12: G1: -0.17 G2: -0.18	
(continued)				Mean Stanford Sleepiness Scale, age group <65 years, change BL to week 1: G1: -0.27 G2: -0.19	
				Mean Stanford Sleepiness Scale, age group >65 years, change BL to week 1: G1: -0.04 G2: -0.03	
				Mean Stanford Sleepiness Scale, age group <65 years, change BL to week 4: G1: -0.14 G2: -0.27	

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Rudy et al., 2006 Rudy et al., 2006 Staskin and Harnett, 2004 (continued)				Mean Stanford Sleepiness Scale, age group >65 years, change from baseline to week 4: G1: -0.23 G2: 0.03	
				Mean Stanford Sleepiness Scale, age group <65 years, change from baseline to week 12: G1: -0.14 G2: -0.22	
				Mean Stanford Sleepiness Scale, age group >65 years, change from baseline to week 12: G1: -0.22 G2: 0.03	
				Mean Stanford Sleepiness Scale, age group < 75 years, change from baseline to week 1: G1: -0.21 G2: -0.14	
				Mean Stanford Sleepiness Scale, age group > 75 years, change from baseline to week 1: G1: -0.02 G2: 0	
				Mean Stanford Sleepiness Scale, age group < 75 years, change from baseline to week 4: G1: 0.14 G2: -0.16	

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Rudy et al., 2006 Rudy et al., 2006 Staskin and Harnett, 2004 (continued)				<p>Mean Stanford Sleepiness Scale, age group > 75 years, change from baseline to week 4: G1: -0.49 G2: -0.09</p> <p>Mean Stanford Sleepiness Scale, age group < 75 years, change from baseline to week 12: G1: 0.15 G2: -0.14</p> <p>Mean Stanford Sleepiness Scale, age group > 75 years, change from baseline to week 12: G1: -0.33 G2: 0.02</p> <p>At least one CNS event: G1: 5.8% G2: 5.2%</p> <p>Somnolence G1: 0.3% G2: 0.6%</p> <p>Sedation: G1: 0 G2: 0.3%</p> <p>Clinically significant increase (≥ 3 points) from baseline to week 12 in SSS score, n (%): G1: 5 (1.5) G2: 8 (2.5)</p>	

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Salvatore et al., 2007</p> <p>Country and setting: Italy, Urogynecology outpatient clinic</p> <p>Enrollment period: January 2004 to October 2005</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Cohort</p> <p>Intervention: Tolterodine SR 4 mg qd</p> <p>Groups: G1: vaginal profile stage 0a or Ia G2: anterior prolapse ≥ stage IIa</p> <p>N at enrollment: G1: 184 G2: 51</p> <p>N at follow-up: NR</p> <p>Women, %: 100</p> <p>Age, median (range): G1: 59 (20, 85) G2: 59 (35, 82)</p> <p>Race/ethnicity: NR</p> <p>Parity, mean (range): G1: 2 (0-6) G2: 2 (0-6)</p> <p>Menopausal, n (%): G1: 132 (72) G2: 7 (18)</p>	<p>Inclusion criteria: Women Urodynamically proven OAB who had no prolapse or pure anterior vaginal prolapse Proven detrusor overactivity</p> <p>Exclusion criteria: All symptoms or signs related to voiding difficulties Prolapse</p>	<p>Previous surgery, n (%): G1: 52 (28) G2: 11 (22)</p> <p>HRT, n (%): G1: 35 (26) G2: 7 (18)</p>	<p>Improvement in condition or cured, n (%): G1: 158 (85.9) G2: 31 (60.8)* P = 0.0002</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: NA</p> <p>Masking: NA</p> <p>Pt selection criteria: +</p> <p>Loss to followup: NR</p> <p>Drop-out rates: NR</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: poor</p> <p>Age: +</p> <p>Baseline OAB status: NR</p> <p>Baseline characteristics: -</p> <p>Length of followup: +</p> <p>Measurement methods: -</p> <p>Measurement reliability: -</p> <p>Intervention description: +</p>

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Sand et al., 2004</p> <p>Country and setting: US, Specialty clinic</p> <p>Enrollment period: Subanalysis of OBJECT trial, women only</p> <p>Funding: ALZA Corp</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: 10mg ER Oxybutynin daily vs 4mg Tolterodine (2mg bid)</p> <p>Groups: G1: 10mg ER Oxybutynin daily x 12 wks G2: 4mg Tolterodine (2mg bid) x 12 wks</p> <p>Stratified by age: a: Age ≤ 64 b: Age 65-74 c: Age ≥ 75</p> <p>N at enrollment: G1: 152 G2: 163</p> <p>N at follow-up: G1: 132 G2: 146</p> <p>Women, %: 100</p> <p>Age, yrs ± SD: G1: 58.4 G2: 58.8</p> <p>Race/ethnicity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • ≥ 7 and ≤ 50 UUI episodes/week • ≥ 10 voids/day • MUI included if predominant UUI <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • UTI • IC • Urethral diverticulum • Bladder tumor • Bladder stone • Delivery within 6 mos • Pelvic, bladder, vaginal surgery in ≤ 6 mos • PVR ≥ 150 mL • CV, renal, pulmonary, GI, endocrine, neurologic, autoimmune, hematological, urological, psychiatric, or hepatic disease • Hematuria • Positive urine culture • Narrow angle glaucoma • Obstructive uropathy • Myasthenia gravis • POP to hymenal ring • Gastrointestinal obstruction • Decreased GI motility • GI narrowing • GI retention • Investigational drugs within 1 month of screening • Hypersensitivity to drugs 	<p>UUI episodes/week, mean: G1: 25.2 G2: 25.1</p> <p>Incontinence episodes/week, mean: G1: 28.1 G2: 28.9</p> <p>voids/week, mean: G1: 91.7 G2: 91.6</p> <p>Naïve to anti-cholinergics, %: G1: 60.5 G2: 60.7</p>	<p>UUI episodes/week, %: G1: 6.2 G2: 8.5 G1a: 5.0 G2a: 8.4 G1b: 5.5 G2b: 7.5 G1c: 8.5 G2c: 11.1 G1/G2: P = 0.038 G1a/G2a: P = 0.005 G1b/G2b: P = 0.337 G1c/G2c: P = 0.568</p> <p>Incontinence episodes/week, mean: G1: 7.3 G2: 10.1 G1a: 5.8 G2a: 10.0 G1b: 6.1 G2b: 9.2 G1c: 10.5 G2c: 12.5 G1/G2: P = 0.030 G1a/G2a: P = 0.005 G1b/G2b: P = 0.164 G1c/G2c: P = 0.714</p> <p>voids/week, mean: G1: 68.0 G2: 71.2 G1a: 63.7 G2a: 71.2 G1b: 73.8 G2b: 71.9 G1c: 66.8 G2c: 65.5 G1/G2: P = 0.272 G1a/G2a: P = 0.024 G1b/G2b: P = 0.706 G1c/G2c: P = 0.838</p>	<p>Quality: Overall quality score: fair INTERNAL VALIDITY: fair Randomization: + Masking: + Pt selection criteria: + Loss to followup: + Drop-out rates: + Power calculation: - Statistical issues: + EXTERNAL VALIDITY: good Age: + Baseline OAB status: + Baseline characteristics: ++ Length of followup: + Measurement methods: + Measurement reliability: + Intervention description: +</p>

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Sand et al., 2004 (continued)		<ul style="list-style-type: none"> • Current drug/ EtOH abuse • Pregnant • Breastfeeding • Inability to follow protocol 		<p>Dry mouth, %: G1: 28.3 G2: 33.7</p> <p>Constipation, %: G1: 8.6 G2: 6.7</p> <p>Retention, %: G1: 4.0 G2: 1.2</p> <p>Blurred vision, %: G1: 2.6 G2: 0.6</p> <p>Dizziness, %: G1: 3.9 G2: 4.3</p> <p>Insomnia, %: G1: 0.7 G2: 1.8</p> <p>Somnolence, %: G1: 3.3 G2: 1.8</p> <p>Nervousness, %: G1: 0 G2: 1.2</p> <p>Headache, %: G1: 9.2 G2: 10.4</p> <p>Dyspepsia, %: G1: 5.3 G2: 6.1</p> <p>Nausea, %: G1: 3.3 G2: 1.8</p> <p>Vomiting, %: G1: 2.0 G2: 1.8</p>	

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Serati et al, 2008</p> <p>Country and setting: NR; Urogynecology unit</p> <p>Enrollment period: January 2005 to December 2006</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: None</p>	<p>Design: Prospective case series</p> <p>Intervention: Tolterodine ER 5 mg qd (prescribed to women with proven pure DO w/o SUI)</p> <p>Groups: G1: Coital incontinence at orgasm G2: Coital incontinence during penetration G3: DO + coital incontinence at orgasm (sub-group of G1) G4: DO w/o coital incontinence (controls)</p> <p>N at enrollment: G1: 49 G2: 83 G3: 34 G4: 53</p> <p>N at follow-up: NR</p> <p>Age, median (range): G1-G2: 51 (22-70) G3: 50 (22-66) G4: 51 (19-68)</p> <p>BMI (kg/m²), median (range): G1-G2: 24.1 (17-43) G3: 25.3 (17-43) G4: 26 (19-35)</p> <p>Race/ethnicity: NR</p> <p>Postmenopausal, n (%): G1-G2: 59 (44.7) G3: 14 (41.2) G4: 19 (35.8)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> Sexually active women with UI <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Documented UTI Previous anti-muscarinic trt POP-Q ≥ Stage 2 Refusal to answer questions about sex life 	<p>Type of UI, n (%):</p> <p>DO: G1: 34 (69.4) G2: 13 (15.7) <i>P</i> < 0.0001</p> <p>SUI: G1: 5 (10.2) G2: 40 (48.2) MUI: G1: 0 Inconclusive: G1: 10 (20.4) G2: 19 (22.9)</p>	<p>Non-responders to trt, n (%): G3: 14 (41.2) G4: 9 (17) <i>P</i> = 0.023</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: NA</p> <p>Method and blinding: NA</p> <p>Pt selection criteria: +</p> <p>Loss to followup: ++</p> <p>Drop-out rates: ++</p> <p>Power calculation: -</p> <p>Statistical issues: -</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: +</p> <p>Baseline OAB status: NR</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: -</p> <p>Intervention description: +</p>

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Serati et al, 2008 (continued)	<p>Parity, median (range): G1-G2: 2 (0-5) G3: 2 (0-5) G4: 2 (0-4)</p> <p>Nulliparous, n (%): G1-G2: 7 (5.3) G3: NR G4: NR</p> <p>Follow-up: 12 weeks</p>				

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Steers et al., 2007</p> <p>Country and setting: Australia, Canada, US; multicenter, 30 centers</p> <p>Enrollment period: July 2001 to September 2003</p> <p>Funding: Eli Lilly</p> <p>FVC - frequency volume chart</p> <p>DOA - detrusor overactivity</p> <p>SU - sensory urge</p> <p>PVR - postvoid residual urine</p> <p>BL - baseline</p> <p>PGI-I- Patient Global Impression of Improvement</p> <p>TEAEs= Treatment emergent adverse events</p>	<p>Design: RCT, placebo-controlled, double-blind, stratified by urodynamic observation (DOA)</p> <p>Intervention: duloxetine vs placebo</p> <p>Groups: G1: placebo x 2 wks, duloxetine 40 mg bid x 4 wks, duloxetine 60 mg bid x bid 8 wks G2: placebo bid x 14 wks</p> <p>N at enrollment: G1: 153 G2: 153</p> <p>N at follow-up, N (%): G1: 90 (59) G2: 120 (78)</p> <p>Age, mean ± SD: G1: 56.0 ± 14.8 G2: 53.3 ± 14.1</p> <p>Women, N (%): 306 (100)</p> <p>BMI, mean kg/m2 ± SD: G1: 29.2 (7.4) G2: 30.2 (7.8)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> women >18 yoa predominant symptoms of OAB ≥ 3 months defined as bothersome urinary urgency and/or urge, abnormal voiding frequency (≤2 h mean daytime voiding interval) UDS observation of either detrusor OA or urgency that limited bladder capacity to <400mL <p>Exclusion criteria:</p> <ul style="list-style-type: none"> -SUI PVR >100 mL Mean 24 h total volume voided of ≥ 3L Urine culture ≥4 UTIs in past yr Regular use of meds for OAB w/in 4wks of enrollment Previous use of duloxetine Continenence surgery ≤ 6 mos Any major surgery ≤ 3 mos Pelvic organ prolapse > ICS Stage II Non-pharm intervention ≤ 3 mos PFM training that had not been stable for 3 mos or would not remain stable during the trial 	<p>Void per 24 hrs: G1: 10.76 G2: 10.49</p> <p>Daytime VI G1: 113.58 G2: 119.63</p> <p>Urinary incontinence episodes(all assessable pts): G1: 1.70 G2: 1.44</p> <p>Urinary incontinence episode (wet OAB): G1: 2.34 G2: 2.07</p> <p>Continenence pads/wk: G1: 7.81 G2: 7.05</p> <p>Volume voided, mL: G1: 175.41 G2: 183.40</p> <p>Nocturia episodes/d: G1: 1.47 G2: 1.63</p> <p>Quality of Life Instrument score, mean ± SD: I-QOL G1: 56.65 ± 24.80 G2: 57.11 ± 23.3</p> <p>U-IIQ: G1: 2.44 ± 1.18 G2: 2.39 ± 1.15</p> <p>U-UDI: G1: 3.18 ± 0.58 G2: 3.15 ± 0.71</p> <p>Bladder compliance (mL/cmH20) G1: 71.8 ± 85.7 G2: 79.3 ± 91.0</p>	<p>VE24 G1: -1.81 G2: -0.62 G1vG2, P<0.001</p> <p>Daytime VI G1: 29.46 G2: 6.51 G1vG2, P<0.001</p> <p>UIE (all assessable pts) G1: -0.74 G2: -0.14 p=0.006</p> <p>UIE (wet OAB) G1: -1.03 G2: -0.24 p=0.032</p> <p>Continenence pads/wk: G1: -1.41 G2: -0.53 P=0.143</p> <p>Volume voided, mL: G1: 8.85 G2: 5.07 P=0.171</p> <p>Nocturia episodes/d: G1: 0.05 G2: -0.18 p=0.004</p> <p>8/12 wks (120 mg/day) G1: 61.6 G2: 42.1 p=0.008</p> <p>Overall: G1: 53.4 G2: 41.9 G=0.052</p> <p>Headache: G1: 13 (8.5) G2: 8 (5.2) p=0.336</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: -</p> <p>Drop-out rates: ++</p> <p>Power calculation: +</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Steers et al., 2007 (continued)			SUI: G1: 0 G2: 0 Maximum cystometric capacity mL G1: 318.3 ± 152.4 G2: 330.4 ± 135.4 Volume threshold for first detrusor contraction mL G1: 226.4 ± 141.8 G2: 250.6 ± 146.0 Moderate or severe bladder condition from PGI-I Scale (%): G1: 84.9 G2: 88.8 I-QOL total score, mean ± SD: G1: 56.6 (24.9) G2: 57.0 (23.2) UIE , mean ± SD: G1: 1.55 (2.08) G2: 1.41 (2.00) VE24 , mean ± SD: G1: 10.8 (3.3) G2: 10.6 (3.6) Symptom of bothersome urgency, %: G1: 99.3 G2: 99.3 Symptoms of urge UI, %: G1: 88.2 G2: 88.9 Symptoms of SUI: G1: 42.5 G2: 50.3 Urodynamic DOA G1: 42.5 G2: 41.8	Diarrhea: G1: 10 (6.5) G2: 5 (3.3) p=0.289 Cough: G1: 7 (4.6) G2: 6 (3.9) p>0.999 Change in QoL Instrument score, mean ± SD: I-QOL G1: 8.37 ± 15.89 G2: 4.87 ± 15.27 G1>G2 (P=.035) U-IIQ G1: -0.44 ± 0.76 G2: -0.22 ± 0.87 P=0.018 U-UDI G1: -0.26± 0.67 G2: -0.19 ± 0.59 P=0.440 Bladder compliance (mL/cmH20) G1: 16.5 ±131.3 G2: 1.0 ± 120.8 P=0.655 Maximum cystometric capacity mL G1: 22.1 ± 116.4 G2: 23.8 ± 108.5 P=0.619 Volume threshold for first detrusor contraction mL G1: 90.0 ± 200.6 G2 : 13.9 ± 112.8 P=0.254 PGI-I score Better 4 wks (80 mg/day) G1: 59.9 G2: 42.9 p=0.005	

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Steers et al., 2007 (continued)			Sensory urgency: G1: 57.5 G2: 58.2	TEAEs reported, % G1: 79.1 G2: 55.6 P<0.001	
			Previous continence surgery, %: G1: 13.1 G2: 11.8	Appetite decreased G1: 6 (3.9) G2: 0 G1>G2, p=0.030	
			Current pelvic-floor muscle training, %: G1: 9.2 G2: 11.8	Arthralgia G1: 6 (3.9) G2: 3 (2.0) p=0.501	
			Previous behavioral therapy, %: G1: 3.3 G2: 2.6	Somnolence: G1: 6 (3.9) G2: 0 p=0.30	
			Previous tolterodine therapy, %: G1: 10.5 G2: 7.2	Sweating increased: G1: 6 (3.9) G2: 2 (1.3) p=0.283	
			Previous oxybutynin therapy, %: G1: 13.0 G2: 9.2	UTI: G1: 6 (3.9) G2: 6 (3.9) p>0.999	
				Anorgasmia G1: 5 (3.3) G2: 0 G1>G2, p=0.060	
				Anxiety: G1: 5 (3.3) G2: 0 G1>G2, p=0.060	
				Tremor: G1: 5 (3.3) G2: 0 G1>G2, p=0.060	
				Upper respiratory infection: G1: 5 (3.3) G2: 4 (2.6) p>0.999	
				Vomiting: G1: 5 (3.3) G2: 3 (2.0) p=0.723	

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Steers et al., 2007 (continued)				<p>Abdominal pain: G1: 4 (2.6) G2: 1 (0.7) p=0.371</p> <p>Back pain: G1: 4 (2.6) G2: 1 (0.7) p=0.371 Note: mean change consistent at 4 wks (duloxetine 80 mg/d) and 8 wks (duloxetine 120 mg/d)</p> <p>Change in I-QOL total score, mean ± SD:</p> <p>Discontinuation due to TEAEs, % G1: 28.1 G2: 5.2 p<0.001</p> <p>Treatment emergent adverse events, n (%):</p> <p>Nausea: G1: 47 (30.7) G2: 7 (4.6) G1>G2, p<0.001</p> <p>Dry mouth: G1: 25 (16.3) G2: 2 (1.3) G1>G2, p<0.001</p> <p>Dizziness: G1: 22 (14.4) G2: 1 (0.07) G1>G2, p<0.001</p> <p>Constipation: G1: 21 (13.7) G2: 5 (3.3) G1>G2, p=0.002</p> <p>Insomnia: G1: 20 (13.1) G2: 5 (3.3) G1>G2, p=0.003</p> <p>Fatigue: G1: 16 (10.5) G2: 3 (2.0) G1>G2, p=0.003</p>	

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Sussman and Garely, 2002</p> <p>Country and setting: US; 2 sites</p> <p>Enrollment period: NR</p> <p>Funding: NR</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: Multicenter RCT</p> <p>Intervention: Tolterodine ER vs. Oxybutynin ER</p> <p>Groups: G1: Tolterodine ER 2 mg qd x 8 wks G2: Tolterodine ER 4 mg qd x 8 wks G3: Oxybutynin ER 5 mg qd x 8 wks G4: Oxybutynin ER 10 mg qd x 8 wks</p> <p>N at enrollment: G1: 333 G2: 336 G3: 313 G4: 307</p> <p>N at follow-up: G1: 313 (86) G2: 316 (88) G3: 286 (81) G4: 285 (79)</p> <p>Age, yrs ± SD: G1: 63.8 ± 15.7 G2: 63.4 ± 16.6 G3: 59.8 ± 16.5 G4: 63.2 ± 15.9</p> <p>Women, N (%): G1: 243 (73) G2: 254 (76) G3: 245 (78) G4: 222 (72)</p> <p>Race/ethnicity, n (%): White: G1: 278 (84) G2: 296 (88) G3: 256 (82) G4: 247 (81) Black: G1: 28 (8) G2: 23 (7) G3: 41 (13) G4: 42 (14)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • 18+ years old • OAB (urinary frequency + urgency) w/ or w/o UI <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Pure SUI • Urinary retention • Gastric retention • Uncontrolled narrow-angle glaucoma • Hepatic or renal dysfunction • Symptomatic or recurrent UTI • ES • BT • Pelvic floor exercise w/in 1 wk of first study visit or expected to start during study • Indwelling catheter • Intermittent self-catherization • Contraindication to antimuscarinic trt • Estrogen therapy started less than 2 mos prior to first visit • Any treatment for UI w/in 1 wk of first visit • Use of anticholinergic drug or potent inhibitors of cytochrome P450 3A4 • Pregnant • Lactating • Childbearing potential w/o adequate contraception 	<p>Presence of UI, n (%): G1: 195 (59) G2: 214 (64) G3: 199 (64) G4: 195 (64)</p> <p>Duration of sx, n (%): <6 mos G1: 38 (11) G2: 48 (14) G3: 62 (20) G4: 43 (14) 6mos-5yrs: G1: 226 (68) G2: 224 (67) G3: 183 (59) G4: 205 (67) >5yrs: G1: 68 (20) G2: 64 (19) G3: 67 (21) G4: 59 (19)</p> <p>Severity of bladder condition, % No problems: G1: 1.3 G2: 0.3 G3: 1.0 G4: 0.7 Some very minor problems: G1: 6 G2: 5 G3: 5 G4: 5 Some minor problems: G1: 18 G2: 16 G3: 21 G4: 15 Moderate problems G1: 45 G2: 50 G3: 41 G4: 47</p>	<p>Bladder Condition Questionnaire</p> <p>Improvement in bladder condition at 8 wks, %:</p> <p>Overall G1: 60 G2: 70 G3: 59 G4: 60 G2 vs G3 p<0.01 G2 vs G4 p<0.01</p> <p>Pts with moderate or severe bladder condition: G2: 77 G4: 65 G2 vs G4 p<0.01</p> <p>Treatment naive G1: 60 G2: 69 G3: 60 G4: 61 p=0.11 for improvement rates p>0.05 for overall difference btw trt arms</p> <p>Treatment experienced G1: 57 G2: 75 G3: 59 G4: 54 p=0.11 for improvement rates p>0.05 for overall difference btw trt arms</p> <p>Adverse Events</p> <p>Withdrawal d/t AE, (%): G2: 6 G4: 13 P=0.001</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Masking: -</p> <p>Pt selection criteria: +</p> <p>Loss to followup: -</p> <p>Drop-out rates: -</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: -</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Sussman and Garely, 2002 (continued)	<p>Hispanic: G1: 19 (6) G2: 15 (5) G3: 15 (5) G4: 16 (5)</p> <p>Other, n (%): G1: 8 (2) G2: 2 (<1) G3: 1 (<1) G4: 2 (<1)</p> <p>Parity: NR</p>		<p>Severe problems: G1: 26 G2: 21 G3: 23 G4: 25</p> <p>Many severe problems: G1: 4 G2: 7 G3: 8 G4: 8</p>	<p>Mean change in severity of dry mouth (visual analogue scale): G1: 2.3 G2: 6.0 G3: 6.3 G4: 11.3 G1 vs G2: p=NS G3 vs G4: p=0.05 G2 vs G4: p=0.03</p>	

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Swift et al., 2003</p> <p>Country and setting: Europe (167 centers), North America (74 centers), Australia and New Zealand (4 centers), University</p> <p>Enrollment period: February 1999 to October 1999</p> <p>Funding: Pharmacia Corp</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT double blind placebo-controlled double dummy, random permuted blocks of 6</p> <p>Intervention: Tolterodine ER vs. Tolterodine IR</p> <p>Groups: G1: Tolterodine ER 4mg daily G2: Tolterodine IR 2 mg BID G3: placebo</p> <p>N at enrollment: G1: 417 G2: 408 G3: 410</p> <p>N at follow-up: Total: 1092</p> <p>Women, %: 100</p> <p>Age, yrs ± SD: G1: 59 ± 14 G2: 59 ± 14 G3: 60 ± 14</p> <p>Race/ethnicity, n (%): White: G1: 396 (95) G2: 389 (95) G3: 383 (93) Black: G1: 15 (4) G2: 12 (3) G3: 20 (5) Asian/Pacific: G1: 5 (1) G2: 4 (1) G3: 2 (1) Mixed: G1: 0 G2: 3 (1) G3: 5 (1) Unknown: G1: 1 (<1) G2: 0 G3: 0</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Women • Age ≥ 18 • ≥ 8 voids/day • > 5 UUI/ week • Symptoms x ≥ 6 months (per voiding diary) <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • SUI • Total daily urine volume > 3 liters • Hepatic/ renal disease • UTI • IC • Hematuria • BOO • Current e-stim • Current bladder training • Indwelling catheter • CIC • Pregnant/ nursing • Childbearing age without BC • Anticholinergic meds • Meds inhibit cytochrome450 	<p>Incontinence episodes/week, mean ± SD: G1: 22.1 ± 22.5 G2: 22.9 ± 21.9 G3: 23.9 ± 21.2</p> <p>Pads/day, mean ± SD: G1: 1.6 ± 2.1 G2: 1.5 ± 2.0 G3: 1.7 ± 2.4</p> <p>voids/day, mean ± SD: G1: 10.8 ± 4.2 G2: 11.1 ± 3.7 G3: 11.2 ± 3.9</p> <p>Voided volume (mL), mean ± SD: G1: 141.2 ± 43.1 G2: 137.2 ± 41.2 G3: 135.7 ± 43.2</p> <p>Previous drug therapy: n (%): G1: 235 (56) G2: 222 (54) G3: 225 (55)</p>	<p>Incontinence episodes/week, mean ± SD: G1: 10.3 ± 17.2 G2: 12.8 ± 19.8 G3: 16.7 ± 19.7 G1/G3: P = 0.001 G2/G3: P = 0.001</p> <p>Pads/day, mean ± SD: G1: 1.0 ± 1.8 G2: 1.0 ± 1.5 G3: 1.5 ± 2.2 G1/G3: P = 0.001 G2/G3: P = 0.001</p> <p>voids/day, mean ± SD: G1: 9.0 ± 3.2 G2: 9.3 ± 4.0 G3: 9.9 ± 3.8 G1/G3: P = 0.001 G2/G3: P = 0.005</p> <p>Voided volume (mL), mean ± SD: G1: 179.1 ± 66.6 G2: 169.7 ± 65.6 G3: 149.0 ± 56.3 G1/G3: P = 0.001 G2/G3: P = 0.001</p> <p>Clinical effectiveness*, dry mouth: G1: 0.53 G2: 0.39 G3: 0.30</p> <p>Dry mouth, n (%): G1: 105 (25.3) G2: 127 (31.2) G3: 33 (8.0) G1/G3: P < 0.01 G2/G3: P < 0.01</p> <p>Abdominal pain, n (%): G1: 18 (4.3) G2: 12 (2.9) G3: 7 (1.7) G1/G3: P = 0.03</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: fair</p> <p>Randomization: +</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: ++</p> <p>Drop-out rates: ++</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Swift et al., 2003 (continued)	BMI, kg/m² ± SD: G1: 28.8 ± 13.8 G2: 29.0 ± 11.0 G3: 28.8 ± 6.7			Constipation, n (%): G1: 27 (6.5) G2: 27 (6.6) G3: 14 (3.4) Dyspepsia, n (%): G1: 11 (2.7) G2: 14 (3.4) G3: 6 (1.5) Nausea, n (%): G1: 7 (1.7) G2: 9 (2.2) G3: 9 (2.2) Diarrhea, n (%): G1: 10 (2.4) G2: 14 (3.4) G3: 9 (2.2) Flatulence, n (%): G1: 8 (1.9) G2: 11 (2.7) G3: 6 (1.5) Xerophthalmia, n (%): G1: 16 (3.9) G2: 8 (2.0) G3: 8 (2.0) Abnormal vision, n (%): G1: 5 (1.2) G2: 4 (1.0) G3: 2 (0.5) Headache, n (%): G1: 29 (7.0) G2: 14 (3.4) G3: 19 (4.6) UTI, n (%): G1: 15 (3.6) G2: 11 (2.7) G3: 19 (4.6) Insomnia, n (%): G1: 7 (1.7) G2: 2 (0.5) G3: 9 (2.2) Somnolence, n (%): G1: 12 (2.9) G2: 11 (2.7) G3: 8 (2.0)	

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Swift et al., 2003 (continued)				<p>Dizziness, n (%): G1: 7 (1.7) G2: 7 (1.7) G3: 4 (1.0)</p> <p>Hypertension, n (%): G1: 6 (1.4) G2: 4 (1.0) G3: 4 (1.0)</p> <p>Sinusitis, n (%): G1: 8 (1.9) G2: 2 (0.5) G3: 3 (0.7)</p> <p>Arthritis, n (%): G1: 1 (0.2) G2: 5 (1.2) G3: 1 (0.2)</p> <p>Dry skin, n (%): G1: 2 (0.5) G2: 5 (1.2) G3: 1 (0.2)</p>	

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Van Kerrebroeck et al., 2002</p> <p>Freeman et al., 2003*</p> <p>Country and setting: North America (74 centers), Australasia (4 centers), Europe (89 centers)</p> <p>Enrollment period: NR</p> <p>Funding: Pharmacia Corporation</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: Tolterodine ER vs Tolterodine IR vs placebo</p> <p>Groups: G1: tolterodine ER 4 mg sid G2: tolterodine IR 2 mg bid G3: placebo</p> <p>N at enrollment: G1: 507 G2: 514 G3: 508</p> <p>N at follow-up: Total: 1442 G1: 398 G3: 374</p> <p>Women, n (%): G1: 417 (82) G2: 408 (79) G3: 410 (81)</p> <p>Age, mean (range): G1: 60 (20, 89) G2: 60 (22, 92) G3: 61 (22, 93)</p> <p>Race/ethnicity, %:* White: G1: 95.7 G3: 94.7 Black: G1: 3.0 G3: 3.5 Asian/Pacific Islander: G1: 1.0 G3: 0.8 Other: G1: 0.3 G3: 1.1</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 18 • Urinary frequency (≥ 8 voids/day) <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • SUI • Total daily urine volume > 3 L • Contra-indications to antimuscarinic treatment • Hepatic or renal disease • UITs • Interstitial cystitis • Hematuria • BOO • Current electrostimulation or bladder training therapy • Indwelling catheter or intermittent self-catheterization • Pregnant or nursing • Women not using reliable contraception • Being treated for OAB with other anticholinergic drugs or drugs that inhibit cytochrome P450 3A4 isoenzymes • Estrogen therapy < 2 months • Treatment w/ investigational drug < 2 months 	<p>Incontinence episodes/week, mean (range): G1: 22.1 (0, 168.0) G2: 23.2 (0, 168.0) G3: 23.3 (0, 168.0)</p> <p>≥ 5 incontinence episodes/week, n (%): G1: 492 (97) G2: 498 (97) G3: 494 (97)</p> <p>Pads/day, mean (range): G1: 1.4 (0-18) G2: 1.4 (0-25) G3: 1.5 (0-22)</p> <p>Voids/day, mean (range): G1: 10.9 (2.3, 51.3) G2: 11.1 (2.0, 48.6) G3: 11.3 (2.0, 37.4)</p> <p>≥ 8 voids/day, n (%): G1: 458 (90) G2: 469 (91) G3: 467 (92)</p> <p>Previous drug therapy, n (%): G1: 270 (53) G2: 276 (54) G3: 263 (52)</p> <p>Poor efficacy, %: G1: 43 G2: 38.4 G3: 40.7</p> <p>Able to finish tasks before visiting a toilet, %:* G1: 5 G3: 6</p>	<p>Urinary Urgency, subjective assessment, 12 wks, n (%):* Improvement: G1: 173 (44) G3: 118 (32) G1/G3: P < 0.001 No change: G1: 201 (51) G3: 212 (57) Deterioration: G1: 22 (6) G3: 44 (12) G1/G3: P < 0.002</p> <p>Urinary urgency, improvement, 12 wks, women only, %:* G1: 46.6 G3: 26.6 G1/G3: P = 0.001 OR 1.81 (95% CI: 1.31, 2.49)</p> <p>Not able to hold urine, 12 wks, %:* G1: 58 G3: 32 G1/G3: P < 0.001</p> <p>Incontinence episodes/week, mean change ± SD (%) G1: -11.8 ± 17.8 G2: -10.6 ± 16.9 G3: -6.9 ± 15.4 G1/G3: P = 0.00001 G2/G3: P = 0.0005 G1/G2: P < 0.05</p> <p>Incontinence episodes/week, median % change: G1: -71 G2: -60 G3: -30 G1/G2: P < 0.05</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: + Masking: + Pt selection criteria: + Loss to followup: + Drop-out rates: - Power calculation: - Statistical issues: + EXTERNAL VALIDITY: good</p> <p>Age: + Baseline OAB status: + Baseline characteristics: ++ Length of followup: + Measurement methods: + Measurement reliability: + Intervention description: +</p>

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Van Kerrebroeck et al., 2002			Voided volume (mL), mean (range):	Pads/day, mean ± SD:	
Freeman et al., 2003* (continued)			G1: 141 (36, 338) G2: 137 (38, 283) G3: 136 (31, 374)	G1: -0.5 ± 1.4 G2: -0.5 ± 1.8 G3: -0.2 ± 1.4 G1/G3: P = 0.0145 G2/G3: P = 0.0035	
				Voids/day, mean ± SD:	
				G1: -3.5 ± 4.9 G2: -3.3 ± 4.4 G3: -2.2 ± 4.0 G1/G3: P = 0.00001 G2/G3: P = 0.0002	
				Voluntary voids/day, mean ± SD:	
				G1: -1.8 ± 3.4 G2: -1.7 ± 3.3 G3: -1.2 ± 2.9 G1 vs G3 G1/G3: P = 0.00047 G2/G3: P = 0.0079	
				Bladder symptoms, improvement, 12 wks, women only, %:*	
				G1: 62.8 G3: 48.4 G1/G3: P = 0.001 OR 1.78 (95% CI: 0.34, 2.37)	
				Treatment benefit, 12 wks, n (%):*	
				Much benefit:	
				G1: 172 (43.2) G3: 88 (23.5) G1/G3: P < 0.001	
				Little benefit	
				G1: 138 (34.7) G3: 118 (31.6)	
				No benefit	
				G1: 88 (22.1) G3: 168 (44.9)	

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Van Kerrebroeck et al., 2002</p> <p>Freeman et al., 2003* (continued)</p>				<p>Able to finish tasks before visiting a toilet, 12 wks, %:* G1: 33 G3: 18 G1/G3: P < 0.001</p> <p>Voided volume (mL), mean change ± SD: G1: +34 ± 51 G2: +29 ± 47 G3: +14 ± 41 G1/G3: P = 0.00001 G2/G3: P = 0.0001</p> <p>Discontinued due to AEs, n (%): G1: 27 (5) G2: 28 (5) G3: 33 (6)</p> <p>Reported serious adverse events, n: G1: 7 G2: 12 G3: 18</p> <p>Parasympathetic Dry mouth, n (%): G1: 118 (23) G2: 156 (30) G3: 39 (8)</p> <p>Xerophthalmia, n (%): G1: 17 (3) G2: 12 (2) G3: 10 (2)</p> <p>Abnormal vision, n (%): G1: 6 (1) G2: 4 (1) G3: 2 (0.5)</p> <p>Dry skin, n (%): G1: 2 (0.5) G2: 6 (1) G3: 1 (0.5)</p>	

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Van Kerrebroeck et al., 2002 Freeman et al., 2003* (continued)				Gastrointestinal Constipation, n (%): G1: 30 (6) G2: 35 (7) G3: 22 (4) Dyspepsia, n (%): G1: 15 (3) G2: 16 (3) G3: 7 (1) Abdominal pain, n (%): G1: 19 (4) G2: 13 (3) G3: 8 (2) Diarrhea, n (%): G1: 10 (2) G2: 16 (3) G3: 11 (2) Flatulence, n (%): G1: 10 (2) G2: 14 (3) G3: 9 (2) Nausea, n (%): G1: 7 (1) G2: 10 (2) G3: 10 (2) Headache, n (%): G1: 32 (6) G2: 19 (4) G3: 23 (5) Somnolence, n (%): G1: 14 (3) G2: 13 (3) G3: 9 (2) Dizziness, n (%): G1: 11 (2) G2: 9 (2) G3: 5 (1) Fatigue, n (%): G1: 11 (2) G2: 6 (1) G3: 4 (1) Insomnia, n (%): G1: 7 (1) G2: 2 (0.5) G3: 9 (2)	

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Van Kerrebroeck et al., 2002				Urinary tract infection, n (%): G1: 16 (3) G2: 13 (3) G3: 20 (4)	
Freeman et al., 2003* (continued)				Dysuria, n (%): G1: 5 (1) G2: 8 (2) G3: 1 (0.5)	
				Peripheral edema, n (%): G1: 7 (1) G2: 7 (1) G3: 4 (1)	

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Zinner et al., 2002</p> <p>[See evidence table for Van Kerrebroeck et al., 2001}</p> <p>Country and setting: North America (74 centers), Australasia (4 centers), Europe (89 centers)</p> <p>Enrollment period: NR</p> <p>Funding: Pharmacia Corporation</p> <p>Author industry relationship disclosures: NR</p>	<p>Design: RCT</p> <p>Intervention: Tolterodine ER vs. placebo</p> <p>Groups: G1: tolterodine ER 4 mg qd < 65 years of age G2: placebo < 65 years of age G3: tolterodine ER 4 mg qd ≥ 65 years of age G4: placebo ≥ 65 years of age</p> <p>N at enrollment: G1: 293 G2: 285 G3: 214 G4: 223</p> <p>N at follow-up: G1: 257 G2: 246 G3: 193 G4: 194</p> <p>Women, n (%): G1: 255 (87)** G2: 247 (87)** G3: 162 (76) G4: 163 (73)</p> <p>Age, mean ± SD: G1: 51 ± 10.5 G2: 51 ± 10 G3: 74 ± 6 G4: 74 ± 6</p> <p>Race/ethnicity: NR</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥ 18 • Urinary frequency (≥ 8 voids/day) <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Demonstrable SUI • Total daily urine volume > 3 L • Contra-indications to antimuscarinic treatment • Hepatic or renal disease • UTI • Interstitial cystitis • Hematuria • BOO • Current electrostimulation or bladder training therapy • Indwelling catheter or intermittent self-catheterization • Pregnant or nursing • Women not using reliable contraception • Being treated for OAB with other anticholinergic drugs or drugs that inhibit cytochrome P450 3A4 isoenzymes • Estrogen therapy < 2 months • Treatment w/ investigational drug < 2 months 	<p>Not able to hold urine upon experiencing urgency, %: G1: 24.9 G2: 29.1 G3: 33.6 G4: 34.5</p> <p>Able to complete tasks before toilet visit in response to urgency, %: G1: 6.5 G2: 7.7 G3: 5.1 G4: 4.9</p> <p>Incontinence episodes/week, mean ± SD: G1: 21.4 ± 22.1 G2: 23.2 ± 22.0 G3: 23.2 ± 22.7 G4: 23.4 ± 18.9</p> <p>Urinary frequency, mean ± SD: G1: 11.0 ± 3.9 G2: 11.4 ± 4.2 G3: 10.8 ± 4.5 G4: 11.0 ± 3.2</p> <p>Voided volume (mL), mean ± SD: G1: 140 ± 41 G2: 137 ± 45 G3: 141 ± 45 G4: 134 ± 39</p> <p>Previous drug therapy, n (%): G1: 148 (50.5) G2: 146 (51.2) G3: 121 (56.5) G4: 117 (52.5)</p> <p>Percentage with poor efficacy, %: G1: 74 (50.0)** G2: 60 (41.1)** G3: 40 (33.1) G4: 46 (39.3)</p>	<p>Not able to hold urine upon experiencing urgency, %: G1: 11.3 G2: 21.1 G3: 15.9 G4: 25.6 G1/G2: P = 0.003 G3/G4: P = 0.007</p> <p>No age-related difference</p> <p>Able to complete tasks before toilet visit in response to urgency, %: G1: 32.8 G2: 16.8 G3: 26.2 G4: 14.8 G1/G2: P = 0.001 G3/G4: P = 0.003</p> <p>No age-related difference</p> <p>Incontinence episodes/week, mean ± SD G1: -12.0 ± 17.6 G2: -7.4 ± 15.6 G3: -11.5 ± 18.2 G4: -6.3 ± 15.0 G1/G2: P = 0.001 G3/G4: P < 0.001</p> <p>No age-related difference</p> <p>Urinary frequency, mean ± SD G1: -2.0 ± 3.1 G2: -1.4 ± 3.1 G3: -1.4 ± 3.7 G4: -0.9 ± 2.6 G1/G2: P = 0.26 G3/G4: P = 0.92</p>	<p>Quality: Overall quality score: poor</p> <p>INTERNAL VALIDITY: poor</p> <p>Randomization: +</p> <p>Masking: +</p> <p>Pt selection criteria: +</p> <p>Loss to followup: +</p> <p>Drop-out rates: NR</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: fair</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: -</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Zinner et al., 2002 (continued)				<p>Perception of bladder condition, improved, %: G1: 60.1 G2: 51.9 G3: 54.2 G4: 31.4 G1/G2: P < 0.05 G3/G4: P < 0.0001</p> <p>Perception of bladder condition, no change, %: G1: 32.4 G2: 38.9 G3: 38.3 G4: 51.1 G4/G2: P < 0.0001</p> <p>Perception of bladder condition, deterioration, %: G1: 7.2 G2: 9.1 G3: 7.5 G4: 17.5</p> <p>Treatment beneficial, %: G1: 78.3 G2: 58.3 G3: 69.8 G4: 46.9 G1/G2: P = 0.001 G3/G4: P = 0.001 No age-related difference</p> <p>Voided volume (mL), mean ± SD: G1: 35 ± 53 G2: 13 ± 41 G3: 33± 47 G4: 16 ± 41 G1/G2: P < 0.001 G3/G4: P < 0.001 No age-related difference</p>	

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Zinner et al., 2002 (continued)				<p>Adverse events, %: G1: 50.7 G2: 50.5 G3: 54.2 G4: 46.0</p> <p>Dry mouth, severe, % G1-G2: 1.7 G3-G4: 1.9</p> <p>Dry mouth, moderate, % G1-G2: 7.6 G3-G4: 6.5</p> <p>Dry mouth, mild, % G1-G2: 13.4 G3-G4: 15.9</p> <p>No dry mouth, % G1-G2: 77.3 G3-G4: 75.7</p>	

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
<p>Author: Zinner et al., 2004</p> <p>Country and setting: US, 51 sites</p> <p>Enrollment period: NR</p> <p>Funding: Indevus Corporation</p> <p>Author industry relationship disclosures: 4 of 5 Alza (1) Indevus (4) Kyowa (1) Lilly (2) Merck (1) Ortho-McNeil (1) Pfizer (2) Pharma (1) Pharmacia (1) Schwarz (1) Watson (2) Yamanouchi (1)</p>	<p>Design: RCT with 2 wk washout period</p> <p>Intervention: Tropium chloride vs placebo</p> <p>Groups: G1: Tropium chloride 20 mg bid G2: placebo</p> <p>N at enrollment: G1: 262 G2: 261</p> <p>N at follow-up: G1: 256 G2: 256</p> <p>Women, n (%): G1: 203 (77.5) G2: 186 (71.3)</p> <p>Age, mean ± SD: G1: 63 ± 0.8 G2: 61.5 ± 0.8</p> <p>Race/ethnicity, n (%): White: G1: 222 (84.7) G2: 225 (86.2) Black: G1: 26 (9.9) G2: 20 (7.7) Hispanic: G1: 10 (3.8) G2: 13 (5.0) Other: G1: 4 (1.6) G2: 3 (1.1)</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age ≥18 years • OAB symptoms ≥ 6 mos • Urinary urgency • ≥ 70 voids per week • ≥ 7 UUI episodes/week <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Stress predominant MUI • Insensate or overflow • Neurogenic bladder disorders • Significant renal disease • Hematuria hematuria and UTI at washout or ≥ twice in the prior year • BOO with PVR ≥ 100 mL • Concurrent use of any anticholinergic drug or other drug therapy for OAB within 21 days before randomization • Bladder surgery ≤ 6 mos • Bladder cancer • Interstitial cystitis • PSA ≥ 10 ng/mL • Diuretic use, estrogen therapy and nonmedical bladder therapy that was not part of a stable, long-term program. 	<p>UUI episodes/day, mean: G1: 3.9 G2: 4.3</p> <p>Urgency episodes/day, mean: G1: 11.29 G2: 11.72</p> <p>Urgency severity score, mean: G1: 1.8 G2: 1.8</p> <p>Voids/day, mean: G1: 12.7 G2: 12.9</p> <p>Daytime voids/day, mean: G1: 10.6 G2: 10.9</p> <p>Nocturia episodes/day, mean: G1: 2.1 G2: 2.0</p> <p>Voided volume (mL), mean: G1: 155.1 G2: 156.1</p> <p>IIQ, mean score (SE): G1: 183.5 (5.6) G2: 195.4 (5.6)</p> <p>IIQ,, women only, mean score (SE): G1: 191.2 (6.3) G2: 201.9 (6.6)</p> <p>Incontinence Impact Questionnaire, travel subscale, score (SE): G1: 52.4 (1.7) G2: 55.6 (1.7)</p>	<p>UUI episodes/day, mean %change: G1: 59.0 G2: 44.2 G1/BL: P < 0.0001</p> <p>Urgency episodes/day, mean change: G1: -0.22 G2: -1.08 G1/BL: P < 0.001</p> <p>Urgency severity score, mean: G1: 1.8 G2: 1.8</p> <p>Voids/day, mean change: G1: -2.37 G2: -1.29 G1/BL: P < 0.0001</p> <p>Daytime voids/day, mean change: G1: -1.90 G2: -0.98 G1/BL: P < 0.0001</p> <p>Nocturia episodes/day, mean change: G1: -0.47 G2: -0.29 G1/BL: P < 0.05</p> <p>Voided volume (mL), mean change: G1: 32.1 G2: 7.7 G1/BL: P < 0.0001</p> <p>IIQ,, LS mean change (SE): G1: -54.0 (5.6) G2: -36.0 (5.6) G1/BL: P ≤ 0.05</p>	<p>Quality: Overall quality score: fair</p> <p>INTERNAL VALIDITY: fair</p> <p>Randomization: +</p> <p>Masking: -</p> <p>Pt selection criteria: +</p> <p>Loss to followup: ++</p> <p>Drop-out rates: +</p> <p>Power calculation: -</p> <p>Statistical issues: +</p> <p>EXTERNAL VALIDITY: good</p> <p>Age: +</p> <p>Baseline OAB status: +</p> <p>Baseline characteristics: ++</p> <p>Length of followup: +</p> <p>Measurement methods: +</p> <p>Measurement reliability: +</p> <p>Intervention description: +</p>

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Zinner et al., 2004 (continued)			IIQ, social relationships, score (SE): G1: 37.8 (1.5) G2: 40.3 (1.5)	IIQ, women only, LS mean change (SE): G1: -59.1 (6.6) G2: -35.7 (6.9) G1/BL: $P \leq 0.05$	
			IIQ, emotional health, mean score (SE): G1: 47.1 (1.6) G2: 49.6 (1.6)	IIQ, travel subscale, LS mean change (SE): G1: -14.9 (1.7) G2: -9.9 (1.7) G1/BL: $P < \leq 0.05$	
			IIQ,, physical activity, mean score (SE): G1: 46.1 (1.6) G2: 50.2 (1.6)	IIQ, social relationships, LS mean change (SE): G1: -10.8 (1.4) G2: -6.3 (1.4) G1/BL: $P \leq 0.05$	
			Prior OAB med, n (%): G1: 135 (51.5) G2: 142 (54.5)	IIQ, emotional health, LS mean change (SE): G1: 14.1 (1.5) G2: -9.2 (1.5) G1/BL: $P \leq 0.05$	
				IIQ, physical activity, LS mean change (SE): G1: -13.5 (1.7) G2: -11.0 (1.7)	
				Dry mouth, n (%): G1: 57 (21.8) G2: 17 (6.5)	
				Constipation, n (%): G1: 25 (9.5) G2: 10 (3.8)	
				Headache, n (%): G1: 17 (6.5) G2: 12 (4.6)	

Evidence Table 7. KQ4 Modifiers of outcomes (continued)

Study Description	Study Design, Interventions, and Population	Inclusion/ Exclusion Criteria	Symptom Characteristics	Outcomes	Quality Rating
Zinner et al., 2004 (continued)				Diarrhea, n (%): G1: 8 (3.1) G2: 14 (5.4) Abdominal pain, n (%): G1: 8 (3.1) G2: 3 (1.1) Discontinuation due to AE, %: G1: 8.8 G2: 5.7	

Appendix D. List of Excluded Studies

Excluded Articles

Full Text Article Exclusion Criteria Codes for Database

X-1A: Not OAB (including post-operative/iatrogenic)

X-1B: Stress or mixed incontinence

X-1C: Isolated nocturia

X-1D: Interstitial cystitis/painful bladder syndrome

X-1E: Pelvic organ prolapse

X-1F: Neurogenic conditions

X-1G: Basic science or anatomy only

X-1H: Imaging/diagnostic study only

X-1I: Other (including questionnaire validation studies, coping strategies, etc.)

X-2: Not original research

X-4: Less than 75% of study population are adult females

X-5: Non-ambulatory population

X-6: Ineligible study type

X-7: N <50

X-8: Non- US cost study/general cost of incontinence (not specifically OAB)

X-9: Treatment not currently available in the US

X-10: Unable to obtain full text of manuscript

List of Excluded Studies

1. Terodiline and oxybutynin in detrusor instability. *Drug Ther Bull.* 1988 May 16;26(10):37-8. X-2
2. Effects of terodiline on urinary incontinence among older non-institutionalized women. Terodiline in the Elderly American Multicenter Study Group. *J Am Geriatr Soc.* 1993 Sep;41(9):915-22. X-9
3. Biofeedback, exercise for urge incontinence. *Health News.* 1999 Jan 5;5(1):7. X-2
4. Overactive bladder. *Harv Womens Health Watch.* 1999 Oct;7(2):6. X-2
5. Sacral nerve stimulation for the treatment of urinary urgency/frequency. *Tecnologica MAP Suppl.* 2000 Jun:12-5. X-2
6. Detrol LA and Diropan XL for overactive bladder. *Med Lett Drugs Ther.* 2001 Apr 2;43(1101):28. X-2
7. Incontinence: an overview. *N2N: Nurse2Nurse.* 2001 Feb;1(9):11-2, 14-5. X-2
8. Don't let elderly patients 'put up with' incontinence. *Senior Care Management.* 2002 Jun;5(6):87-8. X-10
9. Behavior therapy and urge incontinence. *Mayo Clin Health Lett.* 2003 May;21(5):4. X-2
10. Skin patch approved for treatment of overactive bladder. *Mayo Clin Womens Healthsource.* 2003 Aug;7(8):3. X-2
11. Overactive bladder. Relief for urgency, frequency and incontinence. *Mayo Clin Womens Healthsource.* 2004 Mar;8(3):1-2. X-2
12. Trospium chloride (Sanctura): another anticholinergic for overactive bladder. *Med Lett Drugs Ther.* 2004 Aug 2;46(1188):63-4. X-2
13. Innovations in overactive bladder treatment and the nursing home environment. *Long-Term Care Interface.* 2005;6:19-24. X-5
14. Design of the Behavior Enhances Drug Reduction of Incontinence (BE-DRI) study. *Contemp Clin Trials.* 2007 Jan;28(1):48-58. X-11
15. Appropriate pharmacotherapy for nocturia in elderly patients reduces urinary frequency, thereby improving sleep and health-related quality of life. *Drugs & Therapy Perspectives.* 2008;24(2):15-18. X-1C
16. Blander DS, Carpiello VL, Harryhill JF, et al. Extraperitoneal laparoscopic urethropexy with Marlex mesh. *Urology.* 1999 May;53(5):985-9. X-1B, X-7
17. Aagaard J, Reuther K and Stimpel H. A comparison between the combination emepromium bromide/flavoxate and emepromium bromide in the treatment of detrusor instability. *Urol Int.* 1983;38(3):191-2. X-7
18. Ab E, Dik P, Klijn AJ, et al. Detrusor overactivity in spina bifida: how long does it need to be treated? *Neurourol Urodyn.* 2004;23(7):685-8. X-1F, X-4, X-7
19. Abarbanel J and Marcus EL. Impaired detrusor contractility in community-dwelling elderly presenting with lower urinary tract symptoms. *Urology.* 2007 Mar;69(3):436-40. X-1H
20. Abdel-Fattah M, Ramsay I and Barrington JW. A simple visual analogue scale to assess the quality of life in women with urinary incontinence. *Eur J Obstet Gynecol Reprod Biol.* 2007 Jul;133(1):86-9. X-11
21. Abdel-Hady el S and Constantine G. Outcome of the use of tension-free vaginal tape in women with mixed urinary incontinence, previous failed surgery, or low valsalva pressure. *J Obstet Gynaecol Res.* 2005 Feb;31(1):38-42. X-1B
22. Aboseif S, Tamaddon K, Chalfin S, et al. Sacral neuromodulation as an effective treatment for refractory pelvic floor dysfunction. *Urology.* 2002 Jul;60(1):52-6. X-7
23. Abrams P, Blaivas JG, Fowler CJ, et al. The role of neuromodulation in the management of urinary urge incontinence. *BJU Int.* 2003 Mar;91(4):355-9. X-2
24. Abrams P, Kelleher C, Huels J, et al. Clinical relevance of health-related quality of life outcomes with darifenacin. *BJU Int.* 2008 Jul;102(2):208-13. X-2
25. Abrams P and Klevmark B. Frequency volume charts: an indispensable part of lower urinary tract assessment. *Scand J Urol Nephrol Suppl.* 1996;179:47-53. X-1H
26. Abrams P, Malone-Lee J, Jacquetin B, et al. Twelve-month treatment of overactive bladder: efficacy and tolerability of tolterodine. *Drugs Aging.* 2001;18(7):551-60. X-4
27. Abrams P and Swift S. Solifenacin is effective for the treatment of OAB dry patients: a pooled analysis. *Eur Urol.* 2005 Sep;48(3):483-7. X-2
28. Addison R. Catheter valves: a special focus on the Bard Flip-Flo catheter. *Br J Nurs.* 1999 May 13-26;8(9):576-80. X-1I
29. Adekanmi OA, Freeman RM, Reed H, et al. Improving the diagnosis of genuine stress incontinence in symptomatic women with negative cough stress test: the Distal Urethral Electrical Conductance test (DUEC) revisited. *Int Urogynecol J Pelvic Floor Dysfunct.* 2003 Feb;14(1):9-12; discussion 12. X-1H, X-7
30. Agarwal A, Dhirraaj S, Singhal V, et al. Comparison of efficacy of oxybutynin and tolterodine for prevention of catheter related bladder discomfort: a prospective, randomized, placebo-controlled, double-blind study. *Br J Anaesth.* 2006 Mar;96(3):377-80. X-1A
31. Agarwal A, Yadav G, Gupta D, et al. Evaluation of intra-operative tramadol for prevention of catheter-related bladder discomfort: a prospective, randomized, double-blind study. *Br J Anaesth.* 2008 Oct;101(4):506-10. X-1A
32. Agnew G and Byrne P. The evaluation and treatment of female urinary incontinence--a comparison of clinical practice in the Republic of Ireland with the recommendations of the International Continence Society. *Ir Med J.* 2004 Sep;97(8):238-40. X-1H
33. Agwu UM, Umeora OU and Ejikeme BN. Patterns of menopausal symptoms and adaptive ability in a rural population in South-east Nigeria. *J Obstet Gynaecol.* 2008 Feb;28(2):217-21. X-1I
34. Ahlbeck G and Ulmsten U. A urethral valve for bladder drill in patients with motor urge incontinence. *Gynecol Obstet Invest.* 1986;21(2):103-7. X-7
35. Ahlberg J, Edlund C, Wikkelso C, et al. Neurological signs are common in patients with urodynamically verified "idiopathic" bladder overactivity. *Neurourol Urodyn.* 2002;21(1):65-70. X-1F, X-1H
36. Aitchison M, Carter R, Paterson P, et al. Is the treatment of urgency incontinence a placebo response? Results of a five-year follow-up. *Br J Urol.* 1989 Nov;64(5):478-80. X-1I

37. Akkad T, Pelzer A, Mitterberger M, et al. Influence of intravesical potassium on pelvic floor activity in women with overactive bladder syndrome: comparative urodynamics might provide better detection of dysfunctional voiding. *BJU Int.* 2007 Oct;100(4):830-4; discussion 834. X-1H
38. Akpınar H, Cetinel B, Demirkesen O, et al. Long-term results of Burch colposuspension. *Int J Urol.* 2000 Apr;7(4):119-25. X-1B
39. Ala-Ketola L, Kauppila A, Jouppila P, et al. Pre- and postoperative bead chain urethrocytography in female stress urinary incontinence. *Acta Obstet Gynecol Scand.* 1981;60(4):369-74. X-1B
40. Albo ME, Richter HE, Brubaker L, et al. Burch colposuspension versus fascial sling to reduce urinary stress incontinence. *N Engl J Med.* 2007 May 24;356(21):2143-55. X-1B
41. Al-Buheissi S, Khasriya R, Maraj BH, et al. A simple validated scale to measure urgency. *J Urol.* 2008 Mar;179(3):1000-5; discussion 1005. X-1I
42. Alcalay M, Monga A and Stanton SL. Burch colposuspension: a 10-20 year follow up. *Br J Obstet Gynaecol.* 1995 Sep;102(9):740-5. X-1B
43. Alcalay M, Thompson PK and Boone TB. Ball urethroplasty combined with Marshall-Marchetti-Krantz urethropexy versus suburethral sling in patients with intrinsic sphincter deficiency and urethral hypermobility. *Am J Obstet Gynecol.* 2000 Dec;183(6):1348-53; discussion 1353-4. X-1A
44. Allen FJ and de Kock ML. Genito-urinary tuberculosis--experience with 52 urology inpatients. *S Afr Med J.* 1993 Dec;83(12):903-7. X-1I
45. Allen JH, McKenney JM, Stratton MA, et al. Antihypertensive effect of hydrochlorothiazide administered once or twice daily. *Clin Pharm.* 1982 May-Jun;1(3):239-43. X-1A, X-7
46. Alling Moller L, Lose G and Jorgensen T. Risk factors for lower urinary tract symptoms in women 40 to 60 years of age. *Obstet Gynecol.* 2000 Sep;96(3):446-51. X-1I
47. Alloussi S, Laval KU, Eckert R, et al. Trospium chloride (Spasmo-lyt) in patients with motor urge syndrome (detrusor instability): A double-blind, randomised, multicentre, placebo-controlled study. *Journal of Clinical Research.* 1998;1(439-451):439-451. X-10
48. Alloussi S, Loew F, Mast GJ, et al. Treatment of detrusor instability of the urinary bladder by selective sacral blockade. *Br J Urol.* 1984 Oct;56(5):464-7. X-7
49. Allsworth JE, Omicicoli VA, Cunkelman JA, et al. Discussion: 'Reproductive factors associated with nocturia and urgency' by Tikkinen et al. *Am J Obstet Gynecol.* 2008 Aug;199(2):e1-3. X-2
50. Allsworth JE, Omicicoli VA, Cunkelman JA, et al. Reproductive factors associated with nocturia and urgency: Tikkinen et al. *Am J Obstet Gynecol.* 2008 Aug;199(2):205-6. X-2
51. Almeida FG, Bruschini H and Srougi M. Urodynamic and clinical evaluation of 91 female patients with urinary incontinence treated with perineal magnetic stimulation: 1-year followup. *J Urol.* 2004 Apr;171(4):1571-4; discussion 1574-5. X-1B
52. Almeida FG, Bruschini H and Srougi M. Correlation between urethral sphincter activity and Valsalva leak point pressure at different bladder distentions: revisiting the urethral pressure profile. *J Urol.* 2005 Oct;174(4 Pt 1):1312-5; discussion 1315-6. X-1B
53. Al-Mulhim AA, Al-Gazzar SA and Bahnassy AA. Conservative treatment of idiopathic detrusor instability in elderly women. *Saudi Med J.* 2002 May;23(5):543-5. X-7
54. Alnaif B and Drutz HP. The prevalence of urinary and fecal incontinence in Canadian secondary school teenage girls: questionnaire study and review of the literature. *Int Urogynecol J Pelvic Floor Dysfunct.* 2001;12(2):134-137; discussion138. X-4
55. Alperin M, Abrahams-Gessel S and Wakamatsu MM. Development of de novo urge incontinence in women post sling: the role of preoperative urodynamics in assessing the risk. *Neurourol Urodyn.* 2008;27(5):407-11. X-1A
56. Altman D, Ekstrom A, Gustafsson C, et al. Risk of urinary incontinence after childbirth: a 10-year prospective cohort study. *Obstet Gynecol.* 2006 Oct;108(4):873-8. X-6
57. Amarenco G, Ismael SS, Even-Schneider A, et al. Urodynamic effect of acute transcuteaneous posterior tibial nerve stimulation in overactive bladder. *J Urol.* 2003 Jun;169(6):2210-5. X-1F
58. Amarenco G, Ismael SS, Lagauche D, et al. Cough anal reflex: strict relationship between intravesical pressure and pelvic floor muscle electromyographic activity during cough. Urodynamic and electrophysiological study. *J Urol.* 2005 Jan;173(1):149-52. X-1H, X-1I, X-7
59. Amaro JL, Gameiro MO, Kawano PR, et al. Intravaginal electrical stimulation: a randomized, double-blind study on the treatment of mixed urinary incontinence. *Acta Obstet Gynecol Scand.* 2006;85(5):619-22. X-1B, X-7
60. Amaro JL, Gameiro MO and Padovani CR. Effect of intravaginal electrical stimulation on pelvic floor muscle strength. *Int Urogynecol J Pelvic Floor Dysfunct.* 2005 Sep-Oct;16(5):355-8. X-1B, X-7
61. Amaro JL, Oliveira Gameiro MO and Padovani CR. Treatment of urinary stress incontinence by intravaginal electrical stimulation and pelvic floor physiotherapy. *Int Urogynecol J Pelvic Floor Dysfunct.* 2003 Aug;14(3):204-8; discussion 208. X-1B
62. Amayo EO, Kayima J, McLigeyo SO, et al. Autonomic nervous function in patients with chronic renal failure at the Kenyatta National Hospital. *East Afr Med J.* 1994 Apr;71(4):253-5. X-1A
63. Amoroso L, Pelliccioni G, Ghiselli R, et al. Sacral-neuromodulation CT-guided. *Radiol Med (Torino).* 2005 Apr;109(4):421-9. X-7
64. Amundsen C, Lau M, English SF, et al. Do urinary symptoms correlate with urodynamic findings? *J Urol.* 1999 Jun;161(6):1871-4. X-1H
65. Amundsen CL, Flynn BJ and Webster GD. Anatomical correction of vaginal vault prolapse by uterosacral ligament fixation in women who also require a pubovaginal sling. *J Urol.* 2003 May;169(5):1770-4. X-1B, X-7
66. Amundsen CL, Flynn BJ and Webster GD. Urethral erosion after synthetic and nonsynthetic pubovaginal slings: differences in management and continence outcome. *J Urol.* 2003 Jul;170(1):134-7; discussion 137. X-1I

67. Amundsen CL, Guralnick ML and Webster GD. Variations in strategy for the treatment of urethral obstruction after a pubovaginal sling procedure. *J Urol.* 2000 Aug;164(2):434-7. X-II
68. Amundsen CL, Parsons M, Cardozo L, et al. Bladder diary volume per void measurements in detrusor overactivity. *J Urol.* 2006 Dec;176(6 Pt 1):2530-4. X-IH
69. Amundsen CL, Parsons M, Tissot B, et al. Bladder diary measurements in asymptomatic females: functional bladder capacity, frequency, and 24-hr volume. *Neurourol Urodyn.* 2007;26(3):341-9. X-1A
70. Amundsen CL, Romero AA, Jamison MG, et al. Sacral neuromodulation for intractable urge incontinence: are there factors associated with cure? *Urology.* 2005 Oct;66(4):746-50. X-1F
71. Amundsen CL, Visco AG, Ruiz H, et al. Outcome in 104 pubovaginal slings using freeze-dried allograft fascia lata from a single tissue bank. *Urology.* 2000 Dec 4;56(6 Suppl 1):2-8. X-1B
72. Amundsen CL and Webster GD. Sacral neuromodulation in an older, urge-incontinent population. *Am J Obstet Gynecol.* 2002 Dec;187(6):1462-5; discussion 1465. X-7
73. Andersen JR, Lose G, Norgaard M, et al. Terodiline, emepronium bromide or placebo for treatment of female detrusor overactivity? A randomised, double-blind, cross-over study. *Br J Urol.* 1988 Apr;61(4):310-3. X-7
74. Andersen JT and Bradley WE. Abnormalities of detrusor and sphincter function in multiple sclerosis. *Br J Urol.* 1976 Jun;48(3):193-8. X-1F
75. Andersen LF, Agner T, Walter S, et al. Micturition pattern in hyperthyroidism and hypothyroidism. *Urology.* 1987 Feb;29(2):223-4. X-II
76. Anderson RU. Efficacy and safety of oxybutynin for urinary urge incontinence. *Clinical Geriatrics.* 2000;8(8):20. X-2
77. Andrades M, Paul R, Ambreen A, et al. Distribution of lower urinary tract symptoms (LUTS) in adult women. *J Coll Physicians Surg Pak.* 2004 Mar;14(3):132-5. X-6
78. Anger JT, Amundsen CL and Webster GD. Obstruction after Burch colposuspension: a return to retropubic urethrolysis. *Int Urogynecol J Pelvic Floor Dysfunct.* 2006 Sep;17(5):455-9. X-7
79. Anger JT, Litwin MS, Wang Q, et al. Complications of sling surgery among female Medicare beneficiaries. *Obstet Gynecol.* 2007 Mar;109(3):707-14. X-1A
80. Anger JT, Litwin MS, Wang Q, et al. The effect of age on outcomes of sling surgery for urinary incontinence. *J Am Geriatr Soc.* 2007 Dec;55(12):1927-31. X-1A
81. Anger JT, Rodriguez LV, Wang Q, et al. The role of preoperative testing on outcomes after sling surgery for stress urinary incontinence. *J Urol.* 2007 Oct;178(4 Pt 1):1364-8; discussion 1368-9. X-1B
82. Anger JT, Saigal CS, Madison R, et al. Increasing costs of urinary incontinence among female Medicare beneficiaries. *J Urol.* 2006 Jul;176(1):247-51; discussion 251. X-8
83. Apostolidis A, Brady CM, Yiangou Y, et al. Capsaicin receptor TRPV1 in urothelium of neurogenic human bladders and effect of intravesical resiniferatoxin. *Urology.* 2005 Feb;65(2):400-5. X-1A, X-7
84. Apostolidis A, Gonzales GE and Fowler CJ. Effect of intravesical Resiniferatoxin (RTX) on lower urinary tract symptoms, urodynamic parameters, and quality of life of patients with urodynamic increased bladder sensation. *Eur Urol.* 2006 Dec;50(6):1299-305. X-7
85. Apostolidis A, Popat R, Yiangou Y, et al. Decreased sensory receptors P2X3 and TRPV1 in suburothelial nerve fibers following intradetrusor injections of botulinum toxin for human detrusor overactivity. *J Urol.* 2005 Sep;174(3):977-82; discussion 982-3. X-1G, X-7
86. Appell RA. Clinical efficacy and safety of tolterodine in the treatment of overactive bladder: a pooled analysis. *Urology.* 1997 Dec;50(6A Suppl):90-6; discussion 97-9. X-2
87. Appell RA, Chancellor MB, Zobrist RH, et al. Pharmacokinetics, metabolism, and saliva output during transdermal and extended-release oral oxybutynin administration in healthy subjects. *Mayo Clin Proc.* 2003 Jun;78(6):696-702. X-II, X-7
88. Appendino G, De Petrocellis L, Trevisani M, et al. Development of the first ultra-potent "capsaicinoid" agonist at transient receptor potential vanilloid type 1 (TRPV1) channels and its therapeutic potential. *J Pharmacol Exp Ther.* 2005 Feb;312(2):561-70. X-1G
89. Arena MG, Di Rosa AE, Arcudi L, et al. Voiding disorders in patients with cerebrovascular disease. *Funct Neurol.* 1992 Jan-Feb;7(1):47-9. X-1F
90. Arikian N, Soygur T, Selcuki M, et al. Role of magnetic resonance imaging in children with voiding dysfunction: retrospective analysis of 81 patients. *Urology.* 1999 Jul;54(1):157-60; discussion 160-1. X-4
91. Armstrong RB, Dmochowski RR, Sand PK, et al. Safety and tolerability of extended-release oxybutynin once daily in urinary incontinence: combined results from two phase 4 controlled clinical trials. *Int Urol Nephrol.* 2007;39(4):1069-77. X-2
92. Arruda RM, Castro RA, Sartori MG, et al. Clinical and urodynamic evaluation of women with detrusor instability before and after functional pelvic floor electrostimulation. *Clin Exp Obstet Gynecol.* 2003;30(4):220-2. X-7
93. Arunkalaivanan AS and Barrington JW. Randomized trial of porcine dermal sling (Pelvic implant) vs. tension-free vaginal tape (TVT) in the surgical treatment of stress incontinence: a questionnaire-based study. *Int Urogynecol J Pelvic Floor Dysfunct.* 2003 Feb;14(1):17-23; discussion 21-2. X-1B
94. Arunkalaivanan AS, Mahomoud S and Howell M. Does posture affect cystometric parameters and diagnoses? *Int Urogynecol J Pelvic Floor Dysfunct.* 2004 Nov-Dec;15(6):422-4; discussion 424. X-1H
95. Arya LA, Banks C, Gopal M, et al. Development and testing of a new instrument to measure fluid intake, output, and urinary symptoms: the questionnaire-based voiding diary. *Am J Obstet Gynecol.* 2008 May;198(5):559 e1-7. X-1H
96. Arya LA, Myers DL and Jackson ND. Dietary caffeine intake and the risk for detrusor instability: a case-control study. *Obstet Gynecol.* 2000 Jul;96(1):85-9. X-II
97. Ashken MH, Abrams PH and Lawrence WT. Stamey endoscopic bladder neck suspension for stress incontinence. *Br J Urol.* 1984 Dec;56(6):629-34. X-1B

98. Aslan G, Koseoglu H, Sadik O, et al. Sexual function in women with urinary incontinence. *Int J Impot Res.* 2005 May-Jun;17(3):248-51. X-7
99. Atan A, Konety BR, Erickson JR, et al. Tolterodine for overactive bladder: time to onset of action, preferred dosage, and 9-month follow-up. *Tech Urol.* 1999 Jun;5(2):67-70. X-7
100. Atan A, Konety BR, Nangia A, et al. Advantages and risks of ileovesicostomy for the management of neuropathic bladder. *Urology.* 1999 Oct;54(4):636-40. X-1F, X-7
101. Athanasiou S, Anastakis A, Betsi GI, et al. Clinical and urodynamic parameters associated with history of urinary tract infections in women. *Acta Obstet Gynecol Scand.* 2007;86(9):1130-5. X-1I
102. Attanucci CA, Ball HG, Zweizig SL, et al. Differences in symptoms between patients with benign and malignant ovarian neoplasms. *Am J Obstet Gynecol.* 2004 May;190(5):1435-7. X-1A
103. Awad SA, Al-Zahrani HM, Gajewski JB, et al. Long-term results and complications of augmentation ileocystoplasty for idiopathic urge incontinence in women. *Br J Urol.* 1998 Apr;81(4):569-73. X-7
104. Awad SA, Flood HD and Acker KL. The significance of prior anti-incontinence surgery in women who present with urinary incontinence. *J Urol.* 1988 Sep;140(3):514-7. X-1B
105. Awad SA, Gajewski JB, Katz NO, et al. Final diagnosis and therapeutic implications of mixed symptoms of urinary incontinence in women. *Urology.* 1992 Apr;39(4):352-7. X-1B
106. Awad SA, Gajewski JB, Sogbein SK, et al. Relationship between neurological and urological status in patients with multiple sclerosis. *J Urol.* 1984 Sep;132(3):499-502. X-1F
107. Awad SA and McGinnis RH. Factors that influence the incidence of detrusor instability in women. *J Urol.* 1983 Jul;130(1):114-5. X-1H
108. Axelsen SM, Bek KM and Petersen LK. Urodynamic and ultrasound characteristics of incontinence after radical hysterectomy. *Neurourol Urodyn.* 2007;26(6):794-9. X-1A
109. Babu R. Effectiveness of tolterodine in nonneurogenic voiding dysfunction. *Indian Pediatr.* 2006 Nov;43(11):980-3. X-1I, X-4, X-7
110. Bachmann G. The estradiol vaginal ring—a study of existing clinical data. *Maturitas.* 1995 Dec;22 Suppl:S21-9. X-1I
111. Bachmann G. Urogenital ageing: an old problem newly recognized. *Maturitas.* 1995 Dec;22 Suppl:S1-S5. X-1I
112. Backhouse M and Harding L. Do males diagnosed with Duchenne muscular dystrophy (DMD) unnecessarily suffer bowel and bladder problems? *Australian & New Zealand Continence Journal.* 2006;12(3):66. X-1F
113. Badawi JK, Ding A and Bross S. Inhibitory effects of different ATP-sensitive potassium channel openers on electrically generated and carbachol-induced contractions of porcine and human detrusor muscle. *Fundam Clin Pharmacol.* 2008 Feb;22(1):75-86. X-1G
114. Bade JJ, Laseur M, Nieuwenburg A, et al. A placebo-controlled study of intravesical pentosanpolysulphate for the treatment of interstitial cystitis. *Br J Urol.* 1997 Feb;79(2):168-71. X-1D
115. Bael A, Lax H, de Jong TP, et al. The relevance of urodynamic studies for Urge syndrome and dysfunctional voiding: a multicenter controlled trial in children. *J Urol.* 2008 Oct;180(4):1486-93; discussion 1494-5. X-4
116. Baerheim A, Digranes A, Jureen R, et al. Generalized symptoms in adult women with acute uncomplicated lower urinary tract infection: an observational study. *MedGenMed.* 2003 Jul 1;5(3):1. X-1A
117. Bafghi A, Benizri EI, Trastour C, et al. Multifilament polypropylene mesh for urinary incontinence: 10 cases of infections requiring removal of the sling. *BJOG.* 2005 Mar;112(3):376-8. X-1B
118. Bafghi A, Valerio L, Benizri EI, et al. Comparison between monofilament and multifilament polypropylene tapes in urinary incontinence. *Eur J Obstet Gynecol Reprod Biol.* 2005 Oct 1;122(2):232-6. X-1B
119. Bagger PV, Fischer-Rasmussen W and Hansen R. Emepromium carrageenate: clinical effects and urinary excretion in treatment of female urge incontinence. *Scand J Urol Nephrol.* 1985;19(1):31-35. X-7
120. Bagi P and Biering-Sorensen F. Botulinum toxin A for treatment of neurogenic detrusor overactivity and incontinence in patients with spinal cord lesions. *Scand J Urol Nephrol.* 2004;38(6):495-8. X-1F, X-7
121. Bai SW, Jang JB, Lee DY, et al. Uterine arterial embolization for the treatment of uterine leiomyomas. *Yonsei Med J.* 2002 Jun;43(3):346-50. X-1I
122. Bai SW, Jung HJ, Jeon MJ, et al. Surgical repair of anterior wall vaginal defects. *Int J Gynaecol Obstet.* 2007 Aug;98(2):147-50. X-1I
123. Bai SW, Park SH, Chung DJ, et al. The significance of bladder trabeculation in the female lower urinary system: an objective evaluation by urodynamic studies. *Yonsei Med J.* 2005 Oct 31;46(5):673-8. X-1E
124. Baigis-Smith J, Smith DA, Rose M, et al. Managing urinary incontinence in community-residing elderly persons. *Gerontologist.* 1989 Apr;29(2):229-33. X-7
125. Bakas P, Liapis A, Giner M, et al. Quality of life in relation to TVT procedure for the treatment of stress urinary incontinence. *Acta Obstet Gynecol Scand.* 2006;85(6):748-52. X-1B
126. Balat A, Sarica K, Cekmen M, et al. Adrenomedullin and nitric oxide in children with detrusor instability. *Pediatr Nephrol.* 2003 May;18(5):422-5. X-1G, X-4, X-7
127. Ballert KN, Kanofsky JA and Nitti VW. Effect of tension-free vaginal tape and TVT-obturator on lower urinary tract symptoms other than stress urinary incontinence. *Int Urogynecol J Pelvic Floor Dysfunct.* 2008 Mar;19(3):335-40. X-1B
128. Bano F and Barrington JW. Serum calcium and magnesium levels in women with detrusor overactivity. *J Obstet Gynaecol.* 2005 May;25(4):371-3. X-1H, X-7
129. Baraza W, Boereboom C, Shorthouse A, et al. The long-term efficacy of fissurectomy and botulinum toxin injection for chronic anal fissure in females. *Dis Colon Rectum.* 2008 Feb;51(2):239-43. X-1I, X-7

130. Barbaliás GA, Nikiforidis G and Liatsikos EN. Vesicourethral dysfunction associated with multiple sclerosis: clinical and urodynamic perspectives. *J Urol.* 1998 Jul;160(1):106-11. X-1F
131. Barber MD, Gustilo-Ashby AM, Chen CC, et al. Perioperative complications and adverse events of the MONARC transobturator tape, compared with the tension-free vaginal tape. *Am J Obstet Gynecol.* 2006 Dec;195(6):1820-5. X-1B
132. Barber MD, Kleeman S, Karram MM, et al. Transobturator tape compared with tension-free vaginal tape for the treatment of stress urinary incontinence: a randomized controlled trial. *Obstet Gynecol.* 2008 Mar;111(3):611-21. X-1B
133. Barber MD, Visco AG, Wyman JF, et al. Sexual function in women with urinary incontinence and pelvic organ prolapse. *Obstet Gynecol.* 2002 Feb;99(2):281-9. X-II
134. Barnes NM, Dmochowski RR, Park R, et al. Pubovaginal sling and pelvic prolapse repair in women with occult stress urinary incontinence: effect on postoperative emptying and voiding symptoms. *Urology.* 2002 Jun;59(6):856-60. X-1B, X-1E
135. Barrero R, Fijo J, Fernandez-Hurtado M, et al. Vesicoureteral reflux after kidney transplantation in children. *Pediatr Transplant.* 2007 Aug;11(5):498-503. X-II, X-4
136. Barroso U, Jr., Jednak R, Barthold JS, et al. Outcome of ureteral reimplantation in children with the urge syndrome. *J Urol.* 2001 Sep;166(3):1031-5. X-4
137. Bar-Sela S, Levy M, Westin JB, et al. Medical findings in nickel-cadmium battery workers. *Isr J Med Sci.* 1992 Aug-Sep;28(8-9):578-83. X-II
138. Barthold JS, Martin-Crespo R, Kryger JV, et al. Quantitative nuclear cystography does not predict outcome in patients with primary vesicoureteral reflux. *J Urol.* 1999 Sep;162(3 Pt 2):1193-6. X-II, X-4
139. Bary PR, Moisey CU, Stephenson TP, et al. The urodynamic and subjective results of treatment of detrusor instability with oxybutynin chloride. *Prog Clin Biol Res.* 1981;78:313-9. X-7
140. Baselli EC, Brandes SB, Luthin GR, et al. The effect of pregnancy and contractile activity on bladder muscarinic receptor subtypes. *Neurourol Urodyn.* 1999;18(5):511-20. X-1G, X-4
141. Basok EK, Yildirim A, Atsu N, et al. Cadaveric fascia lata versus intravaginal slingplasty for the pubovaginal sling: surgical outcome, overall success and patient satisfaction rates. *Urol Int.* 2008;80(1):46-51. X-1B
142. Basok EK, Yildirim A, Atsu N, et al. The surgical results of the pubovaginal sling procedure using Intravaginal Slingplasty (IVS) for stress urinary incontinence. *Int Urol Nephrol.* 2006;38(3-4):507-12. X-1B
143. Bassi P, De Marco V, Tavolini IM, et al. Pharmacokinetic study of intravesical gemcitabine in carcinoma in situ of the bladder refractory to bacillus Calmette-Guerin therapy. *Urol Int.* 2005;75(4):309-13. X-1A, X-7
144. Bates CP, Whiteside CG and Turner-Warwick R. Synchronous cine-pressure-flow-cysto-urethrography with special reference to stress and urge incontinence. *Br J Urol.* 1970 Dec;42(6):714-23. X-1H
145. Batista JE, Caffaratti J, Arano P, et al. The reliability of cysto-urethrographic signs in the diagnosis of detrusor instability in children. *Br J Urol.* 1998 Jun;81(6):900-4. X-4
146. Batra AK, Mathews R and Lopresti A. Initial experience with the modified vaginal wall sling in the treatment of female stress urinary incontinence. *Int Urogynecol J Pelvic Floor Dysfunct.* 1997;8(4):209-12. X-1B, X-7
147. Bauer SB, Reda EF, Colodny AH, et al. Detrusor instability: a delayed complication in association with the artificial sphincter. *J Urol.* 1986 Jun;135(6):1212-5. X-1A, X-4, X-7
148. Beck RP. Urinary stress incontinence. *S Afr Med J.* 1980 May 24;57(21):853-65. X-1B
149. Beck RP, McCormick S and Nordstrom L. The fascia lata sling procedure for treating recurrent genuine stress incontinence of urine. *Obstet Gynecol.* 1988 Nov;72(5):699-703. X-1B
150. Beck RP, McCormick S and Nordstrom L. A 25-year experience with 519 anterior colporrhaphy procedures. *Obstet Gynecol.* 1991 Dec;78(6):1011-8. X-1B
151. Beck RP, Warren KG and Whitman P. Urodynamic studies in female patients with multiple sclerosis. *Am J Obstet Gynecol.* 1981 Feb 1;139(3):273-6. X-1F
152. Beco J. Interest of retro-anal levator plate myorrhaphy in selected cases of descending perineum syndrome with positive anti-sagging test. *BMC Surg.* 2008;8:13. X-1A
153. Beer AM, Fey S, Walch S, et al. The effect of peat components on endocrine and immunological parameters and on trace elements--results of two pilot studies. *Clin Lab.* 2001;47(3-4):161-7. X-7, X-II
154. Beier-Holgersen R, Kirkeby LT and Nordling J. 'Clam' ileocystoplasty. *Scand J Urol Nephrol.* 1994 Mar;28(1):55-8. X-7
155. Beisland HO and Fossberg E. The effects of terodiline and meladrazine on severe motor urge incontinence in geriatric patients. *J Am Geriatr Soc.* 1985 Jan;33(1):29-32. X-7
156. Bell DA, Collie D and Statham PF. Cauda equina syndrome: what is the correlation between clinical assessment and MRI scanning? *Br J Neurosurg.* 2007 Apr;21(2):201-3. X-1F, X-1A, X-7
157. Belman AB. The clinical significance of cystitis cystica in girls: results of a prospective study. *J Urol.* 1978 May;119(5):661-3. X-1A, X-4, X-7
158. Benassi L, Bocchialini E, Bertelli M, et al. Risk of genital prolapse and urinary incontinence due to pregnancy and delivery. A prospective study. *Minerva Ginecol.* 2002 Aug;54(4):317-24. X-II
159. Benedetti-Panici P, Zullo MA, Plotti F, et al. Long-term bladder function in patients with locally advanced cervical carcinoma treated with neoadjuvant chemotherapy and type 3-4 radical hysterectomy. *Cancer.* 2004 May 15;100(10):2110-7. X-1A
160. Bennett N, O'Leary M, Patel AS, et al. Can higher doses of oxybutynin improve efficacy in neurogenic bladder? *J Urol.* 2004 Feb;171(2 Pt 1):749-51. X-1F, X-7
161. Benson JT. Sacral nerve stimulation results may be improved by electrodiagnostic techniques. *Int Urogynecol J Pelvic Floor Dysfunct.* 2000 Dec;11(6):352-7. X-1H

162. Bent AE, Gousse AE, Hendrix SL, et al. Validation of a two-item quantitative questionnaire for the triage of women with urinary incontinence. *Obstet Gynecol.* 2005 Oct;106(4):767-73. X-II
163. Bent AE, Gousse AE, Hendrix SL, et al. Duloxetine compared with placebo for the treatment of women with mixed urinary incontinence. *Neurourol Urodyn.* 2008;27(3):212-21. X-1B
164. Bent AE, Richardson DA and Ostergard DR. Diagnosis of lower urinary tract disorders in postmenopausal patients. *Am J Obstet Gynecol.* 1983 Jan 15;145(2):218-22. X-1H
165. Bent AE, Sand PK, Ostergard DR, et al. Transvaginal electrical stimulation in the treatment of genuine stress incontinence and detrusor instability... This article has been reprinted from the *International Urogynecology Journal*, Vol. 4, #1, 1993 with permission of Springer Virag London Ltd. *Journal of Urological Nursing.* 1995 1995 Jan-Apr;14(1):930-941. X-2
166. Bergman A and Bader K. Reliability of the patient's history in the diagnosis of urinary incontinence. *Int J Gynaecol Obstet.* 1990 Jul;32(3):255-9. X-1B
167. Bergman A, Ballard CA and Platt LD. Ultrasonic evaluation of urethrovesical junction in women with stress urinary incontinence. *J Clin Ultrasound.* 1988 Jun;16(5):295-300. X-1B, X-1H
168. Bergman A, Bhatia NN and Hasen J. Effect of thyroid releasing hormone on bladder and urethral pressures. *Br J Urol.* 1984 Aug;56(4):397-400. X-1G, X-7
169. Bergman A, Bhatia NN and Hasen J. Urinary urgency and thyrotropin-releasing hormone. *Am J Obstet Gynecol.* 1984 Jan 1;148(1):106-7. X-1G, X-7
170. Bergman A, Koonings PP and Ballard CA. Detrusor instability. Is the bladder the cause or the effect? *J Reprod Med.* 1989 Oct;34(10):834-8. X-6
171. Bergman A, Stanczyk FZ and Lobo RA. The role of prostaglandins in detrusor instability. *Am J Obstet Gynecol.* 1991 Dec;165(6 Pt 1):1833-6. X-1G
172. Bergman J and Elia G. Effects of the menstrual cycle on urodynamic work-up: should we change our practice? *Int Urogynecol J Pelvic Floor Dysfunct.* 1999;10(6):375-7. X-1H, X-7
173. Bernardi R, Perin C, Becker FL, et al. Effect of pilocarpine mouthwash on salivary flow. *Braz J Med Biol Res.* 2002 Jan;35(1):105-10. X-1A, X-7
174. Bharucha AE, Locke GR, Zinsmeister AR, et al. Differences between painless and painful constipation among community women. *Am J Gastroenterol.* 2006 Mar;101(3):604-12. X-1A
175. Bharucha AE, Seide B, Guan Z, et al. Effect of tolterodine on gastrointestinal transit and bowel habits in healthy subjects. *Neurogastroenterol Motil.* 2008 Jun;20(6):643-8. X-1I
176. Bhatia NN and Bergman A. Urinary urgency following administration of thyrotropin releasing hormone during simultaneous urethrocytometry. *Urology.* 1988 Feb;31(2):180-3. X-7
177. Bhattacharya S, Mollison J, Pinion S, et al. A comparison of bladder and ovarian function two years following hysterectomy or endometrial ablation. *Br J Obstet Gynaecol.* 1996 Sep;103(9):898-903. X-1A
178. Bidmead J, Cardozo L, McLellan A, et al. A comparison of the objective and subjective outcomes of colposuspension for stress incontinence in women. *BJOG.* 2001 Apr;108(4):408-13. X-1B
179. Bihrlle W, 3rd and Tarantino AF. Complications of retropubic bladder neck suspension. *Urology.* 1990 Mar;35(3):213-4. X-1A, X-7
180. Billington A. Incontinence in older women. *Elderly Care.* 1999 Nov;11(8):32-3. X-2
181. Bing MH, Moller LA, Jennum P, et al. Pathophysiological aspects of nocturia in a danish population of men and women age 60 to 80 years. *J Urol.* 2007 Aug;178(2):552-7. X-6
182. Birch BR and Miller RA. Primary nocturnal enuresis: a urodynamic study spanning three generations. *Scand J Urol Nephrol.* 1995 Sep;29(3):285-8. X-II
183. Birder LA, Wolf-Johnston A, Griffiths D, et al. Role of urothelial nerve growth factor in human bladder function. *Neurourol Urodyn.* 2007;26(3):405-9. X-1G X-1H, X-7
184. Birns J, Lukkari E and Malone-Lee JG. A randomized controlled trial comparing the efficacy of controlled-release oxybutynin tablets (10 mg once daily) with conventional oxybutynin tablets (5 mg twice daily) in patients whose symptoms were stabilized on 5 mg twice daily of oxybutynin. *BJU Int.* 2000 May;85(7):793-8. X-4
185. Bjornsdottir LT, Geirsson RT and Jonsson PV. Urinary incontinence and urinary tract infections in octogenarian women. *Acta Obstet Gynecol Scand.* 1998 Jan;77(1):105-9. X-5
186. Black N, Griffiths J, Pope C, et al. Impact of surgery for stress incontinence on morbidity: cohort study. *BMJ.* 1997 Dec 6;315(7121):1493-8. X-1B
187. Blackford HN, Murray K, Stephenson TP, et al. Results of transvesical infiltration of the pelvic plexuses with phenol in 116 patients. *Br J Urol.* 1984 Dec;56(6):647-9. X-9
188. Blaivas JG, Bhimani G and Labib KB. Vesicourethral dysfunction in multiple sclerosis. *J Urol.* 1979 Sep;122(3):342-7. X-1F
189. Blaivas JG, Groutz A and Verhaaren M. Does the method of cystometry affect the incidence of involuntary detrusor contractions? A prospective randomized urodynamic study. *Neurourol Urodyn.* 2001;20(2):141-5. X-1H, X-7
190. Blaivas JG, Heritz DM and Romanzi LJ. Early versus late repair of vesicovaginal fistulas: vaginal and abdominal approaches. *J Urol.* 1995 Apr;153(4):1110-2; discussion 1112-3. X-II, X-7
191. Blaivas JG and Jacobs BZ. Pubovaginal fascial sling for the treatment of complicated stress urinary incontinence. *J Urol.* 1991 Jun;145(6):1214-8. X-1B
192. Blaivas JG and Olsson CA. Stress incontinence: classification and surgical approach. *J Urol.* 1988 Apr;139(4):727-31. X-1B
193. Blaivas JG, Panagopoulos G, Weiss JP, et al. Validation of the overactive bladder symptom score. *J Urol.* 2007 Aug;178(2):543-7; discussion 547. X-1H, X-II
194. Blaivas JG, Panagopoulos G, Weiss JP, et al. The urgency perception score: validation and test-retest. *J Urol.* 2007 Jan;177(1):199-202. X-II

195. Blaivas JG, Weiss JP, Desai P, et al. Long-term followup of augmentation enterocystoplasty and continent diversion in patients with benign disease. *J Urol*. 2005 May;173(5):1631-4. X-1A
196. Blander DS, Carpiniello VL, Harryhill JF, et al. Extraperitoneal laparoscopic urethropexy with Marlex mesh. *Urology*. 1999 May;53(5):985-9. X-1B
197. Blatt AH, Titus J and Chan L. Ultrasound measurement of bladder wall thickness in the assessment of voiding dysfunction. *J Urol*. 2008 Jun;179(6):2275-8; discussion 2278-9. X-1H
198. Blok BF, Groen J, Bosch JL, et al. Different brain effects during chronic and acute sacral neuromodulation in urge incontinent patients with implanted neurostimulators. *BJU Int*. 2006 Dec;98(6):1238-43. X-7
199. Blok BF, Sturms LM and Holstege G. Brain activation during micturition in women. *Brain*. 1998 Nov;121 (Pt 11):2033-42. X-1G, X-7
200. Bo K and Borgen JS. Prevalence of stress and urge urinary incontinence in elite athletes and controls. *Med Sci Sports Exerc*. 2001 Nov;33(11):1797-802. X-6
201. Bo K and Talseth T. Long-term effect of pelvic floor muscle exercise 5 years after cessation of organized training. *Obstet Gynecol*. 1996 Feb;87(2):261-5. X-7
202. Bodelsson G, Henriksson L, Osser S, et al. Short term complications of the tension free vaginal tape operation for stress urinary incontinence in women. *BJOG*. 2002 May;109(5):566-9. X-1B
203. Boemers TM, van Gool JD, de Jong TP, et al. Lower urinary tract dysfunction in children with benign sacrococcygeal teratoma. *J Urol*. 1994 Jan;151(1):174-6. X-1A, X-4
204. Bolukbas N, Vural M, Karan A, et al. Effectiveness of functional magnetic versus electrical stimulation in women with urinary incontinence. *Europa Medicophysica*. 2005 Dec;41(4):297-301. X-7
205. Bombier L, Freeman RM, Perkins EP, et al. Why do women have voiding dysfunction and de novo detrusor instability after colposuspension? *BJOG*. 2002 Apr;109(4):402-12. X-1A
206. Bombieri L and Freeman RM. Do bladder neck position and amount of elevation influence the outcome of colposuspension? *BJOG*. 2003 Feb;110(2):197-200. X-1A
207. Bond E, Dorey G, Eckford S, et al. The role of pelvic floor muscle exercises in reducing surgical management of women with stress incontinence: a clinical audit. *Journal of the Association of Chartered Physiotherapists in Women's Health*. 2004;70(17 ref). X-1B
208. Bonney WW, Gupta S, Hunter DR, et al. Bladder dysfunction in schizophrenia. *Schizophr Res*. 1997 Jun 20;25(3):243-9. X-II
209. Borawski KM, Foster RT, Webster GD, et al. Predicting implantation with a neuromodulator using two different test stimulation techniques: A prospective randomized study in urge incontinent women. *Neurourol Urodyn*. 2007;26(1):14-8. X-7
210. Borchert VE, Czyborra P, Fetscher C, et al. Extracts from *Rhois aromatica* and *Solidaginis virgaurea* inhibit rat and human bladder contraction. *Naunyn Schmiedeberg Arch Pharmacol*. 2004 Mar;369(3):281-6. X-1G, X-4, X-5, X-6, X-7
211. Bordman R, Telner D, Jackson B, et al. Step-by-step approach to managing pelvic organ prolapse: information for physicians. *Can Fam Physician*. 2007 Mar;53(3):485-7. X-1E
212. Borello-France D and Burgio KL. Nonsurgical treatment of urinary incontinence. *Clin Obstet Gynecol*. 2004 Mar;47(1):70-82. X-2
213. Borello-France D, Dusi J, O'Leary M, et al. Test-retest reliability of the Urge-Urinary Distress Inventory and Female Sexual Function Index in women with multiple sclerosis. *Urol Nurs*. 2008 Feb;28(1):30-5. X-1F, X-1H
214. Borello-France D, Leng W, O'Leary M, et al. Bladder and sexual function among women with multiple sclerosis. *Mult Scler*. 2004 Aug;10(4):455-61. X-1F
215. Borer JG, Gargollo PC, Kinnamon DD, et al. Bladder growth and development after complete primary repair of bladder exstrophy in the newborn with comparison to staged approach. *J Urol*. 2005 Oct;174(4 Pt 2):1553-7; discussion 1557-8. X-1A, X-II, X-4, X-7
216. Borrie MJ, Campbell AJ, Caradoc-Davies TH, et al. Urinary incontinence after stroke: a prospective study. *Age Ageing*. 1986 May;15(3):177-81. X-1F
217. Borup K and Nielsen JB. Results in 32 women operated for genuine stress incontinence with the pubovaginal sling procedure ad modum Ed McGuire. *Scand J Urol Nephrol*. 2002;36(2):128-33. X-1B, X-7
218. Borzyskowski M and Mundy AR. Videourodynamic assessment of diurnal urinary incontinence. *Arch Dis Child*. 1987 Feb;62(2):128-31. X-1H, X-4
219. Bosch JL and Groen J. Sacral (S3) segmental nerve stimulation as a treatment for urge incontinence in patients with detrusor instability: results of chronic electrical stimulation using an implantable neural prosthesis. *J Urol*. 1995 Aug;154(2 Pt 1):504-7. X-7
220. Bosch JL and Groen J. Neuromodulation: urodynamic effects of sacral (S3) spinal nerve stimulation in patients with detrusor instability or detrusor hyperflexia. *Behav Brain Res*. 1998 May;92(2):141-50. X-7
221. Bosch JL and Groen J. Sacral nerve neuromodulation in the treatment of patients with refractory motor urge incontinence: long-term results of a prospective longitudinal study. *J Urol*. 2000 Apr;163(4):1219-22. X-7
222. Botros SM, Abramov Y, Goldberg RP, et al. Detrusor overactivity and urge urinary incontinence [corrected] following midurethral versus bladder sling procedures. *Am J Obstet Gynecol*. 2005 Dec;193(6):2144-8. X-1A, X-1B
223. Botros SM, Miller JJ, Goldberg RP, et al. Detrusor overactivity and urge urinary incontinence following trans obturator versus midurethral slings. *Neurourol Urodyn*. 2007;26(1):42-5. X-1B
224. Bower WF, Moore KH and Adams RD. A pilot study of the home application of transcutaneous neuromodulation in children with urgency or urge incontinence. *J Urol*. 2001 Dec;166(6):2420-2. X-4
225. Bower WF, Moore KH, Adams RD, et al. A urodynamic study of surface neuromodulation versus sham in detrusor instability and sensory urgency. *J Urol*. 1998 Dec;160(6 Pt 1):2133-6. X-7
226. Bower WF, Sit FK and Yeung CK. Nocturnal enuresis in adolescents and adults is associated with childhood elimination symptoms. *J Urol*. 2006 Oct;176(4 Pt 2):1771-5. X-II, X-4

227. Bower WF, Yip SK and Yeung CK. Dysfunctional elimination symptoms in childhood and adulthood. *J Urol*. 2005 Oct;174(4 Pt 2):1623-7; discussion 1627-8. X-6
228. Boyington AR, Dougherty MC and Phetrasuwan S. Effectiveness of a computer-based system to deliver a continence health promotion intervention. *J Wound Ostomy Continence Nurs*. 2005 Jul-Aug;32(4):246-54. X-1B
229. Bozkurt N, Ozkan S, Korucuoglu U, et al. Urogenital symptoms of postmenopausal women in Turkey. *Menopause*. 2007 Jan-Feb;14(1):150-6. X-1I
230. Bradley CS, Kennedy CM and Nygaard IE. Pelvic floor symptoms and lifestyle factors in older women. *J Womens Health (Larchmt)*. 2005 Mar;14(2):128-36. X-1I
231. Bradshaw HD, Barker AT, Radley SC, et al. The acute effect of magnetic stimulation of the pelvic floor on involuntary detrusor activity during natural filling and overactive bladder symptoms. *BJU Int*. 2003 Jun;91(9):810-3. X-7
232. Bradshaw HD, Radley SC, Rosario DJ, et al. Towards a better understanding of involuntary detrusor activity. *BJU Int*. 2005 Apr;95(6):799-803. X-1H
233. Bradway C, Coyne KS, Irwin D, et al. Lower urinary tract symptoms in women—a common but neglected problem. *J Am Acad Nurse Pract*. 2008 Jun;20(6):311-8. X-2
234. Bramble FJ. The treatment of adult enuresis and urge incontinence by enterocystoplasty. *Br J Urol*. 1982 Dec;54(6):693-6. X-7, X-1I
235. Bratt H, Salvesen KA, Eriksen BC, et al. Long-term effects ten years after maximal electrostimulation of the pelvic floor in women with unstable detrusor and urge incontinence. *Acta Obstet Gynecol Scand Suppl*. 1998;168:22-4. X-7
236. Braun PM, Boschert J, Bross S, et al. Tailored laminectomy: a new technique for neuromodulator implantation. *J Urol*. 1999 Nov;162(5):1607-9. X-7
237. Breen JM, Geer BE and May GE. The fascia lata suburethral sling for treating recurrent urinary stress incontinence. *Am J Obstet Gynecol*. 1997 Dec;177(6):1363-5; discussion 1365-6. X-1B, X-6
238. Briel RC. Follow-up of a new modification of the Marshall-Marchetti-Krantz (MMK) procedure. *Arch Gynecol*. 1986;239(1):1-9. X-1B
239. Britton S. Benefits of inclusive continence assessments for sexual health. *Nurs Times*. 2006 Jan 10-16;102(2):50-1. X-2
240. Brock JW, 3rd. The frequency and frequency dysuria syndromes of childhood: hypercalciuria as a possible etiology. *Urology*. 1994 Sep;44(3):411-2. X-4
241. Brocklehurst JC. Urinary incontinence in the community—analysis of a MORI poll. *BMJ*. 1993 Mar 27;306(6881):832-4. X-1A
242. Bross S, Braun PM, Michel MS, et al. Bladder wall tension during physiological voiding and in patients with an unstable detrusor or bladder outlet obstruction. *BJU Int*. 2003 Oct;92(6):584-8. X-1G, X-1F, X-7
243. Bross S, Honeck P, Kwon ST, et al. Correlation between motor function and lower urinary tract dysfunction in patients with infantile cerebral palsy. *Neurourol Urodyn*. 2007;26(2):222-7. X-1F, X-7
244. Brown JS. Epidemiology and changing demographics of overactive bladder: a focus on the postmenopausal woman. *Geriatrics*. 2002 May;57 Suppl 1:6-12. X-2
245. Brown JS, Bradley CS, Subak LL, et al. The sensitivity and specificity of a simple test to distinguish between urge and stress urinary incontinence. *Ann Intern Med*. 2006 May 16;144(10):715-23. X-1H
246. Brown JS, Grady D, Ouslander JG, et al. Prevalence of urinary incontinence and associated risk factors in postmenopausal women. *Heart & Estrogen/Progesterone Replacement Study (HERS) Research Group. Obstet Gynecol*. 1999 Jul;94(1):66-70. X-6
247. Brown JS, McNaughton KS, Wyman JF, et al. Measurement characteristics of a voiding diary for use by men and women with overactive bladder. *Urology*. 2003 Apr;61(4):802-9. X-1H
248. Brown JS, Posner SF and Stewart AL. Urge incontinence: new health-related quality of life measures. *J Am Geriatr Soc*. 1999 Aug;47(8):980-8. X-7
249. Brown JS, Subak LL, Gras J, et al. Urge incontinence: the patient's perspective. *J Womens Health*. 1998 Dec;7(10):1263-9. X-7
250. Brown JS, Vittinghoff E, Kanaya AM, et al. Urinary tract infections in postmenopausal women: effect of hormone therapy and risk factors. *Obstet Gynecol*. 2001 Dec;98(6):1045-52. X-1I
251. Brown JS, Vittinghoff E, Wyman JF, et al. Urinary incontinence: does it increase risk for falls and fractures? Study of Osteoporotic Fractures Research Group. *J Am Geriatr Soc*. 2000 Jul;48(7):721-5. X-1I
252. Brown K and Hilton P. The incidence of detrusor instability before and after colposuspension: a study using conventional and ambulatory urodynamic monitoring. *BJU Int*. 1999 Dec;84(9):961-5. X-1A, X-1B
253. Brubaker L. Clinical advances in urogynecology. *Minerva Ginecol*. 2006 Aug;58(4):259-64. X-2
254. Brubaker L, Benson JT, Bent A, et al. Transvaginal electrical stimulation for female urinary incontinence. *Am J Obstet Gynecol*. 1997 Sep;177(3):536-40. X-2
255. Brubaker L, Cundiff GW, Fine P, et al. Abdominal sacrocolpopexy with Burch colposuspension to reduce urinary stress incontinence. *N Engl J Med*. 2006 Apr 13;354(15):1557-66. X-1E
256. Brubaker L and FitzGerald MP. Nocturnal polyuria and nocturia relief in patients treated with solifenacin for overactive bladder symptoms. *Int Urogynecol J Pelvic Floor Dysfunct*. 2007 Jul;18(7):737-41. X-2
257. Brubaker L, Kreder K, Richter HE, et al. Refractory urge urinary incontinence and botulinum A injection: the methods of the RUBI Trial. *Journal of Applied Research*. 2006;6(4):260-71. X-6
258. Brubaker L, Richter HE, Visco A, et al. Refractory idiopathic urge urinary incontinence and botulinum A injection. *J Urol*. 2008 Jul;180(1):217-22. X-7
259. Brusa L, Petta F, Pisani A, et al. Central acute D2 stimulation worsens bladder function in patients with mild Parkinson's disease. *J Urol*. 2006 Jan;175(1):202-6; discussion 206-7. X-1F
260. Brusa L, Petta F, Pisani A, et al. Acute vs chronic effects of l-dopa on bladder function in patients with mild Parkinson disease. *Neurology*. 2007 May 1;68(18):1455-9. X-1F, X-7

261. Bruschini H, Almeida FG and Srougi M. Upper and lower urinary tract evaluation of 104 patients with myelomeningocele without adequate urological management. *World J Urol.* 2006 Jun;24(2):224-8. X-II
262. Bryan NP and Chapple CR. Frequency volume charts in the assessment and evaluation of treatment: how should we use them? *Eur Urol.* 2004 Nov;46(5):636-40. X-1H
263. Buchsbaum GM, Chin M, Glantz C, et al. Prevalence of urinary incontinence and associated risk factors in a cohort of nuns. *Obstet Gynecol.* 2002 Aug;100(2):226-9. X-6
264. Bukkapatnam R, Shah S, Raz S, et al. Anterior vaginal wall surgery in elderly patients: outcomes and assessment. *Urology.* 2005 Jun;65(6):1104-8. X-1E
265. Bulmer PJ, James M, Ellis-Jones J, et al. A randomized trial comparing the effectiveness and preference of a touch-screen computer system with a leaflet for providing women with information on urinary symptoms suggestive of detrusor instability. *BJU Int.* 2001 Oct;88(6):532-5. X-II, X-7
266. Bump RC. The urethrodetrusor facilitative reflex in women: results of urethral perfusion studies. *Am J Obstet Gynecol.* 2000 Apr;182(4):794-802; discussion 802-4. X-1H
267. Bump RC, Fantl JA and Hurt WG. Dynamic urethral pressure profilometry pressure transmission ratio determinations after continence surgery: understanding the mechanism of success, failure, and complications. *Obstet Gynecol.* 1988 Dec;72(6):870-4. X-1A, X-7
268. Bump RC, Hurt WG, Elser DM, et al. Understanding lower urinary tract function in women soon after bladder neck surgery. *Continence Program for Women Research Group. Neurourol Urodyn.* 1999;18(6):629-37. X-II
269. Bump RC, Norton PA, Zinner NR, et al. Mixed urinary incontinence symptoms: urodynmic findings, incontinence severity, and treatment response. *Obstet Gynecol.* 2003 Jul;102(1):76-83. X-1B
270. Bump RC, Sugerman HJ, Fantl JA, et al. Obesity and lower urinary tract function in women: effect of surgically induced weight loss. *Am J Obstet Gynecol.* 1992 Aug;167(2):392-7; discussion 397-9. X-7
271. Bunyavejchevin S. Risk factors of female urinary incontinence and overactive bladder in Thai postmenopausal women. *J Med Assoc Thai.* 2005 Sep;88 Suppl 4:S119-23. X-II
272. Bunyavejchevin S. Urinary symptoms and quality of life changes in Thai women with overactive bladder after tolterodine treatment. *J Med Assoc Thai.* 2005 Nov;88(11):1497-501. X-7
273. Bunyavejchevin S. The impact of overactive bladder, stress and mixed urinary incontinence on quality of life in Thai postmenopausal women. *J Med Assoc Thai.* 2006 Mar;89(3):294-8. X-II
274. Bunyavejchevin S and Veeraranapanich S. Quality of life assessment in Thai postmenopausal women with an overactive bladder. *J Med Assoc Thai.* 2005 Aug;88(8):1023-7. X-II
275. Bunyavejchevin S and Wisawasukmongchol W. Five years follow up of laparoscopic burch colposuspension for stress urinary incontinence in Thai women. *J Med Assoc Thai.* 2005 Sep;88(9):1182-6. X-1B, X-7
276. Burgio KL. Behavioral training for stress and urge incontinence in the community. *Gerontology.* 1990;36 Suppl 2:27-34. X-7
277. Burgio KL. Influence of behavior modification on overactive bladder. *Urology.* 2002 Nov;60(5 Suppl 1):72-6; discussion 77. X-7
278. Burgio KL. Behavioral treatment options for urinary incontinence. *Gastroenterology.* 2004 Jan;126(1 Suppl 1):S82-9. X-2
279. Burgio KL, Goode PS, Locher JL, et al. Predictors of outcome in the behavioral treatment of urinary incontinence in women. *Obstet Gynecol.* 2003 Nov;102(5 Pt 1):940-7. X-2
280. Burgio KL, Nygaard IE, Richter HE, et al. Bladder symptoms 1 year after abdominal sacrocolpopexy with and without Burch colposuspension in women without preoperative stress incontinence symptoms. *Am J Obstet Gynecol.* 2007 Dec;197(6):647 e1-6. X-1A, X-1B
281. Burgio KL, Whitehead WE and Engel BT. Urinary incontinence in the elderly. Bladder-sphincter biofeedback and toileting skills training. *Ann Intern Med.* 1985 Oct;103(4):507-15. X-7
282. Burke DM, Shackley DC and O'Reilly PH. The community-based morbidity of flexible cystoscopy. *BJU Int.* 2002 Mar;89(4):347-9. X-II
283. Burkhard FC, Blick N, Hochreiter WW, et al. Urinary urgency and frequency, and chronic urethral and/or pelvic pain in females. Can doxycycline help? *J Urol.* 2004 Jul;172(1):232-5. X-II
284. Burkman RT. Transdermal hormonal contraception: benefits and risks. *Am J Obstet Gynecol.* 2007 Aug;197(2):134 e1-6. X-II
285. Burks FN, Diokno AC, Lajiness MJ, et al. Sacral neuromodulation reprogramming: is it an office burden? *Int Urogynecol J Pelvic Floor Dysfunct.* 2008 Aug;19(8):1137-40. X-7
286. Burrows LJ, Meyn LA, Walters MD, et al. Pelvic symptoms in women with pelvic organ prolapse. *Obstet Gynecol.* 2004 Nov;104(5 Pt 1):982-8. X-1E
287. Burton JR, Pearce KL, Burgio KL, et al. Behavioral training for urinary incontinence in elderly ambulatory patients. *J Am Geriatr Soc.* 1988 Aug;36(8):693-8. X-7
288. Butler RN, Maby JJ, Montella JM, et al. Urinary incontinence: when to refer for procedural therapies. *Geriatrics.* 1999 Dec;54(12):49-54, 56. X-1H, X-2
289. Byck DB, Varner RE and Clough C. Urinary complaints after modified Burch urethropexy: an analysis. *Am J Obstet Gynecol.* 1994 Dec;171(6):1460-2; discussion 1462-4. X-1B
290. Byrne RR, Auge BK, Kourambas J, et al. Routine ureteral stenting is not necessary after ureteroscopy and ureteropyeloscopy: a randomized trial. *J Endourol.* 2002 Feb;16(1):9-13. X-II
291. Cain MP, Wu SD, Austin PF, et al. Alpha blocker therapy for children with dysfunctional voiding and urinary retention. *J Urol.* 2003 Oct;170(4 Pt 2):1514-5; discussion 1516-7. X-1F, X-4
292. Cameron-Strange A and Millard RJ. Management of refractory detrusor instability by transvesical phenol injection. *Br J Urol.* 1988 Oct;62(4):323-5. X-7

293. Cammu H and Van Nysten M. Pelvic floor muscle exercises: 5 years later. *Urology*. 1995 Jan;45(1):113-7; discussion 118. X-1B
294. Candiani GB, Fedele L, Vercellini P, et al. Presacral neurectomy for the treatment of pelvic pain associated with endometriosis: a controlled study. *Am J Obstet Gynecol*. 1992 Jul;167(1):100-3. X-1A
295. Cantor TJ and Bates CP. A comparative study of symptoms and objective urodynamic findings in 214 incontinent women. *Br J Obstet Gynaecol*. 1980 Oct;87(10):889-92. X-1H
296. Cappellano F, Bertapelle P, Spinelli M, et al. Quality of life assessment in patients who undergo sacral neuromodulation implantation for urge incontinence: an additional tool for evaluating outcome. *J Urol*. 2001 Dec;166(6):2277-80. X-1F
297. Caputo RM, Benson JT and McClellan E. Intravaginal maximal electrical stimulation in the treatment of urinary incontinence. *J Reprod Med*. 1993 Sep;38(9):667-71. X-7
298. Caraballo R, Bologna RA, Lukban J, et al. Sacral nerve stimulation as a treatment for urge incontinence and associated pelvic floor disorders at a pelvic floor center: a follow-up study. *Urology*. 2001 Jun;57(6 Suppl 1):121. X-7
299. Carbone A, Gezeroglu H, Aloisi P, et al. Is aging a real risk factor for urological pathologies in men and women? *Arch Esp Urol*. 2001 Jan-Feb;54(1):87-94. X-1H
300. Carbone A, Palleschi G, Conte A, et al. Gabapentin treatment of neurogenic overactive bladder. *Clin Neuropharmacol*. 2006 Jul-Aug;29(4):206-14. X-1F, X-7
301. Carbone A, Palleschi G, Parasciani R, et al. Modulation of viscerosomatic H-reflex during bladder filling: a possible tool in the differential diagnosis of neurogenic voiding dysfunctions. *Eur Urol*. 2002 Sep;42(3):281-8. X-1F, X-1H
302. Cardozo L. Urinary frequency and urgency. *Br Med J (Clin Res Ed)*. 1986 Nov 29;293(6559):1419-23. X-2
303. Cardozo L, Castro-Diaz D, Gittelman M, et al. Reductions in overactive bladder-related incontinence from pooled analysis of phase III trials evaluating treatment with solifenacin. *Int Urogynecol J Pelvic Floor Dysfunct*. 2006 Sep;17(5):512-9. X-2
304. Cardozo L, Chapple CR, Tooze-Hobson P, et al. Efficacy of trospium chloride in patients with detrusor instability: a placebo-controlled, randomized, double-blind, multicentre clinical trial. *BJU Int*. 2000 Apr;85(6):659-64. X-4
305. Cardozo L, Coyne KS and Versi E. Validation of the urgency perception scale. *BJU Int*. 2005 Mar;95(4):591-6. X-1I
306. Cardozo L and Dixon A. Increased warning time with darifenacin: a new concept in the management of urinary urgency. *J Urol*. 2005 Apr;173(4):1214-8. X-4
307. Cardozo L, Hextall A, Bailey J, et al. Colposuspension after previous failed incontinence surgery: a prospective observational study. *Br J Obstet Gynaecol*. 1999 Apr;106(4):340-4. X-1B
308. Cardozo L and Robinson D. Special considerations in premenopausal and postmenopausal women with symptoms of overactive bladder. *Urology*. 2002 Nov;60(5 Suppl 1):64-71; discussion 71. X-2
309. Cardozo LD, Abrams PD, Stanton SL, et al. Idiopathic bladder instability treated by biofeedback. *Br J Urol*. 1978 Dec;50(7):521-3. X-7
310. Cardozo LD and Stanton SL. An objective comparison of the effects of parenterally administered drugs in patients suffering from detrusor instability. *J Urol*. 1979 Jul;122(1):58-9. X-7
311. Cardozo LD and Stanton SL. A comparison between bromocriptine and indomethacin in the treatment of detrusor instability. *J Urol*. 1980 Mar;123(3):399-401. X-1F, X-7
312. Cardozo LD and Stanton SL. Genuine stress incontinence and detrusor instability--a review of 200 patients. *Br J Obstet Gynaecol*. 1980 Mar;87(3):184-90. X-7
313. Cardozo LD, Stanton SL, Robinson H, et al. Evaluation of flurbiprofen in detrusor instability. *Br Med J*. 1980 Feb 2;280(6210):281-2. X-7
314. Cardozo LD, Stanton SL and Williams JE. Detrusor instability following surgery for genuine stress incontinence. *Br J Urol*. 1979 Jun;51(3):204-7. X-1B
315. Carey JM, Chon JK and Leach GE. Urethrolisis with Martius labial fat pad graft for iatrogenic bladder outlet obstruction. *Urology*. 2003 Apr;61(4 Suppl 1):21-5. X-1A
316. Carey M, Fynes M, Murray C, et al. Sacral nerve root stimulation for lower urinary tract dysfunction: overcoming the problem of lead migration. *BJU Int*. 2001 Jan;87(1):15-8. X-7
317. Carey MP, De Jong S, Friedhuber A, et al. A prospective evaluation of the pathogenesis of detrusor instability in women, using electron microscopy and immunohistochemistry. *BJU Int*. 2000 Dec;86(9):970-6. X-1G, X-7
318. Carey MP, Goh JT, Fynes MM, et al. Stress urinary incontinence after delayed primary closure of genitourinary fistula: a technique for surgical management. *Am J Obstet Gynecol*. 2002 May;186(5):948-53. X-1B
319. Carlson KV, Rome S and Nitti VW. Dysfunctional voiding in women. *J Urol*. 2001 Jan;165(1):143-7; discussion 147-8. X-1H, X-7
320. Carr LK, Walsh PJ, Abraham VE, et al. Favorable outcome of pubovaginal slings for geriatric women with stress incontinence. *J Urol*. 1997 Jan;157(1):125-8. X-1B
321. Carr LK and Webster GD. Voiding dysfunction following incontinence surgery: diagnosis and treatment with retropubic or vaginal urethrolisis. *J Urol*. 1997 Mar;157(3):821-3. X-1A
322. Carrico DJ, Peters KM and Diokno AC. Guided imagery for women with interstitial cystitis: results of a prospective, randomized controlled pilot study. *J Altern Complement Med*. 2008 Jan-Feb;14(1):53-60. X-1D
323. Carter JE. Laparoscopic bladder neck suspension. *Endosc Surg Allied Technol*. 1995 Apr-Jun;3(2-3):81-7. X-1B
324. Cartwright R, Elvy S and Cardozo L. Do women with female ejaculation have detrusor overactivity? *J Sex Med*. 2007 Nov;4(6):1655-8. X-1A, X-7
325. Castera R, Podesta ML, Ruarte A, et al. 10-Year experience with artificial urinary sphincter in children and adolescents. *J Urol*. 2001 Jun;165(6 Pt 2):2373-6. X-1I, X-4, X-7

326. Castleden CM, Duffin HM, Asher MJ, et al. Factors influencing outcome in elderly patients with urinary incontinence and detrusor instability. *Age Ageing*. 1985 Sep;14(5):303-7. X-4
327. Castleden CM, Duffin HM and Clarkson EJ. In vivo and in vitro studies on the effect of sodium antagonists on the bladder in man and rat. *Age Ageing*. 1983 Aug;12(3):249-55. X-7
328. Castro NM, Freitas DM, Rodrigues W, Jr., et al. Urodynamic features of the voiding dysfunction in HTLV-1 infected individuals. *Int Braz J Urol*. 2007 Mar-Apr;33(2):238-44; discussion 244-5. X-1F
329. Cawley DM and Hendriks O. Evaluation of the Endomed CV 405 as a treatment for urinary incontinence. *Physiotherapy*. 1992 Jul;78(7):495-8. X-7
330. Centonze D, Petta F, Versace V, et al. Effects of motor cortex rTMS on lower urinary tract dysfunction in multiple sclerosis. *Mult Scler*. 2007 Mar;13(2):269-71. X-1F
331. Cespedes RD. Anterior approach bilateral sacrospinous ligament fixation for vaginal vault prolapse. *Urology*. 2000 Dec 4;56(6 Suppl 1):70-5. X-1E
332. Cespedes RD, Cross CA and McGuire EJ. Modified Ingelman-Sundberg bladder denervation procedure for intractable urge incontinence. *J Urol*. 1996 Nov;156(5):1744-7. X-7
333. Cetinel B, Akpınar H, Tufek I, et al. Bladder involvement in Behcet's syndrome. *J Urol*. 1999 Jan;161(1):52-6. X-1I
334. Cetinel B, Demirkesen O, Onal B, et al. Are there any factors predicting the cure and complication rates of tension-free vaginal tape? *Int Urogynecol J Pelvic Floor Dysfunct*. 2004 May-Jun;15(3):188-93. X-1B
335. Cetinel B, Demirkesen O, Tarcan T, et al. Hidden female urinary incontinence in urology and obstetrics and gynecology outpatient clinics in Turkey: what are the determinants of bothersome urinary incontinence and help-seeking behavior? *Int Urogynecol J Pelvic Floor Dysfunct*. 2007 Jun;18(6):659-64. X-6
336. Chai TC, Gray ML and Steers WD. The incidence of a positive ice water test in bladder outlet obstructed patients: evidence for bladder neural plasticity. *J Urol*. 1998 Jul;160(1):34-8. X-1A, X-1H
337. Chai TC, Zhang C, Warren JW, et al. Percutaneous sacral third nerve root neurostimulation improves symptoms and normalizes urinary HB-EGF levels and antiproliferative activity in patients with interstitial cystitis. *Urology*. 2000 May;55(5):643-6. X-1D, X-7
338. Chaikin DC, Blaivas JG and Blaivas ST. Behavioral therapy for the treatment of refractory interstitial cystitis. *J Urol*. 1993 Jun;149(6):1445-8. X-1A, X-1D
339. Chaikin DC, Blaivas JG, Rosenthal JE, et al. Results of pubovaginal sling for stress incontinence: a prospective comparison of 4 instruments for outcome analysis. *J Urol*. 1999 Nov;162(5):1670-3. X-1B
340. Chaikin DC, Groutz A and Blaivas JG. Predicting the need for anti-incontinence surgery in continent women undergoing repair of severe urogenital prolapse. *J Urol*. 2000 Feb;163(2):531-4. X-1E
341. Chaikin DC, Rosenthal J and Blaivas JG. Pubovaginal fascial sling for all types of stress urinary incontinence: long-term analysis. *J Urol*. 1998 Oct;160(4):1312-6. X-1B
342. Chaisaeng S, Santingamkun A, Opanuraks J, et al. IQOL: translation & reliability for use with urinary incontinence patients in Thailand. *J Med Assoc Thai*. 2006 Sep;89 Suppl 3:S33-9. X-6
343. Chaliha C, Bland JM, Monga A, et al. Pregnancy and delivery: a urodynamic viewpoint. *BJOG*. 2000 Nov;107(11):1354-9. X-1B
344. Chaliha C, Digesu GA, Hutchings A, et al. Changes in urethral function with bladder filling in the presence of urodynamic stress incontinence and detrusor overactivity. *Am J Obstet Gynecol*. 2005 Jan;192(1):60-5. X-1G, X-1H
345. Chaliha C, Digesu GA, Salvatore S, et al. Changes in urethral resistance in the presence of detrusor activity. *Int Urogynecol J Pelvic Floor Dysfunct*. 2006 May;17(3):215-8. X-1G, X-7
346. Chan CL, Scott SM, Williams NS, et al. Rectal hypersensitivity worsens stool frequency, urgency, and lifestyle in patients with urge fecal incontinence. *Dis Colon Rectum*. 2005 Jan;48(1):134-40. X-1I
347. Chan KM, Lim PH, Chee YC, et al. Urinary symptoms and urodynamic diagnosis of patients in one geriatric department. *Ann Acad Med Singapore*. 1992 Nov;21(6):813-7. X-1H
348. Chandi DD, Groenendijk PM and Venema PL. Functional extracorporeal magnetic stimulation as a treatment for female urinary incontinence: 'the chair'. *BJU Int*. 2004 Mar;93(4):539-42. X-7
349. Chandiramani VA, Peterson T, Duthie GS, et al. Urodynamic changes during therapeutic intravesical instillations of capsaicin. *Br J Urol*. 1996 Jun;77(6):792-7. X-7
350. Chandra M, Saharia R, Hill V, et al. Prevalence of diurnal voiding symptoms and difficult arousal from sleep in children with nocturnal enuresis. *J Urol*. 2004 Jul;172(1):311-6. X-1C, X-4
351. Chandra M, Saharia R, Shi Q, et al. Giggle incontinence in children: a manifestation of detrusor instability. *J Urol*. 2002 Nov;168(5):2184-7; discussion 2187. X-4, X-6
352. Chapple C, Steers W, Norton P, et al. A pooled analysis of three phase III studies to investigate the efficacy, tolerability and safety of darifenacin, a muscarinic M3 selective receptor antagonist, in the treatment of overactive bladder. *BJU Int*. 2005 May;95(7):993-1001. X-2
353. Chapple CR and Abrams P. Comparison of darifenacin and oxybutynin in patients with overactive bladder: assessment of ambulatory urodynamics and impact on salivary flow. *Eur Urol*. 2005 Jul;48(1):102-9. X-4
354. Chapple CR, Arano P, Bosch JL, et al. Solifenacin appears effective and well tolerated in patients with symptomatic idiopathic detrusor overactivity in a placebo- and tolterodine-controlled phase 2 dose-finding study. *BJU Int*. 2004 Jan;93(1):71-7. X-4
355. Chapple CR, Parkhouse H, Gardener C, et al. Double-blind, placebo-controlled, cross-over study of flavoxate in the treatment of idiopathic detrusor instability. *Br J Urol*. 1990 Nov;66(5):491-4. X-4, X-7
356. Chapple CR, Patroneva A and Raines SR. Effect of an ATP-sensitive potassium channel opener in subjects with overactive bladder: a randomized, double-blind, placebo-

- controlled study (ZD0947IL/0004). *Eur Urol*. 2006 May;49(5):879-86. X-4
357. Chartier-Kastler EJ, Denys P, Chancellor MB, et al. Urodynamic monitoring during percutaneous sacral nerve neurostimulation in patients with neurogenic detrusor hyperreflexia. *Neurourol Urodyn*. 2001;20(1):61-71. X-1F
358. Chartier-Kastler EJ, Mongiat-Artus P, Bitker MO, et al. Long-term results of augmentation cystoplasty in spinal cord injury patients. *Spinal Cord*. 2000 Aug;38(8):490-4. X-1F
359. Chartier-Kastler EJ, Ruud Bosch JL, Perrigot M, et al. Long-term results of sacral nerve stimulation (S3) for the treatment of neurogenic refractory urge incontinence related to detrusor hyperreflexia. *J Urol*. 2000 Nov;164(5):1476-80. X-1F
360. Cheater FM, Baker R, Gillies C, et al. The nature and impact of urinary incontinence experienced by patients receiving community nursing services: a cross-sectional cohort study. *Int J Nurs Stud*. 2008 Mar;45(3):339-51. X-II
361. Chen CS and Cheng CL. Nephrogenic adenoma of the urinary bladder: clinical experience and review of the literature. *J Chin Med Assoc*. 2006 Apr;69(4):166-8. X-II, X-2
362. Chen CW, Huang KH, Liang HM, et al. A comparison of urodynamic parameters and lower urinary tract symptoms in urodynamic genuine stress incontinence women with or without stress urinary incontinence. *Chang Gung Med J*. 2004 Aug;27(8):594-601. X-1B, X-1H
363. Chen GD, Hu SW, Chen YC, et al. Prevalence and correlations of anal incontinence and constipation in Taiwanese women. *Neurourol Urodyn*. 2003;22(7):664-9. X-1A
364. Chen GD, Lin LY and Gardner JD. The results of laparoscopic adhesiolysis for intractable urinary frequency. *J Urol*. 1997 Nov;158(5):1714-6. X-1A, X-7
365. Chen GD, Lin LY, Wang PH, et al. Urinary tract dysfunction after radical hysterectomy for cervical cancer. *Gynecol Oncol*. 2002 May;85(2):292-7. X-1A
366. Chen YH and Kuo HC. Botulinum A toxin treatment of urethral sphincter pseudodyssynergia in patients with cerebrovascular accidents or intracranial lesions. *Urol Int*. 2004;73(2):156-61; discussion 161-2. X-1F, X-7
367. Chene G, Amblard J, Tardieu AS, et al. Long-term results of tension-free vaginal tape (TVT) for the treatment of female urinary stress incontinence. *Eur J Obstet Gynecol Reprod Biol*. 2007 Sep;134(1):87-94. X-1B
368. Chene G, Cotte B, Tardieu AS, et al. Clinical and ultrasonographic correlations following three surgical anti-incontinence procedures (TOT, TVT and TVT-O). *Int Urogynecol J Pelvic Floor Dysfunct*. 2008 Aug;19(8):1125-31. X-1B
369. Cheon WC, Mak JH and Liu JY. Prospective randomised controlled trial comparing laparoscopic and open colposuspension. *Hong Kong Med J*. 2003 Feb;9(1):10-4. X-1B
370. Chertin B, Koulikov D, Abu-Arafah W, et al. Treatment of nocturnal enuresis in children with attention deficit hyperactivity disorder. *J Urol*. 2007 Oct;178(4 Pt 2):1744-7. X-II, X-4
371. Cheshire WP, Tsuboi Y and Wszolek ZK. Physiologic assessment of autonomic dysfunction in pallidopontonigral degeneration with N279K mutation in the tau gene on chromosome 17. *Auton Neurosci*. 2002 Nov 29;102(1-2):71-7. X-1A
372. Chiaffarino F, Parazzini F, Lavezzari M, et al. Impact of urinary incontinence and overactive bladder on quality of life. *Eur Urol*. 2003 May;43(5):535-8. X-II
373. Chiara G, Piccioni V, Perino M, et al. Psychological investigation in female patients suffering from urinary incontinence. *Int Urogynecol J Pelvic Floor Dysfunct*. 1998;9(2):73-7. X-II
374. Chiarelli P. Urinary stress incontinence and overactive bladder symptoms in older women. *Contemp Nurse*. 2007 Oct;26(2):198-207. X-1H, X-2
375. Chin YK and Stanton SL. A follow up of silastic sling for genuine stress incontinence. *Br J Obstet Gynaecol*. 1995 Feb;102(2):143-7. X-1B
376. Chiu NC, Huang FY and Tsai TC. Urinary tract infections in children. *Zhonghua Min Guo Xiao Er Ke Yi Xue Hui Za Zhi*. 1989 Jul-Aug;30(4):225-32. X-II, X-4
377. Choe JH, Choo MS and Lee KS. Symptom change in women with overactive bladder after extracorporeal magnetic stimulation: a prospective trial. *Int Urogynecol J Pelvic Floor Dysfunct*. 2007 Aug;18(8):875-80. X-7
378. Choe JH, Choo MS and Lee KS. The impact of tension-free vaginal tape on overactive bladder symptoms in women with stress urinary incontinence: significance of detrusor overactivity. *J Urol*. 2008 Jan;179(1):214-9. X-1B
379. Choe JM. Preventing urethral obstruction using the 6-point fixation and weight-adjusted spacing nomogram during sling surgery. *Int Urogynecol J Pelvic Floor Dysfunct*. 2001;12(2):122-8. X-1B
380. Choe JM. Suprapubic sling adjustment: minimally invasive method of curing recurrent stress incontinence after sling surgery. *J Urol*. 2002 Nov;168(5):2059-62. X-1B
381. Choe JM, Gallo ML and Staskin DR. A provocative maneuver to elicit cystometric instability: measuring instability at maximum infusion. *J Urol*. 1999 May;161(5):1541-4. X-1H
382. Choe JM, Ogan K and Battino BS. Antimicrobial mesh versus vaginal wall sling: a comparative outcomes analysis. *J Urol*. 2000 Jun;163(6):1829-34. X-1B
383. Choe JM, Ogan K and Bennett S. Antibacterial mesh sling: a prospective outcome analysis. *Urology*. 2000 Apr;55(4):515-20. X-1B
384. Choe JM and Staskin DR. Gore-Tex patch sling: 7 years later. *Urology*. 1999 Oct;54(4):641-6. X-1B
385. Choi JB, Kim YB, Kim BT, et al. Analysis of heart rate variability in female patients with overactive bladder. *Urology*. 2005 Jun;65(6):1109-12; discussion 1113. X-1G
386. Cholhan HJ and Lotze PM. Voiding function after a modified no-tension pubovaginal sling. *Int Urogynecol J Pelvic Floor Dysfunct*. 2004 Jul-Aug;15(4):249-56. X-1B
387. Cholhan HJ and Lotze PM. Urodynamic changes after tension-free sling procedures: Mycromesh-Plus vs TVT sling. *Int Urogynecol J Pelvic Floor Dysfunct*. 2008 Feb;19(2):217-25. X-1B
388. Chompootaweep S, Nunthapisud P, Trivijitsilp P, et al. The use of two estrogen preparations (a combined contraceptive pill versus conjugated estrogen cream) intravaginally to treat urogenital symptoms in postmenopausal Thai women: a comparative study. *Clin Pharmacol Ther*. 1998 Aug;64(2):204-10. X-II, X-7

389. Choo MS, Song C, Kim JH, et al. Changes in overactive bladder symptoms after discontinuation of successful 3-month treatment with an antimuscarinic agent: a prospective trial. *J Urol*. 2005 Jul;174(1):201-4. X-9
390. Chou EC, Blaivas JG, Chou LW, et al. Urodynamic characteristics of mixed urinary incontinence and idiopathic urge urinary incontinence. *Neurourol Urodyn*. 2008;27(5):376-8. X-1H
391. Chou EC, Flisser AJ, Panagopoulos G, et al. Effective treatment for mixed urinary incontinence with a pubovaginal sling. *J Urol*. 2003 Aug;170(2 Pt 1):494-7. X-1B
392. Chou TP, Gorton E, Stanton SL, et al. Can uroflowmetry patterns in women be reliably interpreted? *Int Urogynecol J Pelvic Floor Dysfunct*. 2000 Jun;11(3):142-7. X-6, X-1H
393. Christman NJ, Oakley MG and Cronin SN. Developing and using preparatory information for women undergoing radiation therapy for cervical or uterine cancer. *Oncol Nurs Forum*. 2001 Jan-Feb;28(1):93-8. X-11
394. Christmas TJ, Smith GL and Rickards D. Diagnosis and management of high-pressure peristaltic contractions in cystoplasties. *Br J Urol*. 1997 Jun;79(6):879-82. X-1A, X-7
395. Christoph F, Moschkowitsch A, Kempkensteffen C, et al. Long-term efficacy of tolterodine and patient compliance in pediatric patients with neurogenic detrusor overactivity. *Urol Int*. 2007;79(1):55-9. X-4
396. Chu PS, Ma WK, Wong SC, et al. The destruction of the lower urinary tract by ketamine abuse: a new syndrome? *BJU Int*. 2008 Dec;102(11):1616-22. X-11
397. Chuang YC, Thomas CA, Tyagi S, et al. Human urine with solifenacin intake but not tolterodine or darifenacin intake blocks detrusor overactivity. *Int Urogynecol J Pelvic Floor Dysfunct*. 2008 Oct;19(10):1353-7. X-11
398. Chung JM, Lee SD, Kang DI, et al. Prevalence and associated factors of overactive bladder in Korean children 5-13 years old: a nationwide multicenter study. *Urology*. 2009 Jan;73(1):63-7; discussion 68-9. X-4
399. Chung SY, Franks M, Smith CP, et al. Technique of combined pubovaginal sling and cystocele repair using a single piece of cadaveric dermal graft. *Urology*. 2002 Apr;59(4):538-41. X-1B
400. Cindolo L, Salzano L, Rota G, et al. Tension-free transobturator approach for female stress urinary incontinence. *Minerva Urol Nefrol*. 2004 Mar;56(1):89-98. X-1B
401. Clarke B. Urethral instability. *Aust N Z J Obstet Gynaecol*. 1992 Aug;32(3):270-5. X-1H, X-11
402. Clarke B. The role of urodynamic assessment in the diagnosis of lower urinary tract disorders. *Int Urogynecol J Pelvic Floor Dysfunct*. 1997;8(4):196-9. X-1H
403. Clarke SJ, Forster DM and Thomas DG. Selective sacral neurectomy in the management of urinary incontinence due to detrusor instability. *Br J Urol*. 1979 Dec;51(6):510-4. X-1F, X-7
404. Clauw DJ, Schmidt M, Radulovic D, et al. The relationship between fibromyalgia and interstitial cystitis. *J Psychiatr Res*. 1997 Jan-Feb;31(1):125-31. X-1D
405. Clemens JQ, Stern JA, Bushman WA, et al. Long-term results of the Stamey bladder neck suspension: direct comparison with the Marshall-Marchetti-Krantz procedure. *J Urol*. 1998 Aug;160(2):372-6. X-1B
406. Clemons JL, Aguilar VC, Sokol ER, et al. Suburethral sling treatment of occult stress incontinence and intrinsic sphincter deficiency in women with severe vaginal prolapse of the anterior vs posterior/apical compartment. *Am J Obstet Gynecol*. 2005 May;192(5):1566-72. X-1B
407. Clemons JL, Aguilar VC, Tillinghast TA, et al. Patient satisfaction and changes in prolapse and urinary symptoms in women who were fitted successfully with a pessary for pelvic organ prolapse. *Am J Obstet Gynecol*. 2004 Apr;190(4):1025-9. X-1E
408. Cliff AM, Heatherwick B, Scoble J, et al. The effect of fasting or desmopressin before treatment on the concentration of mitomycin C during intravesical administration. *BJU Int*. 2000 Oct;86(6):644-7. X-11, X-7
409. Close CE, Carr MC, Burns MW, et al. Interstitial cystitis in children. *J Urol*. 1996 Aug;156(2 Pt 2):860-2. X-1D, X-4, X-7
410. Clow WM. Effect of posture on bladder and urethral function in normal pregnancy. A preliminary report. *Urol Int*. 1975;30(1):9-15. X-1B, X-1H, X-7
411. Clyne OJ, O'Sullivan O and Flood HD. Pubovaginal sling for urodynamic stress incontinence: effect on patient quality of life. *Ir Med J*. 2005 Mar;98(3):75-7. X-1B
412. Coates KW, Harris RL, Cundiff GW, et al. Uroflowmetry in women with urinary incontinence and pelvic organ prolapse. *Br J Urol*. 1997 Aug;80(2):217-21. X-1E, X-1F
413. Cohen BL, Barboglio P and Gousse A. The impact of lower urinary tract symptoms and urinary incontinence on female sexual dysfunction using a validated instrument. *J Sex Med*. 2008 Jun;5(6):1418-23. X-11
414. Cohen BL, Rivera R, Barboglio P, et al. Safety and tolerability of sedation-free flexible cystoscopy for intradetrusor botulinum toxin-A injection. *J Urol*. 2007 Mar;177(3):1006-10; discussion 1010. X-7
415. Cohen BL, Tunuguntla HS and Gousse A. Predictors of success for first stage neuromodulation: motor versus sensory response. *J Urol*. 2006 Jun;175(6):2178-80; discussion 2180-1. X-7
416. Cohen HA, Nussinovitch M, Kauschansky A, et al. Extraordinary daytime urinary frequency in children. *J Fam Pract*. 1993 Jul;37(1):28-9. X-4, X-6, X-7
417. Cole EE, Kaufman MR, Scarpero HM, et al. The effects of isolated posterior compartment defects on lower urinary tract symptoms and urodynamic findings. *BJU Int*. 2006 May;97(5):1024-6. X-11, X-7
418. Cole EE, Scarpero HM and Dmochowski RR. Are patient symptoms predictive of the diagnostic and/or therapeutic value of hydrodistention? *Neurourol Urodyn*. 2005;24(7):638-42. X-7
419. Collas DM and Malone-Lee JG. Age-associated changes in detrusor sensory function in women with lower urinary tract symptoms. *Int Urogynecol J Pelvic Floor Dysfunct*. 1996;7(1):24-9. X-1H
420. Collinet P, Ciofu C, Costa P, et al. The safety of the inside-out transobturator approach for transvaginal tape (TVT-O) treatment in stress urinary incontinence: French registry data on 984 women. *Int Urogynecol J Pelvic Floor Dysfunct*. 2008 May;19(5):711-5. X-1B
421. Colman S, Chapple C, Nitti V, et al. Validation of treatment benefit scale for assessing subjective outcomes in

- treatment of overactive bladder. *Urology*. 2008 Oct;72(4):803-7. X-1I
422. Colombo M, Maggioni A, Zanetta G, et al. Prevention of postoperative urinary stress incontinence after surgery for genitourinary prolapse. *Obstet Gynecol*. 1996 Feb;87(2):266-71. X-1B
423. Colombo M, Scalabrino S, Maggioni A, et al. Burch colposuspension versus modified Marshall-Marchetti-Krantz urethropexy for primary genuine stress urinary incontinence: a prospective, randomized clinical trial. *Am J Obstet Gynecol*. 1994 Dec;171(6):1573-9. X-1B
424. Colombo M, Zanetta G, Vitobello D, et al. The Burch colposuspension for women with and without detrusor overactivity. *Br J Obstet Gynaecol*. 1996 Mar;103(3):255-60. X-1B
425. Combs AJ and Nitti VW. Significance of rectal contractions noted on multichannel urodynamics. *Neurourol Urodyn*. 1995;14(1):73-80. X-1A, X-1H
426. Comperat E, Reitz A, Delcourt A, et al. Histologic features in the urinary bladder wall affected from neurogenic overactivity--a comparison of inflammation, oedema and fibrosis with and without injection of botulinum toxin type A. *Eur Urol*. 2006 Nov;50(5):1058-64. X-1F, X-1G, X-7
427. Congregado Ruiz B, Pena Outeirino XM, Campoy Martinez P, et al. Peripheral afferent nerve stimulation for treatment of lower urinary tract irritative symptoms. *Eur Urol*. 2004 Jan;45(1):65-9. X-7
428. Constantinou CE. Pharmacologic treatment of detrusor incontinence with thiphenamil HCl. *Urol Int*. 1992;48(1):42-7. X-7
429. Contreras Ortiz O, Lombardo RJ and Pellicari A. Non-invasive diagnosis of bladder instability using the Bladder Instability Discriminant Index (BIDI). *Zentralbl Gynakol*. 1993;115(10):446-9. X-1D
430. Cooper MJ, Cario G, Lam A, et al. A review of results in a series of 113 laparoscopic colposuspensions. *Aust N Z J Obstet Gynaecol*. 1996 Feb;36(1):44-8. X-1B
431. Corigliano T, Renella R, Robbiani A, et al. Isolated extraordinary daytime urinary frequency of childhood: a case series of 26 children in Switzerland. *Acta Paediatr*. 2007 Sep;96(9):1347-9. X-1I, X-4
432. Cortesse A, Jacquetin B, Grise P, et al. Prospective multicenter clinical trial of Uretex Sup for surgical treatment of stress urinary incontinence. *Int J Urol*. 2007 Jul;14(7):611-5. X-1B
433. Cosimo O, Pierluigi P, Angelo ZM, et al. A clinical and urodynamic study of patients with varying degrees of cystocele. *Maturitas*. 1997 Jun;27(2):125-32. X-1I
434. Costantini E, Lazzeri M, Bini V, et al. Sensitivity and specificity of one-hour pad test as a predictive value for female urinary incontinence. *Urol Int*. 2008;81(2):153-9. X-1H
435. Costantini E, Mearini L, Mearini E, et al. Assessing outcome after a modified vaginal wall sling for stress incontinence with intrinsic sphincter deficiency. *Int Urogynecol J Pelvic Floor Dysfunct*. 2005 Mar-Apr;16(2):138-46; discussion 146. X-1B
436. Costello AJ, Crowe H and Agarwal D. Supratrigonal cystectomy and ileocystoplasty in management of interstitial cystitis. *Aust N Z J Surg*. 2000 Jan;70(1):34-8. X-1D
437. Cowan JA, McGirt MJ, Woodworth G, et al. The syndrome of hydrocephalus in young and middle-aged adults (SHYMA). *Neurol Res*. 2005 Jul;27(5):540-7. X-1I
438. Coyne K, Revicki D, Hunt T, et al. Psychometric validation of an overactive bladder symptom and health-related quality of life questionnaire: the OAB-q. *Qual Life Res*. 2002 Sep;11(6):563-74. X-1I
439. Coyne KS, Elinoff V, Gordon DA, et al. Relationships between improvements in symptoms and patient assessments of bladder condition, symptom bother and health-related quality of life in patients with overactive bladder treated with tolterodine. *Int J Clin Pract*. 2008 Jun;62(6):925-31. X-1I
440. Coyne KS, Margolis MK, Brewster-Jordan J, et al. Evaluating the impact of overactive bladder on sexual health in women: what is relevant? *J Sex Med*. 2007 Jan;4(1):124-36. X-1I
441. Coyne KS, Margolis MK, Jumadilova Z, et al. Overactive bladder and women's sexual health: what is the impact? *J Sex Med*. 2007 May;4(3):656-66. X-7
442. Coyne KS, Matza LS, Kopp Z, et al. The validation of the patient perception of bladder condition (PPBC): a single-item global measure for patients with overactive bladder. *Eur Urol*. 2006 Jun;49(6):1079-86. X-1I
443. Coyne KS, Matza LS, Thompson C, et al. The responsiveness of the OAB-q among OAB patient subgroups. *Neurourol Urodyn*. 2007;26(2):196-203. X-1I
444. Coyne KS, Matza LS and Thompson CL. The responsiveness of the Overactive Bladder Questionnaire (OAB-q). *Qual Life Res*. 2005 Apr;14(3):849-55. X-1I
445. Coyne KS, Matza LS, Thompson CL, et al. Determining the importance of change in the overactive bladder questionnaire. *J Urol*. 2006 Aug;176(2):627-32; discussion 632. X-2
446. Coyne KS, Payne C, Bhattacharyya SK, et al. The impact of urinary urgency and frequency on health-related quality of life in overactive bladder: results from a national community survey. *Value Health*. 2004 Jul-Aug;7(4):455-63. X-1I
447. Coyne KS, Sexton CC, Irwin DE, et al. The impact of overactive bladder, incontinence and other lower urinary tract symptoms on quality of life, work productivity, sexuality and emotional well-being in men and women: results from the EPIC study. *BJU Int*. 2008 Jun;101(11):1388-95. X-1I
448. Coyne KS, Zhou Z, Bhattacharyya SK, et al. The prevalence of nocturia and its effect on health-related quality of life and sleep in a community sample in the USA. *BJU Int*. 2003 Dec;92(9):948-54. X-1I
449. Coyne KS, Zhou Z, Thompson C, et al. The impact on health-related quality of life of stress, urge and mixed urinary incontinence. *BJU Int*. 2003 Nov;92(7):731-5. X-1A
450. Coyne KS, Zyczynski T, Margolis MK, et al. Validation of an overactive bladder awareness tool for use in primary care settings. *Adv Ther*. 2005 Jul-Aug;22(4):381-94. X-1I
451. Cozzi F, Schiavetti A, Zani A, et al. The functional sequelae of sacrococcygeal teratoma: a longitudinal and cross-sectional follow-up study. *J Pediatr Surg*. 2008 Apr;43(4):658-61. X-1I, X-4

452. Craggs MD. Objective measurement of bladder sensation: use of a new patient-activated device and response to neuromodulation. *BJU Int.* 2005 Sep;96 Suppl 1:29-36. X-1H, X-2
453. Creighton SM, Pearce JM, Robson I, et al. Sensory urgency: how full is your bladder? *Br J Obstet Gynaecol.* 1991 Dec;98(12):1287-9. X-7
454. Creighton SM, Plevnik S and Stanton SL. Distal urethral electrical conductance (DUEC)--a preliminary assessment of its role as a quick screening test for incontinent women. *Br J Obstet Gynaecol.* 1991 Jan;98(1):69-72. X-1H
455. Creighton SM and Stanton SL. Caffeine: does it affect your bladder? *Br J Urol.* 1990 Dec;66(6):613-4. X-1H, X-7
456. Crisp JC, Green NA and Ashken MH. Urodynamic studies in the district general hospital. *Br J Urol.* 1976 Oct;48(5):383-7. X-1G, X-1H
457. Cronje HS, De Beer JA and Bam R. The pathophysiology of an enterocele and its management. *J Obstet Gynaecol.* 2004 Jun;24(4):408-13. X-1A
458. Crooks JE, Khan AB, Meddings RN, et al. Bladder transection revisited. *Br J Urol.* 1995 May;75(5):590-1. X-7
459. Cross CA, Cespedes RD, English SF, et al. Transvaginal urethrolysis for urethral obstruction after anti-incontinence surgery. *J Urol.* 1998 Apr;159(4):1199-201. X-1A
460. Cross CA, Cespedes RD and McGuire EJ. Our experience with pubovaginal slings in patients with stress urinary incontinence. *J Urol.* 1998 Apr;159(4):1195-8. X-1B
461. Cucchi A. Acceleration of flow rate as a screening test for detrusor instability in women with stress incontinence. *Br J Urol.* 1990 Jan;65(1):17-9. X-1H
462. Cucchi A. Bladder contractility and idiopathic detrusor instability in the female. *Br J Urol.* 1993 Nov;72(5 Pt 1):559-65. X-1G, X-1H
463. Cucchi A. A possible link between stress urinary incontinence and detrusor instability in the female--urodynamic (pressure/flow) data and speculative considerations. *Int Urogynecol J Pelvic Floor Dysfunct.* 1998;9(1):3-8. X-1B
464. Cucchi A. Sequential changes in voiding dynamics related to the development of detrusor instability in women with stress urinary incontinence. *Neurourol Urodyn.* 1999;18(2):73-80. X-1B, X-7
465. Cucchi A, Quaglini S, Siracusano S, et al. Urgency degree and bladder contraction velocity: sequential changes in women with idiopathic detrusor overactivity. *Neurourol Urodyn.* 2006;25(2):123-7. X-1H
466. Cucchi A, Siracusano S, Di Benedetto P, et al. Urgency of voiding and abdominal pressure transmission in women with mixed urinary incontinence. *Neurourol Urodyn.* 2004;23(1):43-7. X-1B
467. Cucchi A, Siracusano S, Guarnaschelli C, et al. Voiding urgency and detrusor contractility in women with overactive bladders. *Neurourol Urodyn.* 2003;22(3):223-6. X-1G
468. Cukier JM, Cortina-Borja M and Brading AF. A case-control study to examine any association between idiopathic detrusor instability and gastrointestinal tract disorder, and between irritable bowel syndrome and urinary tract disorder. *Br J Urol.* 1997 Jun;79(6):865-78. X-1I
469. Cundiff GW, Harris RL, Coates KW, et al. Clinical predictors of urinary incontinence in women. *Am J Obstet Gynecol.* 1997 Aug;177(2):262-6; discussion 266-7. X-1H
470. Curran MJ, Kaefer M, Peters C, et al. The overactive bladder in childhood: long-term results with conservative management. *J Urol.* 2000 Feb;163(2):574-7. X-4, X-7
471. Currie CJ, McEwan P, Poole CD, et al. The impact of the overactive bladder on health-related utility and quality of life. *BJU Int.* 2006 Jun;97(6):1267-72. X-1I
472. Dainer MJ and Zorn BH. Burch-Cooper's ligament sling procedure: an augmented incontinence operation. *Tech Urol.* 2000 Sep;6(3):175-7. X-1A, X-7
473. Dallosso HM, McGrother CW, Matthews RJ, et al. The association of diet and other lifestyle factors with overactive bladder and stress incontinence: a longitudinal study in women. *BJU Int.* 2003 Jul;92(1):69-77. X-1A
474. Dalpiaz O, Primus G and Schips L. SPARC sling system for treatment of female stress urinary incontinence in the elderly. *Eur Urol.* 2006 Oct;50(4):826-30; discussion 830-1. X-1B
475. Darai E, Frobert JL, Grisard-Anaf M, et al. Functional results after the suburethral sling procedure for urinary stress incontinence: a prospective randomized multicentre study comparing the retropubic and transobturator routes. *Eur Urol.* 2007 Mar;51(3):795-801; discussion 801-2. X-1B
476. Darai E, Perdu M, Benifla JL, et al. Percutaneous needle colposuspension to Cooper's ligament for the treatment of stress incontinence in women: a report of 82 cases. *Eur J Obstet Gynecol Reprod Biol.* 1997 Jul;74(1):53-5. X-1B
477. Darkow T, Fontes CL and Williamson TE. Costs associated with the management of overactive bladder and related comorbidities. *Pharmacotherapy.* 2005 Apr;25(4):511-9. X-1I
478. Darkow T, Fontes CL, Williamson TE, et al. Drug use issues and actions Non-elderly patients with overactive bladder have significant comorbidities. *Formulary.* 2005 Aug;40(8):255-6. X-1I
479. Darling R and Neilson D. Convenience voids: an important new factor in urinary frequency volume chart analysis. *J Urol.* 2005 Feb;173(2):487-9. X-1A
480. Das AK, Carlson AM and Hull M. Improvement in depression and health-related quality of life after sacral nerve stimulation therapy for treatment of voiding dysfunction. *Urology.* 2004 Jul;64(1):62-8. X-6
481. Das S. Dynamic suburethral suspension with pedicled external oblique aponeurosis in the management of female urinary incontinence. *J Urol.* 1999 Aug;162(2):469-73. X-1A, X-7
482. Dattilo J. A long-term study of patient outcomes with pelvic muscle re-education for urinary incontinence. *J Wound Ostomy Continence Nurs.* 2001 Jul;28(4):199-205. X-7
483. Davies MC, Crouch NS, Woodhouse CR, et al. Congenital adrenal hyperplasia and lower urinary tract symptoms. *BJU Int.* 2005 Jun;95(9):1263-6. X-1I, X-7
484. Davila GW, Bernier F, Franco J, et al. Bladder dysfunction in sexual abuse survivors. *J Urol.* 2003 Aug;170(2 Pt 1):476-9. X-1I

485. Davis TL, Lukacz ES, Lubner KM, et al. Determinants of patient satisfaction after the tension-free vaginal tape procedure. *Am J Obstet Gynecol.* 2004 Jul;191(1):176-81. X-1B
486. Day PL. Findings of a three-year retrospective study to investigate prevalence and incidence of urinary incontinence and overactive bladder in a typical managed care setting. *Pharm Pract Manag Q.* 2000 Apr;20(1):1-11. X-6
487. de Araujo MP, Faria AC, Takano CC, et al. Urodynamic study and quality of life in patients with fibromyalgia and lower urinary tract symptoms. *Int Urogynecol J Pelvic Floor Dysfunct.* 2008 Aug;19(8):1103-7. X-1F
488. De Bruin VS, Machado C, Howard RS, et al. Nocturnal and respiratory disturbances in Steele-Richardson-Olszewski syndrome (progressive supranuclear palsy). *Postgrad Med J.* 1996 May;72(847):293-6. X-1I
489. De Cuyper EM, Ismail R and Maher CF. Laparoscopic Burch colposuspension after failed suburethral tape procedures: a retrospective audit. *Int Urogynecol J Pelvic Floor Dysfunct.* 2008 May;19(5):681-5. X-1B
490. De Gennaro M, Capitanucci ML, Mastracci P, et al. Percutaneous tibial nerve neuromodulation is well tolerated in children and effective for treating refractory vesical dysfunction. *J Urol.* 2004 May;171(5):1911-3. X-4, X-7
491. de Kort LM, Nesselaar CH, van Gool JD, et al. The influence of colonic enema irrigation on urodynamic findings in patients with neurogenic bladder dysfunction. *Br J Urol.* 1997 Nov;80(5):731-3. X-1H
492. De Muylder X, Claes H, Neven P, et al. Usefulness of urodynamic investigations in female incontinence. *Eur J Obstet Gynecol Reprod Biol.* 1992 May 13;44(3):205-8. X-1H
493. De Paepe H, Hoebeke P, Renson C, et al. Pelvic-floor therapy in girls with recurrent urinary tract infections and dysfunctional voiding. *Br J Urol.* 1998 May;81 Suppl 3:109-13. X-1I, X-7
494. De Ridder D, Chandiramani V, Dasgupta P, et al. Intravesical capsaicin as a treatment for refractory detrusor hyperreflexia: a dual center study with long-term followup. *J Urol.* 1997 Dec;158(6):2087-92. X-1F, X-7
495. De Ridder D, Vermeulen C, Ketelaer P, et al. Pelvic floor rehabilitation in multiple sclerosis. *Acta Neurol Belg.* 1999 Mar;99(1):61-4. X-1F, X-7
496. de Seze M, Gallien P, Denys P, et al. Intravesical glucidic capsaicin versus glucidic solvent in neurogenic detrusor overactivity: a double blind controlled randomized study. *Neurourol Urodyn.* 2006;25(7):752-7. X-1F, X-7
497. de Seze M, Wiart L, Ferriere J, et al. Intravesical instillation of capsaicin in urology: A review of the literature. *Eur Urol.* 1999 Oct;36(4):267-77. X-2
498. de Seze M, Wiart L, Joseph PA, et al. Capsaicin and neurogenic detrusor hyperreflexia: a double-blind placebo-controlled study in 20 patients with spinal cord lesions. *Neurourol Urodyn.* 1998;17(5):513-23. X-1A, X-7
499. de Tayrac R, Chevalier N, Chauveaud-Lambling A, et al. Is vaginal hysterectomy a risk factor for urinary incontinence at long-term follow-up? *Eur J Obstet Gynecol Reprod Biol.* 2007 Feb;130(2):258-61. X-1A
500. De Wachter S, Vermandel A, De Moerloose K, et al. Value of increase in bladder capacity in treatment of refractory monosymptomatic nocturnal enuresis in children. *Urology.* 2002 Dec;60(6):1090-4. X-1I, X-4
501. De Wachter S and Wyndaele JJ. How sudden is a compelling desire to void? An observational cystometric study on the suddenness of this sensation. *BJU Int.* 2008 Apr;101(8):1000-3. X-1H
502. Deans CL, Morris AR, O'Sullivan R, et al. Voiding difficulty and other adverse events after tension free vaginal tape. *Australian & New Zealand Continence Journal.* 2004 Jun;10(2):34-6, 38-41. X-1B
503. Debodinance P, Delporte P, Engrand JB, et al. Tension-free vaginal tape (TVT) in the treatment of urinary stress incontinence: 3 years experience involving 256 operations. *Eur J Obstet Gynecol Reprod Biol.* 2002 Oct 10;105(1):49-58. X-1B
504. Deffieux X, Hubeaux K, Porcher R, et al. Abnormal pelvic response to cough in women with stress urinary incontinence. *Neurourol Urodyn.* 2008;27(4):291-6. X-1B, X-1G, X-7
505. Defreitas GA, Lemack GE, Zimmern PE, et al. Distinguishing neurogenic from non-neurogenic detrusor overactivity: a urodynamic assessment of lower urinary tract symptoms in patients with and without Parkinson's disease. *Urology.* 2003 Oct;62(4):651-5. X-7
506. del Campo-Rodriguez M, Batista-Miranda JE, Errando-Smet C, et al. Outcome of colposuspension in patients with stress urinary incontinence and abnormal cystometry. *Arch Esp Urol.* 1999 Sep;52(7):810-4. X-1B
507. Del Carro U, Riva D, Comi GC, et al. Neurophysiological evaluation in detrusor instability. *Neurourol Urodyn.* 1993;12(5):455-62. X-1G
508. Del Pizzo JJ, Chew BH, Jacobs SC, et al. Treatment of radiation induced hemorrhagic cystitis with hyperbaric oxygen: long-term followup. *J Urol.* 1998 Sep;160(3 Pt 1):731-3. X-1I
509. Delaere KP, Debruyne FM and Moonen WA. The use of indomethacin in the treatment of idiopathic bladder instability. *Urol Int.* 1981;36(2):124-7. X-7
510. Delaere KP, Michiels HG, Debruyne FM, et al. Flavoxate hydrochloride in the treatment of detrusor instability. *Urol Int.* 1977;32(5):377-81. X-7
511. Demain S, Smith JF, Hiller L, et al. Comparison of group and individual physiotherapy for female urinary incontinence in primary care: pilot study. *Physiotherapy.* 2001 May;87(5):235-42. X-7
512. Demirci F and Yucel O. Comparison of pubovaginal sling and burch colposuspension procedures in type I/II genuine stress incontinence. *Arch Gynecol Obstet.* 2001 Nov;265(4):190-4. X-1B, X-7
513. Demirci F, Yucel O, Eren S, et al. Long-term results of Burch colposuspension. *Gynecol Obstet Invest.* 2001;51(4):243-7. X-1B
514. Dennis PJ, Rohner TJ, Jr., Hu TW, et al. Simple urodynamic evaluation of incontinent elderly female nursing home patients. A descriptive analysis. *Urology.* 1991 Feb;37(2):173-9. X-1I, X-5
515. Densmore M. Incontinence in the female patient. *J Fam Pract.* 1982 May;14(5):935-6, 938, 940 passim. X-2, X-4, X-6, X-7

516. Deutekom M, Dobben AC, Terra MP, et al. Clinical presentation of fecal incontinence and anorectal function: what is the relationship? *Am J Gastroenterol.* 2007 Feb;102(2):351-61. X-II
517. Deval B, Ferchaux J, Berry R, et al. Objective and subjective cure rates after trans-obturator tape (OBTAPE) treatment of female urinary incontinence. *Eur Urol.* 2006 Feb;49(2):373-7. X-1B
518. Deval B, Levardon M, Samain E, et al. A French multicenter clinical trial of SPARC for stress urinary incontinence. *Eur Urol.* 2003 Aug;44(2):254-8; discussion 258-9. X-1B
519. Di Rovasenda E, Podesta E, Scarsi P, et al. Treatment and follow-up of vesico-ureteral reflux in patients with neuropathic bladder. *Eur J Pediatr Surg.* 1992 Jun;2(3):152-3. X-1F
520. Diefenbach K, Arold G, Wollny A, et al. Effects on sleep of anticholinergics used for overactive bladder treatment in healthy volunteers aged > or = 50 years. *BJU Int.* 2005 Feb;95(3):346-9. X-II, X-7
521. Diefenbach K, Jaeger K, Wollny A, et al. Effect of tolterodine on sleep structure modulated by CYP2D6 genotype. *Sleep Med.* 2008 Jul;9(5):579-82. X-II
522. Dietz HP, Barry C, Lim YN, et al. Two-dimensional and three-dimensional ultrasound imaging of suburethral slings. *Ultrasound Obstet Gynecol.* 2005 Aug;26(2):175-9. X-1B
523. Dietz HP, Barry C, Rane A, et al. Is the cough test necessary at TVT insertion? A case control series Tension-free vaginal tape (TVT). *Australian & New Zealand Continence Journal.* 2006 Winter;12(2):50-3. X-1B
524. Dietz HP and Clarke B. Is the irritable bladder associated with anterior compartment relaxation? A critical look at the 'integral theory of pelvic floor dysfunction'. *Aust N Z J Obstet Gynaecol.* 2001 Aug;41(3):317-9. X-1G, X-1H, X-6
525. Dietz HP, Foote AJ, Mak HL, et al. TVT and Sparc suburethral slings: a case-control series. *Int Urogynecol J Pelvic Floor Dysfunct.* 2004 Mar-Apr;15(2):129-31; discussion 131. X-1B
526. Dietz HP, Mouritsen L, Ellis G, et al. How important is TVT location? *Acta Obstet Gynecol Scand.* 2004 Oct;83(10):904-8. X-1B
527. Dietz HP and Simpson JM. Levator trauma is associated with pelvic organ prolapse. *BJOG.* 2008 Jul;115(8):979-84. X-1E
528. Dietz HP and Wilson PD. Colposuspension success and failure: a long-term objective follow-up study. *Int Urogynecol J Pelvic Floor Dysfunct.* 2000 Dec;11(6):346-51. X-1B
529. Dietz HP and Wilson PD. Laparoscopic colposuspension versus urethropexy: a case-control series. *Int Urogynecol J Pelvic Floor Dysfunct.* 2005 Jan-Feb;16(1):15-8; discussion 18. X-1A
530. Dietz HP, Wilson PD, Clarke B, et al. Irritative symptoms after colposuspension: are they due to distortion or overelevation of the anterior vaginal wall and trigone? *Int Urogynecol J Pelvic Floor Dysfunct.* 2001;12(4):232-5; discussion 235-6. X-1A
531. Digesu GA, Athanasiou S, Chaliha C, et al. Urethral retro-resistance pressure and urodynamic diagnoses in women with lower urinary tract symptoms. *BJOG.* 2006 Jan;113(1):34-8. X-1H
532. Digesu GA, Chaliha C, Khullar V, et al. The relationship of urethral resistance pressure and pressure flow parameters in women with lower urinary tract symptoms. *Int Urogynecol J Pelvic Floor Dysfunct.* 2007 May;18(5):493-7. X-1H
533. Digesu GA, Hutchings A, Salvatore S, et al. Pressure flow study: a useful diagnostic test of female lower urinary tract symptoms. *Neurourol Urodyn.* 2004;23(2):104-8. X-1H
534. Digesu GA, Khullar V, Cardozo L, et al. Overactive bladder symptoms: do we need urodynamics? *Neurourol Urodyn.* 2003;22(2):105-8. X-1H
535. Digesu GA, Khullar V, Cardozo L, et al. The open bladder neck: a significant finding? *Int Urogynecol J Pelvic Floor Dysfunct.* 2004 Sep-Oct;15(5):336-9. X-1I
536. Digesu GA, Khullar V, Cardozo L, et al. Preoperative pressure-flow studies: useful variables to predict the outcome of continence surgery. *BJU Int.* 2004 Dec;94(9):1296-9. X-1B
537. Digesu GA, Khullar V, Panayi D, et al. Should we explain lower urinary tract symptoms to patients? *Neurourol Urodyn.* 2008;27(5):368-71. X-1I
538. Digesu GA, Salvatore S, Chaliha C, et al. Do overactive bladder symptoms improve after repair of anterior vaginal wall prolapse? *Int Urogynecol J Pelvic Floor Dysfunct.* 2007 Dec;18(12):1439-43. X-1E
539. Digesu GA, Salvatore S, Fernando R, et al. Mixed urinary symptoms: what are the urodynamic findings? *Neurourol Urodyn.* 2008;27(5):372-5. X-1B, X-1H
540. Diggs C, Meyer WA, Langenberg P, et al. Assessing urgency in interstitial cystitis/painful bladder syndrome. *Urology.* 2007 Feb;69(2):210-4. X-1D
541. Dik P, Klijn AJ, van Gool JD, et al. Early start to therapy preserves kidney function in spina bifida patients. *Eur Urol.* 2006 May;49(5):908-13. X-1I, X-4
542. Diokno AC, Catipay JR and Steinert BW. Office assessment of patient outcome of pharmacologic therapy for urge incontinence. *Int Urogynecol J Pelvic Floor Dysfunct.* 2002;13(5):334-8. X-7
543. Diokno AC, Homma Y, Sekiguchi Y, et al. Interstitial cystitis, gynecologic pelvic pain, prostatitis, and their epidemiology. *Int J Urol.* 2003 Oct;10 Suppl:S3-6. X-1D, X-2
544. Diokno AC, Sand PK, Macdiarmid S, et al. Perceptions and behaviours of women with bladder control problems. *Fam Pract.* 2006 Oct;23(5):568-77. X-6
545. Diokno AC, Vinson RK and McGillicuddy J. Treatment of the severe uninhibited neurogenic bladder by selective sacral rhizotomy. *J Urol.* 1977 Aug;118(2):299-301. X-1F
546. Diokno AC, Wells TJ and Brink CA. Urinary incontinence in elderly women: urodynamic evaluation. *J Am Geriatr Soc.* 1987 Oct;35(10):940-6. X-1H
547. Dmochowski R. Urinary frequency: event measure, outcome significance? *Curr Urol Rep.* 2005 Mar;6(2):87-8. X-2

548. Dmochowski R, Abrams P, Marschall-Kehrel D, et al. Efficacy and tolerability of tolterodine extended release in male and female patients with overactive bladder. *Eur Urol*. 2007 Apr;51(4):1054-64; discussion 1064. X-4
549. Dmochowski RR. Management of postoperative overactive bladder complications. *Geriatrics*. 2002 May;57 Suppl 1:18-23. X-2
550. Dmochowski RR, Avon M, Ross J, et al. Transvaginal radio frequency treatment of the endopelvic fascia: a prospective evaluation for the treatment of genuine stress urinary incontinence. *J Urol*. 2003 Mar;169(3):1028-32. X-1B
551. Dmochowski RR, Miklos JR, Norton PA, et al. Duloxetine versus placebo for the treatment of North American women with stress urinary incontinence. *J Urol*. 2003 Oct;170(4 Pt 1):1259-63. X-1B
552. Dmochowski RR and Newman DK. Impact of overactive bladder on women in the United States: results of a national survey. *Curr Med Res Opin*. 2007 Jan;23(1):65-76. X-6
553. Dmochowski RR, Nitti V, Staskin D, et al. Transdermal oxybutynin in the treatment of adults with overactive bladder: combined results of two randomized clinical trials. *World J Urol*. 2005 Sep;23(4):263-70. X-2
554. Dockter M, Kolstad AM, Martin KA, et al. Prevalence of urinary incontinence: a comparative study of collegiate female athletes [sic] and non-athletic controls. *Journal of Women's Health Physical Therapy*. 2007 Spring;31(1):12-7. X-6
555. Domingo S, Alama P, Ruiz N, et al. Transobturator tape procedure outcome: a clinical and quality of life analysis of a 1-year follow-up. *Int Urogynecol J Pelvic Floor Dysfunct*. 2007 Aug;18(8):895-900. X-1B
556. Dong D, Xu Z, Shi B, et al. Urodynamic study in the neurogenic bladder dysfunction caused by intervertebral disk hernia. *Neurourol Urodyn*. 2006;25(5):446-50. X-1F
557. Doo CK, Hong B, Chung BJ, et al. Five-year outcomes of the tension-free vaginal tape procedure for treatment of female stress urinary incontinence. *Eur Urol*. 2006 Aug;50(2):333-8. X-1B
558. Dooley Y, Lowenstein L, Kenton K, et al. Mixed incontinence is more bothersome than pure incontinence subtypes. *Int Urogynecol J Pelvic Floor Dysfunct*. 2008 Oct;19(10):1359-62. X-1H
559. Dorflinger A, Gorton E, Stanton S, et al. Urethral pressure profile: is it affected by position? *Neurourol Urodyn*. 2002;21(6):553-7. X-1G, X-1H
560. Dorschner W, Stolzenburg JU, Griebenow R, et al. Efficacy and cardiac safety of propiverine in elderly patients - a double-blind, placebo-controlled clinical study. *Eur Urol*. 2000 Jun;37(6):702-8. X-9
561. Dowd T, Kolcaba K and Steiner R. Correlations among measures of bladder function and comfort. *J Nurs Meas*. 2002 Spring-Summer;10(1):27-38. X-1I
562. Drake MJ, Gardner BP and Brading AF. Innervation of the detrusor muscle bundle in neurogenic detrusor overactivity. *BJU Int*. 2003 May;91(7):702-10. X-1G, X-7
563. Drake MJ, Harvey IJ, Gillespie JJ, et al. Localized contractions in the normal human bladder and in urinary urgency. *BJU Int*. 2005 May;95(7):1002-5. X-1G, X-7
564. Drake MJ, Hedlund P, Andersson KE, et al. Morphology, phenotype and ultrastructure of fibroblastic cells from normal and neuropathic human detrusor: absence of myofibroblast characteristics. *J Urol*. 2003 Apr;169(4):1573-6. X-1G, X-4
565. Drake NL, Flynn MK, Romero AA, et al. Nocturnal polyuria in women with overactive bladder symptoms and nocturia. *Am J Obstet Gynecol*. 2005 May;192(5):1682-6. X-1I
566. Dryden MS, Wilkinson M, Redman M, et al. Detection of Chlamydia trachomatis in general practice urine samples. *Br J Gen Pract*. 1994 Mar;44(380):114-7. X-1I
567. D'Souza AO, Smith MJ, Miller LA, et al. Persistence, adherence, and switch rates among extended-release and immediate-release overactive bladder medications in a regional managed care plan. *J Manag Care Pharm*. 2008 Apr;14(3):291-301. X-1I
568. DuBeau CE. Urinary incontinence. *Clinical Geriatrics*. 2001;9(6):31. X-2, X-4
569. DuBeau CE, Bent AE, Dmochowski RR, et al. Addressing the unmet needs of geriatric patients with overactive bladder: challenges and controversies. *Clinical Geriatrics*. 2003;11(12):16. X-2
570. DuBeau CE, Khullar V and Versi E. "Unblinding" in randomized controlled drug trials for urinary incontinence: Implications for assessing outcomes when adverse effects are evident. *Neurourol Urodyn*. 2005;24(1):13-20. X-1I
571. DuBeau CE, Kiely DK and Resnick NM. Quality of life impact of urge incontinence in older persons: a new measure and conceptual structure. *J Am Geriatr Soc*. 1999 Aug;47(8):989-94. X-1H
572. DuBeau CE, Levy B, Mangione CM, et al. The impact of urge urinary incontinence on quality of life: importance of patients' perspective and explanatory style. *J Am Geriatr Soc*. 1998 Jun;46(6):683-92. X-7
573. Duckett J, Aggarwal I, Basu M, et al. The value of cystoscopy and bladder biopsy taken at the time of tension-free vaginal tape insertion. *J Obstet Gynaecol*. 2007 Apr;27(3):297-9. X-1B
574. Duckett J, Aggarwal I, Patil A, et al. Effect of tension-free vaginal tape position on the resolution of irritative bladder symptoms in women with mixed incontinence. *Int Urogynecol J Pelvic Floor Dysfunct*. 2008 Feb;19(2):237-9. X-1B
575. Duckett JR and Basu M. The predictive value of preoperative pressure-flow studies in the resolution of detrusor overactivity and overactive bladder after tension-free vaginal tape insertion. *BJU Int*. 2007 Jun;99(6):1439-42. X-1B
576. Duckett JR and Tamilselvi A. Effect of tension-free vaginal tape in women with a urodynamic diagnosis of idiopathic detrusor overactivity and stress incontinence. *BJOG*. 2006 Jan;113(1):30-3. X-1B
577. Duckett JR, Vella M, Kavalakuntla G, et al. Tolerability and efficacy of duloxetine in a nontrial situation. *BJOG*. 2007 May;114(5):543-7. X-1B
578. Duldulao KE, Diokno AC and Mitchell B. Value of urinary cytology in women presenting with urge incontinence and/or irritative voiding symptoms. *J Urol*. 1997 Jan;157(1):113-6. X-1H

579. Dunn M, Smith JC and Ardran GM. Prolonged bladder distension as a treatment of urgency and urge incontinence of urine. *Br J Urol.* 1974 Dec;46(6):645-52. X-7
580. Dunn MD, Portis AJ, Kahn SA, et al. Clinical effectiveness of new stent design: randomized single-blind comparison of tail and double-pigtail stents. *J Endourol.* 2000 Mar;14(2):195-202. X-1A
581. Durufle A, Petrilli S, Nicolas B, et al. Effects of pregnancy and child birth on urinary symptoms and urodynamics in women with multiple sclerosis. *Int Urogynecol J Pelvic Floor Dysfunct.* 2006 Jun;17(4):352-5. X-1F
582. Dwyer PL and Desmedt E. Impaired bladder emptying in women. *Aust N Z J Obstet Gynaecol.* 1994 Feb;34(1):73-8. X-1G, X-1H
583. Dwyer PL, Lee ET and Hay DM. Obesity and urinary incontinence in women. *Br J Obstet Gynaecol.* 1988 Jan;95(1):91-6. X-1I
584. Dykstra D, Enriquez A and Valley M. Treatment of overactive bladder with botulinum toxin type B: a pilot study. *Int Urogynecol J Pelvic Floor Dysfunct.* 2003 Dec;14(6):424-6. X-7
585. Eastwood HD and Warrell R. Urinary incontinence in the elderly female: prediction in diagnosis and outcome of management. *Age Ageing.* 1984 Jul;13(4):230-4. X-1H
586. Eckford SD, Swami KS, Jackson SR, et al. Desmopressin in the treatment of nocturia and enuresis in patients with multiple sclerosis. *Br J Urol.* 1994 Dec;74(6):733-5. X-1I
587. Edlund C, Dijkema HE, Hassouna MM, et al. Sacral nerve stimulation for refractory urge symptoms in elderly patients. *Scand J Urol Nephrol.* 2004;38(2):131-5. X-7
588. Edlund C, Hellstrom M, Peeker R, et al. First Scandinavian experience of electrical sacral nerve stimulation in the treatment of the overactive bladder. *Scand J Urol Nephrol.* 2000 Dec;34(6):366-76. X-7
589. Edlund C, Peeker R and Fall M. Clam ileocystoplasty: successful treatment of severe bladder overactivity. *Scand J Urol Nephrol.* 2001 Jun;35(3):190-5. X-7
590. Edlund C, Peeker R and Fall M. Lidocaine cystometry in the diagnosis of bladder overactivity. *Neurourol Urodyn.* 2001;20(2):147-55. X-7
591. Ehren I, Hosseini A, Lundberg JO, et al. Nitric oxide: a useful gas in the detection of lower urinary tract inflammation. *J Urol.* 1999 Aug;162(2):327-9. X-1I
592. Ehren I, Volz D, Farrelly E, et al. Efficacy and impact of botulinum toxin A on quality of life in patients with neurogenic detrusor overactivity: a randomised, placebo-controlled, double-blind study. *Scand J Urol Nephrol.* 2007;41(4):335-40. X-1F, X-7
593. Eiberg H, Shaumburg HL, Von Gontard A, et al. Linkage study of a large Danish 4-generation family with urge incontinence and nocturnal enuresis. *J Urol.* 2001 Dec;166(6):2401-3. X-1I, X-6
594. Ekman G, Andersson KE, Rud T, et al. A double-blind, cross-over study of the effects of terodiline in women with unstable bladder. *Acta Pharmacol Toxicol (Copenh).* 1980;46 Suppl 1:39-43. X-7
595. Ekstrom A, Altman D, Wiklund I, et al. Planned cesarean section versus planned vaginal delivery: comparison of lower urinary tract symptoms. *Int Urogynecol J Pelvic Floor Dysfunct.* 2008 Apr;19(4):459-65. X-6
596. Ekstrom B, Andersson KE and Mattiasson A. Urodynamic effects of intravesical instillation of atropine and phentolamine in patients with detrusor hyperactivity. *J Urol.* 1993 Jan;149(1):155-8. X-7
597. Elbadawi A, Hailemariam S, Yalla SV, et al. Structural basis of geriatric voiding dysfunction. VI. Validation and update of diagnostic criteria in 71 detrusor biopsies. *J Urol.* 1997 May;157(5):1802-13. X-1G
598. Elbadawi A, Hailemariam S, Yalla SV, et al. Structural basis of geriatric voiding dysfunction. VII. Prospective ultrastructural/urodynamic evaluation of its natural evolution. *J Urol.* 1997 May;157(5):1814-22. X-1G
599. Elbadawi A, Resnick NM, Dorsam J, et al. Structural basis of neurogenic bladder dysfunction. I. Methods of prospective ultrastructural study and overview of the findings. *J Urol.* 2003 Feb;169(2):540-6. X-1F, X-1G
600. Elbadawi A, Yalla SV and Resnick NM. Structural basis of geriatric voiding dysfunction. I. Methods of a prospective ultrastructural/urodynamic study and an overview of the findings. *J Urol.* 1993 Nov;150(5 Pt 2):1650-6. X-1B, X-1G
601. Elbadawi A, Yalla SV and Resnick NM. Structural basis of geriatric voiding dysfunction. II. Aging detrusor: normal versus impaired contractility. *J Urol.* 1993 Nov;150(5 Pt 2):1657-67. X-1B, X-1G
602. Elbadawi A, Yalla SV and Resnick NM. Structural basis of geriatric voiding dysfunction. III. Detrusor overactivity. *J Urol.* 1993 Nov;150(5 Pt 2):1668-80. X-1B, X-1G
603. Elbadawi A, Yalla SV and Resnick NM. Structural basis of geriatric voiding dysfunction. IV. Bladder outlet obstruction. *J Urol.* 1993 Nov;150(5 Pt 2):1681-95. X-1G, X-1I
604. Elder DD and Stephenson TP. An assessment of the Frewen regime in the treatment of detrusor dysfunction in females. *Br J Urol.* 1980 Dec;52(6):467-71. X-7
605. Elder DD, Stephenson TP and Bary PR. An assessment of the Frewen regime in the treatment of detrusor dysfunction in females. *Prog Clin Biol Res.* 1981;78:321-8. X-7
606. Elder JS. Bladder rehabilitation, the effect of a cognitive training programme on urge incontinence. *J Urol.* 1997 Oct;158(4):1642. X-2
607. Elder JS, Snyder HM, Peters C, et al. Variations in practice among urologists and nephrologists treating children with vesicoureteral reflux. *J Urol.* 1992 Aug;148(2 Pt 2):714-7. X-1I, X-4
608. Elem B, Patil PS and Lucas SB. Haematuria frequency syndrome in patients with positive HIV serology: observations in Zambia. *Br J Urol.* 1991 Feb;67(2):146-9. X-1A, X-7
609. Elgamasy AN, Lewis V, Hassouna ME, et al. Effect of transvaginal stimulation in the treatment of detrusor instability. *Urol Nurs.* 1996 Dec;16(4):127-30. X-7
610. Elhilali MM, Khaled SM, Kashiwabara T, et al. Sacral neuromodulation: long-term experience of one center. *Urology.* 2005 Jun;65(6):1114-7. X-7

611. Elkins TE, Ghosh TS, Tagoe GA, et al. Transvaginal mobilization and utilization of the anterior bladder wall to repair vesicovaginal fistulas involving the urethra. *Obstet Gynecol.* 1992 Mar;79(3):455-60. X-II
612. Elliott RA, Norman RI, Parker SG, et al. Effect of treatment with calcium antagonists in vitro and in vivo on the contractile response of isolated rat and human detrusor muscle. *Clin Sci (Lond).* 1996 Oct;91(4):467-74. X-1G, X-4, X-6
613. Ellsworth PI, Borgstein NG, Nijman RJ, et al. Use of tolterodine in children with neurogenic detrusor overactivity: relationship between dose and urodynamic response. *J Urol.* 2005 Oct;174(4 Pt 2):1647-51; discussion 1651. X-1F, X-4
614. El-Nahas AR, El-Assmy AM, Shoma AM, et al. Self-retaining ureteral stents: analysis of factors responsible for patients' discomfort. *J Endourol.* 2006 Jan;20(1):33-7. X-1A, X-7
615. Elser DM, Wyman JF, McClish DK, et al. The effect of bladder training, pelvic floor muscle training, or combination training on urodynamic parameters in women with urinary incontinence. *Continence Program for Women Research Group. NeuroUrol Urodyn.* 1999;18(5):427-36. X-7
616. el-Toukhy TA and Davies AE. The efficacy of laparoscopic mesh colposuspension: results of a prospective controlled study. *BJU Int.* 2001 Sep;88(4):361-6. X-1B
617. Elzevier HW, Venema PL and Lycklama a Nijeholt AA. Sexual function after tension-free vaginal tape (TVT) for stress incontinence: results of a mailed questionnaire. *Int Urogynecol J Pelvic Floor Dysfunct.* 2004 Sep-Oct;15(5):313-8. X-1B
618. Engel AF, Kamm MA, Bartram CI, et al. Relationship of symptoms in faecal incontinence to specific sphincter abnormalities. *Int J Colorectal Dis.* 1995;10(3):152-5. X-II
619. English PJ and Fowler JW. Videourodynamic assessment of the Stamey procedure for stress incontinence. *Br J Urol.* 1988 Dec;62(6):550-2. X-1B
620. Erdogru T, Kocak T, Serdaroglu P, et al. Evaluation and therapeutic approaches of voiding and erectile dysfunction in neurological Behcet's syndrome. *J Urol.* 1999 Jul;162(1):147-53. X-II
621. Eregie CO. Observations on urinary frequency in exclusively breastfed neonates. *East Afr Med J.* 1998 Oct;75(10):576-8. X-II, X-4
622. Eriksen BC, Bergmann S and Mjølnerod OK. Effect of anal electrostimulation with the 'Incontan' device in women with urinary incontinence. *Br J Obstet Gynaecol.* 1987 Feb;94(2):147-56. X-7
623. Eriksen BC, Hagen B, Eik-Nes SH, et al. Long-term effectiveness of the Burch colposuspension in female urinary stress incontinence. *Acta Obstet Gynecol Scand.* 1990;69(1):45-50. X-1B
624. Eriksen BC and Mjølnerod OK. Changes in urodynamic measurements after successful anal electrostimulation in female urinary incontinence. *Br J Urol.* 1987 Jan;59(1):45-9. X-7
625. Ertberg P, Moller LA and Lose G. A comparison of three methods to evaluate maximum bladder capacity: cystometry, uroflowmetry and a 24-h voiding diary in women with urinary incontinence. *Acta Obstet Gynecol Scand.* 2003 Apr;82(4):374-7. X-1H
626. Esa A, Kiwamoto H, Sugiyama T, et al. Functional electrical stimulation in the management of incontinence: studies of urodynamics. *Int Urol Nephrol.* 1991;23(2):135-41. X-7
627. Espuna Pons M and Puig Clota M. Coital urinary incontinence: impact on quality of life as measured by the King's Health Questionnaire. *Int Urogynecol J Pelvic Floor Dysfunct.* 2008 May;19(5):621-5. X-1A
628. Espuna-Pons M, Dilla T, Castro D, et al. Analysis of the value of the ICIQ-UI SF questionnaire and stress test in the differential diagnosis of the type of urinary incontinence. *NeuroUrol Urodyn.* 2007;26(6):836-41. X-1H
629. Essenhigh DM and Ryan DW. An appraisal of S3 blocks in the management of incontinence. *Br J Urol.* 1982 Dec;54(6):697-9. X-7
630. Esteghamati A, Rashidi A, Nikfallah A, et al. The association between urodynamic findings and microvascular complications in patients with long-term type 2 diabetes but without voiding symptoms. *Diabetes Res Clin Pract.* 2007 Oct;78(1):42-50. X-1A, X-7
631. Ethans KD, Nance PW, Bard RJ, et al. Efficacy and safety of tolterodine in people with neurogenic detrusor overactivity. *J Spinal Cord Med.* 2004;27(3):214-8. X-1F, X-7
632. Everaert K, De Ridder D, Baert L, et al. Patient satisfaction and complications following sacral nerve stimulation for urinary retention, urge incontinence and perineal pain: a multicenter evaluation. *Int Urogynecol J Pelvic Floor Dysfunct.* 2000;11(4):231-5; discussion 236. X-7
633. Ewing R, Bultitude MI and Shuttleworth KE. Subtrigonal phenol injection for urge incontinence secondary to detrusor instability in females. *Br J Urol.* 1982 Dec;54(6):689-92. X-7
634. Ewing R, Bultitude MI and Shuttleworth KE. Subtrigonal phenol injection therapy for incontinence in female patients with multiple sclerosis. *Lancet.* 1983 Jun 11;1(8337):1304-6. X-1F
635. Ezzat IM. Sling operations in the treatment of stress urinary incontinence: how to adjust sling tension. *J Obstet Gynaecol Res.* 2003 Dec;29(6):374-9. X-1B
636. Faber P and Heidenreich J. Effect of intravenous butylscopolamine infusion on urge incontinence. *Urol Int.* 1977;32(2-3):218-20. X-9
637. Faerber GJ. Urethral diverticulectomy and pubovaginal sling for simultaneous treatment of urethral diverticulum and intrinsic sphincter deficiency. *Tech Urol.* 1998 Dec;4(4):192-7. X-II
638. Fall M. Does electrostimulation cure urinary incontinence? *J Urol.* 1984 Apr;131(4):664-7. X-7
639. Fall M. Conservative management of chronic interstitial cystitis: transcutaneous electrical nerve stimulation and transurethral resection. *J Urol.* 1985 May;133(5):774-8. X-1D, X-7
640. Fall M, Ahlstrom K, Carlsson CA, et al. Contelle: pelvic floor stimulator for female stress-urge incontinence. A multicenter study. *Urology.* 1986 Mar;27(3):282-7. X-7

641. Fall M, Erlandson BE, Carlsson CA, et al. Effects of electrical intravaginal stimulation on bladder volume: an experimental and clinical study. *Urol Int*. 1978;33(6):440-2. X-7
642. Fall M, Erlandson BE, Nilson AE, et al. Long-term intravaginal electrical stimulation in urge and stress incontinence. *Scand J Urol Nephrol Suppl*. 1977(44):55-63. X-7
643. Fall M, Erlandson BE and Pettersson S. Evaluation of history and simple supine cystometry as a preoperative test in stress urinary incontinence. *Acta Obstet Gynecol Scand*. 1984;63(3):241-4. X-1H, X-7
644. Fall M, Erlandson BE, Sundin T, et al. Intravaginal electrical stimulation. Clinical experiments on bladder inhibition. *Scand J Urol Nephrol Suppl*. 1977(44):41-7. X-7
645. Fall M, Ohlsson BL and Carlsson CA. The neurogenic overactive bladder. Classification based on urodynamics. *Br J Urol*. 1989 Oct;64(4):368-73. X-1F, X-2
646. Fall M and Pettersson S. The simplified Lapidus' operation for stress incontinence. *Scand J Urol Nephrol*. 1983;17(1):27-30. X-1B
647. Fantl JA. Behavioral intervention for community-dwelling individuals with urinary incontinence. *Urology*. 1998 Feb;51(2A Suppl):30-4. X-2
648. Fantl JA, Bump RC, Robinson D, et al. Efficacy of estrogen supplementation in the treatment of urinary incontinence. The Continence Program for Women Research Group. *Obstet Gynecol*. 1996 Nov;88(5):745-9. X-7
649. Fantl JA, Harkins SW, Wyman JF, et al. Fluid loss quantitation test in women with urinary incontinence: a test-retest analysis. *Obstet Gynecol*. 1987 Nov;70(5):739-43. X-1H, X-7
650. Fantl JA, Hurt WG, Beachley MC, et al. Bead-chain cystourethrogram: an evaluation. *Obstet Gynecol*. 1981 Aug;58(2):237-40. X-1B, X-1H
651. Fantl JA, Wyman JF, Anderson RL, et al. Postmenopausal urinary incontinence: comparison between non-estrogen-supplemented and estrogen-supplemented women. *Obstet Gynecol*. 1988 Jun;71(6 Pt 1):823-8. X-1H
652. Fantl JA, Wyman JF, McClish DK, et al. Urinary incontinence in community-dwelling women: clinical, urodynamic, and severity characteristics. *Am J Obstet Gynecol*. 1990 Apr;162(4):946-51; discussion 951-2. X-7
653. Fantl JA, Wyman JF, McClish DK, et al. Efficacy of bladder training in older women with urinary incontinence. *JAMA*. 1991 Feb 6;265(5):609-13. X-1B, X-7
654. Farghaly SA, Hindmarsh JR and Worth PH. Post-hysterectomy urethral dysfunction: evaluation and management. *Br J Urol*. 1986 Jun;58(3):299-302. X-1A
655. Farrar DJ and Osborne JL. The use of bromocriptine in the treatment of the unstable bladder. *Br J Urol*. 1976 Aug;48(4):235-8. X-7
656. Farrell SA, Baydock S, Amir B, et al. Effectiveness of a new self-positioning pessary for the management of urinary incontinence in women. *Am J Obstet Gynecol*. 2007 May;196(5):474 e1-8. X-7
657. Farrell SA, Singh B and Aldakhil L. Continence pessaries in the management of urinary incontinence in women. *J Obstet Gynaecol Can*. 2004 Feb;26(2):113-7. X-1B
658. Farrell SA and Tynski G. The effect of urethral pressure variation on detrusor activity in women. *Int Urogynecol J Pelvic Floor Dysfunct*. 1996;7(2):87-93. X-1H
659. Fehrling M, Fall M and Pecker R. Maximal functional electrical stimulation as a single treatment: is it cost-effective? *Scand J Urol Nephrol*. 2007;41(2):132-7. X-7
660. Fehrmann-Zumpe P, Karbe K and Blessman G. Using flavoxate as primary medication for patients suffering from urge symptomatology. *Int Urogynecol J Pelvic Floor Dysfunct*. 1999;10(2):91-5. X-4
661. Felsen D, Frye S, Trimble LA, et al. Inflammatory mediator profile in urine and bladder wash fluid of patients with interstitial cystitis. *J Urol*. 1994 Aug;152(2 Pt 1):355-61. X-1D, X-1G
662. Fernando RJ, Thakar R, Sultan AH, et al. Effect of vaginal pessaries on symptoms associated with pelvic organ prolapse. *Obstet Gynecol*. 2006 Jul;108(1):93-9. X-1E
663. Fernie GR, Jewett MA, Halsall P, et al. Urodynamic characterization of incontinence in the elderly by bladder volume. *J Urol*. 1983 Apr;129(4):772-4. X-1G, X-1H
664. Ferrara P, Costa S, Rigante D, et al. Intramedullary epidermoid cyst presenting with abnormal urological manifestations. *Spinal Cord*. 2003 Nov;41(11):645-8. X-1F, X-2, X-4, X-6, X-7
665. Ferrara P, Rigante D, D'Aleo C, et al. Preliminary data on monosymptomatic nocturnal enuresis in children and adolescents with type 1 diabetes. *Scand J Urol Nephrol*. 2006;40(3):238-40. X-1A, X-4
666. Ferrie BG, Smith JS, Logan D, et al. Experience with bladder training in 65 patients. *Br J Urol*. 1984 Oct;56(5):482-4. X-1I
667. Fielding JR, Lee JH, Dubeau CE, et al. Voiding cystourethrography findings in elderly women with urge incontinence. *J Urol*. 2000 Apr;163(4):1216-8. X-1G
668. Fieler VK. Side effects and quality of life in patients receiving high-dose rate brachytherapy. *Oncol Nurs Forum*. 1997 Apr;24(3):545-53. X-1A
669. Fihn SD, Johnson C, Roberts PL, et al. Trimethoprim-sulfamethoxazole for acute dysuria in women: a single-dose or 10-day course. A double-blind, randomized trial. *Ann Intern Med*. 1988 Mar;108(3):350-7. X-1I
670. Finazzi Agro E, Campagna A, Sciobica F, et al. Posterior tibial nerve stimulation: is the once-a-week protocol the best option? *Minerva Urol Nefrol*. 2005 Jun;57(2):119-23. X-7
671. Fine PM, Mauldin JE and Harte EH. Stress incontinence in women. A practical method of evaluation. *J Reprod Med*. 1988 Sep;33(9):749-56. X-1B, X-1H
672. Fink D, Perucchini D, Schaer GN, et al. The role of the frequency-volume chart in the differential diagnostic of female urinary incontinence. *Acta Obstet Gynecol Scand*. 1999 Mar;78(3):254-7. X-1H, X-7
673. Finkelstein K, Glosner S, Sanchez RJ, et al. Prevalence of probable overactive bladder in a private obstetrics and gynecology group practice. *Curr Med Res Opin*. 2008 Apr;24(4):1083-90. X-6
674. Finnerty P. Overactive bladder. *World of Irish Nursing & Midwifery*. 2006;14(6):47-47. X-2

675. Fischer W and Linde A. Pelvic floor findings in urinary incontinence--results of conditioning using vaginal cones. *Acta Obstet Gynecol Scand*. 1997 May;76(5):455-60. X-1B, X-7
676. Fischer-Rasmussen W. Evaluation of long-term safety and clinical benefit of terodiline in women with urgency/urge incontinence. A multicentre study. *Scand J Urol Nephrol Suppl*. 1984;87:35-47. X-9
677. Fischer-Rasmussen W. Transvaginal needle bladder neck suspension for stress urinary incontinence: practicable methods but not optimal results. *Acta Obstet Gynecol Scand Suppl*. 1998;168:38-43. X-1B
678. Fischer-Rasmussen W, Hansen RI and Stage P. Predictive values of diagnostic tests in the evaluation of female urinary stress incontinence. *Acta Obstet Gynecol Scand*. 1986;65(4):291-4. X-1B
679. Fitzgerald MP, Ayuste D and Brubaker L. How do urinary diaries of women with an overactive bladder differ from those of asymptomatic controls? *BJU Int*. 2005 Aug;96(3):365-7. X-7
680. FitzGerald MP and Brubaker L. Urinary incontinence symptom scores and urodynamic diagnoses. *Neurourol Urodyn*. 2002;21(1):30-5. X-1H
681. FitzGerald MP and Brubaker L. Variability of 24-hour voiding diary variables among asymptomatic women. *J Urol*. 2003 Jan;169(1):207-9. X-1H
682. FitzGerald MP, Butler N, Shott S, et al. Bother arising from urinary frequency in women. *Neurourol Urodyn*. 2002;21(1):36-40; discussion 41. X-1A
683. FitzGerald MP, Jaffar J and Brubaker L. Risk factors for an elevated postvoid residual urine volume in women with symptoms of urinary urgency, frequency and urge incontinence. *Int Urogynecol J Pelvic Floor Dysfunct*. 2001;12(4):237-9; discussion 239-40. X-1I
684. FitzGerald MP, Kenton KS and Brubaker L. Localization of the urge to void in patients with painful bladder syndrome. *Neurourol Urodyn*. 2005;24(7):633-7. X-1A, X-7
685. FitzGerald MP, Manaves V, Martin AF, et al. Myosin isoforms in female human detrusor. *Neurourol Urodyn*. 2001;20(1):23-9. X-1G
686. Fjorback MV, Rijkhoff N, Petersen T, et al. Event driven electrical stimulation of the dorsal penile/clitoral nerve for management of neurogenic detrusor overactivity in multiple sclerosis. *Neurourol Urodyn*. 2006;25(4):349-55. X-1F, X-6, X-7
687. Fjorback MV, Van Rey FS, Rijkhoff NJ, et al. Electrical stimulation of sacral dermatomes in multiple sclerosis patients with neurogenic detrusor overactivity. *Neurourol Urodyn*. 2007;26(4):525-30. X-1F
688. Fjorback MV, van Rey FS, van der Pal F, et al. Acute urodynamic effects of posterior tibial nerve stimulation on neurogenic detrusor overactivity in patients with MS. *Eur Urol*. 2007 Feb;51(2):464-70; discussion 471-2. X-1F, X-7
689. Fletcher S. Interstitial cystitis: painful bladder syndrome. *Nurse Pract*. 1988 Oct;13(10):7, 10-2. X-1D, X-2
690. Fletcher SG and Lemack GE. Clarifying the role of urodynamics in the preoperative evaluation of stress urinary incontinence. *ScientificWorldJournal*. 2008;8:1259-68. X-1B, X-1H
691. Fliegner JR and Glenning PP. Seven years experience in the evaluation and management of patients with urge incontinence of urine. *Aust N Z J Obstet Gynaecol*. 1979 Feb;19(1):42-4. X-9
692. Flisser AJ and Blaivas JG. Role of cystometry in evaluating patients with overactive bladder. *Urology*. 2002 Nov;60(5 Suppl 1):33-42; discussion 42. X-2
693. Flood HD, Malhotra SJ, O'Connell HE, et al. Long-term results and complications using augmentation cystoplasty in reconstructive urology. *Neurourol Urodyn*. 1995;14(4):297-309. X-7
694. Flynn BJ and Yap WT. Pubovaginal sling using allograft fascia lata versus autograft fascia for all types of stress urinary incontinence: 2-year minimum followup. *J Urol*. 2002 Feb;167(2 Pt 1):608-12. X-1B
695. Flynn L, Cell P and Luisi E. Effectiveness of pelvic muscle exercises in reducing urge incontinence among community residing elders. *J Gerontol Nurs*. 1994 May;20(5):23-7. X-7
696. Fonda D, Brimage PJ and D'Astoli M. Simple screening for urinary incontinence in the elderly: comparison of simple and multichannel cystometry. *Urology*. 1993 Nov;42(5):536-40. X-1H
697. Foote J, Glavind K, Kralidis G, et al. Treatment of overactive bladder in the older patient: pooled analysis of three phase III studies of darifenacin, an M3 selective receptor antagonist. *Eur Urol*. 2005 Sep;48(3):471-7. X-2
698. Forman A, Andersson KE, Henriksson L, et al. Effects of nifedipine on the smooth muscle of the human urinary tract in vitro and in vivo. *Acta Pharmacol Toxicol (Copenh)*. 1978 Aug;43(2):111-8. X-1G
699. Fortunato P, Schettini M and Gallucci M. Diverticula of the female urethra. *Br J Urol*. 1997 Oct;80(4):628-32. X-1A, X-7
700. Fossberg E, Beisland HO and Sander S. Sensory urgency in females: treatment with phenylpropanolamine. *Eur Urol*. 1981;7(3):157-60. X-7
701. Foster MC and O'Reilly PH. Early experience of the gittes "no-incision" pubovaginal suspension for stress urinary incontinence. *Br J Urol*. 1989 Dec;64(6):590-3. X-1A, X-7
702. Foster RT, Sr., Anoaia EJ, Webster GD, et al. In patients undergoing neuromodulation for intractable urge incontinence a reduction in 24-hr pad weight after the initial test stimulation best predicts long-term patient satisfaction. *Neurourol Urodyn*. 2007;26(2):213-7. X-4
703. Foster RT, Sr., Barber MD, Parasio MF, et al. A prospective assessment of overactive bladder symptoms in a cohort of elderly women who underwent transvaginal surgery for advanced pelvic organ prolapse. *Am J Obstet Gynecol*. 2007 Jul;197(1):82 e1-4. X-1E
704. Fowler CJ, Beck RO, Gerrard S, et al. Intravesical capsaicin for treatment of detrusor hyperreflexia. *J Neurol Neurosurg Psychiatry*. 1994 Feb;57(2):169-73. X-1F, X-7
705. Fox SD and Stanton SL. Vault prolapse and rectocele: assessment of repair using sacrocolpopexy with mesh interposition. *BJOG*. 2000 Nov;107(11):1371-5. X-1E
706. Franco I, Landau-Dyer L, Isom-Batz G, et al. The use of botulinum toxin A injection for the management of external sphincter dyssynergia in neurologically normal children. *J Urol*. 2007 Oct;178(4 Pt 2):1775-9; discussion 1779-80. X-1I, X-4

707. Franco N, Shobeiri SA and Echols KT. Medium-term follow-up of transvaginal suburethral slings: variance in outcome success using two different evaluation methods. *Urology*. 2002 Oct;60(4):607-10; discussion 610-1. X-1B
708. Frauscher F, Helweg G, Strasser H, et al. Intraurethral ultrasound: diagnostic evaluation of the striated urethral sphincter in incontinent females. *Eur Radiol*. 1998;8(1):50-3. X-1B, X-1H, X-7
709. Frazer MI and Haylen BT. Trigonal sensitivity testing in women. *J Urol*. 1989 Feb;141(2):356-8. X-1H
710. Frederick RW and Leach GE. Cadaveric prolapse repair with sling: intermediate outcomes with 6 months to 5 years of followup. *J Urol*. 2005 Apr;173(4):1229-33. X-1B
711. Freeman RM. Female urinary incontinence: aetiology and pathophysiology. *Hosp Med*. 2000 Feb;61(2):84-9. X-2
712. Freeman RM, Adekanmi O, Waterfield MR, et al. The effect of cannabis on urge incontinence in patients with multiple sclerosis: a multicentre, randomised placebo-controlled trial (CAMS-LUTS). *Int Urogynecol J Pelvic Floor Dysfunct*. 2006 Nov;17(6):636-41. X-1F
713. Freeman RM, McPherson FM and Baxby K. Psychological features of women with idiopathic detrusor instability. *Urol Int*. 1985;40(5):257-9. X-1I
714. Freiha FS, Faysal MH and Stamey TA. The surgical treatment of intractable interstitial cystitis. *J Urol*. 1980 May;123(5):632-4. X-1D
715. Frenkl TL, Rackley RR, Vasavada SP, et al. Management of iatrogenic foreign bodies of the bladder and urethra following pelvic floor surgery. *Neurourol Urodyn*. 2008;27(6):491-5. X-1I
716. Frewen WK. An objective assessment of the unstable bladder of psychosomatic origin. *Br J Urol*. 1978 Jun;50(4):246-9. X-7
717. Frewen WK. The management of urgency and frequency of micturition. *Br J Urol*. 1980 Oct;52(5):367-9. X-7
718. Frigerio L, Ferrari A and Candiani GB. The significance of the stop test in female urinary incontinence. *Diagn Gynecol Obstet*. 1981 Winter;3(4):301-4. X-1H
719. Friis E, Hjortrup A, Nielsen JE, et al. Urinary incontinence and genital prolapse: a prospective blind study of the value of urodynamic evaluation. *J Urol*. 1982 Oct;128(4):764-5. X-1H
720. Fritel X, Zabak K, Pigne A, et al. Predictive value of urethral mobility before suburethral tape procedure for urinary stress incontinence in women. *J Urol*. 2002 Dec;168(6):2472-5. X-1B
721. Frohlich G, Bulitta M and Strosser W. Trosipium chloride in patients with detrusor overactivity: meta-analysis of placebo-controlled, randomized, double-blind, multi-center clinical trials on the efficacy and safety of 20 mg trosipium chloride twice daily. *Int J Clin Pharmacol Ther*. 2002 Jul;40(7):295-303. X-2
722. Fujishiro T, Takahashi S, Enomoto H, et al. Magnetic stimulation of the sacral roots for the treatment of urinary frequency and urge incontinence: an investigational study and placebo controlled trial. *J Urol*. 2002 Sep;168(3):1036-9. X-7
723. Fulford SC, Flynn R, Barrington J, et al. An assessment of the surgical outcome and urodynamic effects of the pubovaginal sling for stress incontinence and the associated urge syndrome. *J Urol*. 1999 Jul;162(1):135-7. X-1B
724. Fulmer BR, Sakamoto K, Turk TM, et al. Acute and long-term outcomes of radio frequency bladder neck suspension. *J Urol*. 2002 Jan;167(1):141-5. X-1B
725. Gajewski JB and Awad SA. Oxybutynin versus propantheline in patients with multiple sclerosis and detrusor hyperreflexia. *J Urol*. 1986 May;135(5):966-8. X-1F, X-7
726. Galejs LE and Diokno AC. Frequency urgency syndrome following urinary bladder suspension. *Geriatr Nephrol Urol*. 1998;8(1):25-8. X-1A
727. Galloway NT, El-Galley RE, Sand PK, et al. Extracorporeal magnetic innervation therapy for stress urinary incontinence. *Urology*. 1999 Jun;53(6):1108-11. X-1B
728. Galloway NT, El-Galley RE, Sand PK, et al. Update on extracorporeal magnetic innervation (EXMI) therapy for stress urinary incontinence. *Urology*. 2000 Dec 4;56(6 Suppl 1):82-6. X-1B
729. Gamble TL, Botros SM, Beaumont JL, et al. Predictors of persistent detrusor overactivity after transvaginal sling procedures. *Am J Obstet Gynecol*. 2008 Dec;199(6):696 e1-7. X-1B
730. Game X, Castel-Lacanal E, Bentaleb Y, et al. Botulinum toxin A detrusor injections in patients with neurogenic detrusor overactivity significantly decrease the incidence of symptomatic urinary tract infections. *Eur Urol*. 2008 Mar;53(3):613-8. X-1F
731. Gandini R, Chiochi M, Konda D, et al. Transcatheter foam sclerotherapy of symptomatic female varicocele with sodium-tetradecyl-sulfate foam. *Cardiovasc Intervent Radiol*. 2008 Jul-Aug;31(4):778-84. X-1I
732. Ganesan T, Khadra MH, Wallis J, et al. Vitamin B12 malabsorption following bladder reconstruction or diversion with bowel segments. *ANZ J Surg*. 2002 Jul;72(7):479-82. X-1I
733. Ganio E, Luc AR, Clerico G, et al. Sacral nerve stimulation for treatment of fecal incontinence: a novel approach for intractable fecal incontinence. *Dis Colon Rectum*. 2001 May;44(5):619-29; discussion 629-31. X-1I
734. Ganio E, Masin A, Ratto C, et al. Short-term sacral nerve stimulation for functional anorectal and urinary disturbances: results in 40 patients: evaluation of a new option for anorectal functional disorders. *Dis Colon Rectum*. 2001 Sep;44(9):1261-7. X-1A, X-7
735. Gardy M, Kozminski M, DeLancey J, et al. Stress incontinence and cystoceles. *J Urol*. 1991 Jun;145(6):1211-3. X-1I, X-1B
736. Garnick MB, Schade D, Israel M, et al. Intravesical doxorubicin for prophylaxis in the management of recurrent superficial bladder carcinoma. *J Urol*. 1984 Jan;131(1):43-6. X-1I, X-7
737. Gateau T, Faramarzi-Roques R, Le Normand L, et al. Clinical and urodynamic repercussions after TVT procedure and how to diminish patient complaints. *Eur Urol*. 2003 Sep;44(3):372-6; discussion 376. X-1B

738. Gaudenz R and Weil A. Motor urge incontinence: diagnosis and treatment. *Urol Int.* 1980;35(1):1-12. X-7
739. Gebhart JB, Dixon DA, Trabuco EC, et al. Three-year outcomes of Uretex Urethral Support System for treatment of stress urinary incontinence. *Int Urogynecol J Pelvic Floor Dysfunct.* 2008 Aug;19(8):1075-9. X-1B
740. Gee AS and Durdey P. Urge incontinence of faeces is a marker of severe external anal sphincter dysfunction. *Br J Surg.* 1995 Sep;82(9):1179-82. X-1I
741. Geirsson G and Fall M. Maximal functional electrical stimulation in routine practice. *Neurourol Urodyn.* 1997;16(6):559-65. X-4, X-7
742. Geirsson G, Lindstrom S and Fall M. The bladder cooling reflex in man--characteristics and sensitivity to temperature. *Br J Urol.* 1993 Jun;71(6):675-80. X-1G
743. Geirsson G, Lindstrom S and Fall M. Pressure, volume and infusion speed criteria for the ice-water test. *Br J Urol.* 1994 May;73(5):498-503. X-1H
744. Gelenberg AJ, Kane JM, Keller MB, et al. Comparison of standard and low serum levels of lithium for maintenance treatment of bipolar disorder. *N Engl J Med.* 1989 Nov 30;321(22):1489-93. X-1I
745. George J, Tharion G, Richar J, et al. The effectiveness of intravesical oxybutynin, propantheline, and capsaicin in the management of neuropathic bladder following spinal cord injury. *ScientificWorldJournal.* 2007;7:1683-90. X-1F
746. George NJ, Dunn M, Dounis A, et al. The late symptomatic and functional results of enterocystoplasty. *Br J Urol.* 1978 Dec;50(7):517-20. X-1D, X-7
747. George VK, Russell GL, Shutt A, et al. Clam ileocystoplasty. *Br J Urol.* 1991 Nov;68(5):487-9. X-7
748. Gerharz EW, Kohl UN, Weingartner K, et al. Experience with the Mainz modification of ureterosigmoidostomy. *Br J Surg.* 1998 Nov;85(11):1512-6. X-1A, X-7
749. Gerstenberg TC, Klarskov P, Ramirez D, et al. Terodiline in the treatment of women with urgency and motor urge incontinence. A clinical and urodynamic double-blind cross-over study. *Br J Urol.* 1986 Apr;58(2):129-33. X-7
750. Gerten KA, Richter HE, Burgio KL, et al. Impact of urinary incontinence in morbidly obese women versus women seeking urogynecologic care. *Urology.* 2007 Dec;70(6):1082-5. X-1I
751. Getsios D, Caro JJ, Ishak KJ, et al. Canadian economic comparison of extended-release oxybutynin and immediate-release tolterodine in the treatment of overactive bladder. *Clin Ther.* 2004 Mar;26(3):431-8. X-8
752. Getsios D, Caro JJ, Ishak KJ, et al. Oxybutynin Extended Release and Tolterodine Immediate Release: A Health Economic Comparison. *Clinical Drug Investigation.* 2004;24(2):81-88. X-8
753. Ghalayini IF and Al-Ghazo MA. Intradetrusor injection of botulinum-A toxin in patients with idiopathic and neurogenic detrusor overactivity: urodynamic outcome and patient satisfaction. *Neurourol Urodyn.* 2007;26(4):531-6. X-7
754. Ghei M and Malone-Lee J. Using the circumstances of symptom experience to assess the severity of urgency in the overactive bladder. *J Urol.* 2005 Sep;174(3):972-6. X-1H
755. Ghei M, Maraj BH, Miller R, et al. Effects of botulinum toxin B on refractory detrusor overactivity: a randomized, double-blind, placebo controlled, crossover trial. *J Urol.* 2005 Nov;174(5):1873-7; discussion 1877. X-7
756. Ghezzi F, Serati M, Cromi A, et al. Tension-free vaginal tape for the treatment of urodynamic stress incontinence with intrinsic sphincteric deficiency. *Int Urogynecol J Pelvic Floor Dysfunct.* 2006 Jun;17(4):335-9. X-1B
757. Ghoniem GM, Lapeyrolerie J, Sood OP, et al. Tulane experience with management of urinary incontinence after placement of an artificial urinary sphincter. *World J Urol.* 1994;12(6):333-6. X-1I
758. Giannantoni A, Costantini E, Di Stasi SM, et al. Botulinum A toxin intravesical injections in the treatment of painful bladder syndrome: a pilot study. *Eur Urol.* 2006 Apr;49(4):704-9. X-1A, X-7
759. Giannantoni A, Di Stasi SM, Nardicchi V, et al. Botulinum-A toxin injections into the detrusor muscle decrease nerve growth factor bladder tissue levels in patients with neurogenic detrusor overactivity. *J Urol.* 2006 Jun;175(6):2341-4. X-1F, X-7
760. Giannantoni A, Di Stasi SM, Stephen RL, et al. Intravesical resiniferatoxin versus botulinum-A toxin injections for neurogenic detrusor overactivity: a prospective randomized study. *J Urol.* 2004 Jul;172(1):240-3. X-1F, X-7
761. Giannantoni A, Mearini E, Di Stasi SM, et al. New therapeutic options for refractory neurogenic detrusor overactivity. *Minerva Urol Nefrol.* 2004 Mar;56(1):79-87. X-1F
762. Giannantoni A, Porena M, Costantini E, et al. Botulinum A toxin intravesical injection in patients with painful bladder syndrome: 1-year followup. *J Urol.* 2008 Mar;179(3):1031-4. X-1A, X-7
763. Giddens JL, Radomski SB, Hirshberg ED, et al. Urodynamic findings in adults with the tethered cord syndrome. *J Urol.* 1999 Apr;161(4):1249-54. X-1F
764. Gilja I. Tansvaginal needle suspension operation: the way we do it. Clinical and urodynamic study: long-term results. *Eur Urol.* 2000 Mar;37(3):325-30. X-1B
765. Gilleran JP, Lemack GE and Zimmern PE. Reduction of moderate-to-large cystocele during urodynamic evaluation using a vaginal gauze pack: 8-year experience. *BJU Int.* 2006 Feb;97(2):292-5. X-1E
766. Gillespie WA, Henderson EP, Linton KB, et al. Microbiology of the urethral (frequency and dysuria) syndrome. A controlled study with 5-year review. *Br J Urol.* 1989 Sep;64(3):270-4. X-1G
767. Gillon G, Engelstein D and Servadio C. Risk factors and their effect on the results of Burch colposuspension for urinary stress incontinence. *Isr J Med Sci.* 1992 Jun;28(6):354-6. X-1B
768. Gillon G, Savir A, Nissenkorn I, et al. Results and relevance of urodynamic studies in the management of urinary incontinence in women over eighty. *Isr J Med Sci.* 1989 Jan;25(1):25-7. X-7
769. Giramonti KM, Kogan BA and Halpern LF. The effects of anticholinergic drugs on attention span and short-term memory skills in children. *Neurourol Urodyn.* 2008;27(4):315-8. X-4

770. Girard JM, Deruelle P, Collinet P, et al. Efficacy and complications in the surgical treatment of stress urinary incontinence by insertion of a silicone-coated polyester tape (Lift). *Eur J Obstet Gynecol Reprod Biol.* 2006 May 1;126(1):107-12. X-1A
771. Gittes RF and Nakamura RM. Female urethral syndrome. A female prostatitis? *West J Med.* 1996 May;164(5):435-8. X-1A
772. Giuseppe PG, Pace G and Vicentini C. Sexual function in women with urinary incontinence treated by pelvic floor transvaginal electrical stimulation. *J Sex Med.* 2007 May;4(3):702-7. X-7
773. Gladh G, Mattsson S and Lindstrom S. Anogenital electrical stimulation as treatment of urge incontinence in children. *BJU Int.* 2001 Mar;87(4):366-71. X-4
774. Gladh G, Mattsson S and Lindstrom S. Outcome of the bladder cooling test in children with nonneurogenic bladder problems. *J Urol.* 2004 Sep;172(3):1095-8. X-1G, X-1H, X-4
775. Gladman MA, Scott SM, Chan CL, et al. Rectal hyposensitivity: prevalence and clinical impact in patients with intractable constipation and fecal incontinence. *Dis Colon Rectum.* 2003 Feb;46(2):238-46. X-11
776. Glavind K and Larsen EH. Results and complications of tension-free vaginal tape (TVT) for surgical treatment of female stress urinary incontinence. *Int Urogynecol J Pelvic Floor Dysfunct.* 2001;12(6):370-2. X-1B
777. Glazer HI, Romanzi L and Polaneczky M. Pelvic floor muscle surface electromyography. Reliability and clinical predictive validity. *J Reprod Med.* 1999 Sep;44(9):779-82. X-1H
778. Glazier DB, Ankem MK, Ferlise V, et al. Utility of biofeedback for the daytime syndrome of urinary frequency and urgency of childhood. *Urology.* 2001 Apr;57(4):791-3; discussion 793-4. X-4
779. Glazier DB, Murphy DP, Fleisher MH, et al. Evaluation of the utility of video-urodynamics in children with urinary tract infection and voiding dysfunction. *Br J Urol.* 1997 Nov;80(5):806-8. X-1H, X-4, X-6
780. Glezerman M, Glasner M, Bar-David J, et al. The setting-up of an ambulatory urologic-gynecologic service. Management of patients and results. *Eur J Obstet Gynecol Reprod Biol.* 1986 Mar;21(3):125-33. X-1B, X-7
781. Glia A, Gylin M, Akerlund JE, et al. Biofeedback training in patients with fecal incontinence. *Dis Colon Rectum.* 1998 Mar;41(3):359-64. X-1I, X-7
782. Godec CJ. Detrusor hyperreflexia inhibited by anal dilatation. *Urology.* 1980 Mar;15(3):321-4. X-11
783. Godec CJ and Cass AS. Cystometric variations during postural changes and functional electrical stimulation of the pelvic floor muscles. *J Urol.* 1980 May;123(5):722-5. X-1G
784. Godec CJ, Cass AS and Ruiz E. Another aspect of acute urinary retention in young patients. *Ann Emerg Med.* 1982 Sep;11(9):471-4. X-1A, X-7
785. Godec CJ, Esho J and Cass AS. Correlation among cystometry, urethral pressure profilometry and pelvic floor electromyography in the evaluation of female patients with voiding dysfunction symptoms. *J Urol.* 1980 Nov;124(5):678-82. X-1H
786. Goepel M, Hoffmann JA, Piro M, et al. Prevalence and physician awareness of symptoms of urinary bladder dysfunction. *Eur Urol.* 2002 Mar;41(3):234-9. X-6
787. Goff BA, Mandel LS, Drescher CW, et al. Development of an ovarian cancer symptom index: possibilities for earlier detection. *Cancer.* 2007 Jan 15;109(2):221-7. X-11
788. Goff BA, Mandel LS, Melancon CH, et al. Frequency of symptoms of ovarian cancer in women presenting to primary care clinics. *JAMA.* 2004 Jun 9;291(22):2705-12. X-1A
789. Goffin JR, Rajan R and Souhami L. Tolerance of radiotherapy and chemotherapy in elderly patients with bladder cancer. *Am J Clin Oncol.* 2004 Apr;27(2):172-7. X-11
790. Gold RS, Groutz A, Pauzner D, et al. Bladder perforation during tension-free vaginal tape surgery: does it matter? *J Reprod Med.* 2007 Jul;52(7):616-8. X-1B
791. Goldberg RP, Kwon C, Gandhi S, et al. Urinary incontinence after multiple gestation and delivery: impact on quality of life. *Int Urogynecol J Pelvic Floor Dysfunct.* 2005 Sep-Oct;16(5):334-6. X-11
792. Goldberg RP, Kwon C, Gandhi S, et al. Urinary incontinence among mothers of multiples: the protective effect of cesarean delivery. *Am J Obstet Gynecol.* 2003 Jun;188(6):1447-50; discussion 1450-3. X-1A
793. Goldman HB, Amundsen CL, Mangel J, et al. Dorsal genital nerve stimulation for the treatment of overactive bladder symptoms. *Neurourol Urodyn.* 2008;27(6):499-503. X-7
794. Goldman HB and Dmochowski RR. Lower urinary tract dysfunction in patients with gastroparesis. *J Urol.* 1997 May;157(5):1823-5. X-1A
795. Gomes CM, Hisano M, Machado LR, et al. Urological manifestations of chronic schistosomal myeloradiculopathy. *BJU Int.* 2005 Oct;96(6):853-6. X-1F
796. Gopal M, Haynes K, Bellamy SL, et al. Discontinuation rates of anticholinergic medications used for the treatment of lower urinary tract symptoms. *Obstet Gynecol.* 2008 Dec;112(6):1311-8. X-11
797. Gordon D, Gold R, Pauzner D, et al. Tension-free vaginal tape in the elderly: is it a safe procedure? *Urology.* 2005 Mar;65(3):479-82. X-1B
798. Gordon D, Gold RS, Pauzner D, et al. Combined genitourinary prolapse repair and prophylactic tension-free vaginal tape in women with severe prolapse and occult stress urinary incontinence: preliminary results. *Urology.* 2001 Oct;58(4):547-50. X-1E
799. Gordon D, Groutz A, Ascher-Landsberg J, et al. Double-blind, placebo-controlled study of magnesium hydroxide for treatment of sensory urgency and detrusor instability: preliminary results. *Br J Obstet Gynaecol.* 1998 Jun;105(6):667-9. X-7
800. Gordon D, Groutz A, Goldman G, et al. Anal incontinence: prevalence among female patients attending a urogynecologic clinic. *Neurourol Urodyn.* 1999;18(3):199-204. X-1F
801. Gordon D, Groutz A, Sinai T, et al. Sexual function in women attending a urogynecology clinic. *Int Urogynecol J Pelvic Floor Dysfunct.* 1999;10(5):325-8. X-11
802. Gormley EA, Griffiths DJ, McCracken PN, et al. Polypharmacy and its effect on urinary incontinence in a geriatric population. *Br J Urol.* 1993 Mar;71(3):265-9. X-1A

803. Gorton E and Stanton S. Ambulatory urodynamics: do they help clinical management? *BJOG*. 2000 Mar;107(3):316-9. X-7
804. Graham CA and Mallett VT. Race as a predictor of urinary incontinence and pelvic organ prolapse. *Am J Obstet Gynecol*. 2001 Jul;185(1):116-20. X-1H
805. Graham G, Gupta S and Aarons L. Determination of an optimal dosage regimen using a Bayesian decision analysis of efficacy and adverse effect data. *J Pharmacokinet Pharmacodyn*. 2002 Feb;29(1):67-88. X-1I
806. Granville L. The burden of illness for overactive bladder. *Manag Care Interface*. 2005;18 Suppl B:16-9. X-2
807. Graves PL and Thomas CB. Habits of nervous tension and suicide. *Suicide Life Threat Behav*. 1991 Summer;21(2):91-105. X-1A, 808. Gravina GL, Brandetti F, Martini P, et al. Measurement of the thickness of the urethrovaginal space in women with or without vaginal orgasm. *J Sex Med*. 2008 Mar;5(3):610-8. X-II
809. Gray M, Albo M and Huffstutler S. Interstitial cystitis: a guide to recognition, evaluation, and management for nurse practitioners. *J Wound Ostomy Continence Nurs*. 2002 Mar;29(2):93-102. X-1D, X-2
810. Gray M, McClain R, Peruggia M, et al. A model for predicting motor urge urinary incontinence. *Nurs Res*. 2001 Mar-Apr;50(2):116-22. X-1H
811. Gray R, Wagg A and Malone-Lee JG. Differences in detrusor contractile function in women with neuropathic and idiopathic detrusor instability. *Br J Urol*. 1997 Aug;80(2):222-6. X-1G, X-1H
812. Green M, Filippou A, Sant G, et al. Expression of intercellular adhesion molecules in the bladder of patients with interstitial cystitis. *Urology*. 2004 Apr;63(4):688-93. X-1D, X-6, X-7
813. Green SA, Alon A, Ianus J, et al. Efficacy and safety of a neurokinin-1 receptor antagonist in postmenopausal women with overactive bladder with urge urinary incontinence. *J Urol*. 2006 Dec;176(6 Pt 1):2535-40; discussion 2540. X-9
814. Greenberg P, Brown J, Yates T, et al. Voiding urges perceived by patients with interstitial cystitis/painful bladder syndrome. *Neurourol Urodyn*. 2008;27(4):287-90. X-1D
815. Greenfield S, Friedland G, Scifers S, et al. Protocol management of dysuria, urinary frequency, and vaginal discharge. *Ann Intern Med*. 1974 Oct;81(4):452-7. X-II
816. Greenfield S, Friedland G, Scifers S, et al. Protocol management of dysuria, urinary frequency, and vaginal discharge. *J Fam Pract*. 1975 Jun;2(3):179-84. X-II
817. Griffiths D. Clinical studies of cerebral and urinary tract function in elderly people with urinary incontinence. *Behav Brain Res*. 1998 May;92(2):151-5. X-II
818. Griffiths D, Derbyshire S, Stenger A, et al. Brain control of normal and overactive bladder. *J Urol*. 2005 Nov;174(5):1862-7. X-1G, X-7
819. Griffiths D, Tadic SD, Schaefer W, et al. Cerebral control of the bladder in normal and urge-incontinent women. *Neuroimage*. 2007 Aug 1;37(1):1-7. X-1H, X-7
820. Griffiths DJ. Urodynamic assessment of bladder function. *Br J Urol*. 1977 Feb;49(1):29-36. X-1G, X-1H
821. Griffiths DJ, McCracken PN and Harrison GM. Incontinence in the elderly: objective demonstration and quantitative assessment. *Br J Urol*. 1991 May;67(5):467-71. X-1H
822. Griffiths DJ, McCracken PN, Harrison GM, et al. Characteristics of urinary incontinence in elderly patients studied by 24-hour monitoring and urodynamic testing. *Age Ageing*. 1992 May;21(3):195-201. X-1H
823. Griffiths DJ, McCracken PN, Harrison GM, et al. Relationship of fluid intake to voluntary micturition and urinary incontinence in geriatric patients. *Neurourol Urodyn*. 1993;12(1):1-7. X-II, X-5
824. Griffiths DJ, McCracken PN, Harrison GM, et al. Cerebral aetiology of urinary urge incontinence in elderly people. *Age Ageing*. 1994 May;23(3):246-50. X-1A
825. Griffiths DJ, McCracken PN, Harrison GM, et al. Urge incontinence and impaired detrusor contractility in the elderly. *Neurourol Urodyn*. 2002;21(2):126-31. X-1H
826. Griffiths DJ, McCracken PN, Harrison GM, et al. Urinary incontinence in the elderly: the brain factor. *Scand J Urol Nephrol Suppl*. 1994;157:83-8. X-1F, X-1H
827. Griffiths DJ, McCracken PN, Harrison GM, et al. Urge incontinence in elderly people: factors predicting the severity of urine loss before and after pharmacological treatment. *Neurourol Urodyn*. 1996;15(1):53-7. X-7
828. Griffiths DJ and Scholtmeijer RJ. Place of the free flow curve in the urodynamic investigation of children. *Br J Urol*. 1984 Oct;56(5):474-7. X-4
829. Grigoleit U, Murtz G, Laschke S, et al. Efficacy, tolerability and safety of propiverine hydrochloride in children and adolescents with congenital or traumatic neurogenic detrusor overactivity—a retrospective study. *Eur Urol*. 2006 Jun;49(6):1114-20; discussion 1120-1. X-1F, X-4
830. Grimby A, Milsom I, Molander U, et al. The influence of urinary incontinence on the quality of life of elderly women. *Age Ageing*. 1993 Mar;22(2):82-9. X-6
831. Grischke EM, Anton H, Stolz W, et al. Urodynamic assessment and lateral urethrocytography. A comparison of two diagnostic procedures for female urinary incontinence. *Acta Obstet Gynecol Scand*. 1991;70(3):225-9. X-1H
832. Groen J, Amiel C and Bosch JL. Chronic pudendal nerve neuromodulation in women with idiopathic refractory detrusor overactivity incontinence: results of a pilot study with a novel minimally invasive implantable mini-stimulator. *Neurourol Urodyn*. 2005;24(3):226-30. X-7
833. Groen J, Klijn AJ, Bosch JL, et al. Diagnosis and grading of detrusor instability using a computerized algorithm. *J Urol*. 1998 May;159(5):1669-74. X-4
834. Groen J, Ruud Bosch JL and van Mastrigt R. Sacral neuromodulation in women with idiopathic detrusor overactivity incontinence: decreased overactivity but unchanged bladder contraction strength and urethral resistance during voiding. *J Urol*. 2006 Mar;175(3 Pt 1):1005-9; discussion 1009. X-7
835. Groen J, van Mastrigt R and Bosch JL. Computerized assessment of detrusor instability in patients treated with sacral neuromodulation. *J Urol*. 2001 Jan;165(1):169-73. X-1H, X-7

836. Groenendijk PM, Halilovic M, Chandi DD, et al. Extracorporeal magnetic innervation therapy: assessment of clinical efficacy in relation to urodynamic parameters. *Scand J Urol Nephrol.* 2008;42(5):433-6. X-7
837. Groenendijk PM, Heesakkers JP and Lycklama ANAA. Urethral instability and sacral nerve stimulation-a better parameter to predict efficacy? *J Urol.* 2007 Aug;178(2):568-72; discussion 572. X-7
838. Grosse J, Kramer G and Stohrer M. Success of repeat detrusor injections of botulinum a toxin in patients with severe neurogenic detrusor overactivity and incontinence. *Eur Urol.* 2005 May;47(5):653-9. X-1F
839. Grossklau DJ and Franke JJ. Treatment of elderly women with urge incontinence in middle tennessee: a single institution practice-based study. *Tenn Med.* 2000 Dec;93(12):457-60. X-7
840. Groutz A, Blaivas JG, Hyman MJ, et al. Pubovaginal sling surgery for simple stress urinary incontinence: analysis by an outcome score. *J Urol.* 2001 May;165(5):1597-600. X-1B
841. Groutz A, Blaivas JG, Kesler SS, et al. Outcome results of transurethral collagen injection for female stress incontinence: assessment by urinary incontinence score. *J Urol.* 2000 Dec;164(6):2006-9. X-1B
842. Groutz A, Chaikin DC, Theusen E, et al. Use of cadaveric solvent-dehydrated fascia lata for cystocele repair--preliminary results. *Urology.* 2001 Aug;58(2):179-83. X-1I
843. Groutz A, Gold R, Pauzner D, et al. Tension-free vaginal tape (TVT) for the treatment of occult stress urinary incontinence in women undergoing prolapse repair: a prospective study of 100 consecutive cases. *Neurourol Urodyn.* 2004;23(7):632-5. X-1B
844. Groutz A, Gordon D, Wolman I, et al. Persistent postpartum urinary retention in contemporary obstetric practice. Definition, prevalence and clinical implications. *J Reprod Med.* 2001 Jan;46(1):44-8. X-1A
845. Groutz A, Gordon D, Wolman I, et al. Tension-free vaginal tape for stress urinary incontinence: Is there a learning curve? *Neurourol Urodyn.* 2002;21(5):470-2. X-1B
846. Groutz A, Samandarov A, Gold R, et al. Role of urethroscopy in the evaluation of refractory idiopathic detrusor instability. *Urology.* 2001 Oct;58(4):544-6. X-1H
847. Gruneberger A. Treatment of motor urge incontinence with clenbuterol and flavoxate hydrochloride. *Br J Obstet Gynaecol.* 1984 Mar;91(3):275-8. X-7
848. Gu FL, Reng ZY, Shang GZ, et al. Treatment of urgency and urge incontinence with flavoxate in the People's Republic of China. *J Int Med Res.* 1987 Sep-Oct;15(5):312-8. X-4
849. Gu J, Restorick JM, Blank MA, et al. Vasoactive intestinal polypeptide in the normal and unstable bladder. *Br J Urol.* 1983 Dec;55(6):645-7. X-1G
850. Guarisi T, Pinto-Neto AM, Herrmann V, et al. Urodynamics in climacteric women with urinary incontinence: correlation with route of delivery. *Int Urogynecol J Pelvic Floor Dysfunct.* 2002 Nov;13(6):366-71; discussion 371. X-1G
851. Guerette NL, Bena JF and Davila GW. Transobturator slings for stress incontinence: using urodynamic parameters to predict outcomes. *Int Urogynecol J Pelvic Floor Dysfunct.* 2008 Jan;19(1):97-102. X-1B
852. Guerra LA, Pike J, Milks J, et al. Outcome in patients who underwent tethered cord release for occult spinal dysraphism. *J Urol.* 2006 Oct;176(4 Pt 2):1729-32. X-II, X-4
853. Guerrero K, Emery S, Owen L, et al. Intravesical oxybutynin: practicalities of clinical use. *J Obstet Gynaecol.* 2006 Feb;26(2):141-3. X-7
854. Gunnarsson M and Mattiasson A. Circumvaginal surface electromyography in women with urinary incontinence and in healthy volunteers. *Scand J Urol Nephrol Suppl.* 1994;157:89-95. X-1G, X-7
855. Haab F, Richard F, Amarenco G, et al. Comprehensive evaluation of bladder and urethral dysfunction symptoms: development and psychometric validation of the Urinary Symptom Profile (USP) questionnaire. *Urology.* 2008 Apr;71(4):646-56. X-1I
856. Haab F, Sananes S, Amarenco G, et al. Results of the tension-free vaginal tape procedure for the treatment of type II stress urinary incontinence at a minimum followup of 1 year. *J Urol.* 2001 Jan;165(1):159-62. X-1B
857. Haarala M, Alanen A, Hietarinta M, et al. Lower urinary tract symptoms in patients with Sjogren's syndrome and systemic lupus erythematosus. *Int Urogynecol J Pelvic Floor Dysfunct.* 2000;11(2):84-6. X-1F
858. Haeusler G, Hanzal E, Joura E, et al. Differential diagnosis of detrusor instability and stress-incontinence by patient history: the Gaudenz-Incontinence-Questionnaire revisited. *Acta Obstet Gynecol Scand.* 1995 Sep;74(8):635-7. X-1H
859. Haferkamp A, Dorsam J and Elbadawi A. Ultrastructural diagnosis of neuropathic detrusor overactivity: validation of a common myogenic mechanism. *Adv Exp Med Biol.* 2003;539(Pt A):281-91. X-1F, X-1H
860. Haferkamp A, Dorsam J, Resnick NM, et al. Structural basis of neurogenic bladder dysfunction. II. Myogenic basis of detrusor hyperreflexia. *J Urol.* 2003 Feb;169(2):547-54. X-1F, X-1G
861. Haferkamp A, Freund T, Wagener N, et al. Distribution of neuropeptide Y-containing nerves in the neurogenic and non-neurogenic detrusor. *BJU Int.* 2006 Feb;97(2):393-9. X-1G, X-7
862. Haferkamp A, Mundhenk J, Bastian PJ, et al. Increased expression of connexin 43 in the overactive neurogenic detrusor. *Eur Urol.* 2004 Dec;46(6):799-805. X-1G, X-7
863. Haferkamp A, Schurch B, Reitz A, et al. Lack of ultrastructural detrusor changes following endoscopic injection of botulinum toxin type a in overactive neurogenic bladder. *Eur Urol.* 2004 Dec;46(6):784-91. X-1F
864. Hagglund D, Walker-Engstrom ML, Larsson G, et al. Quality of life and seeking help in women with urinary incontinence. *Acta Obstet Gynecol Scand.* 2001 Nov;80(11):1051-5. X-1I

865. Hains G, Hains F, Descarreaux M, et al. Urinary incontinence in women treated by ischemic compression over the bladder area: a pilot study. *Journal of Chiropractic Medicine*. 2007 Dec;6(4):132-40. X-1B, X-7
866. Hajebrahimi S, Altaweel W, Cadoret J, et al. Efficacy of botulinum-A toxin in adults with neurogenic overactive bladder: initial results. *Can J Urol*. 2005 Feb;12(1):2543-6. X-1F, X-7
867. Hallen B, Bogentoft S, Sandquist S, et al. Tolerability and steady-state pharmacokinetics of terodiline and its main metabolites in elderly patients with urinary incontinence. *Eur J Clin Pharmacol*. 1989;36(5):487-93. X-7
868. Hampel C, Wienhold D, Benken N, et al. Definition of overactive bladder and epidemiology of urinary incontinence. *Urology*. 1997 Dec;50(6A Suppl):4-14; discussion 15-7. X-2
869. Hancock R, Bender P, Dayhoff N, et al. Factors associated with nursing interventions to reduce incontinence in hospitalized older adults. *Urol Nurs*. 1996 Sep;16(3):79-85. X-1A, X-2, X-5, X-6
870. Handa VL, Harvey L, Fox HE, et al. Parity and route of delivery: does cesarean delivery reduce bladder symptoms later in life? *Am J Obstet Gynecol*. 2004 Aug;191(2):463-9. X-6
871. Hanley HG. Urge incontinence. *Br J Urol*. 1965 Dec;37(6):678-80. X-1I
872. Hannestad YS, Lie RT, Rortveit G, et al. Familial risk of urinary incontinence in women: population based cross sectional study. *BMJ*. 2004 Oct 16;329(7471):889-91. X-1I
873. Hanno PM. Analysis of long-term Elmiron therapy for interstitial cystitis. *Urology*. 1997 May;49(5A Suppl):93-9. X-1D
874. Hanno PM, Buehler J and Wein AJ. Use of amitriptyline in the treatment of interstitial cystitis. *J Urol*. 1989 Apr;141(4):846-8. X-1D
875. Hanno PM, Landis JR, Matthews-Cook Y, et al. The diagnosis of interstitial cystitis revisited: lessons learned from the National Institutes of Health Interstitial Cystitis Database study. *J Urol*. 1999 Feb;161(2):553-7. X-1D
876. Hansen BM, Bonnesen T, Hvidberg JE, et al. Changes in symptoms and colpo-cystourethrography in 35 patients before and after total abdominal hysterectomy: a prospective study. *Urol Int*. 1985;40(4):224-6. X-1A, X-7
877. Hansen J, Borau A, Rodriguez A, et al. Urethral sphincter EMG as event detector for Neurogenic detrusor overactivity. *IEEE Trans Biomed Eng*. 2007 Jul;54(7):1212-9. X-1F, X-1H
878. Hansen J, Media S, Nohr M, et al. Treatment of neurogenic detrusor overactivity in spinal cord injured patients by conditional electrical stimulation. *J Urol*. 2005 Jun;173(6):2035-9. X-1F
879. Hanson LA, Schulz JA, Flood CG, et al. Vaginal pessaries in managing women with pelvic organ prolapse and urinary incontinence: patient characteristics and factors contributing to success. *Int Urogynecol J Pelvic Floor Dysfunct*. 2006 Feb;17(2):155-9. X-1E
880. Hansson S, Hjalmas K, Jodal U, et al. Lower urinary tract dysfunction in girls with untreated asymptomatic or covert bacteriuria. *J Urol*. 1990 Feb;143(2):333-5. X-1A, X-4
881. Harding C, Dorkin TJ and Thorpe AC. Is low bladder compliance predictive of detrusor overactivity? *Neurourol Urodyn*. 2009;28(1):74-7. X-1H
882. Harding G, Cella D, Robinson D, Jr., et al. Symptom burden among patients with renal cell carcinoma (RCC): content for a symptom index. *Health Qual Life Outcomes*. 2007;5:34. X-1I
883. Hardy BG, Shulman KI and Zucchero C. Gradual discontinuation of lithium augmentation in elderly patients with unipolar depression. *J Clin Psychopharmacol*. 1997 Feb;17(1):22-6. X-1I, X-7
884. Harms L, Emons G, Bader W, et al. Funneling before and after anti-incontinence surgery--a prognostic indicator? Part 2: tension-free vaginal tape. *Int Urogynecol J Pelvic Floor Dysfunct*. 2007 Mar;18(3):289-94. X-1B
885. Harpe SE, Szeinbach SL, Caswell RJ, et al. The relative importance of health related quality of life and prescription insurance coverage in the decision to pharmacologically manage symptoms of overactive bladder. *J Urol*. 2007 Dec;178(6):2532-6; discussion 2536. X-1I
886. Harper M, Popat RB, Dasgupta R, et al. A minimally invasive technique for outpatient local anaesthetic administration of intradetrusor botulinum toxin in intractable detrusor overactivity. *BJU Int*. 2003 Aug;92(3):325-6. X-2
887. Harris RL, Cundiff GW, Coates KW, et al. Urinary incontinence and pelvic organ prolapse in nulliparous women. *Obstet Gynecol*. 1998 Dec;92(6):951-4. X-1H
888. Harris RL, Cundiff GW, Theofrastous JP, et al. Bladder compliance in neurologically intact women. *Neurourol Urodyn*. 1996;15(5):483-8. X-1H
889. Harrison GL and Memel DS. Urinary incontinence in women: its prevalence and its management in a health promotion clinic. *Br J Gen Pract*. 1994 Apr;44(381):149-52. X-6
890. Hartigan-Go K, Bateman DN, Daly AK, et al. Stereoselective cardiotoxic effects of terodiline. *Clin Pharmacol Ther*. 1996 Jul;60(1):89-98. X-1G
891. Hartnett NM and Saver BG. Is extended-release oxybutynin (Ditropan XL) or tolterodine (Detrol) more effective in the treatment of an overactive bladder? *J Fam Pract*. 2001 Jul;50(7):571. X-2
892. Harvey MA, Baker K and Wells GA. Tolterodine versus oxybutynin in the treatment of urge urinary incontinence: a meta-analysis. *Am J Obstet Gynecol*. 2001 Jul;185(1):56-61. X-2
893. Hasan ST, Marshall C, Robson WA, et al. Clinical outcome and quality of life following enterocystoplasty for idiopathic detrusor instability and neurogenic bladder dysfunction. *Br J Urol*. 1995 Nov;76(5):551-7. X-7
894. Hasan ST, Robson WA, Pridie AK, et al. Transcutaneous electrical nerve stimulation and temporary S3 neuromodulation in idiopathic detrusor instability. *J Urol*. 1996 Jun;155(6):2005-11. X-7
895. Hasan ST, Robson WA, Ramsden PD, et al. Outcome of endoscopic bladder transection. *Br J Urol*. 1995 May;75(5):592-6. X-7
896. Hashim H and Abrams P. Is the bladder a reliable witness for predicting detrusor overactivity? *J Urol*. 2006 Jan;175(1):191-4; discussion 194-5. X-1H, X-6

897. Hashim H and Abrams P. How should patients with an overactive bladder manipulate their fluid intake? *BJU Int.* 2008 Jul;102(1):62-6. X-7
898. Hashim H, Malmberg L, Graugaard-Jensen C, et al. Desmopressin, as a "designer-drug," in the treatment of overactive bladder syndrome. *Neurourol Urodyn.* 2009;28(1):40-6. X-4, X-7
899. Hassouna ME and Ghoniem GM. Long-term outcome and quality of life after modified pubovaginal sling for intrinsic sphincteric deficiency. *Urology.* 1999 Feb;53(2):287-91. X-1B
900. Hastie KJ and Moisey CU. Are urodynamics necessary in female patients presenting with stress incontinence? *Br J Urol.* 1989 Feb;63(2):155-6. X-1B
901. Hastwell GB. Psychogenic pelvic pain or occult prolapse syndrome? *Med J Aust.* 1986 Apr 14;144(8):405-7. X-1E
902. Hattori T, Yasuda K, Sakakibara R, et al. Micturitional disturbance in ossification of the posterior longitudinal ligament in the cervical spine. *J Spinal Disord.* 1990 Dec;3(4):285-7. X-1F
903. Haylen BT, Chetty N, Logan V, et al. Is sensory urgency part of the same spectrum of bladder dysfunction as detrusor overactivity? *Int Urogynecol J Pelvic Floor Dysfunct.* 2007 Feb;18(2):123-8. X-1I
904. Haylen BT, Krishnan S, Schulz S, et al. Has the true prevalence of voiding difficulty in urogynecology patients been underestimated? *Int Urogynecol J Pelvic Floor Dysfunct.* 2007 Jan;18(1):53-6. X-1A, X-1H
905. Haylen BT, Parys BT, Anyaegbunam WI, et al. Urine flow rates in male and female urodynamic patients compared with the Liverpool nomograms. *Br J Urol.* 1990 May;65(5):483-7. X-1H, X-1G
906. Haylen BT, Sutherst JR and Frazer MI. Is the investigation of most stress incontinence really necessary? *Br J Urol.* 1989 Aug;64(2):147-9. X-1B, X-1H, X-6
907. Hebjorn S. Treatment of detrusor hyperreflexia in multiple sclerosis: a double-blind, crossover clinical trial comparing methantheline bromide (Banthine), flavoxate chloride (Urispas) and meladrazine tartrate (Lisidonil). *Urol Int.* 1977;32(2-3):209-17. X-1F
908. Hebjorn S, Andersen JT, Walter S, et al. Detrusor hyperreflexia. A survey on its etiology and treatment. *Scand J Urol Nephrol.* 1976;10(2):103-9. X-1I
909. Heckerling PS, Canaris GJ, Flach SD, et al. Predictors of urinary tract infection based on artificial neural networks and genetic algorithms. *Int J Med Inform.* 2007 Apr;76(4):289-96. X-1A
910. Hedlund H, Schultz A, Talseth T, et al. Sacral neuromodulation in Norway: clinical experience of the first three years. *Scand J Urol Nephrol Suppl.* 2002(210):87-95. X-7
911. Heilenkötter K, Bachmann C, Janhsen E, et al. Prospective evaluation of inpatient and outpatient bladder training in children with functional urinary incontinence. *Urology.* 2006 Jan;67(1):176-80. X-1I, X-4
912. Hellerstein S and Linebarger JS. Voiding dysfunction in pediatric patients. *Clin Pediatr (Phila).* 2003 Jan-Feb;42(1):43-9. X-1I, X-4
913. Hellerstein S and Zguta AA. Outcome of overactive bladder in children. *Clin Pediatr (Phila).* 2003 Jul-Aug;42(6):553-6. X-4
914. Hellstrom AL, Hjalmas K and Jodal U. Terodiline in the treatment of children with unstable bladders. *Br J Urol.* 1989 Apr;63(4):358-62. X-4
915. Hellstrom PA, Tammela TL, Kontturi MJ, et al. The bladder cooling test for urodynamic assessment: analysis of 400 examinations. *Br J Urol.* 1991 Mar;67(3):275-9. X-1G, X-1H
916. Henningssohn L, Steven K, Kallestrup EB, et al. Distressful symptoms and well-being after radical cystectomy and orthotopic bladder substitution compared with a matched control population. *J Urol.* 2002 Jul;168(1):168-74; discussion 174-5. X-1A
917. Herbst F, Kamm MA, Morris GP, et al. Gastrointestinal transit and prolonged ambulatory colonic motility in health and faecal incontinence. *Gut.* 1997 Sep;41(3):381-9. X-1A, X-7
918. Herwig R, Bruns F, Strasser H, et al. Late urologic effects after adjuvant irradiation in stage I endometrial carcinoma. *Urology.* 2004 Feb;63(2):354-8. X-1I, X-7
919. Heslington K and Hilton P. Ambulatory monitoring and conventional cystometry in asymptomatic female volunteers. *Br J Obstet Gynaecol.* 1996 May;103(5):434-41. X-1H, X-7
920. Hicks A, McCafferty GP, Riedel E, et al. GW427353 (solabegron), a novel, selective beta3-adrenergic receptor agonist, evokes bladder relaxation and increases micturition reflex threshold in the dog. *J Pharmacol Exp Ther.* 2007 Oct;323(1):202-9. X-1G X-4
921. Higson RH, Smith JC and Hills W. Intravesical lignocaine and detrusor instability. *Br J Urol.* 1979 Dec;51(6):500-3. X-1B, X-1H, X-7
922. Higson RH, Smith JC and Whelan P. Bladder rupture: an acceptable complication of distension therapy? *Br J Urol.* 1978 Dec;50(7):529-34. X-1A
923. Hill J, Corson RJ, Brandon H, et al. History and examination in the assessment of patients with idiopathic fecal incontinence. *Dis Colon Rectum.* 1994 May;37(5):473-7. X-1I, X-1H
924. Hilton P. Urinary incontinence during sexual intercourse: a common, but rarely volunteered, symptom. *Br J Obstet Gynaecol.* 1988 Apr;95(4):377-81. X-1I
925. Hilton P. A clinical and urodynamic study comparing the Stamey bladder neck suspension and suburethral sling procedures in the treatment of genuine stress incontinence. *Br J Obstet Gynaecol.* 1989 Feb;96(2):213-20. X-1B
926. Hilton P. Urodynamic findings in patients with urogenital fistulae. *Br J Urol.* 1998 Apr;81(4):539-42. X-1I
927. Hilton P and Mayne CJ. The Stamey endoscopic bladder neck suspension: a clinical and urodynamic investigation, including actuarial follow-up over four years. *Br J Obstet Gynaecol.* 1991 Nov;98(11):1141-9. X-1B
928. Hilton P and Stanton SL. The use of desmopressin (DDAVP) in nocturnal urinary frequency in the female. *Br J Urol.* 1982 Jun;54(3):252-5. X-1C
929. Hinata N, Shirakawa T, Okada H, et al. Quantitative analysis of the levels of expression of muscarinic receptor subtype RNA in the detrusor muscle of patients with overactive bladder. *Mol Diagn.* 2004;8(1):17-22. X-1G
930. Hindmarsh JR, Essenhigh DM and Yeates WK. Indications for bladder transection. *Urol Int.* 1978;33(5):322-6. X-7

931. Hindmarsh JR, Gosling PT and Deane AM. Bladder instability. Is the primary defect in the urethra? *Br J Urol.* 1983 Dec;55(6):648-51. X-1G, X-4, X-7
932. Hirao Y, Okajima E, Ozono S, et al. A prospective randomized study of prophylaxis of tumor recurrence following transurethral resection of superficial bladder cancer--intravesical thio-TEPA versus oral UFT. *Cancer Chemother Pharmacol.* 1992;30 Suppl:S26-30. X-11
933. Hirst GR, Watkins AJ, Guerrero K, et al. Botulinum toxin B is not an effective treatment of refractory overactive bladder. *Urology.* 2007 Jan;69(1):69-73. X-7
934. Hitchcock RJ, Duffy PG and Malone PS. Uretrocystoplasty: the 'bladder' augmentation of choice. *Br J Urol.* 1994 May;73(5):575-9. X-11, X-4
935. Hjalmas K, Hellstrom AL, Mogren K, et al. The overactive bladder in children: a potential future indication for tolterodine. *BJU Int.* 2001 Apr;87(6):569-74. X-4, X-7
936. Ho CH, Chen SC, Chung SD, et al. Determining the appropriate length of a double-pigtail ureteral stent by both stent configurations and related symptoms. *J Endourol.* 2008 Jul;22(7):1427-31. X-11
937. Hodgkinson CP and Drukker BH. Infravesical nerve resection for detrusor dyssynergia. The Ingelman-Sundberg operation. *Acta Obstet Gynecol Scand.* 1977;56(4):401-8. X-2
938. Hoebeke P, De Caestecker K, Vande Walle J, et al. The effect of botulinum-A toxin in incontinent children with therapy resistant overactive detrusor. *J Urol.* 2006 Jul;176(1):328-30; discussion 330-1. X-4, X-7
939. Hoebeke P, Van Laecke E, Everaert K, et al. Transcutaneous neuromodulation for the urge syndrome in children: a pilot study. *J Urol.* 2001 Dec;166(6):2416-9. X-4
940. Hoebeke P, Van Laecke E, Renson C, et al. Pelvic floor spasms in children: an unknown condition responding well to pelvic floor therapy. *Eur Urol.* 2004 Nov;46(5):651-4; discussion 654. X-4, X-7
941. Hoebeke P, Van Laecke E, Van Camp C, et al. One thousand video-urodynamic studies in children with non-neurogenic bladder sphincter dysfunction. *BJU Int.* 2001 Apr;87(6):575-80. X-11, X-4
942. Hoebeke PB, Van Gool JD, Van Laecke E, et al. The after-contraction in paediatric urodynamics. *Br J Urol.* 1996 Nov;78(5):780-2. X-1H, X-4
943. Hoekx L, Vermandel A and Wyndaele JJ. Functional bladder capacity after bladder biofeedback predicts long-term outcome in children with nocturnal enuresis. *Scand J Urol Nephrol.* 2003;37(2):120-3. X-1C
944. Hoekx L, Wyndaele JJ and Vermandel A. The role of bladder biofeedback in the treatment of children with refractory nocturnal enuresis associated with idiopathic detrusor instability and small bladder capacity. *J Urol.* 1998 Sep;160(3 Pt 1):858-60. X-11, X-4
945. Holmes DM, Montz FJ and Stanton SL. Oxybutynin versus propantheline in the management of detrusor instability. A patient-regulated variable dose trial. *Br J Obstet Gynaecol.* 1989 May;96(5):607-12. X-7
946. Holmes DM, Plevnik S and Stanton SL. Bladder neck electrical conductivity in female urinary urgency and urge incontinence. *Br J Obstet Gynaecol.* 1989 Jul;96(7):816-20. X-1H, X-7
947. Holmes DM, Plevnik S and Stanton SL. Bladder neck electrical conductivity in the treatment of detrusor instability with biofeedback. *Br J Obstet Gynaecol.* 1989 Jul;96(7):821-6. X-7
948. Holmes NM, Coplen DE, Strand W, et al. Is bladder dysfunction and incontinence associated with ureteroceles congenital or acquired? *J Urol.* 2002 Aug;168(2):718-9. X-11
949. Holmgren C, Nilsson S, Lanner L, et al. Frequency of de novo urgency in 463 women who had undergone the tension-free vaginal tape (TVT) procedure for genuine stress urinary incontinence--a long-term follow-up. *Eur J Obstet Gynecol Reprod Biol.* 2007 May;132(1):121-5. X-1A
950. Holschneider CH, Solh S, Leberz TB, et al. The modified Pereyra procedure in recurrent stress urinary incontinence: a 15-year review. *Obstet Gynecol.* 1994 Apr;83(4):573-8. X-1B
951. Homayoon K, Chen JJ, Cummings JM, et al. Voiding dysfunction: outcome in infants with congenital vesicoureteral reflux. *Urology.* 2005 Nov;66(5):1091-4; discussion 1094. X-11, X-4
952. Homma Y and Kawabe K. Health-related quality of life of Japanese patients with overactive bladder treated with extended-release tolterodine or immediate-release oxybutynin: a randomized, placebo-controlled trial. *World J Urol.* 2004 Oct;22(4):251-6. X-4
953. Homma Y, Kondo Y, Takahashi S, et al. Reproducibility of cystometry in overactive detrusor. *Eur Urol.* 2000 Dec;38(6):681-5. X-1H, X-7
954. Homma Y, Paick JS, Lee JG, et al. Clinical efficacy and tolerability of extended-release tolterodine and immediate-release oxybutynin in Japanese and Korean patients with an overactive bladder: a randomized, placebo-controlled trial. *BJU Int.* 2003 Nov;92(7):741-7. X-4
955. Homma Y and Uemura S. Use of the short form of King's Health Questionnaire to measure quality of life in patients with an overactive bladder. *BJU Int.* 2004 May;93(7):1009-13. X-11
956. Homma Y, Yamaguchi T and Yamaguchi O. A randomized, double-blind, placebo-controlled phase II dose-finding study of the novel anti-muscarinic agent imidafenacin in Japanese patients with overactive bladder. *Int J Urol.* 2008 Sep;15(9):809-15. X-4
957. Homma Y, Yoshida M, Seki N, et al. Symptom assessment tool for overactive bladder syndrome--overactive bladder symptom score. *Urology.* 2006 Aug;68(2):318-23. X-11
958. Hooper P, Tincello DG and Richmond DH. The use of salivary stimulant pastilles to improve compliance in women taking oxybutynin hydrochloride for detrusor instability: a pilot study. *Br J Urol.* 1997 Sep;80(3):414-6. X-7
959. Horstmann M, Schaefer T, Aguilar Y, et al. Neurogenic bladder treatment by doubling the recommended antimuscarinic dosage. *Neurourol Urodyn.* 2006;25(5):441-5. X-1F

960. Hougaard C, Fraser MO, Chien C, et al. A positive modulator of K Ca 2 and K Ca 3 channels, 4,5-dichloro-1,3-diethyl-1,3-dihydro-benzoimidazol-2-one (NS4591), inhibits bladder afferent firing in vitro and bladder overactivity in vivo. *J Pharmacol Exp Ther*. 2009 Jan;328(1):28-39. X-11
961. Hoverd PA and Fowler CJ. Desmopressin in the treatment of daytime urinary frequency in patients with multiple sclerosis. *J Neurol Neurosurg Psychiatry*. 1998 Nov;65(5):778-80. X-1F, X-7
962. Ho-Yin PL, Man-Wah P and Shing-Kai Y. Effects of aging on generic SF-36 quality of life measurements in Hong Kong Chinese women with urinary incontinence. *Acta Obstet Gynecol Scand*. 2003 Mar;82(3):275-9. X-11
963. Hsieh CH, Chen HY, Hsu CS, et al. Risk factors for urinary frequency in Taiwanese women aged 20-59 years. *Taiwan J Obstet Gynecol*. 2006 Dec;45(4):329-32. X-11
964. Hsieh TF, Yu KJ and Lin SY. Possible application of Raman microspectroscopy to verify the interstitial cystitis diagnosis after potassium sensitivity test: phenylalanine or tryptophan as a biomarker. *Dis Markers*. 2007;23(3):147-52. X-1D, X-1H
965. Hu TW and Wagner TH. Health-related consequences of overactive bladder: an economic perspective. *BJU Int*. 2005 Sep;96 Suppl 1:43-5. X-2
966. Hu TW, Wagner TH, Bentkover JD, et al. Costs of urinary incontinence and overactive bladder in the United States: a comparative study. *Urology*. 2004 Mar;63(3):461-5. X-2
967. Huang KH, Kung FT, Liang HM, et al. Management of polypropylene mesh erosion after intravaginal midurethral sling operation for female stress urinary incontinence. *Int Urogynecol J Pelvic Floor Dysfunct*. 2005 Nov-Dec;16(6):437-40. X-1B
968. Huang KH, Kung FT, Liang HM, et al. Concomitant pelvic organ prolapse surgery with TVT procedure. *Int Urogynecol J Pelvic Floor Dysfunct*. 2006 Jan;17(1):60-5. X-1B
969. Huang KH, Kung FT, Liang HM, et al. Concomitant surgery with tension-free vaginal tape. *Acta Obstet Gynecol Scand*. 2003 Oct;82(10):948-53. X-1B
970. Hubeaux K, Deffieux X, Ismael SS, et al. Autonomic nervous system activity during bladder filling assessed by heart rate variability analysis in women with idiopathic overactive bladder syndrome or stress urinary incontinence. *J Urol*. 2007 Dec;178(6):2483-7. X-7
971. Hughes DA and Dubois D. Cost-effectiveness analysis of extended-release formulations of oxybutynin and tolterodine for the management of urge incontinence. *Pharmacoeconomics*. 2004;22(16):1047-59. X-8
972. Hui-Chi H. A checklist for assessing the risk of falls among the elderly. *J Nurs Res*. 2004 Jun;12(2):131-42. X-1A
973. Hullfish KL, Bovbjerg VE and Steers WD. Colpocleisis for pelvic organ prolapse: patient goals, quality of life, and satisfaction. *Obstet Gynecol*. 2007 Aug;110(2 Pt 1):341-5. X-1E
974. Hullfish KL, Fenner DE, Sorser SA, et al. Postpartum depression, urge urinary incontinence, and overactive bladder syndrome: is there an association? *Int Urogynecol J Pelvic Floor Dysfunct*. 2007 Oct;18(10):1121-6. X-11
975. Hung MJ, Ho ES, Shen PS, et al. Urgency is the core symptom of female overactive bladder syndrome, as demonstrated by a statistical analysis. *J Urol*. 2006 Aug;176(2):636-40. X-1H
976. Hutchings A, Griffiths J and Black NA. Surgery for stress incontinence: factors associated with a successful outcome. *Br J Urol*. 1998 Nov;82(5):634-41. X-1B
977. Huwyler M, Schurch B, Knapp PA, et al. Can the ice-water test predict the outcome of intradetrusor injections of botulinum toxin in patients with neurogenic bladder dysfunction? *World J Urol*. 2007 Dec;25(6):613-7. X-1F, X-7
978. Huwyler M, Springer J, Kessler TM, et al. A safe and simple solution for intravesical tension-free vaginal tape erosion: removal by standard transurethral resection. *BJU Int*. 2008 Aug 5;102(5):582-5. X-1G, X-7 (5)
979. Iglesias X and Espuna M. Surgical treatment of urinary stress incontinence using a method for postoperative adjustment of sling tension (Remeex System). *Int Urogynecol J Pelvic Floor Dysfunct*. 2003 Nov;14(5):326-30; discussion 330. X-1B
980. Ijland MM, Fischer DC, Kieback DG, et al. Midline intravaginal slingplasty for treatment of urinary stress incontinence: results of an independent audit up to 2 years after surgery. *Int Urogynecol J Pelvic Floor Dysfunct*. 2005 Nov-Dec;16(6):447-54. X-1F
981. Ikeda K, Kobayashi S, Suzuki M, et al. M(3) receptor antagonism by the novel antimuscarinic agent solifenacin in the urinary bladder and salivary gland. *Naunyn Schmiedebergs Arch Pharmacol*. 2002 Aug;366(2):97-103. X-1G
982. Ilker Y, Tarcan T, Ozveren B, et al. Anterior urethropexy according to lapides: how effective is it in the treatment of genuine female stress incontinence? *Arch Esp Urol*. 2003 Jan-Feb;56(1):95-9. X-1B
983. Imanishi M, Tomishima Y, Itou S, et al. Discovery of a novel series of biphenyl benzoic acid derivatives as potent and selective human beta3-adrenergic receptor agonists with good oral bioavailability. Part I. *J Med Chem*. 2008 Mar 27;51(6):1925-44. X-1G
984. Ingelman-Sundberg A. Urge incontinence in women. *Acta Obstet Gynecol Scand*. 1975;54(2):153-6. X-1F
985. Iosif S, Henriksson L and Ulmsten U. Urethrocytometry as a routine method for the objective evaluation of women with urinary incontinence. *Arch Gynecol*. 1980;230(1):41-7. X-1H
986. Iosif S, Henriksson L and Ulmsten U. The frequency of disorders of the lower urinary tract, urinary incontinence in particular, as evaluated by a questionnaire survey in a gynecological health control population. *Acta Obstet Gynecol Scand*. 1981;60(1):71-6. X-6
987. Irwin DE, Abrams P, Milsom I, et al. Understanding the elements of overactive bladder: questions raised by the EPIC study. *BJU Int*. 2008 Jun;101(11):1381-7. X-11
988. Irwin DE, Milsom I, Kopp Z, et al. Symptom bother and health care-seeking behavior among individuals with overactive bladder. *Eur Urol*. 2008 May;53(5):1029-37. X-11
989. Irwin DE, Milsom I, Kopp Z, et al. Impact of overactive bladder symptoms on employment, social interactions and emotional well-being in six European countries. *BJU Int*. 2006 Jan;97(1):96-100. X-11

990. Irwin P and Samsudin A. Reinvestigation of patients with a diagnosis of interstitial cystitis: common things are sometimes common. *J Urol.* 2005 Aug;174(2):584-7; discussion 587. X-1D
991. Irwin PP, Hammonds WD and Galloway NT. Lumbar epidural blockade for management of pain in interstitial cystitis. *Br J Urol.* 1993 Apr;71(4):413-6. X-1D
992. Ishigooka M, Hashimoto T, Hayami S, et al. Ice water test in patients with overactive bladder due to cerebrovascular accidents and bladder outlet obstruction. *Urol Int.* 1997;58(2):84-7. X-1A, X-7
993. Ishigooka M, Hashimoto T, Izumiya K, et al. Electrical pelvic floor stimulation in the management of urinary incontinence due to neuropathic overactive bladder. *Front Med Biol Eng.* 1993;5(1):1-10. X-1F
994. Ishigooka M, Hashimoto T, Sasagawa I, et al. Terodiline in the treatment of nocturnal enuresis in children. *Int Urol Nephrol.* 1992;24(5):509-13. X-II, X-4
995. Ishigooka M, Hashimoto T, Sasagawa I, et al. Electrical pelvic floor stimulation by percutaneous implantable electrode. *Br J Urol.* 1994 Aug;74(2):191-4. X-7
996. Ishigooka M, Zermann DH, Doggweiler R, et al. Sacral nerve stimulation and diurnal urine volume. *Eur Urol.* 1999 Nov;36(5):421-6. X-7
997. Ishikawa N, Suda S, Sasaki T, et al. Development of a non-invasive treatment system for urinary incontinence using a functional continuous magnetic stimulator (FCMS). *Med Biol Eng Comput.* 1998 Nov;36(6):704-10. X-II, X-4, X-5
998. Ishiko O, Hirai K, Sumi T, et al. The urinary incontinence score in the diagnosis of female urinary incontinence. *Int J Gynaecol Obstet.* 2000 Feb;68(2):131-7. X-II
999. Ishiko O, Sumi T, Hirai K, et al. Classification of female urinary incontinence by the scored incontinence questionnaire. *Int J Gynaecol Obstet.* 2000 Jun;69(3):255-60. X-1G
1000. Ismael SS, Epstein T, Bayle B, et al. Bladder cooling reflex in patients with multiple sclerosis. *J Urol.* 2000 Oct;164(4):1280-4. X-1G, X-7
1001. Ismail SI. Recurrent prolapse after sacrocolpopexy for post-hysterectomy vaginal vault prolapse. *J Obstet Gynaecol.* 2007 Apr;27(3):292-6. X-1E
1002. Ito T, Sakakibara R, Yamamoto T, et al. Urinary dysfunction and autonomic control in amyloid neuropathy. *Clin Auton Res.* 2006 Feb;16(1):66-71. X-1F
1003. Ito T, Sakakibara R, Yasuda K, et al. Incomplete emptying and urinary retention in multiple-system atrophy: when does it occur and how do we manage it? *Mov Disord.* 2006 Jun;21(6):816-23. X-1F
1004. Ito T, Ueda T, Honma Y, et al. Recent trends in patient characteristics and therapeutic choices for interstitial cystitis: analysis of 282 Japanese patients. *Int J Urol.* 2007 Dec;14(12):1068-70. X-1D
1005. Ivil KD and Suresh G. Review of augmentation 'clam' cystoplasty in a district general hospital setting. *Int Urol Nephrol.* 2002;34(1):129-32. X-7
1006. Jackson RA, Vittinghoff E, Kanaya AM, et al. Urinary incontinence in elderly women: findings from the Health, Aging, and Body Composition Study. *Obstet Gynecol.* 2004 Aug;104(2):301-7. X-6
1007. Jackson SL, Scholes D, Boyko EJ, et al. Urinary incontinence and diabetes in postmenopausal women. *Diabetes Care.* 2005 Jul;28(7):1730-8. X-1B
1008. Jacobs CD and van der Pas M. A multicenter maintenance study of oral pilocarpine tablets for radiation-induced xerostomia. *Oncology (Williston Park).* 1996 Mar;10(3 Suppl):16-20. X-II
1009. Jacoby K and Rowbotham RK. Double balloon positive pressure urethrography is a more sensitive test than voiding cystourethrography for diagnosing urethral diverticulum in women. *J Urol.* 1999 Dec;162(6):2066-9. X-II, X-7
1010. Jacome EG, Tutera G and Mattox FT. Laparoscopic Burch urethropexy in a private clinical practice. *J Am Assoc Gynecol Laparosc.* 1999 Feb;6(1):39-44. X-1B, X-1E
1011. James ED and Niblett PG. Provisional study to quantify and compare parameters of urine leakage in stress and instability incontinence. *Br J Urol.* 1988 Sep;62(3):223-7. X-1H
1012. James M, Jackson S, Shepherd A, et al. Pure stress leakage symptomatology: is it safe to discount detrusor instability? *Br J Obstet Gynaecol.* 1999 Dec;106(12):1255-8. X-1B, X-1H
1013. Janknegt RA, Weil EH and Eerdmans PH. Improving neuromodulation technique for refractory voiding dysfunctions: two-stage implant. *Urology.* 1997 Mar;49(3):358-62. X-7
1014. Jankovic J, Gilden JL, Hiner BC, et al. Neurogenic orthostatic hypotension: a double-blind, placebo-controlled study with midodrine. *Am J Med.* 1993 Jul;95(1):38-48. X-II
1015. Jansen L and Forbes D. The psychometric testing of a urinary incontinence nursing assessment instrument. *J Wound Ostomy Continence Nurs.* 2006 Jan-Feb;33(1):69-76. X-II
1016. Jarvis GJ. The management of urinary incontinence due to primary vesical sensory urgency by bladder drill. *Br J Urol.* 1982 Aug;54(4):374-6. X-7
1017. Jarvis GJ and Millar DR. The treatment of incontinence due to detrusor instability by bladder drill. *Prog Clin Biol Res.* 1981;78:341-3. X-7
1018. Jeffery S, Fynes M, Lee F, et al. Efficacy and complications of intradetrusor injection with botulinum toxin A in patients with refractory idiopathic detrusor overactivity. *BJU Int.* 2007 Dec;100(6):1302-6. X-7
1019. Jensen LN, Gerstenberg T, Kallestrup EB, et al. Urodynamic evaluation of patients with autosomal dominant pure spastic paraplegia linked to chromosome 2p21-p24. *J Neurol Neurosurg Psychiatry.* 1998 Nov;65(5):693-6. X-1F
1020. Jeon MJ, Chung da J, Park JH, et al. Surgical therapeutic index of tension-free vaginal tape and transobturator tape for stress urinary incontinence. *Gynecol Obstet Invest.* 2008;65(1):41-6. X-1B
1021. Jha S, Moran P, Greenham H, et al. Sexual function following surgery for urodynamic stress incontinence. *Int Urogynecol J Pelvic Floor Dysfunct.* 2007 Aug;18(8):845-50. X-1B

1022. Jha S, Toozs-Hobson P, Parsons M, et al. Does preoperative urodynamics change the management of prolapse? *J Obstet Gynaecol*. 2008 Apr;28(3):320-2. X-1B, X-1E, X-1H
1023. Jitapunkul S and Khovichunkit W. Urinary incontinence in Thai elderly living in Klong Toey slum. *J Med Assoc Thai*. 1998 Mar;81(3):160-8. X-6
1024. Johannesson M, O'Connor RM, Kobelt-Nguyen G, et al. Willingness to pay for reduced incontinence symptoms. *Br J Urol*. 1997 Oct;80(4):557-62. X-1I
1025. Johansen J, Claudi T and Holtedahl K. Insulin treatment for poorly regulated diabetic patients in general practice. Better regulation and symptom relief? *Scand J Prim Health Care*. 1999 Dec;17(4):244-9. X-1I
1026. Johnson DN, Vaglianti RM, Huber SJ, et al. Genitourinary applications of sacral neuromodulation. *W V Med J*. 2003 May-Jun;99(3):111-3. X-2, X-6, X-7
1027. Johnstone JM, Ardran GM and Ramsden PD. A preliminary assessment of bladder distension in the treatment of enuretic children. *Br J Urol*. 1977 Feb;49(1):43-9. X-1A, X-4
1028. Jolleys JV. Reported prevalence of urinary incontinence in women in a general practice. *Br Med J (Clin Res Ed)*. 1988 May 7;296(6632):1300-2. X-1A
1029. Jolleys JV. Diagnosis and management of female urinary incontinence in general practice. *J R Coll Gen Pract*. 1989 Jul;39(324):277-9. X-7
1030. Jones JS, Rackley RR, Berglund R, et al. Porcine small intestinal submucosa as a percutaneous mid-urethral sling: 2-year results. *BJU Int*. 2005 Jul;96(1):103-6. X-1B
1031. Jorgensen L, Lose G and Molsted-Pedersen L. Vaginal repair in female motor urge incontinence. *Eur Urol*. 1987;13(6):382-5. X-7
1032. Jorgensen L, Mortensen SO, Colstrup H, et al. Bladder distension in the management of detrusor instability. *Scand J Urol Nephrol*. 1985;19(2):101-4. X-7
1033. Jose CC, Price A, Norman A, et al. Hypofractionated radiotherapy for patients with carcinoma of the bladder. *Clin Oncol (R Coll Radiol)*. 1999;11(5):330-3. X-1I
1034. Jozwik M, Jr., Jozwik M and Lotocki W. Stress urinary incontinence in women. III. Different tissue biochemistry in patients with concomitant detrusor instability. Preliminary report. *Int Urol Nephrol*. 1998;30(3):279-82. X-1B
1035. Juma S. Anterior vaginal suspension for vaginal vault prolapse. *Tech Urol*. 1995 Fall;1(3):150-6. X-1E
1036. Juma S and Brito CG. Transobturator tape (TOT): Two years follow-up. *Neurourol Urodyn*. 2007;26(1):37-41. X-1B
1037. Juma S, Little NA and Raz S. Vaginal wall sling: four years later. *Urology*. 1992 May;39(5):424-8. X-1B
1038. Jumadilova Z, Varadharajan S, Girase P, et al. Retrospective evaluation of outcomes in patients with overactive bladder receiving tolterodine versus oxybutynin. *Am J Health Syst Pharm*. 2006 Dec 1;63(23):2357-64. X-8
1039. Jundt K, Scheer I, Schiessl B, et al. Physical and sexual abuse in patients with overactive bladder: is there an association? *Int Urogynecol J Pelvic Floor Dysfunct*. 2007 Apr;18(4):449-53. X-1I
1040. Junemann KP, Halaska M, Rittstein T, et al. Propiverine versus tolterodine: efficacy and tolerability in patients with overactive bladder. *Eur Urol*. 2005 Sep;48(3):478-82. X-9
1041. Junemann KP, Hessdorfer E, Unamba-Oparah I, et al. Propiverine hydrochloride immediate and extended release: comparison of efficacy and tolerability in patients with overactive bladder. *Urol Int*. 2006;77(4):334-9. X-9
1042. Kaat LD, Boon AJ, Kamphorst W, et al. Frontal presentation in progressive supranuclear palsy. *Neurology*. 2007 Aug 21;69(8):723-9. X-1A
1043. Kabay SC, Kabay S, Yucel M, et al. Acute urodynamic effects of percutaneous posterior tibial nerve stimulation on neurogenic detrusor overactivity in patients with Parkinson's disease. *Neurourol Urodyn*. 2009;28(1):62-7. X-1F
1044. Kabay SC, Yucel M and Kabay S. Acute effect of posterior tibial nerve stimulation on neurogenic detrusor overactivity in patients with multiple sclerosis: urodynamic study. *Urology*. 2008 Apr;71(4):641-5. X-1F, X-7
1045. Kadar N. The value of bladder filling in the clinical detection of urine loss and selection of patients for urodynamic testing. *Br J Obstet Gynaecol*. 1988 Jul;95(7):698-704. X-1H
1046. Kafri R, Langer R, Dvir Z, et al. Rehabilitation vs drug therapy for urge urinary incontinence: short-term outcome. *Int Urogynecol J Pelvic Floor Dysfunct*. 2007 Apr;18(4):407-11. X-7
1047. Kafri R, Shames J, Raz M, et al. Rehabilitation versus drug therapy for urge urinary incontinence: long-term outcomes. *Int Urogynecol J Pelvic Floor Dysfunct*. 2008 Jan;19(1):47-52. X-7
1048. Kagan G. Emepronium bromide in the management of nightly urinary frequency and urinary incontinence. *Practitioner*. 1970 May;204(223):705-10. X-7
1049. Kajiwaru M, Inoue K, Kato M, et al. Nocturnal enuresis and overactive bladder in children: an epidemiological study. *Int J Urol*. 2006 Jan;13(1):36-41. X-4
1050. Kajiwaru M, Inoue K, Mutaguchi K, et al. The prevalence of overactive bladder and nocturnal enuresis in Japanese early adolescents: a questionnaire survey. *Hinyokika Kyo*. 2006 Feb;52(2):107-11. X-4
1051. Kajiwaru M, Kato M, Mutaguchi K, et al. Overactive bladder in children should be strictly differentiated from monosymptomatic nocturnal enuresis. *Urol Int*. 2008;80(1):57-61. X-1A, X-4
1052. Kajiwaru M and Mutaguchi K. Clinical efficacy and tolerability of gosha-jinki-gan, Japanese traditional herbal medicine, in females with overactive bladder. *Hinyokika Kyo*. 2008 Feb;54(2):95-9. X-7
1053. Kalita J, Shah S, Kapoor R, et al. Bladder dysfunction in acute transverse myelitis: magnetic resonance imaging and neurophysiological and urodynamic correlations. *J Neurol Neurosurg Psychiatry*. 2002 Aug;73(2):154-9. X-1H, X-1I, X-7
1054. Kallestrup EB, Jorgensen SS, Nordling J, et al. Treatment of interstitial cystitis with Cystistat: a hyaluronic acid product. *Scand J Urol Nephrol*. 2005;39(2):143-7. X-1D

1055. Kalsi V, Apostolidis A, Gonzales G, et al. Early effect on the overactive bladder symptoms following botulinum neurotoxin type A injections for detrusor overactivity. *Eur Urol*. 2008 Jul;54(1):181-7. X-7
1056. Kalsi V, Apostolidis A, Popat R, et al. Quality of life changes in patients with neurogenic versus idiopathic detrusor overactivity after intradetrusor injections of botulinum neurotoxin type A and correlations with lower urinary tract symptoms and urodynamic changes. *Eur Urol*. 2006 Mar;49(3):528-35. X-7
1057. Kalsi V, Gonzales G, Popat R, et al. Botulinum injections for the treatment of bladder symptoms of multiple sclerosis. *Ann Neurol*. 2007 Nov;62(5):452-7. X-1F, X-7
1058. Kalsi V, Popat RB, Apostolidis A, et al. Cost-consequence analysis evaluating the use of botulinum neurotoxin-A in patients with detrusor overactivity based on clinical outcomes observed at a single UK centre. *Eur Urol*. 2006 Mar;49(3):519-27. X-7
1059. Kamm MA, Hawley PR and Lennard-Jones JE. Lateral division of the puborectalis muscle in the management of severe constipation. *Br J Surg*. 1988 Jul;75(7):661-3. X-11
1060. Kanayama N, Kanari C, Masuda Y, et al. Drug-drug interactions in the metabolism of imidafenacin: role of the human cytochrome P450 enzymes and UDP-glucuronic acid transferases, and potential of imidafenacin to inhibit human cytochrome P450 enzymes. *Xenobiotica*. 2007 Feb;37(2):139-54. X-1G, X-6
1061. Kang HS, Wang KC, Kim KM, et al. Prognostic factors affecting urologic outcome after untethering surgery for lumbosacral lipoma. *Childs Nerv Syst*. 2006 Sep;22(9):1111-21. X-1F, X-4
1062. Kaplan HJ and Mamo GJ. Pubovaginal sling technique utilizing a unique bone anchor instrumentation system. *Can J Urol*. 2000 Oct;7(5):1116-21. X-1B
1063. Kaplan SA, Santarosa RP and Te AE. Comparison of fascial and vaginal wall slings in the management of intrinsic sphincter deficiency. *Urology*. 1996 Jun;47(6):885-9. X-11
1064. Kaplan SA, Te AE, Young GP, et al. Prospective analysis of 373 consecutive women with stress urinary incontinence treated with a vaginal wall sling: the Columbia-Cornell University experience. *J Urol*. 2000 Nov;164(5):1623-7. X-1B
1065. Karademir K, Baykal K, Sen B, et al. A peripheric neuromodulation technique for curing detrusor overactivity: Stoller afferent neurostimulation. *Scand J Urol Nephrol*. 2005;39(3):230-3. X-7
1066. Karaman MI, Kaya C, Caskurlu T, et al. Urodynamic findings in children with cerebral palsy. *Int J Urol*. 2005 Aug;12(8):717-20. X-1F, X-4
1067. Karantanis E, Fynes MM and Stanton SL. The tension-free vaginal tape in older women. *BJOG*. 2004 Aug;111(8):837-41. X-1B
1068. Karon S. A team approach to bladder retraining: a pilot study. *Urol Nurs*. 2005 Aug;25(4):269-76. X-4
1069. Karram MM and Bhatia NN. Management of coexistent stress and urge urinary incontinence. *Obstet Gynecol*. 1989 Jan;73(1):4-7. X-1B
1070. Karram MM, Segal JL, Vassallo BJ, et al. Complications and untoward effects of the tension-free vaginal tape procedure. *Obstet Gynecol*. 2003 May;101(5 Pt 1):929-32. X-1B
1071. Karsenty G, Chartier-Kastler E, Mozer P, et al. A novel technique to achieve cutaneous continent urinary diversion in spinal cord-injured patients unable to catheterize through native urethra. *Spinal Cord*. 2008 Apr;46(4):305-10. X-1F, X-7
1072. Karsenty G, Elzayat E, Delapparent T, et al. Botulinum toxin type a injections into the trigone to treat idiopathic overactive bladder do not induce vesicoureteral reflux. *J Urol*. 2007 Mar;177(3):1011-4. X-7
1073. Kasirga E, Akil I, Yilmaz O, et al. Evaluation of voiding dysfunctions in children with chronic functional constipation. *Turk J Pediatr*. 2006 Oct-Dec;48(4):340-3. X-4
1074. Kato K, Furuhashi K, Suzuki K, et al. Overactive bladder and glaucoma: a survey at outpatient clinics in Japan. *Int J Urol*. 2007 Jul;14(7):595-7. X-11
1075. Kaufman JM. Urodynamics in stress urinary incontinence. *J Urol*. 1979 Dec;122(6):778-82. X-1B
1076. Kauppila A, Alavaikko P and Kujansuu E. Detrusor instability score in the evaluation of stress urinary incontinence. *Acta Obstet Gynecol Scand*. 1982;61(2):137-41. X-1B
1077. Kavallaris A, Kohler C, Diebolder H, et al. Repair of prolapse with vaginal sacrocolporectomy: technique and results. *Eur J Obstet Gynecol Reprod Biol*. 2005 Oct 1;122(2):237-42. X-1E
1078. Kawabe K, Abe S, Kanda T, et al. Clinical reevaluation of the effect of oxybutynin chloride on uninhibited neurogenic and reflex neurogenic bladder. *Urol Int*. 1986;41(1):16-20. X-1F
1079. Kay G, Crook T, Rekeda L, et al. Differential effects of the antimuscarinic agents darifenacin and oxybutynin ER on memory in older subjects. *Eur Urol*. 2006 Aug;50(2):317-26. X-4
1080. Kaya C, Ilktac A, Gunes M, et al. The efficacy of trospium chloride in women with an overactive bladder. *Marmara Medical Journal*. 2008;21(2):127-132. X-7
1081. Kaya H, Sezik M, Ozbasar D, et al. Intrafascial versus extrafascial abdominal hysterectomy: effects on urinary urge incontinence. *Int Urogynecol J Pelvic Floor Dysfunct*. 2004 May-Jun;15(3):171-4. X-1A
1082. Kayigil O, Atahan O and Metin A. Experiences with clam ileocystoplasty. *Int Urol Nephrol*. 1998;30(1):45-8. X-7
1083. Kayigil O, Aytac B, Cakar KS, et al. Clam ileocystoplasty in adult nocturnal enuresis. *Int Urol Nephrol*. 2001;32(4):647-9. X-1C, X-7
1084. Kayigil O, Metin A and Atmaca AF. Obstructive urodynamic findings in idiopathic detrusor overactivity. *Int Urol Nephrol*. 2007;39(2):445-8. X-1G
1085. Kebapci N, Yenilmez A, Efe B, et al. Bladder dysfunction in type 2 diabetic patients. *Neurourol Urodyn*. 2007;26(6):814-9. X-7
1086. Keeler JR, Hurst CG and Dunn MA. Pyridostigmine used as a nerve agent pretreatment under wartime conditions. *JAMA*. 1991 Aug 7;266(5):693-5. X-11

1087. Kelleher C, Cardozo L, Kobashi K, et al. Solifenacin: as effective in mixed urinary incontinence as in urge urinary incontinence. *Int Urogynecol J Pelvic Floor Dysfunct.* 2006 Jun;17(4):382-8. X-2
1088. Kelleher CJ, Cardozo LD, Khullar V, et al. A medium-term analysis of the subjective efficacy of treatment for women with detrusor instability and low bladder compliance. *Br J Obstet Gynaecol.* 1997 Sep;104(9):988-93. X-1I
1089. Kelleher CJ, Cardozo LD, Khullar V, et al. A new questionnaire to assess the quality of life of urinary incontinent women. *Br J Obstet Gynaecol.* 1997 Dec;104(12):1374-9. X-1I
1090. Kelleher CJ, Pleil AM, Reese PR, et al. How much is enough and who says so? *BJOG.* 2004 Jun;111(6):605-12. X-1I
1091. Kelleher CJ, Tubaro A, Wang JT, et al. Impact of fesoterodine on quality of life: pooled data from two randomized trials. *BJU Int.* 2008 Jul;102(1):56-61. X-2
1092. Kelly JD, Kernohan RM and Keane PF. Symptomatic outcome following clam ileocystoplasty. *Eur Urol.* 1997;32(1):30-3. X-7
1093. Kempton CL, Kurtin PJ, Inwards DJ, et al. Malignant lymphoma of the bladder: evidence from 36 cases that low-grade lymphoma of the MALT-type is the most common primary bladder lymphoma. *Am J Surg Pathol.* 1997 Nov;21(11):1324-33. X-1I
1094. Kendall AR and Stein BS. Practical approach to stress urinary incontinence. *Geriatrics.* 1983 May;38(5):69-70, 73-4, 79. X-1B
1095. Kennelly MJ and Kang J. Botulinum-A toxin injections as a treatment for refractory detrusor hyperreflexia. *Topics in Spinal Cord Injury Rehabilitation.* 2003;8(3):46-53. X-1F
1096. Kenton K and Brubaker L. Relationship between levator ani contraction and motor unit activation in the urethral sphincter. *Am J Obstet Gynecol.* 2002 Aug;187(2):403-6. X-1G
1097. Kenton K, Fitzgerald MP and Brubaker L. What is a clinician to do-believe the patient or her urinary diary? *J Urol.* 2006 Aug;176(2):633-5; discussion 635. X-1H
1098. Kenton K, Fitzgerald MP, Shott S, et al. Role of urethral electromyography in predicting outcome of Burch retropubic urethropexy. *Am J Obstet Gynecol.* 2001 Jul;185(1):51-5. X-1B
1099. Kenton K, Lowenstein L, Simmons J, et al. Aging and overactive bladder may be associated with loss of urethral sensation in women. *Neurourol Urodyn.* 2007;26(7):981-4. X-1H
1100. Kenton K, Oldham L and Brubaker L. Open Burch urethropexy has a low rate of perioperative complications. *Am J Obstet Gynecol.* 2002 Jul;187(1):107-10. X-1B
1101. Kersey J, Martin MR and Mishra P. A further assessment of the gauze hammock sling operation in the treatment of stress incontinence. *Br J Obstet Gynaecol.* 1988 Apr;95(4):382-5. X-1B
1102. Kessler R and Constantinou CE. Internal urethrotomy in girls and its impact on the urethral intrinsic and extrinsic continence mechanisms. *J Urol.* 1986 Dec;136(6):1248-53. X-1I, X-4
1103. Kessler TM, Burkhard FC, Madersbacher H, et al. Safety of prolonged sacral neuromodulation timed lead testing. *Curr Med Res Opin.* 2008 Feb;24(2):343-7. X-1H, X-7
1104. Kessler TM, Burkhard FC, Z'Brun S, et al. Effect of thalamic deep brain stimulation on lower urinary tract function. *Eur Urol.* 2008 Mar;53(3):607-12. X-7
1105. Kessler TM, Danuser H, Schumacher M, et al. Botulinum A toxin injections into the detrusor: an effective treatment in idiopathic and neurogenic detrusor overactivity? *Neurourol Urodyn.* 2005;24(3):231-6. X-7
1106. Kessler TM, Lackner J, Kiss G, et al. Predictive value of initial urodynamic pattern on urinary continence in patients with myelomeningocele. *Neurourol Urodyn.* 2006;25(4):361-7. X-1A
1107. Kessler TM, Madersbacher H and Kiss G. Prolonged sacral neuromodulation testing using permanent leads: a more reliable patient selection method? *Eur Urol.* 2005 May;47(5):660-5. X-7
1108. Khan Z, Starer P and Bhola A. Urinary incontinence in female Parkinson disease patients. Pitfalls of diagnosis. *Urology.* 1989 Jun;33(6):486-9. X-1F, X-1H
1109. Khan Z, Starer P, Singh VK, et al. Role of detrusor instability in primary enuresis. *Urology.* 1993 Feb;41(2):189-91. X-1I, X-4
1110. Khoury AE, Dave S, Peralta-Del Valle MH, et al. Severe bladder trabeculation obviates the need for bladder outlet procedures during augmentation cystoplasty in incontinent patients with neurogenic bladder. *BJU Int.* 2008 Jan;101(2):223-6. X-1F, X-7
1111. Khullar V, Cardozo LD, Salvatore S, et al. Ultrasound: a noninvasive screening test for detrusor instability. *Br J Obstet Gynaecol.* 1996 Sep;103(9):904-8. X-1H
1112. Khullar V, Damiano R, Tooze-Hobson P, et al. Prevalence of faecal incontinence among women with urinary incontinence. *Br J Obstet Gynaecol.* 1998 Nov;105(11):1211-3. X-1H
1113. Khullar V, Hill S, Laval KU, et al. Treatment of urge-predominant mixed urinary incontinence with tolterodine extended release: a randomized, placebo-controlled trial. *Urology.* 2004 Aug;64(2):269-74; discussion 274-5. X-1B
1114. Khullar V, Rovner ES, Dmochowski R, et al. Fesoterodine dose response in subjects with overactive bladder syndrome. *Urology.* 2008 May;71(5):839-43. X-2
1115. Kibar Y, Demir E, Irkilata C, et al. Effect of biofeedback treatment on spinning top urethra in children with voiding dysfunction. *Urology.* 2007 Oct;70(4):781-4; discussion 784-5. X-1I, X-4
1116. Kilic N, Balkan E, Akgoz S, et al. Comparison of the effectiveness and side-effects of tolterodine and oxybutynin in children with detrusor instability. *Int J Urol.* 2006 Feb;13(2):105-8. X-4
1117. Kim HL, Gerber GS, Patel RV, et al. Practice patterns in the treatment of female urinary incontinence: a postal and internet survey. *Urology.* 2001 Jan;57(1):45-8. X-6
1118. Kim JC, Park EY, Seo SI, et al. Nerve growth factor and prostaglandins in the urine of female patients with overactive bladder. *J Urol.* 2006 May;175(5):1773-6; discussion 1776. X-1G, X-1H

1119. Kim JW, Kim MJ, Noh JY, et al. Extracorporeal pelvic floor magnetic stimulation in children with voiding dysfunction. *BJU Int.* 2005 Jun;95(9):1310-3. X-4, X-7
1120. Kim MD, Won JW, Lee DY, et al. Uterine artery embolization for adenomyosis without fibroids. *Clin Radiol.* 2004 Jun;59(6):520-6. X-11
1121. Kim SW, Song SH and Ku JH. Bladder training versus combination of propiverine with bladder training for female urinary frequency. A prospective, randomized, comparative study. *Gynecol Obstet Invest.* 2008;65(2):123-7. X-7
1122. Kim Y, Yoshimura N, Masuda H, et al. Intravesical instillation of human urine after oral administration of tropsium, tolterodine and oxybutynin in a rat model of detrusor overactivity. *BJU Int.* 2006 Feb;97(2):400-3. X-1G, X-4
1123. Kim YH, Kattan MW and Boone TB. Correlation of urodynamic results and urethral coaptation with success after transurethral collagen injection. *Urology.* 1997 Dec;50(6):941-8. X-1B
1124. Kim YH, Seo JT and Yoon H. The effect of overactive bladder syndrome on the sexual quality of life in Korean young and middle aged women. *Int J Impot Res.* 2005 Mar-Apr;17(2):158-63. X-1I, X-6E
1125. Kim YT, Kwon DD, Kim J, et al. Gabapentin for overactive bladder and nocturia after anticholinergic failure. *Int Braz J Urol.* 2004 Jul-Aug;30(4):275-8. X-7
1126. Kinn AC and Nilsson BY. Urethral sensitivity in incontinent women. *Eur Urol.* 2005 Jul;48(1):116-20. X-1H
1127. Kjolhede P. Long-term efficacy of Burch colposuspension: a 14-year follow-up study. *Acta Obstet Gynecol Scand.* 2005 Aug;84(8):767-72. X-1A
1128. Kjolhede P and Ryden G. Clinical and urodynamic characteristics of women with recurrent urinary incontinence after Burch colposuspension. *Acta Obstet Gynecol Scand.* 1997 May;76(5):461-7. X-1A, X-1D
1129. Kjolhede P, Wahlstrom J and Wingren G. Pelvic floor dysfunction after Burch colposuspension--a comprehensive study. Part I. *Acta Obstet Gynecol Scand.* 2005 Sep;84(9):894-901. X-1A
1130. Kjolseth D, Madsen B, Knudsen LM, et al. Biofeedback treatment of children and adults with idiopathic detrusor instability. *Scand J Urol Nephrol.* 1994 Sep;28(3):243-7. X-4, X-7
1131. Klaphajone J, Kitisomprayoonkul W and Sriplakit S. Botulinum toxin type A injections for treating neurogenic detrusor overactivity combined with low-compliance bladder in patients with spinal cord lesions. *Arch Phys Med Rehabil.* 2005 Nov;86(11):2114-8. X-1F, X-7
1132. Klarskov N and Lose G. Urethral pressure reflectometry vs urethral pressure profilometry in women: a comparative study of reproducibility and accuracy. *BJU Int.* 2007 Aug;100(2):351-6. X-1G, X-1H
1133. Klarskov P, Gerstenberg TC and Hald T. Bladder training and terodiline in females with idiopathic urge incontinence and stable detrusor function. *Scand J Urol Nephrol.* 1986;20(1):41-6. X-7
1134. Klausner AP, Sharma S, Fletcher S, et al. Does oxybutynin alter plaques, amyloid beta peptides and behavior in a mouse model of Alzheimer's disease? *J Urol.* 2008 Mar;179(3):1173-7. X-1G, X-4, X-5, X-6, X-7
1135. Klay M and Marfyak K. Use of a continence nurse specialist in an extended care facility. *Urol Nurs.* 2005 Apr;25(2):101-2, 107-8. X-1I, X-5
1136. Kleeman S, Vassallo B, Segal J, et al. The ability of history and a negative cough stress test to detect occult stress incontinence in patients undergoing surgical repair of advanced pelvic organ prolapse. *Int Urogynecol J Pelvic Floor Dysfunct.* 2006 Jan;17(1):27-9. X-1E
1137. Klein LA. Measurement of trigonal sensitivity as a test of bladder function. *J Urol.* 1987 Feb;137(2):245-8. X-1H
1138. Klevan JL and De Jong AR. Urinary tract symptoms and urinary tract infection following sexual abuse. *Am J Dis Child.* 1990 Feb;144(2):242-4. X-1I
1139. Klijn AJ, Uiterwaal CS, Vijverberg MA, et al. Home uroflowmetry biofeedback in behavioral training for dysfunctional voiding in school-age children: a randomized controlled study. *J Urol.* 2006 Jun;175(6):2263-8; discussion 2268. X-4
1140. Klingele CJ, Carley ME and Hill RF. Patient characteristics that are associated with urodynamically diagnosed detrusor instability and genuine stress incontinence. *Am J Obstet Gynecol.* 2002 May;186(5):866-8. X-1H
1141. Klingler HC, Pycha A, Schmidbauer J, et al. Use of peripheral neuromodulation of the S3 region for treatment of detrusor overactivity: a urodynamic-based study. *Urology.* 2000 Nov 1;56(5):766-71. X-7
1142. Klotz T, Bruggenjurgen B, Burkart M, et al. The economic costs of overactive bladder in Germany. *Eur Urol.* 2007 Jun;51(6):1654-62; discussion 1662-3. X-8
1143. Klovning A, Hunskaar S and Eriksen BC. Validity of a scored urological history in detecting detrusor instability in female urinary incontinence. *Acta Obstet Gynecol Scand.* 1996 Nov;75(10):941-5. X-1I
1144. Klutke C, Siegel S, Carlin B, et al. Urinary retention after tension-free vaginal tape procedure: incidence and treatment. *Urology.* 2001 Nov;58(5):697-701. X-1B
1145. Klutke JJ, Bergman J and Klutke CG. Transvaginal bladder neck suspension with Cooper's ligament fixation. Long-term urodynamic results. *J Reprod Med.* 2000 Jul;45(7):541-5. X-1B
1146. Klutke JJ and Ramos S. Urodynamic outcome after surgery for severe prolapse and potential stress incontinence. *Am J Obstet Gynecol.* 2000 Jun;182(6):1378-81. X-1B, X-1E
1147. Kobashi KC and Govier FE. Management of vaginal erosion of polypropylene mesh slings. *J Urol.* 2003 Jun;169(6):2242-3. X-1I
1148. Kobashi KC, Leach GE, Chon J, et al. Continued multicenter followup of cadaveric prolapse repair with sling. *J Urol.* 2002 Nov;168(5):2063-8. X-1E
1149. Kobashi KC, Mee SL and Leach GE. A new technique for cystocele repair and transvaginal sling: the cadaveric prolapse repair and sling (CAPS). *Urology.* 2000 Dec 4;56(6 Suppl 1):9-14. X-1E
1150. Kobelt G. Economic considerations and outcome measurement in urge incontinence. *Urology.* 1997 Dec;50(6A Suppl):100-7; discussion 108-10. X-6

1151. Kochakarn W, Pummangura N, Kijvikai K, et al. Reliability of a Thai version of King's Health Questionnaire in Thai females with overactive bladder symptoms. *J Med Assoc Thai*. 2005 Nov;88(11):1526-34. X-1H
1152. Kockelbergh RC, Tan JB, Bates CP, et al. Clam enterocystoplasty in general urological practice. *Br J Urol*. 1991 Jul;68(1):38-41. X-7
1153. Koduri S, Goldberg RP and Sand PK. Transvaginal therapy of genuine stress incontinence. *Urology*. 2000 Dec 4;56(6 Suppl 1):23-7. X-1B
1154. Koefoot RB, Jr. and Webster GD. Urodynamic evaluation in women with frequency, urgency symptoms. *Urology*. 1983 Jun;21(6):648-51. X-1H
1155. Koelbl H and Bernaschek G. A new method for sonographic urethrocytography and simultaneous pressure-flow measurements. *Obstet Gynecol*. 1989 Sep;74(3 Pt 1):417-22. X-1G, X-1H
1156. Koelbl H, Bernaschek G and Deutinger J. Assessment of female urinary incontinence by introital sonography. *J Clin Ultrasound*. 1990 May;18(4):370-4. X-1B, X-7
1157. Koh CJ, DeFilippo RE, Borer JG, et al. Bladder and external urethral sphincter function after prenatal closure of myelomeningocele. *J Urol*. 2006 Nov;176(5):2232-6. X-4, X-7
1158. Kok AL, Burger CW, van de Weijer PH, et al. Micturition complaints in postmenopausal women treated with continuously combined hormone replacement therapy: a prospective study. *Maturitas*. 1999 Jan 4;31(2):143-9. X-7
1159. Komine S, Yoshida H, Fujiyama C, et al. Voiding dysfunction in patients with human T-lymphotropic-virus-type-1-associated myelopathy. *Urol Int*. 1991;47 Suppl 1:67-8. X-1F
1160. Komine S, Yoshida K, Yamashita H, et al. Voiding dysfunction in patients with human T-lymphotropic virus type-1-associated myelopathy (HAM). *Paraplegia*. 1989 Jun;27(3):217-21. X-1F
1161. Kondo A, Isobe Y, Kimura K, et al. Efficacy, safety and hospital costs of tension-free vaginal tape and pubovaginal sling in the surgical treatment of stress incontinence. *J Obstet Gynaecol Res*. 2006 Dec;32(6):539-44. X-1B
1162. Kondo A, Kobayashi M, Otani T, et al. Children with unstable bladder: clinical and urodynamic observation. *J Urol*. 1983 Jan;129(1):88-91. X-4
1163. Kondo Y, Homma Y, Takahashi S, et al. Transvaginal ultrasound of urethral sphincter at the mid urethra in continent and incontinent women. *J Urol*. 2001 Jan;165(1):149-52. X-1H
1164. Kong KH, Chan KF, Lim AC, et al. Detrusor hyperreflexia in strokes. *Ann Acad Med Singapore*. 1994 May;23(3):319-21. X-1F, X-7
1165. Konturi MJ, Hellstrom PA, Tammela TL, et al. Colocystoplasty for the treatment of severe interstitial cystitis. *Urol Int*. 1991;46(1):50-4. X-1D
1166. Koonings P, Bergman A and Ballard CA. Combined detrusor instability and stress urinary incontinence: where is the primary pathology? *Gynecol Obstet Invest*. 1988;26(3):250-6. X-1B
1167. Kopp JB, Miller KD, Mican JA, et al. Crystalluria and urinary tract abnormalities associated with indinavir. *Ann Intern Med*. 1997 Jul 15;127(2):119-25. X-1A
1168. Korda A, Cooper M and Hunter P. Coital urinary incontinence in an Australian population. *Asia Oceania J Obstet Gynaecol*. 1989 Dec;15(4):313-5. X-1H
1169. Korda A, Ferry J and Hunter P. Colposuspension for the treatment of female urinary incontinence. *Aust N Z J Obstet Gynaecol*. 1989 May;29(2):146-9. X-1B
1170. Korda A, Peat B and Hunter P. Experience with silastic slings for female urinary incontinence. *Aust N Z J Obstet Gynaecol*. 1989 May;29(2):150-4. X-1B
1171. Kowalczyk JJ. Office evaluation of the patient with an overactive urinary bladder. *J Am Osteopath Assoc*. 2000 Mar;100(3 Suppl):S1-4. X-2
1172. Krauwinkel WJ, Smulders RA, Mulder H, et al. Effect of age on the pharmacokinetics of solifenacin in men and women. *Int J Clin Pharmacol Ther*. 2005 May;43(5):227-38. X-II, X-7
1173. Krogh RA, Neumann GA, Lauszus FF, et al. Hysterectomy is associated with stress incontinence in women who previously had a transcervical endometrial resection. *Gynecol Obstet Invest*. 2007;63(3):121-5. X-1A, X-1B
1174. Krue S, Jensen H, Agger AO, et al. The influence of infant birth weight on post partum stress incontinence in obese women. *Arch Gynecol Obstet*. 1997;259(3):143-5. X-1B
1175. Ku JH, Oh JG, Shin JW, et al. Age is not a limiting factor for midurethral sling procedures in the elderly with urinary incontinence. *Gynecol Obstet Invest*. 2006;61(4):194-9. X-1B
1176. Kuhn A, Hildebrand R and Birkhauser M. Do transsexuals have micturition disorders? *Eur J Obstet Gynecol Reprod Biol*. 2007 Apr;131(2):226-30. X-1A, X-7
1177. Kuhn A, Vits K, Kuhn P, et al. Do women with urinary incontinence really know where all the toilets are? The toilet paper. *Eur J Obstet Gynecol Reprod Biol*. 2006 Nov;129(1):65-8. X-II
1178. Kuipers M, Smulders R, Krauwinkel W, et al. Open-label study of the safety and pharmacokinetics of solifenacin in subjects with hepatic impairment. *J Pharmacol Sci*. 2006 Dec;102(4):405-12. X-7
1179. Kulseng-Hanssen S. Prevalence and pattern of unstable urethral pressure in one hundred seventy-four gynecologic patients referred for urodynamic investigation. *Am J Obstet Gynecol*. 1983 Aug 15;146(8):895-900. X-1H
1180. Kulseng-Hanssen S. The development of a national database of the results of surgery for urinary incontinence in women. *BJOG*. 2003 Nov;110(11):975-82. X-1B, X-6
1181. Kulseng-Hanssen S and Borstad E. The development of a questionnaire to measure the severity of symptoms and the quality of life before and after surgery for stress incontinence. *BJOG*. 2003 Nov;110(11):983-8. X-1B, X-6
1182. Kulseng-Hanssen S, Husby H and Schiotz HA. The tension free vaginal tape operation for women with mixed incontinence: Do preoperative variables predict the outcome? *Neurourol Urodyn*. 2007;26(1):115-21; discussion 122. X-1B
1183. Kulseng-Hanssen S, Husby H and Schiotz HA. Follow-up of TVT operations in 1,113 women with mixed urinary incontinence at 7 and 38 months. *Int Urogynecol J Pelvic Floor Dysfunct*. 2008 Mar;19(3):391-6. X-1B

1184. Kulseng-Hanssen S, Kristoffersen M and Larsen E. Evaluation of the subjective and objective effect of maximal electrical stimulation in patients complaining of urge incontinence. *Acta Obstet Gynecol Scand Suppl.* 1998;168:12-5. X-7
1185. Kumar SP and Abrams PH. Detrusor myectomy: long-term results with a minimum follow-up of 2 years. *BJU Int.* 2005 Aug;96(3):341-4. X-7
1186. Kunkle JC, Payne CK and Whitmore KE. Urinary incontinence: combined biofeedback and electrical stimulation treatment. *Journal of Urological Nursing.* 1993;12(3):537-543. X-7
1187. Kuo HC. Videourodynamic results after pubovaginal sling procedure for stress urinary incontinence. *Urology.* 1999 Nov;54(5):802-6; discussion 806-7. X-1B
1188. Kuo HC. Videourodynamic study for diagnosis of bladder outlet obstruction in women. *J Formos Med Assoc.* 2000 May;99(5):386-92. X-1I
1189. Kuo HC. Anatomical and functional results of pubovaginal sling procedure using polypropylene mesh for the treatment of stress urinary incontinence. *J Urol.* 2001 Jul;166(1):152-7. X-1B, X-7
1190. Kuo HC. Comparison of video urodynamic results after the pubovaginal sling procedure using rectus fascia and polypropylene mesh for stress urinary incontinence. *J Urol.* 2001 Jan;165(1):163-8. X-1B
1191. Kuo HC. Sonographic evaluation of anatomic results after the pubovaginal sling procedure for stress urinary incontinence. *J Ultrasound Med.* 2001 Jul;20(7):739-47. X-1B
1192. Kuo HC. Efficacy of desmopressin in treatment of refractory nocturia in patients older than 65 years. *Urology.* 2002 Apr;59(4):485-9. X-1C, X-7
1193. Kuo HC. Effectiveness of intravesical resiniferatoxin for anticholinergic treatment refractory detrusor overactivity due to nonspinal cord lesions. *J Urol.* 2003 Sep;170(3):835-9. X-7
1194. Kuo HC. Videourodynamic results in stress urinary incontinence patients after pelvic floor muscle training. *J Formos Med Assoc.* 2003 Jan;102(1):23-9. X-1I
1195. Kuo HC. Urodynamic evidence of effectiveness of botulinum A toxin injection in treatment of detrusor overactivity refractory to anticholinergic agents. *Urology.* 2004 May;63(5):868-72. X-7
1196. Kuo HC. Clinical effects of suburothelial injection of botulinum A toxin on patients with nonneurogenic detrusor overactivity refractory to anticholinergics. *Urology.* 2005 Jul;66(1):94-8. X-7
1197. Kuo HC. Long-term surgical results of pubovaginal sling procedure using polypropylene mesh in the treatment of stress urinary incontinence. *Urol Int.* 2005;74(2):147-52. X-1B
1198. Kuo HC. Preliminary results of suburothelial injection of botulinum A toxin in the treatment of chronic interstitial cystitis. *Urol Int.* 2005;75(2):170-4. X-1D
1199. Kuo HC. Videourodynamic characteristics and lower urinary tract symptoms of female bladder outlet obstruction. *Urology.* 2005 Nov;66(5):1005-9. X-1H, X-1I
1200. Kuo HC. Therapeutic effects of suburothelial injection of botulinum A toxin for neurogenic detrusor overactivity due to chronic cerebrovascular accident and spinal cord lesions. *Urology.* 2006 Feb;67(2):232-6. X-1F
1201. Kuo HC. Will suburothelial injection of small dose of botulinum A toxin have similar therapeutic effects and less adverse events for refractory detrusor overactivity? *Urology.* 2006 Nov;68(5):993-7; discussion 997-8. X-7
1202. Kuo HC. Comparison of effectiveness of detrusor, suburothelial and bladder base injections of botulinum toxin A for idiopathic detrusor overactivity. *J Urol.* 2007 Oct;178(4 Pt 1):1359-63. X-7
1203. Kuo HC. Effect of detrusor function on the therapeutic outcome of a suburethral sling procedure using a polypropylene sling for stress urinary incontinence in women. *Scand J Urol Nephrol.* 2007;41(2):138-43. X-1B
1204. Kuo HC. Therapeutic satisfaction and dissatisfaction in patients with spinal cord lesions and detrusor sphincter dyssynergia who received detrusor botulinum toxin A injection. *Urology.* 2008 Nov;72(5):1056-60. X-1F
1205. Kuo HC and Liu HT. Investigation of dysfunctional voiding in children with urgency frequency syndrome and urinary incontinence. *Urol Int.* 2006;76(1):72-6. X-1H, X-4, X-7
1206. Kuo HC, Liu HT and Yang WC. Therapeutic effect of multiple resiniferatoxin intravesical instillations in patients with refractory detrusor overactivity: a randomized, double-blind, placebo controlled study. *J Urol.* 2006 Aug;176(2):641-5. X-4, X-7
1207. Kupelian V, Rosen RC, Link CL, et al. Association of urological symptoms and chronic illness in men and women: contributions of symptom severity and duration--results from the BACH Survey. *J Urol.* 2009 Feb;181(2):694-700. X-8
1208. Kuriyama N, Tokuda T, Kondo M, et al. Evaluation of autonomic malfunction in idiopathic normal pressure hydrocephalus. *Clin Auton Res.* 2008 Aug;18(4):213-20. X-1I
1209. Kurstjens GA, Borau A, Rodriguez A, et al. Intraoperative recording of electroneurographic signals from cuff electrodes on extradural sacral roots in spinal cord injured patients. *J Urol.* 2005 Oct;174(4 Pt 1):1482-7. X-1F, X-7
1210. Kurstjens M, Rijkhoff NJ, Borau A, et al. Intraoperative recording of sacral root nerve signals in humans. *Artif Organs.* 2005 Mar;29(3):242-5. X-1F, X-7
1211. Kuruba R, Almahmeed T, Martinez F, et al. Bariatric surgery improves urinary incontinence in morbidly obese individuals. *Surg Obes Relat Dis.* 2007 Nov-Dec;3(6):586-90; discussion 590-1. X-7
1212. Kuschel S, Werner M, Schmid DM, et al. Botulinum toxin-A for idiopathic overactivity of the vesical detrusor: a 2-year follow-up. *Int Urogynecol J Pelvic Floor Dysfunct.* 2008 Jul;19(7):905-9. X-7
1213. Kushner L, Chiu PY, Brettschneider N, et al. Urinary substance P concentration correlates with urinary frequency and urgency in interstitial cystitis patients treated with intravesical dimethyl sulfoxide and not intravesical anesthetic cocktail. *Urology.* 2001 Jun;57(6 Suppl 1):129. X-1D
1214. Kutzenberger J. Surgical therapy of neurogenic detrusor overactivity (hyperreflexia) in paraplegic patients by sacral deafferentation and implant driven micturition by sacral anterior root stimulation: methods, indications, results, complications, and future prospects. *Acta Neurochir Suppl.* 2007;97(Pt 1):333-9. X-1F

1215. Kuuva N and Nilsson CG. Tension-free vaginal tape procedure: an effective minimally invasive operation for the treatment of recurrent stress urinary incontinence? *Gynecol Obstet Invest.* 2003;56(2):93-8. X-1B
1216. Kwak KW and Park KH. Clinical inconsistency of lower urinary tract symptoms between questionnaire and bladder diary in children with nocturnal enuresis. *J Urol.* 2008 Sep;180(3):1085-9; discussion 1089-90. X-1A, X-4
1217. Lackner TE, Wyman JF, McCarthy TC, et al. Randomized, placebo-controlled trial of the cognitive effect, safety, and tolerability of oral extended-release oxybutynin in cognitively impaired nursing home residents with urge urinary incontinence. *J Am Geriatr Soc.* 2008 May;56(5):862-70. X-5
1218. Lagro-Janssen AL, Debruyne FM, Smits AJ, et al. The effects of treatment of urinary incontinence in general practice. *Fam Pract.* 1992 Sep;9(3):284-9. X-7
1219. Lagro-Janssen AL, Debruyne FM and van Weel C. Value of the patient's case history in diagnosing urinary incontinence in general practice. *Br J Urol.* 1991 Jun;67(6):569-72. X-1H
1220. Lagro-Janssen AL, Debruyne FM and Van Weel C. Psychological aspects of female urinary incontinence in general practice. *Br J Urol.* 1992 Nov;70(5):499-502. X-1I
1221. Lagro-Janssen T, Smits A and Van Weel C. Urinary incontinence in women and the effects on their lives. *Scand J Prim Health Care.* 1992 Sep;10(3):211-6. X-1I
1222. Lagro-Janssen T and van Weel C. Long-term effect of treatment of female incontinence in general practice. *Br J Gen Pract.* 1998 Nov;48(436):1735-8. X-7
1223. Lajer H, Thranov IR, Bagi P, et al. Evaluation of urologic morbidity after radiotherapy for cervical carcinoma by urodynamic examinations and patient voiding schemes: a prospective study. *Int J Radiat Oncol Biol Phys.* 2002 Dec 1;54(5):1362-8. X-1A, X-1I, X-7
1224. Lalos O, Berglund AL and Lalos A. Impact of urinary and climacteric symptoms on social and sexual life after surgical treatment of stress urinary incontinence in women: a long-term outcome. *J Adv Nurs.* 2001 Feb;33(3):316-27. X-1B, X-7
1225. Lalos O and Bjerle P. Bladder wall mechanics and micturition before and after subtotal and total hysterectomy. *Eur J Obstet Gynecol Reprod Biol.* 1986 Mar;21(3):143-50. X-1A, X-7
1226. Lammers RL, Gibson S, Kovacs D, et al. Comparison of test characteristics of urine dipstick and urinalysis at various test cutoff points. *Ann Emerg Med.* 2001 Nov;38(5):505-12. X-1G, X-1I
1227. Lang EW, Chesnut RM and Hennerici M. Urinary retention and space-occupying lesions of the frontal cortex. *Eur Neurol.* 1996;36(1):43-7. X-1F
1228. Langer R, Golan A, Arad D, et al. Effects of induced menopause on Burch colposuspension for urinary stress incontinence. *J Reprod Med.* 1992 Dec;37(12):956-8. X-1I
1229. Langer R, Golan A, Neuman M, et al. The effect of large uterine fibroids on urinary bladder function and symptoms. *Am J Obstet Gynecol.* 1990 Oct;163(4 Pt 1):1139-41. X-1I, X-7
1230. Langer R, Golan A, Ron-El R, et al. Colposuspension for urinary stress incontinence in premenopausal and postmenopausal women. *Surg Gynecol Obstet.* 1990 Jul;171(1):13-6. X-1B
1231. Langer R, Lipshitz Y, Halperin R, et al. Long-Term (10-15 years) follow-up after Burch colposuspension for urinary stress incontinence. *Int Urogynecol J Pelvic Floor Dysfunct.* 2001;12(5):323-6; discussion 326-7. X-1A
1232. Langer R, Ron-El R, Bukovsky I, et al. Colposuspension in patients with combined stress incontinence and detrusor instability. *Eur Urol.* 1988;14(6):437-9. X-1B
1233. Langer R, Ron-el R, Newman M, et al. Detrusor instability following colposuspension for urinary stress incontinence. *Br J Obstet Gynaecol.* 1988 Jun;95(6):607-10. X-1B
1234. Lapitan MC and Chye PL. The epidemiology of overactive bladder among females in Asia: a questionnaire survey. *Int Urogynecol J Pelvic Floor Dysfunct.* 2001;12(4):226-31. X-6
1235. Larsson G, Hallen B and Nilvebrant L. Tolterodine in the treatment of overactive bladder: analysis of the pooled phase II efficacy and safety data. *Urology.* 1999 May;53(5):990-8. X-1F
1236. Latini JM, Alipour M and Kreder KJ, Jr. Efficacy of sacral neuromodulation for symptomatic treatment of refractory urinary urge incontinence. *Urology.* 2006 Mar;67(3):550-3; discussion 553-4. X-7
1237. Lau DT, Morlock RJ and Hill CD. Psychometric evaluation of the medical outcomes study-sleep scale in persons with overactive bladder. *Clin Ther.* 2006 Dec;28(12):2119-32. X-1I
1238. Laurikainen E and Kiilholma P. The tension-free vaginal tape procedure for female urinary incontinence without preoperative urodynamic evaluation. *J Am Coll Surg.* 2003 Apr;196(4):579-83. X-1B
1239. Laurikainen E, Rosti J, Pitkanen Y, et al. The Rosti sling: a new, minimally invasive, tension-free technique for the surgical treatment of female urinary incontinence-the first 217 patients. *J Urol.* 2004 Apr;171(4):1576-80; discussion 1580. X-1B
1240. Laurikainen E, Valpas A, Kivela A, et al. Retropubic compared with transobturator tape placement in treatment of urinary incontinence: a randomized controlled trial. *Obstet Gynecol.* 2007 Jan;109(1):4-11. X-1B
1241. Laurikkala J and Juhola M. A genetic-based machine learning system to discover the diagnostic rules for female urinary incontinence. *Comput Methods Programs Biomed.* 1998 Mar;55(3):217-28. X-1B, X-1H, X-6, X-7
1242. Lawrence M, Guay DR, Benson SR, et al. Immediate-release oxybutynin versus tolterodine in detrusor overactivity: a population analysis. *Pharmacotherapy.* 2000 Apr;20(4):470-5. X-1I
1243. Lawton V and Smith AR. Laparoscopic colposuspension. *Semin Laparosc Surg.* 1999 Jun;6(2):90-9. X-1B
1244. Lazzeri M, Beneforti P and Turini D. Urodynamic effects of intravesical resiniferatoxin in humans: preliminary results in stable and unstable detrusor. *J Urol.* 1997 Dec;158(6):2093-6. X-7
1245. Lazzeri M, Calo G, Spinelli M, et al. Urodynamic effects of intravesical nociceptin/orphanin FQ in neurogenic detrusor overactivity: a randomized, placebo-controlled, double-blind study. *Urology.* 2003 May;61(5):946-50. X-1F

1246. Lazzeri M, Calo G, Spinelli M, et al. Daily intravesical instillation of 1 mg nociceptin/orphanin FQ for the control of neurogenic detrusor overactivity: a multicenter, placebo controlled, randomized exploratory study. *J Urol*. 2006 Nov;176(5):2098-102. X-1F, X-7
1247. Learman LA, Summitt RL, Jr., Varner RE, et al. Hysterectomy versus expanded medical treatment for abnormal uterine bleeding: clinical outcomes in the medicine or surgery trial. *Obstet Gynecol*. 2004 May;103(5 Pt 1):824-33. X-11
1248. Leboeuf L, Miles RA, Kim SS, et al. Grade 4 cystocele repair using four-defect repair and porcine xenograft acellular matrix (Pelvicol): outcome measures using SEAPI. *Urology*. 2004 Aug;64(2):282-6. X-11, X-7
1249. Lee HC, Liu CS, Chiao C, et al. Hyperbaric oxygen therapy in hemorrhagic radiation cystitis: a report of 20 cases. *Undersea Hyperb Med*. 1994 Sep;21(3):321-7. X-1D, X-7
1250. Lee JC, Yokoyama T, Hwang HJ, et al. Clinical application of Clostridium botulinum type A neurotoxin purified by a simple procedure for patients with urinary incontinence caused by refractory detrusor overactivity. *FEMS Immunol Med Microbiol*. 2007 Oct;51(1):201-11. X-7
1251. Lee KA and DeJoseph JF. Sleep disturbances, vitality, and fatigue among a select group of employed childbearing women. *Birth*. 1992 Dec;19(4):208-13. X-11
1252. Lee KS, Chan CJ, Merriman A, et al. Clinical profile of elderly urinary incontinence in Singapore: a community-based study. *Ann Acad Med Singapore*. 1991 Nov;20(6):736-9. X-7
1253. Lee KS, Han DH, Choi YS, et al. A prospective trial comparing tension-free vaginal tape and transobturator vaginal tape inside-out for the surgical treatment of female stress urinary incontinence: 1-year followup. *J Urol*. 2007 Jan;177(1):214-8. X-1B
1254. Lee PE, Kung RC and Drutz HP. Periurethral autologous fat injection as treatment for female stress urinary incontinence: a randomized double-blind controlled trial. *J Urol*. 2001 Jan;165(1):153-8. X-1B
1255. Lee SS, Liu YC, Wann SR, et al. Once daily isepamicin treatment in complicated urinary tract infections. *J Microbiol Immunol Infect*. 1999 Jun;32(2):105-10. X-11
1256. Lee WC, Wu HP, Tai TY, et al. Investigation of urodynamic characteristics and bladder sensory function in the early stages of diabetic bladder dysfunction in women with type 2 diabetes. *J Urol*. 2009 Jan;181(1):198-203. X-1H, X-11
1257. Lehtoranta K, Tainio H, Lukkari-Lax E, et al. Pharmacokinetics, efficacy, and safety of intravesical formulation of oxybutynin in patients with detrusor overactivity. *Scand J Urol Nephrol*. 2002 Feb;36(1):18-24. X-7
1258. Leibovici D, Cooper A, Lindner A, et al. Ureteral stents: morbidity and impact on quality of life. *Isr Med Assoc J*. 2005 Aug;7(8):491-4. X-11
1259. Leiby BE, Landis JR, Propert KJ, et al. Discovery of morphological subgroups that correlate with severity of symptoms in interstitial cystitis: a proposed biopsy classification system. *J Urol*. 2007 Jan;177(1):142-8. X-1A, X-1D
1260. Lekka E and Lee LK. Successful treatment with intradetrusor Botulinum-A toxin for urethral urinary leakage (catheter bypassing) in patients with end-staged multiple sclerosis and indwelling suprapubic catheters. *Eur Urol*. 2006 Oct;50(4):806-9; discussion 809-10. X-1F
1261. Leksukulchai O and Dietz HP. Normal values for detrusor wall thickness in nulligravid women. *Australian & New Zealand Continence Journal*. 2007 Autumn;13(1):4-6. X-1G
1262. Leksukulchai O and Dietz HP. Detrusor wall thickness as a test for detrusor overactivity in women. *Ultrasound Obstet Gynecol*. 2008 Sep;32(4):535-9. X-1H
1263. Lemack GE. Reversing the tide: emerging options for treating overactive bladder symptoms in women. *Tex Med*. 1999 Nov;95(11):78-81. X-2
1264. Lemack GE. Management and treatment of overactive bladder in the older female patient. *Clinical Geriatrics*. 2002;10(11):32. X-2
1265. Lemack GE, Frohman E and Ramnarayan P. Women with voiding dysfunction secondary to bladder outlet dyssynergia in the setting of multiple sclerosis do not demonstrate significantly elevated intravesical pressures. *Urology*. 2007 May;69(5):893-7. X-1F
1266. Lemack GE, Frohman EM, Zimmern PE, et al. Urodynamic distinctions between idiopathic detrusor overactivity and detrusor overactivity secondary to multiple sclerosis. *Urology*. 2006 May;67(5):960-4. X-1F, X-1G
1267. Lemack GE, Hawker K and Frohman E. Incidence of upper tract abnormalities in patients with neurovesical dysfunction secondary to multiple sclerosis: analysis of risk factors at initial urologic evaluation. *Urology*. 2005 May;65(5):854-7. X-1H
1268. Lemack GE, Xu Y, Brubaker L, et al. Clinical and demographic factors associated with valsalva leak point pressure among women undergoing burch bladder neck suspension or autologous rectus fascial sling procedures. *Neurourol Urodyn*. 2007;26(3):392-6. X-1B
1269. Lemack GE and Zimmern PE. Predictability of urodynamic findings based on the Urogenital Distress Inventory-6 questionnaire. *Urology*. 1999 Sep;54(3):461-6. X-1A
1270. Lemack GE and Zimmern PE. Identifying patients who require urodynamic testing before surgery for stress incontinence based on questionnaire information and surgical history. *Urology*. 2000 Apr;55(4):506-11. X-1B
1271. Leng WW, Davies BJ, Tarin T, et al. Delayed treatment of bladder outlet obstruction after sling surgery: association with irreversible bladder dysfunction. *J Urol*. 2004 Oct;172(4 Pt 1):1379-81. X-1A, X-7
1272. Lentz GM, Bavendam T, Stenchever MA, et al. Hormonal manipulation in women with chronic, cyclic irritable bladder symptoms and pelvic pain. *Am J Obstet Gynecol*. 2002 Jun;186(6):1268-71; discussion 1271-3. X-1D, X-6, X-7
1273. Leonardo CR, Filgueiras MF, Vasconcelos MM, et al. Risk factors for renal scarring in children and adolescents with lower urinary tract dysfunction. *Pediatr Nephrol*. 2007 Nov;22(11):1891-6. X-11, X-4
1274. Leppilahti M, Sairanen J, Tammela TL, et al. Prevalence of clinically confirmed interstitial cystitis in women: a population based study in Finland. *J Urol*. 2005 Aug;174(2):581-3. X-1D

1275. Leroi AM, Michot F, Grise P, et al. Effect of sacral nerve stimulation in patients with fecal and urinary incontinence. *Dis Colon Rectum*. 2001 Jun;44(6):779-89. X-1I, X-7
1276. Letourneau R, Sant GR, el-Mansoury M, et al. Activation of bladder mast cells in interstitial cystitis. *Int J Tissue React*. 1992;14(6):307-12. X-1D
1277. Lettgen B, von Gontard A, Olbing H, et al. Urge incontinence and voiding postponement in children: somatic and psychosocial factors. *Acta Paediatr*. 2002;91(9):978-84; discussion 895-6. X-1G, X-4
1278. Leung VY, Chu WC, Yeung CK, et al. Ureteric jet Doppler waveform and bladder wall thickness in children with nocturnal enuresis. *Pediatr Res*. 2006 Nov;60(5):582-6. X-1G, X-1H, X-4
1279. LeVeque FG, Montgomery M, Potter D, et al. A multicenter, randomized, double-blind, placebo-controlled, dose-titration study of oral pilocarpine for treatment of radiation-induced xerostomia in head and neck cancer patients. *J Clin Oncol*. 1993 Jun;11(6):1124-31. X-1I
1280. Levin I, Groutz A, Gold R, et al. Surgical complications and medium-term outcome results of tension-free vaginal tape: a prospective study of 313 consecutive patients. *Neurourol Urodyn*. 2004;23(1):7-9. X-1B
1281. Lev-Toaff AS, Karasick S and Toaff ME. Hysterosalpingography before and after myomectomy: clinical value and imaging findings. *AJR Am J Roentgenol*. 1993 Apr;160(4):803-7. X-1I, X-7
1282. Lewicky-Gaupp C, Cao DC and Culbertson S. Urinary and anal incontinence in African American teenaged gravidas during pregnancy and the puerperium. *J Pediatr Adolesc Gynecol*. 2008 Feb;21(1):21-6. X-4
1283. Lewis JB, Ng AV, O'Connor RC, et al. Are there differences between women with urge predominant and stress predominant mixed urinary incontinence? *Neurourol Urodyn*. 2007;26(2):204-7. X-1B
1284. Liang CC, Chang YL, Chang SD, et al. Pessary test to predict postoperative urinary incontinence in women undergoing hysterectomy for prolapse. *Obstet Gynecol*. 2004 Oct;104(4):795-800. X-1B, X-1E
1285. Liang CC, Tseng LH, Horng SG, et al. Correlation of pelvic organ prolapse quantification system scores with obstetric parameters and lower urinary tract symptoms in primiparae postpartum. *Int Urogynecol J Pelvic Floor Dysfunct*. 2007 May;18(5):537-41. X-1A
1286. Liao KK, Chen JT, Lai KL, et al. Effect of sacral-root stimulation on the motor cortex in patients with idiopathic overactive bladder syndrome. *Neurophysiol Clin*. 2008 Feb;38(1):39-43. X-7
1287. Liao YM, Dougherty MC, Biemer PP, et al. Factors related to lower urinary tract symptoms among a sample of employed women in Taipei. *Neurourol Urodyn*. 2008;27(1):52-9. X-6, X-6E
1288. Liapis A, Bakas P and Creatsas G. The efficacy of bladder distention therapy in the treatment of frequency and urgency. *Eur J Obstet Gynecol Reprod Biol*. 2001 Mar;95(1):97-9. X-7
1289. Liapis A, Bakas P and Creatsas G. Burch colposuspension and tension-free vaginal tape in the management of stress urinary incontinence in women. *Eur Urol*. 2002 Apr;41(4):469-73. X-1B
1290. Liapis A, Bakas P and Creatsas G. Assessment of TVT efficacy in the management of patients with genuine stress incontinence with the use of epidural vs intravenous anesthesia. *Int Urogynecol J Pelvic Floor Dysfunct*. 2007 Oct;18(10):1197-200. X-1B
1291. Liapis A, Bakas P and Creatsas G. Long-term efficacy of tension-free vaginal tape in the management of stress urinary incontinence in women: efficacy at 5- and 7-year follow-up. *Int Urogynecol J Pelvic Floor Dysfunct*. 2008 Nov;19(11):1509-12. X-1A
1292. Liapis A, Bakas P, Pafiti A, et al. Changes in the quantity of collagen type I in women with genuine stress incontinence. *Urol Res*. 2000 Oct;28(5):323-6. X-1B, X-1G
1293. Liatsikos EN, Gershbaum D, Kapoor R, et al. Comparison of symptoms related to positioning of double-pigtail stent in upper pole versus renal pelvis. *J Endourol*. 2001 Apr;15(3):299-302. X-1A, X-7
1294. Liberman JN, Hunt TL, Stewart WF, et al. Health-related quality of life among adults with symptoms of overactive bladder: results from a U.S. community-based survey. *Urology*. 2001 Jun;57(6):1044-50. X-1I
1295. Lilly JD and Parsons CL. Bladder surface glycosaminoglycans is a human epithelial permeability barrier. *Surg Gynecol Obstet*. 1990 Dec;171(6):493-6. X-1G
1296. Lim J, Cornish A and Carey MP. Clinical and quality-of-life outcomes in women treated by the TVT-O procedure. *BJOG*. 2006 Nov;113(11):1315-20. X-1B
1297. Lim JR, Bak CW and Lee JB. Comparison of anxiety between patients with mixed incontinence and those with stress urinary incontinence. *Scand J Urol Nephrol*. 2007;41(5):403-6. X-1B
1298. Lin AD, Lin AT, Chen KK, et al. Nocturnal enuresis in older adults. *J Chin Med Assoc*. 2004 Mar;67(3):136-40. X-1I
1299. Lin HH, Sheu BC, Lo MC, et al. Abnormal urodynamic findings after radical hysterectomy or pelvic irradiation for cervical cancer. *Int J Gynaecol Obstet*. 1998 Nov;63(2):169-74. X-1I
1300. Lin HH, Torng PL, Sheu BC, et al. Urodynamically age-specific prevalence of urinary incontinence in women with urinary symptoms. *Neurourol Urodyn*. 2003;22(1):29-32. X-1G, X-1H
1301. Lin HH, Yu HJ, Sheu BC, et al. Importance of urodynamic study before radical hysterectomy for cervical cancer. *Gynecol Oncol*. 2001 May;81(2):270-2. X-1H, X-7
1302. Lin LY, Yeh NH, Lin CY, et al. Comparisons of urodynamic characteristics between female patients with overactive bladder and overactive bladder plus stress urinary incontinence. *Urology*. 2004 Nov;64(5):945-9. X-1H
1303. Lin TL, Ng SC, Chen YC, et al. What affects the occurrence of nocturia more: menopause or age? *Maturitas*. 2005 Feb 14;50(2):71-7. X-1C
1304. Linder A, Leach GE and Raz S. Augmentation cystoplasty in the treatment of neurogenic bladder dysfunction. *J Urol*. 1983 Mar;129(3):491-3. X-1F
1305. Lindholm P and Lose G. Terbutaline (Bricanyl) in the treatment of female urge incontinence. *Urol Int*. 1986;41(2):158-60. X-7

1306. Lindseth PD and Lindseth GN. Assessing for preflight predictors of airsickness. *Aviat Space Environ Med.* 1992 Oct;63(10):908-13. X-1A, X-4
1307. Linn JF, Hohenfellner M, Roth S, et al. Treatment of interstitial cystitis: comparison of subtrigonal and supratrigonal cystectomy combined with orthotopic bladder substitution. *J Urol.* 1998 Mar;159(3):774-8. X-1D
1308. Lipton RB, Kolodner K and Wesnes K. Assessment of cognitive function of the elderly population: effects of darifenacin. *J Urol.* 2005 Feb;173(2):493-8. X-4
1309. Litwiller SE, Nelson RS, Fone PD, et al. Vaginal wall sling: long-term outcome analysis of factors contributing to patients satisfaction and surgical success. *J Urol.* 1997 Apr;157(4):1279-82. X-1B
1310. Liu CY. Laparoscopic retropubic colposuspension (Burch procedure). A review of 58 cases. *J Reprod Med.* 1993 Jul;38(7):526-30. X-1B
1311. Liu HT, Chancellor MB and Kuo HC. Urinary nerve growth factor level could be a biomarker in the differential diagnosis of mixed urinary incontinence in women. *BJU Int.* 2008 Nov;102(10):1440-4. X-1G
1312. Liu HT and Kuo HC. Increased expression of transient receptor potential vanilloid subfamily 1 in the bladder predicts the response to intravesical instillations of resiniferatoxin in patients with refractory idiopathic detrusor overactivity. *BJU Int.* 2007 Nov;100(5):1086-90. X-1G, X-7
1313. Liu HT and Kuo HC. Urinary nerve growth factor level could be a potential biomarker for diagnosis of overactive bladder. *J Urol.* 2008 Jun;179(6):2270-4. X-1G
1314. Liu HT and Kuo HC. Urinary nerve growth factor levels are elevated in patients with overactive bladder and do not significantly increase with bladder distention. *Neurourol Urodyn.* 2009;28(1):78-81. X-1G, X-7
1315. Liu L, Mansfield KJ, Kristiana I, et al. The molecular basis of urgency: regional difference of vanilloid receptor expression in the human urinary bladder. *Neurourol Urodyn.* 2007;26(3):433-8; discussion 439; discussion 451-3. X-1G
1316. Lloyd SN, Lloyd SM, Rogers K, et al. Is there still a place for prolonged bladder distension? *Br J Urol.* 1992 Oct;70(4):382-6. X-1I
1317. Lo SK, Naidu J and Cao Y. Additive effect of interferential therapy over pelvic floor exercise alone in the treatment of female urinary stress and urge incontinence: a randomized controlled trial. *Hong Kong Physiotherapy Journal.* 2003;21:37-42. X-7
1318. Lo TS, Chang TC, Chao AS, et al. Tension-free vaginal tape procedure on genuine stress incontinent women with coexisting genital prolapse. *Acta Obstet Gynecol Scand.* 2003 Nov;82(11):1049-53. X-1B
1319. Lo TS, Horng SG, Chang CL, et al. Tension-free vaginal tape procedure after previous failure in incontinence surgery. *Urology.* 2002 Jul;60(1):57-61. X-1B
1320. Lo TS, Huang HJ, Chang CL, et al. Use of intravenously anesthesia for tension-free vaginal tape therapy in elderly women with genuine stress incontinence. *Urology.* 2002 Mar;59(3):349-53. X-1B
1321. Lo TS, Lin CT, Huang HJ, et al. The use of general anesthesia for the tension-free vaginal tape procedure and concomitant surgery. *Acta Obstet Gynecol Scand.* 2003 Apr;82(4):367-73. X-1B
1322. Locher JL, Goode PS, Roth DL, et al. Reliability assessment of the bladder diary for urinary incontinence in older women. *J Gerontol A Biol Sci Med Sci.* 2001 Jan;56(1):M32-5. X-1H
1323. Lockhart JL, Ellis GF, Helal M, et al. Combined cystourethropexy for the treatment of type 3 and complicated female urinary incontinence. *J Urol.* 1990 Apr;143(4):722-5. X-1B
1324. Lockhart JL, Shessel F, Weinstein D, et al. Urodynamics in women with stress and urge incontinence. *Urology.* 1982 Sep;20(3):333-6. X-1G, X-1H
1325. Loddenkemper T, Foldvary N, Raja S, et al. Ictal urinary urge: further evidence for lateralization to the nondominant hemisphere. *Epilepsia.* 2003 Jan;44(1):124-6. X-1F, X-7
1326. Long CY, Hsu SC, Chen YH, et al. The effects on vesicourethral function following laparoscopic hysterectomy. *Kaohsiung J Med Sci.* 2001 Nov;17(11):564-9. X-1I, X-7
1327. Long CY, Hsu SC, Sun DJ, et al. Abnormal clinical and urodynamic findings in women with severe genitourinary prolapse. *Kaohsiung J Med Sci.* 2002 Dec;18(12):593-7. X-4
1328. Long CY, Hsu SC, Wu TP, et al. Effect of laparoscopic hysterectomy on bladder neck and urinary symptoms. *Aust N Z J Obstet Gynaecol.* 2003 Feb;43(1):65-9. X-1I
1329. Long CY, Hsu SC, Wu TP, et al. Urodynamic comparison of continent and incontinent women with severe uterovaginal prolapse. *J Reprod Med.* 2004 Jan;49(1):33-7. X-1E, X-1G
1330. Long CY, Jang MY, Chen SC, et al. Changes in vesicourethral function following laparoscopic hysterectomy versus abdominal hysterectomy. *Aust N Z J Obstet Gynaecol.* 2002 Aug;42(3):259-63. X-1A
1331. Long CY, Liu CM, Hsu SC, et al. A randomized comparative study of the effects of oral and topical estrogen therapy on the lower urinary tract of hysterectomized postmenopausal women. *Fertil Steril.* 2006 Jan;85(1):155-60. X-1A
1332. Loomis ME and Conco D. Patients' perceptions of health, chronic illness, and nursing diagnoses. *Nurs Diagn.* 1991 Oct-Dec;2(4):162-70. X-1A, X-1H, X-7
1333. Lopez Pereira P, Miguelez C, Caffarati J, et al. Trospium chloride for the treatment of detrusor instability in children. *J Urol.* 2003 Nov;170(5):1978-81. X-4
1334. Lorenzo AJ, Wallis MC, Cook A, et al. What is the variability in urodynamic parameters with position change in children? Analysis of a prospectively enrolled cohort. *J Urol.* 2007 Dec;178(6):2567-70. X-1H, X-4
1335. Lose G, Jorgensen L and Thunedborg P. Doxepin in the treatment of female detrusor overactivity: a randomized double-blind crossover study. *J Urol.* 1989 Oct;142(4):1024-6. X-7
1336. Lowenstein L, FitzGerald MP, Kenton K, et al. Validation of a real-time urodynamic measure of urinary sensation. *Am J Obstet Gynecol.* 2008 Jun;198(6):661 e1-4; discussion 661 e4-5. X-1H
1337. Lowenstein L, Kenton K, Brubaker L, et al. The relationship between obstructive sleep apnea, nocturia, and daytime overactive bladder syndrome in women. *Am J Obstet Gynecol.* 2008 May;198(5):598 e1-5. X-1A, X-7

1338. Lowenstein L, Kenton K, Dooley Y, et al. Women who experience detrusor overactive at lower bladder volumes report greater bother. *Neurourol Urodyn.* 2008;27(1):45-7. X-II
1339. Lowenstein L, Kenton K, FitzGerald MP, et al. Clinically useful measures in women with mixed urinary incontinence. *Am J Obstet Gynecol.* 2008 Jun;198(6):664 e1-3; discussion 664 e3-4. X-1A, X-1B, X-7
1340. Lubeck DP, Prebil LA, Peebles P, et al. A health related quality of life measure for use in patients with urge urinary incontinence: a validation study. *Qual Life Res.* 1999 Jun;8(4):337-44. X-II
1341. Lubner KM and Wolde-Tsadik G. Efficacy of functional electrical stimulation in treating genuine stress incontinence: a randomized clinical trial. *Neurourol Urodyn.* 1997;16(6):543-51. X-1B
1342. Lucas MG and Thomas DG. Endoscopic bladder transection for detrusor instability. *Br J Urol.* 1987 Jun;59(6):526-8. X-7
1343. Lucas MG, Thomas DG, Clarke S, et al. Long-term follow-up of selective sacral neurectomy. *Br J Urol.* 1988 Mar;61(3):218-20. X-7
1344. Lucioni A, Rapp DE, Gong EM, et al. Intravesical botulinum type A toxin injection in patients with overactive bladder: Trigone versus trigone-sparing injection. *Can J Urol.* 2006 Oct;13(5):3291-5. X-7
1345. Lukacz ES, Lawrence JM, Buckwalter JG, et al. Epidemiology of prolapse and incontinence questionnaire: validation of a new epidemiologic survey. *Int Urogynecol J Pelvic Floor Dysfunct.* 2005 Jul-Aug;16(4):272-84. X-1H
1346. Lutgendorf SK, Kreder KJ, Rothrock NE, et al. Diurnal cortisol variations and symptoms in patients with interstitial cystitis. *J Urol.* 2002 Mar;167(3):1338-43. X-1D
1347. Luukinen H, Koski K, Kivela SL, et al. Social status, life changes, housing conditions, health, functional abilities and life-style as risk factors for recurrent falls among the home-dwelling elderly. *Public Health.* 1996 Mar;110(2):115-8. X-1A
1348. Maahe C, Landman M, Wang X, et al. Expression of smoothelin in the normal and the overactive human bladder. *J Urol.* 2006 Mar;175(3 Pt 1):1152-7. X-1G, X-7
1349. Macaulay AJ, Stern RS and Stanton SL. Psychological aspects of 211 female patients attending a urodynamic unit. *J Psychosom Res.* 1991;35(1):1-10. X-1I
1350. MacDiarmid SA, Anderson RU, Armstrong RB, et al. Efficacy and safety of extended release oxybutynin for the treatment of urge incontinence: an analysis of data from 3 flexible dosing studies. *J Urol.* 2005 Oct;174(4 Pt 1):1301-5; discussion 1305. X-2
1351. Macfarlane JR and Tolley DA. The effect of terodiline on patients with detrusor instability. *Scand J Urol Nephrol Suppl.* 1984;87:51-4. X-7
1352. Madeiro AP, Rufino AC, Sartori MG, et al. The effects of bethanechol and cisapride on urodynamic parameters in patients undergoing radical hysterectomy for cervical cancer. A randomized, double-blind, placebo-controlled study. *Int Urogynecol J Pelvic Floor Dysfunct.* 2006 May;17(3):248-52. X-1H
1353. Madersbacher H, Halaska M, Voigt R, et al. A placebo-controlled, multicentre study comparing the tolerability and efficacy of propiverine and oxybutynin in patients with urgency and urge incontinence. *BJU Int.* 1999 Oct;84(6):646-51. X-9
1354. Madersbacher S, Pycha A, Klingler CH, et al. The International Prostate Symptom score in both sexes: a urodynamics-based comparison. *Neurourol Urodyn.* 1999;18(3):173-82. X-1H
1355. Madersbacher S, Pycha A, Schatzl G, et al. The aging lower urinary tract: a comparative urodynamic study of men and women. *Urology.* 1998 Feb;51(2):206-12. X-1H
1356. Madjar S, Covington-Nichols C and Secrest CL. New periurethral bulking agent for stress urinary incontinence: modified technique and early results. *J Urol.* 2003 Dec;170(6 Pt 1):2327-9. X-1F
1357. Mahady IW and Begg BM. Long-term symptomatic and cystometric cure of the urge incontinence syndrome using a technique of bladder re-education. *Br J Obstet Gynaecol.* 1981 Oct;88(10):1038-43. X-7
1358. Mahajan ST, Elkadry EA, Kenton KS, et al. Patient-centered surgical outcomes: the impact of goal achievement and urge incontinence on patient satisfaction one year after surgery. *Am J Obstet Gynecol.* 2006 Mar;194(3):722-8. X-1A
1359. Mahawong P, Chaiyaprasithi B, Soontrapa S, et al. A role of intravesical capsaicin instillation in benign prostatic hyperplasia with overactive bladder symptoms: the first reported study in the literature. *J Med Assoc Thai.* 2007 Sep;90(9):1821-7. X-4
1360. Maher C, Carey M, Dwyer P, et al. Pubovaginal or vicryl mesh rectus fascia sling in intrinsic sphincter deficiency. *Int Urogynecol J Pelvic Floor Dysfunct.* 2001;12(2):111-6. X-1B
1361. Maher C, Dwyer P, Carey M, et al. The Burch colposuspension for recurrent urinary stress incontinence following retropubic continence surgery. *Br J Obstet Gynaecol.* 1999 Jul;106(7):719-24. X-1B
1362. Maher CF, Dwyer PL, Carey MP, et al. Colposuspension or sling for low urethral pressure stress incontinence? *Int Urogynecol J Pelvic Floor Dysfunct.* 1999;10(6):384-9. X-1B
1363. Mahony DT, Laferte RO and Blais DJ. Incontinence of urine due to instability of micturition reflexes. Part II. Pudendal nucleus instability. *Urology.* 1980 Apr;15(4):379-88. X-1G, X-2, X-4, X-6, X-7
1364. Mahony DT, Laferte RO and Blais DJ. Incontinence of urine due to instability of micturition reflexes: Part I. Detrusor reflex instability. *Urology.* 1980 Mar;15(3):229-39. X-1G, X-2, X-4, X-6, X-7
1365. Major H, Culligan P and Heit M. Urethral sphincter morphology in women with detrusor instability. *Obstet Gynecol.* 2002 Jan;99(1):63-8. X-1G
1366. Makinen JI, Pitkanen YA, Salmi TA, et al. Transdermal estrogen for female stress urinary incontinence in postmenopause. *Maturitas.* 1995 Nov;22(3):233-8. X-1B

1367. Malm-Buatsi E, Nepple KG, Boyt MA, et al. Efficacy of transcutaneous electrical nerve stimulation in children with overactive bladder refractory to pharmacotherapy. *Urology*. 2007 Nov;70(5):980-3. X-4, X-7
1368. Malone RP, Delaney MA, Luebbert JF, et al. A double-blind placebo-controlled study of lithium in hospitalized aggressive children and adolescents with conduct disorder. *Arch Gen Psychiatry*. 2000 Jul;57(7):649-54. X-II, X-4, X-5
1369. Malone-Lee J, Shaffu B, Anand C, et al. Tolterodine: superior tolerability than and comparable efficacy to oxybutynin in individuals 50 years old or older with overactive bladder: a randomized controlled trial. *J Urol*. 2001 May;165(5):1452-6. X-4
1370. Malone-Lee JG, Walsh JB and Maugour MF. Tolterodine: a safe and effective treatment for older patients with overactive bladder. *J Am Geriatr Soc*. 2001 Jun;49(6):700-5. X-4
1371. Malouf AJ, Vaizey CJ, Nicholls RJ, et al. Permanent sacral nerve stimulation for fecal incontinence. *Ann Surg*. 2000 Jul;232(1):143-8. X-1A, X-7
1372. Manca A, Sculpher MJ, Ward K, et al. A cost-utility analysis of tension-free vaginal tape versus colposuspension for primary urodynamic stress incontinence. *BJOG*. 2003 Mar;110(3):255-62. X-1B
1373. Manning J, Eyers AA, Korda A, et al. Is there an association between fecal incontinence and lower urinary dysfunction? *Dis Colon Rectum*. 2001 Jun;44(6):790-8. X-II
1374. Manonai J, Chittacharoen A, Sarit-ahirak S, et al. Lower urinary tract symptoms in Thai women attending the menopause clinic: prevalence and associated factors. *J Med Assoc Thai*. 2004 Nov;87(11):1265-9. X-6
1375. Manonai J, Songchitsomboon S, Chanda K, et al. The effect of a soy-rich diet on urogenital atrophy: a randomized, cross-over trial. *Maturitas*. 2006 May 20;54(2):135-40. X-7
1376. Mansfield KJ, Liu L, Moore KH, et al. Molecular characterization of M2 and M3 muscarinic receptor expression in bladder from women with refractory idiopathic detrusor overactivity. *BJU Int*. 2007 Jun;99(6):1433-8. X-1G
1377. Marinkovic SP and Stanton SL. Triple compartment prolapse: sacrocolpopexy with anterior and posterior mesh extensions. *BJOG*. 2003 Mar;110(3):323-6. X-1E
1378. Mark SD, McRae CU, Arnold EP, et al. Clam cystoplasty for the overactive bladder: a review of 23 cases. *Aust N Z J Surg*. 1994 Feb;64(2):88-90. X-7
1379. Martin MR and James ED. Treatment of unstable detrusor (irritable bladder) using fluphenazine and nortriptyline: progress report. *Prog Clin Biol Res*. 1981;78:335-8. X-7
1380. Martin MR and Schiff AA. Fluphenazine/nortriptyline in the irritable bladder syndrome. A double-blind placebo controlled study. *Br J Urol*. 1984 Apr;56(2):178-9. X-7
1381. Martinez JM, Russell JM and Hirschfeld RM. Tolerability of oral loading of divalproex sodium in the treatment of acute mania. *Depress Anxiety*. 1998;7(2):83-6. X-II
1382. Mascarenhas F, Cocuzza M, Gomes CM, et al. Trigonal injection of botulinum toxin-A does not cause vesicoureteral reflux in neurogenic patients. *Neurourol Urodyn*. 2008;27(4):311-4. X-1F, X-7
1383. Mason RC and Roach M. Modified pubovaginal sling for treatment of intrinsic sphincteric deficiency. *J Urol*. 1996 Dec;156(6):1991-4. X-1B
1384. Massey JA and Abrams P. Dose titration in clinical trials. An example using emepromium carrageenate in detrusor instability. *Br J Urol*. 1986 Apr;58(2):125-8. X-9
1385. Mastropietro MA, Geary W, Fuller E, et al. Detrusor biopsy as a potential clinical tool. *Int Urogynecol J Pelvic Floor Dysfunct*. 2001;12(6):355-60. X-1G, X-1H, X-7
1386. Masunaga K, Yoshida M, Inadome A, et al. Pharmacological effects of solifenacin on human isolated urinary bladder. *Pharmacology*. 2008;82(1):43-52. X-1G
1387. Matharu G, Donaldson MM, McGrother CW, et al. Relationship between urinary symptoms reported in a postal questionnaire and urodynamic diagnosis. *Neurourol Urodyn*. 2005;24(2):100-5. X-1H
1388. Matharu GS, Assassa RP, Williams KS, et al. Objective assessment of urinary incontinence in women: comparison of the one-hour and 24-hour pad tests. *Eur Urol*. 2004 Feb;45(2):208-12. X-1H
1389. Matharu GS, Assassa RP, Williams KS, et al. Continence nurse treatment of women's urinary symptoms. *Br J Nurs*. 2004 Feb 12-25;13(3):140-3. X-II
1390. Mattiasson A, Ekstrom B and Andersson KE. Effects of intravesical instillation of verapamil in patients with detrusor hyperactivity. *J Urol*. 1989 Jan;141(1):174-7. X-7
1391. Mattiasson A and Teleman P. Abnormal urethral motor function is common in female stress, mixed, and urge incontinence. *Neurourol Urodyn*. 2006;25(7):703-8. X-II
1392. Mattox TF and Bhatia NN. The prevalence of urinary incontinence or prolapse among white and Hispanic women. *Am J Obstet Gynecol*. 1996 Feb;174(2):646-8. X-6
1393. Mattsson S and Gladh G. Urethrovaginal reflux--a common cause of daytime incontinence in girls. *Pediatrics*. 2003 Jan;111(1):136-9. X-II, X-4
1394. Matza LS, Thompson CL, Krasnow J, et al. Test-retest reliability of four questionnaires for patients with overactive bladder: the overactive bladder questionnaire (OAB-q), patient perception of bladder condition (PPBC), urgency questionnaire (UQ), and the primary OAB symptom questionnaire (POSQ). *Neurourol Urodyn*. 2005;24(3):215-25. X-II
1395. Mayo ME. Pelvic floor function in women with detrusor instability. *Prog Clin Biol Res*. 1981;78:91-9. X-1H, X-7
1396. Mazur D, Wehnert J, Dorschner W, et al. Clinical and urodynamic effects of propiverine in patients suffering from urgency and urge incontinence. A multicentre dose-optimizing study. *Scand J Urol Nephrol*. 1995 Sep;29(3):289-94. X-9
1397. McAchrans SE and Daneshgari F. Sacral neuromodulation in the older woman. *Clin Obstet Gynecol*. 2007 Sep;50(3):735-44. X-2

1398. McCahy PJ and Styles RA. Prolonged bladder distension: experience in the treatment of detrusor overactivity and interstitial cystitis. *Eur Urol.* 1995;28(4):325-7. X-7
1399. McCannel DA and Haile W. Urethral narrowing and its treatment: a possible solution for the problems of recurrent urinary infection and the "irritable bladder". *Int Urol Nephrol.* 1982;14(4):407-14. X-II
1400. McClish DK, Fantl JA, Wyman JF, et al. Bladder training in older women with urinary incontinence: relationship between outcome and changes in urodynamic observations. *Obstet Gynecol.* 1991 Feb;77(2):281-6. X-7
1401. McClish DK, Wyman JF, Sale PG, et al. Use and costs of incontinence pads in female study volunteers. Continence Program for Women Research Group. *J Wound Ostomy Continence Nurs.* 1999 Jul;26(4):207-8, 210-3. X-II
1402. McCrery R and Appell R. Transvaginal urethrolisis for obstruction after antiincontinence surgery. *Int Urogynecol J Pelvic Floor Dysfunct.* 2007 Jun;18(6):627-33. X-1A
1403. McDowell BJ, Burgio KL, Dombrowski M, et al. An interdisciplinary approach to the assessment and behavioral treatment of urinary incontinence in geriatric outpatients. *J Am Geriatr Soc.* 1992 Apr;40(4):370-4. X-7
1404. McGuire EJ. Symptoms that depress the doctor. Urge incontinence. *Br J Hosp Med.* 1984 Feb;31(2):149-54. X-2
1405. McInerney PD, Harris SA, Pritchard A, et al. Night studies for primary diurnal and nocturnal enuresis and preliminary results of the "clam" ileocystoplasty. *Br J Urol.* 1991 Jan;67(1):42-3. X-II
1406. McInerney PD, Vanner TF, Harris SA, et al. Ambulatory urodynamics. *Br J Urol.* 1991 Mar;67(3):272-4. X-1H, X-7
1407. McInerney PD, Vanner TF, Matenhelia S, et al. Assessment of the long-term results of subtrigonal phenolisation. *Br J Urol.* 1991 Jun;67(6):586-7. X-7
1408. McKenna SP, Whalley D, Renck-Hooper U, et al. The development of a quality of life instrument for use with post-menopausal women with urogenital atrophy in the UK and Sweden. *Qual Life Res.* 1999 Aug;8(5):393-8. X-II
1409. McLennan MT. The role of electrodiagnostic techniques in the reprogramming of patients with a delayed suboptimal response to sacral nerve stimulation. *Int Urogynecol J Pelvic Floor Dysfunct.* 2003 Jun;14(2):98-103. X-7
1410. McLennan MT and Bent AE. Fascia lata suburethral sling vs. Burch retropubic urethropexy. A comparison of morbidity. *J Reprod Med.* 1998 Jun;43(6):488-94. X-1B, X-6
1411. McLennan MT, Melick C and Bent AE. Urethral instability: clinical and urodynamic characteristics. *Neurourol Urodyn.* 2001;20(6):653-60. X-1G, X-1H
1412. McPherson K, Herbert A, Judge A, et al. Self-reported bladder function five years post-hysterectomy. *J Obstet Gynaecol.* 2005 Jul;25(5):469-75. X-1A
1413. McQuire WA. Electrotherapy and exercises for stress incontinence and urinary frequency. *Physiotherapy.* 1975 Oct;61(10):305-7. X-1B
1414. McRae P, Murray KH, Nurse DE, et al. Clam enterocystoplasty in the neuropathic bladder. *Br J Urol.* 1987 Dec;60(6):523-5. X-1F
1415. Meade-D'Alisera P, Merriweather T and Wentland M. Impact of commercial marketing on patient demand. *Urol Nurs.* 2001 Dec;21(6):406-7, 410. X-4
1416. Medel R, Ruarte AC, Castera R, et al. Primary enuresis: a urodynamic evaluation. *Br J Urol.* 1998 May;81 Suppl 3:50-2. X-4
1417. Medina-Bombardo D, Segui-Diaz M, Roca-Fusalba C, et al. What is the predictive value of urinary symptoms for diagnosing urinary tract infection in women? *Fam Pract.* 2003 Apr;20(2):103-7. X-1H, X-1I
1418. Menarini M, Del Popolo G, Di Benedetto P, et al. Trosipium chloride in patients with neurogenic detrusor overactivity: is dose titration of benefit to the patients? *Int J Clin Pharmacol Ther.* 2006 Dec;44(12):623-32. X-1F
1419. Merlini E, Sangiorgio L and Seymandi P. The lazy bladder syndrome: a possible urodynamic evolution in patients with idiopathic detrusor and pelvic floor overactivity. *Pediatr Med Chir.* 2004 May-Jun;26(3):187-90. X-II, X-4, X-7
1420. Meschia M, Pifarotti P, Spennacchio M, et al. A randomized comparison of tension-free vaginal tape and endopelvic fascia plication in women with genital prolapse and occult stress urinary incontinence. *Am J Obstet Gynecol.* 2004 Mar;190(3):609-13. X-1B
1421. Messing E, Pauk D, Schaeffer A, et al. Associations among cystoscopic findings and symptoms and physical examination findings in women enrolled in the Interstitial Cystitis Data Base (ICDB) Study. *Urology.* 1997 May;49(5A Suppl):81-5. X-1D
1422. Metcalfe A, Bick D, Tohill S, et al. A prospective cohort study of repair and non-repair of second-degree perineal trauma: results and issues for future research. *Evidence Based Midwifery.* 2006 Oct;4(2):60-4. X-1I
1423. Metello J, Nogueira B, Torgal M, et al. Comparison of the efficacy and tolerability of solifenacin succinate with or without previous use of trospium chloride. *Int Urogynecol J Pelvic Floor Dysfunct.* 2007 Sep;18(9):1021-5. X-7
1424. Meyhoff HH, Gerstenberg TC and Nordling J. Placebo--the drug of choice in female motor urge incontinence? *Br J Urol.* 1983 Feb;55(1):34-7. X-7
1425. Meyhoff HH, Walter S, Gerstenberg T, et al. Incontinence surgery in females with motor urge incontinence. *Prog Clin Biol Res.* 1981;78:347-50. X-7
1426. Meyhoff HH, Walter S, Gerstenberg TC, et al. Incontinence surgery in female motor urge incontinence. *Acta Obstet Gynecol Scand.* 1983;62(4):365-8. X-7
1427. Michel MC, de la Rosette JJ, Piro M, et al. Does concomitant stress incontinence alter the efficacy of tolterodine in patients with overactive bladder? *J Urol.* 2004 Aug;172(2):601-4. X-1B
1428. Milani R, Merlo E, Scalabrino S, et al. A double-blind trial of fentonium bromide in the treatment of incontinent unstable bladder. *Int J Clin Pharmacol Ther Toxicol.* 1986 Aug;24(8):421-4. X-7

1429. Milani R, Scalabrino S, Carrera S, et al. Comparison of flavoxate hydrochloride in daily dosages of 600 versus 1200 mg for the treatment of urgency and urge incontinence. *J Int Med Res.* 1988 May-Jun;16(3):244-8. X-7
1430. Milani R, Scalabrino S, Carrera S, et al. Flavoxate hydrochloride for urinary urgency after pelvic radiotherapy: comparison of 600 mg versus 1200 mg daily dosages. *J Int Med Res.* 1988 Jan-Feb;16(1):71-4. X-1A, X-7
1431. Milisen K, Dejaeger E, Braes T, et al. Process evaluation of a nurse-led multifactorial intervention protocol for risk screening and assessment of fall problems among community-dwelling older persons: a pilot-study. *J Nutr Health Aging.* 2006 Sep-Oct;10(5):446-52. X-11
1432. Millard RJ and Halaska M. Efficacy of solifenacin in patients with severe symptoms of overactive bladder: a pooled analysis. *Curr Med Res Opin.* 2006 Jan;22(1):41-8. X-2
1433. Milleman M, Langenstroer P and Guralnick ML. Post-void residual urine volume in women with overactive bladder symptoms. *J Urol.* 2004 Nov;172(5 Pt 1):1911-4. X-1H
1434. Miller EA, Amundsen CL, Toh KL, et al. Preoperative urodynamic evaluation may predict voiding dysfunction in women undergoing pubovaginal sling. *J Urol.* 2003 Jun;169(6):2234-7. X-1A
1435. Miller KL, DuBeau CE, Bergmann M, et al. Quest for a detrusor overactivity index. *J Urol.* 2002 Feb;167(2 Pt 1):578-84; discussion 584-5. X-1H
1436. Mills IW, Greenland JE, McMurray G, et al. Studies of the pathophysiology of idiopathic detrusor instability: the physiological properties of the detrusor smooth muscle and its pattern of innervation. *J Urol.* 2000 Feb;163(2):646-51. X-1G
1437. Minaglia S, Ozel B, Bizhang R, et al. Increased prevalence of interstitial cystitis in women with detrusor overactivity refractory to anticholinergic therapy. *Urology.* 2005 Oct;66(4):702-6. X-1H, X-7
1438. Minaglia S, Ozel B, Nguyen JN, et al. Validation of Spanish version of Pelvic Pain and Urgency/Frequency (PUF) patient symptom scale. *Urology.* 2005 Apr;65(4):664-9. X-11
1439. Minardi D, Piloni V, Amadi A, et al. Correlation between urodynamics and perineal ultrasound in female patients with urinary incontinence. *Neurourol Urodyn.* 2007;26(2):176-82; discussion 183-4. X-1H
1440. Minassian VA, Lovatsis D, Pascali D, et al. Effect of childhood dysfunctional voiding on urinary incontinence in adult women. *Obstet Gynecol.* 2006 Jun;107(6):1247-51. X-11
1441. Misawa T, Kamimura M, Kinoshita T, et al. Neurogenic bladder in patients with cervical compressive myelopathy. *J Spinal Disord Tech.* 2005 Aug;18(4):315-20. X-1F, X-1H
1442. Mishail A, Shahsavari M, Kim J, et al. Deficits in urological knowledge among medical students and primary care providers: potential for impact on urological care. *J Urol.* 2008 Nov;180(5):2140-7. X-11
1443. Mitsui T, Tanaka H, Moriya K, et al. Outcomes of lower urinary and bowel function in meningomyelocele patients with augmentation enterocystoplasty. *Spinal Cord.* 2008 Jun;46(6):432-7. X-1F, X-4, X-7
1444. Miyamae K, Yoshida M, Murakami S, et al. Pharmacological effects of darifenacin on human isolated urinary bladder. *Pharmacology.* 2003 Dec;69(4):205-11. X-1G, X-7
1445. Moen M and Stien R. Urge incontinence. *Ann Chir Gynaecol.* 1982;71(4):203-7. X-2
1446. Moisey CU, Stephenson TP and Brendler CB. The urodynamic and subjective results of treatment of detrusor instability with oxybutynin chloride. *Br J Urol.* 1980 Dec;52(6):472-5. X-7
1447. Moiyadi AV, Devi BI and Nair KP. Urinary disturbances following traumatic brain injury: clinical and urodynamic evaluation. *NeuroRehabilitation.* 2007;22(2):93-8. X-1F
1448. Molander U, Milsom I, Ekelund P, et al. An epidemiological study of urinary incontinence and related urogenital symptoms in elderly women. *Maturitas.* 1990 Apr;12(1):51-60. X-11
1449. Mold JW. Pharmacotherapy of urinary incontinence. *Am Fam Physician.* 1996 Aug;54(2):673-80, 683-5. X-2
1450. Monga AK, Marrero JM, Stanton SL, et al. Is there an irritable bladder in the irritable bowel syndrome? *Br J Obstet Gynaecol.* 1997 Dec;104(12):1409-12. X-11, X-7
1451. Montz FJ and Stanton SL. Q-Tip test in female urinary incontinence. *Obstet Gynecol.* 1986 Feb;67(2):258-60. X-1H
1452. Monz B, Chartier-Kastler E, Hampel C, et al. Patient characteristics associated with quality of life in European women seeking treatment for urinary incontinence: results from PURE. *Eur Urol.* 2007 Apr;51(4):1073-81; discussion 1081-2. X-11
1453. Moore K, Griffiths D, Latimer G, et al. Twenty four-hour monitoring of incontinence and bladder function in a community hospital. *J ET Nurs.* 1993 Jul-Aug;20(4):163-8. X-1H
1454. Moore KH, Foote A, Siva S, et al. The use of the bladder neck support prosthesis in combined genuine stress incontinence and detrusor instability. *Aust N Z J Obstet Gynaecol.* 1997 Nov;37(4):440-5. X-1B
1455. Moore KH, Gilpin SA, Dixon JS, et al. Increase in presumptive sensory nerves of the urinary bladder in idiopathic detrusor instability. *Br J Urol.* 1992 Oct;70(4):370-2. X-1H, X-7
1456. Moore KH, Lam DS, Lynch W, et al. The tachykinin NK-2 receptor antagonist SR48968 does not block noncholinergic contractions in unstable human bladder. *Peptides.* 2002 Jun;23(6):1155-60. X-1G
1457. Moore KH, O'Sullivan RJ, Simons A, et al. Randomised controlled trial of urose continence advisor therapy compared with standard urogynaecology regimen for conservative incontinence treatment: efficacy, costs and two year follow up. *BJOG.* 2003 Jul;110(7):649-57. X-7
1458. Moore KH, Richmond DH and Parys BT. Sex distribution of adult idiopathic detrusor instability in relation to childhood bedwetting. *Br J Urol.* 1991 Nov;68(5):479-82. X-11
1459. Moore KH, Simons A, Mukerjee C, et al. The relative incidence of detrusor instability and bacterial cystitis detected on the urodynamic-test day. *BJU Int.* 2000 May;85(7):786-92. X-1H

1460. Morabito F, Rossi R, Graziano ME, et al. Multicenter study on the use of gemcitabine to prevent recurrence of multiple-recurring superficial bladder tumors following intravesical antitubercular agents and/or BCG: evaluation of tolerance. *Arch Ital Urol Androl.* 2006 Mar;78(1):1-4. X-II
1461. Moran PA, Dwyer PL, Carey MP, et al. Oral methotrexate in the management of refractory interstitial cystitis. *Aust N Z J Obstet Gynaecol.* 1999 Nov;39(4):468-71. X-1D
1462. Moran PA, Ward KL, Johnson D, et al. Tension-free vaginal tape for primary genuine stress incontinence: a two-centre follow-up study. *BJU Int.* 2000 Jul;86(1):39-42. X-1B
1463. Morgan JE, Heritz DM, Stewart FE, et al. The polypropylene pubovaginal sling for the treatment of recurrent stress urinary incontinence. *J Urol.* 1995 Sep;154(3):1013-4; discussion 1015-6. X-1B
1464. Morgan TO, Jr., Westney OL and McGuire EJ. Pubovaginal sling: 4-YEAR outcome analysis and quality of life assessment. *J Urol.* 2000 Jun;163(6):1845-8. X-1B
1465. Morita T, Iizuka H, Iwata T, et al. Function and distribution of beta3-adrenoceptors in rat, rabbit and human urinary bladder and external urethral sphincter. *J Smooth Muscle Res.* 2000 Feb;36(1):21-32. X-1G
1466. Morkved S and Bo K. Effect of postpartum pelvic floor muscle training in prevention and treatment of urinary incontinence: a one-year follow up. *BJOG.* 2000 Aug;107(8):1022-8. X-1B
1467. Morris AR, O'Sullivan R, Dunkley P, et al. Extracorporeal magnetic stimulation is of limited clinical benefit to women with idiopathic detrusor overactivity: a randomized sham controlled trial. *Eur Urol.* 2007 Sep;52(3):876-81. X-7
1468. Morris AR, Westbrook JI and Moore KH. A longitudinal study over 5 to 10 years of clinical outcomes in women with idiopathic detrusor overactivity. *BJOG.* 2008 Jan;115(2):239-46. X-11
1469. Morrison LM, Eadie AS, McAlister A, et al. Personality testing in 226 patients with urinary incontinence. *Br J Urol.* 1986 Aug;58(4):387-9. X-11
1470. Moser F, Bjelic-Radisic V and Tamussino K. Needle suspension of the bladder neck for stress urinary incontinence: objective results at 11 to 16 years. *Int Urogynecol J Pelvic Floor Dysfunct.* 2006 Nov;17(6):611-4. X-1B
1471. Moskowitz MO, Byrne DS, Callahan HJ, et al. Decreased expression of a glycoprotein component of bladder surface mucin (GP1) in interstitial cystitis. *J Urol.* 1994 Feb;151(2):343-5. X-1D, X-1G
1472. Mosley LJ, Lee GP, Hughes ML, et al. Analysis of symptoms, functional impairments, and participation in occupational therapy for individuals with multiple sclerosis. *Occupational Therapy in Health Care.* 2003;17(3/4):27-43. X-1F, X-7
1473. Mukerji G, Waters J, Chessell IP, et al. Pain during ice water test distinguishes clinical bladder hypersensitivity from overactivity disorders. *BMC Urol.* 2006;6:31. X-1H, X-7
1474. Mulholland SG. UTI in women: how best to treat acute and recurrent infections... urinary tract infection (UTI). *Consultant (00107069).* 1999;39(5):1457. X-II
1475. Muller SC, Frohneberg D, Schwab R, et al. Selective sacral nerve blockade for the treatment of unstable bladders. *Eur Urol.* 1986;12(6):408-12. X-7
1476. Mulvany NJ and Riley CB. Granulosa cell tumors of unilocular cystic type. *Pathology.* 1997 Nov;29(4):348-53. X-1A, X-1G, X-7
1477. Munding M, Wessells H, Thornberry B, et al. Use of tolterodine in children with dysfunctional voiding: an initial report. *J Urol.* 2001 Mar;165(3):926-8. X-II, X-4
1478. Mundy AR. Bladder transection for urge incontinence associated with detrusor instability. *Br J Urol.* 1980 Dec;52(6):480-3. X-7
1479. Mundy AR. The surgical treatment of urge incontinence of urine. *J Urol.* 1982 Sep;128(3):481-3. X-7
1480. Mundy AR and Stephenson TP. "Clam" ileocystoplasty for the treatment of refractory urge incontinence. *Br J Urol.* 1985 Dec;57(6):641-6. X-7
1481. Murakami S, Yoshida M, Iwashita H, et al. Pharmacological effects of KRP-197 on the human isolated urinary bladder. *Urol Int.* 2003;71(3):290-8. X-1G
1482. Murphy EL, Glynn SA, Frider J, et al. Increased prevalence of infectious diseases and other adverse outcomes in human T lymphotropic virus types I- and II-infected blood donors. *Retrovirus Epidemiology Donor Study (REDS) Study Group. J Infect Dis.* 1997 Dec;176(6):1468-75. X-1G, X-II
1483. Murray C, Goh JT, Fynes M, et al. Urinary and faecal incontinence following delayed primary repair of obstetric genital fistula. *BJOG.* 2002 Jul;109(7):828-32. X-11
1484. Murray K, Mundy AR, Blackford HN, et al. Transvesical phenolisation of the pelvic plexuses: a simple technique for the treatment of refractory detrusor instability and hyperreflexia. *Urol Int.* 1986;41(3):202-6. X-9
1485. Murray KH. Effect of naloxone-induced opioid blockade on idiopathic detrusor instability. *Urology.* 1983 Sep;22(3):329-31. X-7
1486. Musa AA. Use of double-J stents prior to extracorporeal shock wave lithotripsy is not beneficial: results of a prospective randomized study. *Int Urol Nephrol.* 2008;40(1):19-22. X-1A
1487. Muskat Y, Bukovsky I, Schneider D, et al. The use of scopolamine in the treatment of detrusor instability. *J Urol.* 1996 Dec;156(6):1989-90. X-7
1488. Musselman DM, Ford AP, Gennevois DJ, et al. A randomized crossover study to evaluate Ro 115-1240, a selective alpha1A/1L-adrenoceptor partial agonist in women with stress urinary incontinence. *BJU Int.* 2004 Jan;93(1):78-83. X-1B
1489. Mustafa M and Wadie BS. Placard-shaped in situ vaginal wall sling for the treatment of stress urinary incontinence. *Int J Urol.* 2006 Feb;13(2):132-4. X-1B
1490. Myren CJ, Beuke HP and Thybo E. Functional results after ileocystoplasty. *Int Urol Nephrol.* 1988;20(3):269-74. X-1A, X-7
1491. Nabi G, Singh I, Ansari MS, et al. Primary small cell neuroendocrine carcinoma of urinary bladder: an uncommon entity to be recognized. *Int Urol Nephrol.* 2001;33(4):637-40. X-II
1492. Nager CW, Albo ME, Fitzgerald MP, et al. Reference urodynamic values for stress incontinent women. *NeuroUrol Urodyn.* 2007;26(3):333-40. X-1B

1493. Nager CW, FitzGerald M, Kraus SR, et al. Urodynamic measures do not predict stress continence outcomes after surgery for stress urinary incontinence in selected women. *J Urol*. 2008 Apr;179(4):1470-4. X-1B
1494. Naglie G, Radomski SB, Brymer C, et al. A randomized, double-blind, placebo controlled crossover trial of nimodipine in older persons with detrusor instability and urge incontinence. *J Urol*. 2002 Feb;167(2 Pt 1):586-90. X-9
1495. Nakamura M, Sakurai T, Sugao H, et al. Maximum electrical stimulation for urge incontinence. *Urol Int*. 1987;42(4):285-7. X-7
1496. Nakamura M, Sakurai T, Tsujimoto Y, et al. Bladder inhibition by electrical stimulation of the perianal skin. *Urol Int*. 1986;41(1):62-3. X-7
1497. Nama G. Toilet tied. *Nursing in the Community*. 2007;8(6):17-19. X-10
1498. Naoemova I, De Wachter S, Wuyts FL, et al. Do sensation-related bladder diaries differ between patients with urodynamically confirmed and non-objectivised urinary incontinence? *Int Urogynecol J Pelvic Floor Dysfunct*. 2008 Feb;19(2):213-6. X-1A
1499. Naoemova I, De Wachter S, Wuyts FL, et al. Reliability of the 24-h sensation-related bladder diary in women with urinary incontinence. *Int Urogynecol J Pelvic Floor Dysfunct*. 2008 Jul;19(7):955-9. X-1H
1500. Naoemova I, De Wachter S and Wyndaele JJ. Comparison of sensation-related voiding patterns between continent and incontinent women: a study with a 3-day sensation-related bladder diary (SR-BD). *NeuroUrol Urodyn*. 2008;27(6):511-4. X-1H
1501. Natsume O. Detrusor contractility and overactive bladder in patients with cerebrovascular accident. *Int J Urol*. 2008 Jun;15(6):505-10; discussion 510. X-1F, X-7
1502. Nauth MA and Funfgeld C. Correction of cystocele and stress incontinence with anterior transobturator mesh. *Eur J Obstet Gynecol Reprod Biol*. 2008 Feb;136(2):249-53. X-1B
1503. Nazarko L. Management of a patient with diabetes and a hypotonic bladder. *Nurs Times*. 2005 Nov 22-28;101(47):63-4. X-1F, X-7
1504. Nazarko L. Treating the person with an overactive bladder. *Nursing & Residential Care*. 2006;8(7):303-305. X-2
1505. Nazarko L. Stroke and continence: the benefits of assessment. *Nursing & Residential Care*. 2007 May;9(5):203-6. X-1I
1506. Nazir M, Stien R, Carlsen E, et al. Early evaluation of bowel symptoms after primary repair of obstetric perineal rupture is misleading: an observational cohort study. *Dis Colon Rectum*. 2003 Sep;46(9):1245-50. X-1I
1507. Ndamba J, Nyazema N, Makaza N, et al. Traditional herbal remedies used for the treatment of urinary schistosomiasis in Zimbabwe. *J Ethnopharmacol*. 1994 Apr;42(2):125-32. X-1I
1508. N'Dow J, Leung HY, Marshall C, et al. Bowel dysfunction after bladder reconstruction. *J Urol*. 1998 May;159(5):1470-4; discussion 1474-5. X-7
1509. Nejat F, Radmanesh F, Ansari S, et al. Spina bifida occulta: is it a predictor of underlying spinal cord abnormality in patients with lower urinary tract dysfunction? *J Neurosurg Pediatrics*. 2008 Feb;1(2):114-7. X-1F
1510. Nelson MS. Acute urinary retention secondary to an incarcerated gravid uterus. *Am J Emerg Med*. 1986 May;4(3):231-2. X-1I, X-7
1511. Neuhaus J, Pfeiffer F, Wolburg H, et al. Alterations in connexin expression in the bladder of patients with urge symptoms. *BJU Int*. 2005 Sep;96(4):670-6. X-1G
1512. Neumann G, Olesen PG, Hansen V, et al. The short-term prevalence of de novo urinary symptoms after different modes of hysterectomy. *Int Urogynecol J Pelvic Floor Dysfunct*. 2004 Jan-Feb;15(1):14-9; discussion 19. X-1A
1513. Neumann GA, Lauszus FF, Ljungstrom B, et al. Incidence and remission of urinary incontinence after hysterectomy--a 3-year follow-up study. *Int Urogynecol J Pelvic Floor Dysfunct*. 2007 Apr;18(4):379-82. X-6
1514. Neveus T, Hetta J, Cnattingius S, et al. Depth of sleep and sleep habits among enuretic and incontinent children. *Acta Paediatr*. 1999 Jul;88(7):748-52. X-4
1515. Newman DK. Report of a mail survey of women with bladder control disorders. *Urol Nurs*. 2004 Dec;24(6):499-507. X-1I
1516. Newton M, Kosier JH and Smith D. Treatments for overactive bladder. *Urol Nurs*. 2000 Aug;20(4):267-8. X-2
1517. Ney P, Pandita RK, Newgreen DT, et al. Pharmacological characterization of a novel investigational antimuscarinic drug, fesoterodine, in vitro and in vivo. *BJU Int*. 2008 Apr;101(8):1036-42. X-1G
1518. Ng CC, Lee LC and Han WH. Use of three-dimensional ultrasound scan to assess the clinical importance of midurethral placement of the tension-free vaginal tape (TVT) for treatment of incontinence. *Int Urogynecol J Pelvic Floor Dysfunct*. 2005 May-Jun;16(3):220-5. X-1B
1519. Ng S, Tee YT, Tsui KP, et al. Is the role of Burch colposuspension fading away in this epoch for treating female urinary incontinence? *Int Urogynecol J Pelvic Floor Dysfunct*. 2007 Aug;18(8):937-42. X-1B
1520. Ng SC, Lin TL, Chang SJ, et al. Nursing intervention to enhance efficacy of home practice of pelvic floor muscle exercises in treating mixed urinary incontinence. *Int Urogynecol J Pelvic Floor Dysfunct*. 2008 May;19(5):637-42. X-1B
1521. Nguyen A, Mahoney S, Minor L, et al. A simple objective method of adjusting sling tension. *J Urol*. 1999 Nov;162(5):1674-6. X-1B
1522. Nguyen JK and Bhatia NN. Resolution of motor urge incontinence after surgical repair of pelvic organ prolapse. *J Urol*. 2001 Dec;166(6):2263-6. X-1A
1523. Nguyen JK and Bhatia NN. Risk of recurrent stress incontinence in women undergoing the combined modified pereyra procedure and transvaginal sacrospinous ligament vault suspension. *Urology*. 2001 Dec;58(6):947-52. X-1B
1524. Nickel JC, Tripp D, Teal V, et al. Sexual function is a determinant of poor quality of life for women with treatment refractory interstitial cystitis. *J Urol*. 2007 May;177(5):1832-6. X-1D

1525. Nicolson P, Kopp Z, Chapple CR, et al. It's just the worry about not being able to control it! A qualitative study of living with overactive bladder. *Br J Health Psychol*. 2008 May;13(Pt 2):343-59. X-II
1526. Nigro DA, Wein AJ, Foy M, et al. Associations among cystoscopic and urodynamic findings for women enrolled in the Interstitial Cystitis Data Base (ICDB) Study. *Urology*. 1997 May;49(5A Suppl):86-92. X-1D
1527. Nijman RJ, Borgstein NG, Ellsworth P, et al. Tolterodine treatment for children with symptoms of urinary urge incontinence suggestive of detrusor overactivity: results from 2 randomized, placebo controlled trials. *J Urol*. 2005 Apr;173(4):1334-9. X-4
1528. Nijman RJ, Borgstein NG, Ellsworth P, et al. Long-term tolerability of tolterodine extended release in children 5-11 years of age: results from a 12-month, open-label study. *Eur Urol*. 2007 Nov;52(5):1511-6. X-4
1529. Nilsson CG, Lukkari E, Haarala M, et al. Comparison of a 10-mg controlled release oxybutynin tablet with a 5-mg oxybutynin tablet in urge incontinent patients. *Neurourol Urodyn*. 1997;16(6):533-42. X-7
1530. Nilvebrant L, Andersson KE, Gillberg PG, et al. Tolterodine--a new bladder-selective antimuscarinic agent. *Eur J Pharmacol*. 1997 May 30;327(2-3):195-207. X-1G, X-4
1531. Nissenkorn I and De Jong PR. A novel surgical technique for implanting a new electrostimulation system for treating female overactive bladder: a preliminary report. *BJU Int*. 2005 Jun;95(9):1253-8. X-7
1532. Nissenkorn I, Katz A, Rubinstein R, et al. Selective effect of nortriptyline on smooth muscle as an anticholinergic drug: a pharmacological and clinical study. *Eur Urol*. 1986;12(2):109-12. X-1G, X-7
1533. Nissenkorn I, Shalev M, Radziszewski P, et al. Patient-adjusted intermittent electrostimulation for treating stress and urge urinary incontinence. *BJU Int*. 2004 Jul;94(1):105-9. X-1B, X-7
1534. Nitahara KS, Aboseif S and Tanagho EA. Long-term results of colpocystourethropey for persistent or recurrent stress urinary incontinence. *J Urol*. 1999 Jul;162(1):138-41. X-1B
1535. Nitti VW and Combs AJ. Correlation of Valsalva leak point pressure with subjective degree of stress urinary incontinence in women. *J Urol*. 1996 Jan;155(1):281-5. X-1B, X-1H
1536. Nitti VW, Dmochowski R, Appell RA, et al. Efficacy and tolerability of tolterodine extended-release in continent patients with overactive bladder and nocturia. *BJU Int*. 2006 Jun;97(6):1262-6. X-4
1537. Nitti VW, Tu LM and Gitlin J. Diagnosing bladder outlet obstruction in women. *J Urol*. 1999 May;161(5):1535-40. X-1I
1538. Nixon A, Colman S, Sabounjian L, et al. A validated patient reported measure of urinary urgency severity in overactive bladder for use in clinical trials. *J Urol*. 2005 Aug;174(2):604-7. X-1I
1539. Noblett KL, Jensen JK and Ostergard DR. The relationship of body mass index to intra-abdominal pressure as measured by multichannel cystometry. *Int Urogynecol J Pelvic Floor Dysfunct*. 1997;8(6):323-6. X-1I
1540. Nordling J and van Ophoven A. Intravesical glycosaminoglycan replenishment with chondroitin sulphate in chronic forms of cystitis. A multi-national, multi-centre, prospective observational clinical trial. *Arzneimittelforschung*. 2008;58(7):328-35. X-1D
1541. Norris RD, Sur RL, Springhart WP, et al. A prospective, randomized, double-blinded placebo-controlled comparison of extended release oxybutynin versus phenazopyridine for the management of postoperative ureteral stent discomfort. *Urology*. 2008 May;71(5):792-5. X-1I
1542. Norton C and Kamm MA. Outcome of biofeedback for faecal incontinence. *Br J Surg*. 1999 Sep;86(9):1159-63. X-1I
1543. Norton P, Karram M, Wall LL, et al. Randomized double-blind trial of terodiline in the treatment of urge incontinence in women. *Obstet Gynecol*. 1994 Sep;84(3):386-91. X-9
1544. Nuhoglu B, Fidan V, Ayyildiz A, et al. Stoller afferent nerve stimulation in woman with therapy resistant over active bladder; a 1-year follow up. *Int Urogynecol J Pelvic Floor Dysfunct*. 2006 May;17(3):204-7. X-7
1545. Nuotio M, Jylha M, Luukkaala T, et al. Urgency, urge incontinence and voiding symptoms in men and women aged 70 years and over. *BJU Int*. 2002 Mar;89(4):350-5. X-2
1546. Nuotio M, Tammela TL, Luukkaala T, et al. Predictors of institutionalization in an older population during a 13-year period: the effect of urge incontinence. *J Gerontol A Biol Sci Med Sci*. 2003 Aug;58(8):756-62. X-1I
1547. Nygaard I, Kreder K, Mueller E, et al. Does urethral competence affect urodynamic voiding parameters in women with prolapse? *Neurourol Urodyn*. 2007;26(7):1030-5. X-1H
1548. Nygaard I and Linder M. Thirst at work--an occupational hazard? *Int Urogynecol J Pelvic Floor Dysfunct*. 1997;8(6):340-3. X-1I
1549. Nygaard IE. Does prolonged high-impact activity contribute to later urinary incontinence? A retrospective cohort study of female Olympians. *Obstet Gynecol*. 1997 Nov;90(5):718-22. X-1I
1550. Nygaard IE, Kreder KJ, Lepic MM, et al. Efficacy of pelvic floor muscle exercises in women with stress, urge, and mixed urinary incontinence. *Am J Obstet Gynecol*. 1996 Jan;174(1 Pt 1):120-5. X-7
1551. Obrink A and Bunne G. Treatment of urgency by instillation of emepromium bromide in the urinary bladder. *Scand J Urol Nephrol*. 1978;12(3):215-8. X-7
1552. Obrink A and Bunne G. Urethrocystometric findings in urgency. *Urol Int*. 1978;33(6):377-80. X-1G
1553. Odegard PS and Setter SM. Geriatric pharmacotherapy update: new medications, recent releases, and coming attractions. *Annals of Long Term Care*. 2006;14(5):20-27. X-2
1554. Oga M, Shono H, Kohara M, et al. Chronological changes in subjective symptoms during pregnancy in nulliparous and multiparous women. *Acta Obstet Gynecol Scand*. 1995 Nov;74(10):784-7. X-1A

1555. Ogundipe A, Rosenzweig BA, Karram MM, et al. Modified suburethral sling procedure for treatment of recurrent or severe stress urinary incontinence. *Surg Gynecol Obstet.* 1992 Aug;175(2):173-6. X-1B
1556. Oh SJ, Choo MS, Kim HS, et al. Generic and disease-specific health-related quality of life in women with coital incontinence: a prospective, multicenter study. *Gynecol Obstet Invest.* 2008;65(1):62-7. X-1A
1557. Oh SJ and Ku JH. Impact of self-perceived bothersomeness on health-related quality of life in women with overactive bladder symptoms. *Urol Int.* 2007;78(1):63-9. X-1H
1558. Oh SJ and Ku JH. Impact of stress urinary incontinence and overactive bladder on micturition patterns and health-related quality of life. *Int Urogynecol J Pelvic Floor Dysfunct.* 2007 Jan;18(1):65-71. X-II
1559. Oh SJ, Ku JH, Choo MS, et al. Health-related quality of life and sexual function in women with stress urinary incontinence and overactive bladder. *Int J Urol.* 2008 Jan;15(1):62-7; discussion 67. X-II
1560. Oh SJ, Ku JH, Hong SK, et al. Factors influencing self-perceived disease severity in women with stress urinary incontinence combined with or without urge incontinence. *Neurourol Urodyn.* 2005;24(4):341-7. X-1B
1561. Oh SJ, Son H, Jeong JY, et al. Patients' experience with ambulatory urodynamics. A prospective study. *Scand J Urol Nephrol.* 2006;40(5):391-6. X-1H, X-7
1562. Oh SJ, Son H, Jeong JY, et al. Anticholinergics [corrected] in patients with overactive bladder: Assessment of ambulatory urodynamics and patient perception. *Neurourol Urodyn.* 2007;26(6):789-93. X-1H, X-7
1563. Oh SJ, Son H, Jeong JY, et al. Anticholinergics in patients with overactive bladder: Assessment of ambulatory urodynamics and patient perception. *Neurourol Urodyn.* 2007;26(6):789-93. X-7
1564. Ohlsson BL, Fall M and Frankenberg-Sommar S. Effects of external and direct pudendal nerve maximal electrical stimulation in the treatment of the uninhibited overactive bladder. *Br J Urol.* 1989 Oct;64(4):374-80. X-7
1565. Okada H, Sengoku J, Gohji K, et al. Clinical effect of propiverine in patients with urge or stress incontinence. Kobe University Incontinence Study Group. *Hinyokika Kyo.* 1998 Jan;44(1):65-9. X-7
1566. Okada N, Igawa Y, Ogawa A, et al. Transcutaneous electrical stimulation of thigh muscles in the treatment of detrusor overactivity. *Br J Urol.* 1998 Apr;81(4):560-4. X-7
1567. Okamura K, Hasegawa T, Nojiri Y, et al. Diagnosis and treatment of lower urinary tract symptoms in the elderly by general practitioners. *Geriatr Gerontol Int.* 2008 Jun;8(2):119-25. X-II, X-4, X-7
1568. Okamura K, Nojiri Y, Hirano M, et al. Lower urinary tract symptoms (LUTS) in people aged 50 or older consulting general practitioners. *Geriatrics & Gerontology International.* 2007 Jun;7(2):147-53. X-6
1569. Okorochoa I, Cumming G and Gould I. Female urodynamics and lower urinary tract infection. *BJU Int.* 2002 Jun;89(9):863-7. X-1A, X-1H
1570. Oldenburg B and Millard RJ. Predictors of long term outcome following a bladder re-training programme. *J Psychosom Res.* 1986;30(6):691-8. X-1A
1571. O'Leary M, Erickson JR, Smith CP, et al. Effect of controlled-release oxybutynin on neurogenic bladder function in spinal cord injury. *J Spinal Cord Med.* 2003 Summer;26(2):159-62. X-1F, X-7
1572. Olesen KP and Walter S. Posterior bladder suspension defects in the female. A radiological classification with urodynamic and clinical evaluation. *Acta Obstet Gynecol Scand.* 1980;59(6):543-8. X-1E, X-1G
1573. Olesen KP, Walter S, Frimodt-Moller C, et al. Morphology and function of the bladder and urethra in female urinary incontinence. *Int Urol Nephrol.* 1975;7(4):303-13. X-1H, X-2
1574. Oliver S, Fowler C, Mundy A, et al. Measuring the sensations of urge and bladder filling during cystometry in urge incontinence and the effects of neuromodulation. *Neurourol Urodyn.* 2003;22(1):7-16. X-1H
1575. Olsson B and Landgren BM. The effect of tolterodine on the pharmacokinetics and pharmacodynamics of a combination oral contraceptive containing ethinyl estradiol and levonorgestrel. *Clin Ther.* 2001 Nov;23(11):1876-88. X-1A, X-7
1576. Olsson B and Szamosi J. Food does not influence the pharmacokinetics of a new extended release formulation of tolterodine for once daily treatment of patients with overactive bladder. *Clin Pharmacokinet.* 2001;40(2):135-43. X-7
1577. Olsson B and Szamosi J. Multiple dose pharmacokinetics of a new once daily extended release tolterodine formulation versus immediate release tolterodine. *Clin Pharmacokinet.* 2001;40(3):227-35. X-7
1578. Ommert A, Albrecht K, Wenger F, et al. Stapled transanal rectal resection (STARR): a new option in the treatment of obstructive defecation syndrome. *Langenbecks Arch Surg.* 2006 Feb;391(1):32-7. X-II
1579. Opisso E, Borau A, Rodriguez A, et al. Patient controlled versus automatic stimulation of pudendal nerve afferents to treat neurogenic detrusor overactivity. *J Urol.* 2008 Oct;180(4):1403-8. X-1F
1580. Opsomer RJ, Klarskov P, Holm-Bentzen M, et al. Long term results of superselective sacral nerve resection for motor urge incontinence. *Scand J Urol Nephrol.* 1984;18(2):101-5. X-7
1581. O'Reilly BA, Kosaka AH, Knight GF, et al. P2X receptors and their role in female idiopathic detrusor instability. *J Urol.* 2002 Jan;167(1):157-64. X-1G
1582. Ornan D, White R, Pollak J, et al. Pelvic embolization for intractable postpartum hemorrhage: long-term follow-up and implications for fertility. *Obstet Gynecol.* 2003 Nov;102(5 Pt 1):904-10. X-1A, X-7
1583. Oskay UY, Beji NK and Yalcin O. A study on urogenital complaints of postmenopausal women aged 50 and over. *Acta Obstet Gynecol Scand.* 2005 Jan;84(1):72-8. X-6
1584. Osman T. Stress incontinence surgery for patients presenting with mixed incontinence and a normal cystometrogram. *BJU Int.* 2003 Dec;92(9):964-8. X-1B
1585. Osorio AV and Alon US. The relationship between urinary calcium, sodium, and potassium excretion and the role of potassium in treating idiopathic hypercalciuria. *Pediatrics.* 1997 Oct;100(4):675-81. X-II

1586. O'Sullivan R, Karantanis E, Stevermuer TL, et al. Definition of mild, moderate and severe incontinence on the 24-hour pad test. *BJOG*. 2004 Aug;111(8):859-62. X-1A
1587. Ouslander J, Staskin D, Raz S, et al. Clinical versus urodynamic diagnosis in an incontinent geriatric female population. *J Urol*. 1987 Jan;137(1):68-71. X-1H
1588. Ouslander JG. Geriatric considerations in the diagnosis and management of overactive bladder. *Urology*. 2002 Nov;60(5 Suppl 1):50-5; discussion 55. X-6, X-7
1589. Ouslander JG, Blaustein J, Connor A, et al. Pharmacokinetics and clinical effects of oxybutynin in geriatric patients. *J Urol*. 1988 Jul;140(1):47-50. X-7
1590. Ouslander JG, Schnelle JF, Uman G, et al. Does oxybutynin add to the effectiveness of prompted voiding for urinary incontinence among nursing home residents? A placebo-controlled trial. *J Am Geriatr Soc*. 1995 Jun;43(6):610-7. X-5
1591. Ozel SK, Dokumcu Z, Akyildiz C, et al. Factors affecting renal scar development in children with spina bifida. *Urol Int*. 2007;79(2):133-6. X-1F, X-4, X-6, X-7
1592. Ozkan KU, Bauer SB, Khoshbin S, et al. Neurogenic bladder dysfunction after sacrococcygeal teratoma resection. *J Urol*. 2006 Jan;175(1):292-6; discussion 296. X-1A, X-1F, X-4
1593. Ozono S, Okajima E, Hirao Y, et al. Transcatheter arterial embolization of vesical artery in the treatment of invasive bladder cancer. *Eur Urol*. 1988;15(3-4):176-9. X-1I
1594. Pace G, Aceto G, Cormio L, et al. Nocturnal enuresis can be caused by absorptive hypercalciuria. *Scand J Urol Nephrol*. 1999 Apr;33(2):111-4. X-1C
1595. Pages IH, Jahr S, Schaufele MK, et al. Comparative analysis of biofeedback and physical therapy for treatment of urinary stress incontinence in women. *Am J Phys Med Rehabil*. 2001 Jul;80(7):494-502. X-1B, X-7
1596. Paick JS, Kim SW, Ku JH, et al. Preoperative maximal flow rate may be a predictive factor for the outcome of tension-free vaginal tape procedure for stress urinary incontinence. *Int Urogynecol J Pelvic Floor Dysfunct*. 2004 Nov-Dec;15(6):413-7; discussion 417. X-1B
1597. Paick JS, Ku JH, Kim SW, et al. Tension-free vaginal tape procedure for the treatment of mixed urinary incontinence: significance of maximal urethral closure pressure. *J Urol*. 2004 Sep;172(3):1001-5. X-1B
1598. Paick JS, Ku JH, Shin JW, et al. Significance of pad test loss for the evaluation of women with urinary incontinence. *Neurourol Urodyn*. 2005;24(1):39-43. X-1B
1599. Palleschi G, Pastore AL, Stocchi F, et al. Correlation between the Overactive Bladder questionnaire (OAB-q) and urodynamic data of Parkinson disease patients affected by neurogenic detrusor overactivity during antimuscarinic treatment. *Clin Neuropharmacol*. 2006 Jul-Aug;29(4):220-9. X-1F, X-7
1600. Palma P, Riccetto C, Herrmann V, et al. Transobturator crossover readjustable sling for severe female incontinence: technique and preliminary results. *Int Urogynecol J Pelvic Floor Dysfunct*. 2007 Nov;18(11):1277-82. X-1B
1601. Palma PC, Thiel M, Riccetto CL, et al. Resiniferatoxin for detrusor instability refractory to anticholinergics. *Int Braz J Urol*. 2004 Jan-Feb;30(1):53-8. X-7
1602. Palmer J. Report of a double-blind crossover study of flurbiprofen and placebo in detrusor instability. *J Int Med Res*. 1983;11 Suppl 2:11-7. X-7
1603. Panayi DC, Duckett J, Digesu GA, et al. Pre-operative opening detrusor pressure is predictive of detrusor overactivity following TVT in patients with pre-operative mixed urinary incontinence. *Neurourol Urodyn*. 2009;28(1):82-5. X-1A
1604. Pannek J and Pieper P. Clinical usefulness of ambulatory urodynamics in the diagnosis and treatment of lower urinary tract dysfunction. *Scand J Urol Nephrol*. 2008;42(5):428-32. X-1H, X-4, X-7
1605. Papa Petros PE. Detrusor instability and low compliance may represent different levels of disturbance in peripheral feedback control of the micturition reflex. *Neurourol Urodyn*. 1999;18(2):81-91. X-1G
1606. Papa Petros PE and Ulmsten U. An anatomical classification--a new paradigm for management of urinary dysfunction in the female. *Int Urogynecol J Pelvic Floor Dysfunct*. 1999;10(1):29-35. X-1G
1607. Parazzini F, Lavezzari M and Arbitani W. Prevalence of overactive bladder and urinary incontinence. *J Fam Pract*. 2002 Dec;51(12):1072-5. X-6
1608. Parekh DJ, Pope JCT, Adams MC, et al. The use of radiography, urodynamic studies and cystoscopy in the evaluation of voiding dysfunction. *J Urol*. 2001 Jan;165(1):215-8. X-1I, X-4
1609. Parekh DJ, Pope JI, Adams MC, et al. The role of hypercalciuria in a subgroup of dysfunctional voiding syndromes of childhood. *J Urol*. 2000 Sep;164(3 Pt 2):1008-10. X-4
1610. Parkin DE, Davis JA and Symonds RP. Long-term bladder symptomatology following radiotherapy for cervical carcinoma. *Radiother Oncol*. 1987 Jul;9(3):195-9. X-1I
1611. Parkin DE, Davis JA and Symonds RP. Urodynamic findings following radiotherapy for cervical carcinoma. *Br J Urol*. 1988 Mar;61(3):213-7. X-1I
1612. Parsons CL, Dell J, Stanford EJ, et al. Increased prevalence of interstitial cystitis: previously unrecognized urologic and gynecologic cases identified using a new symptom questionnaire and intravesical potassium sensitivity. *Urology*. 2002 Oct;60(4):573-8. X-1D
1613. Parsons CL, Greenberger M, Gabal L, et al. The role of urinary potassium in the pathogenesis and diagnosis of interstitial cystitis. *J Urol*. 1998 Jun;159(6):1862-6; discussion 1866-7. X-1D
1614. Parsons CL and Koprowski PF. Interstitial cystitis: successful management by increasing urinary voiding intervals. *Urology*. 1991 Mar;37(3):207-12. X-1D, X-7
1615. Parsons M, Amundsen CL, Cardozo L, et al. Bladder diary patterns in detrusor overactivity and urodynamic stress incontinence. *Neurourol Urodyn*. 2007;26(6):800-6. X-1B
1616. Parys BT, Woolfenden KA and Parsons KF. Bladder dysfunction after simple hysterectomy: urodynamic and neurological evaluation. *Eur Urol*. 1990;17(2):129-33. X-1H

1617. Patel AS, O'Leary ML, Stein RJ, et al. The relationship between overactive bladder and sexual activity in women. *Int Braz J Urol.* 2006 Jan-Feb;32(1):77-87. X-11
1618. Paterson J. Outcomes achieved by an elderly community dwelling population who undertook conservative continence management programs. *Geriacton.* 2004 Sep;22(3):5-16. X-10
1619. Patki P, Hamid R, Somayaji S, et al. Long-term urological outcomes in paediatric spinal cord injury. *Spinal Cord.* 2006 Dec;44(12):729-33. X-1F, X-4
1620. Patki PS, Hamid R, Arumugam K, et al. Botulinum toxin-type A in the treatment of drug-resistant neurogenic detrusor overactivity secondary to traumatic spinal cord injury. *BJU Int.* 2006 Jul;98(1):77-82. X-1F, X-7
1621. Pauls RN, Marinkovic SP, Silva WA, et al. Effects of sacral neuromodulation on female sexual function. *Int Urogynecol J Pelvic Floor Dysfunct.* 2007 Apr;18(4):391-5. X-1I, X-7
1622. Payne C and Allee T. Goal achievement provides new insights into interstitial cystitis/painful bladder syndrome symptoms and outcomes. *Neurourol Urodyn.* 2009;28(1):13-7. X-1D, X-7
1623. Payne CK, Joyce GF, Wise M, et al. Interstitial cystitis and painful bladder syndrome. *J Urol.* 2007 Jun;177(6):2042-9. X-1D
1624. Payne CK, Mosbaugh PG, Forrest JB, et al. Intravesical resiniferatoxin for the treatment of interstitial cystitis: a randomized, double-blind, placebo controlled trial. *J Urol.* 2005 May;173(5):1590-4. X-1D
1625. Peacock LM, Wiskind AK and Wall LL. Clinical features of urinary incontinence and urogenital prolapse in a black inner-city population. *Am J Obstet Gynecol.* 1994 Dec;171(6):1464-9; discussion 1469-71. X-1H
1626. Peattie AB, Plevnik S and Stanton SL. The use of bladder neck electric conductance (BNEC) in the investigation and management of sensory urge incontinence in the female. *J R Soc Med.* 1988 Aug;81(8):442-4. X-7
1627. Peattie AB and Stanton SL. The Stamey operation for correction of genuine stress incontinence in the elderly woman. *Br J Obstet Gynaecol.* 1989 Aug;96(8):983-6. X-1B
1628. Peeker R, Aldenborg F and Fall M. Nephrogenic adenoma--a study with special reference to clinical presentation. *Br J Urol.* 1997 Oct;80(4):539-42. X-1I
1629. Peeker R, Aldenborg F and Fall M. Complete transurethral resection of ulcers in classic interstitial cystitis. *Int Urogynecol J Pelvic Floor Dysfunct.* 2000;11(5):290-5. X-1D
1630. Peeker R, Haghsheno MA, Holmang S, et al. Intravesical bacillus Calmette-Guerin and dimethyl sulfoxide for treatment of classic and nonulcer interstitial cystitis: a prospective, randomized double-blind study. *J Urol.* 2000 Dec;164(6):1912-5; discussion 1915-6. X-1D, X-7
1631. Pelosi MA, 3rd and Pelosi MA. Stress urinary incontinence. Laparoscopically assisted transvaginal needle suspension. *J Reprod Med.* 1997 Jun;42(6):319-24. X-1B
1632. Pelosi MA, 3rd and Pelosi MA. Pubic bone suburethral stabilization sling: laparoscopic assessment of a transvaginal operation for the treatment of stress urinary incontinence. *J Laparoendosc Adv Surg Tech A.* 1999 Feb;9(1):45-50. X-1B
1633. Pelsang RE and Bonney WW. Voiding cystourethrography in female stress incontinence. *AJR Am J Roentgenol.* 1996 Mar;166(3):561-5. X-1B
1634. Pengelly AW and Booth CM. A prospective trial of bladder training as treatment for detrusor instability. *Br J Urol.* 1980 Dec;52(6):463-6. X-7
1635. Pengelly AW, Stephenson TP, Milroy EJ, et al. Results of prolonged bladder distension as treatment for detrusor instability. *Br J Urol.* 1978 Jun;50(4):243-5. X-7
1636. Peretto I, Forlani R, Fossati C, et al. Discovery of diaryl imidazolidin-2-one derivatives, a novel class of muscarinic M3 selective antagonists (Part 1). *J Med Chem.* 2007 Apr 5;50(7):1571-83. X-1G, X-4
1637. Perez-Medrano A, Brune ME, Buckner SA, et al. Structure-activity studies of novel cyanoguanidine ATP-sensitive potassium channel openers for the treatment of overactive bladder. *J Med Chem.* 2007 Nov 29;50(24):6265-73. X-1G
1638. Persson J, Bergqvist CE and Wolner-Hanssen P. An ultra-short perineal pad-test for evaluation of female stress urinary incontinence treatment. *Neurourol Urodyn.* 2001;20(3):277-85. X-1H
- X-1B , 1639. Persson-Junemann C, Seemann O, Kohrmann KU, et al. Comparison of urodynamic findings and response to oxybutynin in nocturnal enuresis. *Eur Urol.* 1993;24(1):92-6. X-1C, X-4
1640. Pesce F, Castellano V, Finazzi Agro E, et al. Voiding dysfunction in patients with spinal cord lesions at the thoracolumbar vertebral junction. *Spinal Cord.* 1997 Jan;35(1):37-9. X-1F
1641. Peschers UM, Jundt K and Dimpfl T. Differences between cough and Valsalva leak-point pressure in stress incontinent women. *Neurourol Urodyn.* 2000;19(6):677-81. X-1B
1642. Peters D. Terodiline in the treatment of urinary frequency and motor urge incontinence. A controlled multicentre trial. *Scand J Urol Nephrol Suppl.* 1984;87:21-33. X-9
1643. Peters KM, Carey JM and Konstant D. Sacral neuromodulation for the treatment of refractory interstitial cystitis: outcomes based on technique. *Int Urogynecol J Pelvic Floor Dysfunct.* 2003 Oct;14(4):223-8; discussion 228. X-1D
1644. Peters KM, Diokno AC, Steinert BW, et al. The efficacy of intravesical bacillus Calmette-Guerin in the treatment of interstitial cystitis: long-term followup. *J Urol.* 1998 May;159(5):1483-6; discussion 1486-7. X-1D
1645. Peters KM and Konstant D. Sacral neuromodulation decreases narcotic requirements in refractory interstitial cystitis. *BJU Int.* 2004 Apr;93(6):777-9. X-1D
1646. Petersen T, Chandiramani V and Fowler CJ. The ice-water test in detrusor hyper-reflexia and bladder instability. *Br J Urol.* 1997 Feb;79(2):163-7. X-1H
1647. Petersen T and Pedersen E. Neurourodynamic evaluation of voiding dysfunction in multiple sclerosis. *Acta Neurol Scand.* 1984 Jun;69(6):402-11. X-1F
1648. Petros PE and Ulmsten U. Bladder instability in women: a premature activation of the micturition reflex. *Neurourol Urodyn.* 1993;12(3):235-9. X-1H

1649. Petros PE and Ulmsten UI. The combined intravaginal sling and tuck operation. An ambulatory procedure for cure of stress and urge incontinence. *Acta Obstet Gynecol Scand Suppl.* 1990;153:53-9. X-1B
1650. Petros PE and Ulmsten UI. Cure of urge incontinence by the combined intravaginal sling and tuck operation. *Acta Obstet Gynecol Scand Suppl.* 1990;153:61-2. X-7
1651. Petros PE and Ulmsten UI. Non stress non urge female urinary incontinence--diagnosis and cure: a preliminary report. *Acta Obstet Gynecol Scand Suppl.* 1990;153:69-70. X-1I
1652. Petros PP. The intravaginal slingplasty operation, a minimally invasive technique for cure of urinary incontinence in the female. *Aust N Z J Obstet Gynaecol.* 1996 Nov;36(4):453-61. X-7
1653. Petros PP and Ulmsten U. Urge incontinence history is an accurate predictor of urge incontinence. *Acta Obstet Gynecol Scand.* 1992 Oct;71(7):537-9. X-1H
1654. Petrou SP, Brown JA and Blaivas JG. Suprameatal transvaginal urethrolysis. *J Urol.* 1999 Apr;161(4):1268-71. X-7
1655. Petrou SP and Young PR. Rate of recurrent stress urinary incontinence after retropubic urethrolysis. *J Urol.* 2002 Feb;167(2 Pt 1):613-5. X-1B
1656. Pettersen R, Haig Y, Nakstad PH, et al. Subtypes of urinary incontinence after stroke: relation to size and location of cerebrovascular damage. *Age Ageing.* 2008 May;37(3):324-7. X-1I
1657. Pettersen R, Stien R and Wyller TB. Post-stroke urinary incontinence with impaired awareness of the need to void: clinical and urodynamic features. *BJU Int.* 2007 May;99(5):1073-7. X-1F, X-5
1658. Pfister C, Dacher JN, Gaucher S, et al. The usefulness of a minimal urodynamic evaluation and pelvic floor biofeedback in children with chronic voiding dysfunction. *BJU Int.* 1999 Dec;84(9):1054-7. X-4
1659. Pfisterer MH, Griffiths DJ, Rosenberg L, et al. The impact of detrusor overactivity on bladder function in younger and older women. *J Urol.* 2006 May;175(5):1777-83; discussion 1783. X-1G, X-1H
1660. Pfisterer MH, Griffiths DJ, Rosenberg L, et al. Parameters of bladder function in pre-, peri-, and postmenopausal continent women without detrusor overactivity. *Neurourol Urodyn.* 2007;26(3):356-61. X-1I, X-7
1661. Pfisterer MH, Griffiths DJ, Schaefer W, et al. The effect of age on lower urinary tract function: a study in women. *J Am Geriatr Soc.* 2006 Mar;54(3):405-12. X-1H
1662. Philip J, Willmott S and Irwin P. Interstitial cystitis versus detrusor overactivity: a comparative, randomized, controlled study of cystometry using saline and 0.3 M potassium chloride. *J Urol.* 2006 Feb;175(2):566-70; discussion 570-1. X-1H, X-7
1663. Philip J, Willmott S, Owen D, et al. A double-blind, randomized controlled trial of cystometry using saline versus 0.3 M potassium chloride infusion in women with overactive bladder syndrome. *Neurourol Urodyn.* 2007;26(1):110-4. X-1H, X-7
1664. Philp T, Shah PJ and Worth PH. Acupuncture in the treatment of bladder instability. *Br J Urol.* 1988 Jun;61(6):490-3. X-7
1665. Piauxt E, Evans CJ, Espindle D, et al. Development and validation of the Overactive Bladder Satisfaction (OAB-S) Questionnaire. *Neurourol Urodyn.* 2008;27(3):179-90. X-1I
1666. Pidcock FS, Krishnan C, Crawford TO, et al. Acute transverse myelitis in childhood: center-based analysis of 47 cases. *Neurology.* 2007 May 1;68(18):1474-80. X-1F, X-4, X-7
1667. Pieri S, Agresti P, Morucci M, et al. Percutaneous treatment of pelvic congestion syndrome. *Radiol Med (Torino).* 2003 Jan-Feb;105(1-2):76-82. X-1A, X-7
1668. Pikaart DP, Miklos JR and Moore RD. Laparoscopic removal of pubovaginal polypropylene tension-free tape slings. *JSLs.* 2006 Apr-Jun;10(2):220-5. X-1B
1669. Pilsgaard K and Mouritsen L. Follow-up after repair of vaginal vault prolapse with abdominal colposacropexy. *Acta Obstet Gynecol Scand.* 1999 Jan;78(1):66-70. X-1E, X-7
1670. Pleil AM, Coyne KS, Reese PR, et al. The validation of patient-rated global assessments of treatment benefit, satisfaction, and willingness to continue--the BSW. *Value Health.* 2005 Nov-Dec;8 Suppl 1:S25-34. X-1I
1671. Plevnik S and Janez J. Urethral pressure variations. *Urology.* 1983 Feb;21(2):207-9. X-1H
1672. Podesta ML, Castera R and Ruarde AC. Videourodynamic findings in young infants with severe primary reflux. *J Urol.* 2004 Feb;171(2 Pt 1):829-33; discussion 833. X-1I, X-4
1673. Podnar S, Trsinar B and Vodusek DB. Bladder dysfunction in patients with cauda equina lesions. *Neurourol Urodyn.* 2006;25(1):23-31. X-1I
1674. Poggesi A, Pracucci G, Chabriat H, et al. Urinary complaints in nondisabled elderly people with age-related white matter changes: the Leukoaraiosis And Disability (LADIS) Study. *J Am Geriatr Soc.* 2008 Sep;56(9):1638-43. X-1H, X-4
1675. Ponsky LE, Sharma S, Pandrangi L, et al. Screening and monitoring for bladder cancer: refining the use of NMP22. *J Urol.* 2001 Jul;166(1):75-8. X-1A
1676. Poole CJ and Lightman SL. Inhibition of vasopressin secretion during migraine. *J Neurol Neurosurg Psychiatry.* 1988 Nov;51(11):1441-4. X-1G, X-7
1677. Poon CI and Zimmern PE. Is there a role for periurethral collagen injection in the management of urodynamically proven mixed urinary incontinence? *Urology.* 2006 Apr;67(4):725-9; discussion 729-30. X-1B
1678. Popat R, Apostolidis A, Kalsi V, et al. A comparison between the response of patients with idiopathic detrusor overactivity and neurogenic detrusor overactivity to the first intradetrusor injection of botulinum-A toxin. *J Urol.* 2005 Sep;174(3):984-9. X-7
1679. Porru D and Usai E. Standard and extramural ambulatory urodynamic investigation for the diagnosis of detrusor instability-correlated incontinence and micturition disorders. *Neurourol Urodyn.* 1994;13(3):237-42. X-1H
1680. Posey JA, 3rd, Saif MW, Carlisle R, et al. Phase 1 study of weekly polyethylene glycol-camptothecin in patients with advanced solid tumors and lymphomas. *Clin Cancer Res.* 2005 Nov 1;11(21):7866-71. X-1A, X-7
1681. Potts JM, Ward AM and Rackley RR. Association of chronic urinary symptoms in women and Ureaplasma urealyticum. *Urology.* 2000 Apr;55(4):486-9. X-1A, X-1H

1682. Pow-Sang JM, Lockhart JL, Suarez A, et al. Female urinary incontinence: preoperative selection, surgical complications and results. *J Urol*. 1986 Oct;136(4):831-3. X-1B
1683. Prasopsanti K, Santi-Ngamkun A and Pornprasit K. Estimated cost of overactive bladder in Thailand. *J Med Assoc Thai*. 2007 Nov;90(11):2316-20. X-6
1684. Prati A, Cantoni C and Potenzoni D. Use of rociverine in patients with motor urge incontinence: A cystomanometric study of dose-effect correlations. *Current Therapeutic Research, Clinical & Experimental*. 1992;51(6):933-936. X-7
1685. Pregazzi R, Sartore A, Bortoli P, et al. Immediate postpartum perineal examination as a predictor of puerperal pelvic floor dysfunction. *Obstet Gynecol*. 2002 Apr;99(4):581-4. X-1A
1686. Pregazzi R, Sartore A, Troiano L, et al. Postpartum urinary symptoms: prevalence and risk factors. *Eur J Obstet Gynecol Reprod Biol*. 2002 Jul 10;103(2):179-82. X-6
1687. Preminger GM, Kettelhut MC, Elkins SL, et al. Ureteral stenting during extracorporeal shock wave lithotripsy: help or hindrance? *J Urol*. 1989 Jul;142(1):32-6. X-1I
1688. Pressman MR, Figueroa WG, Kendrick-Mohamed J, et al. Nocturia. A rarely recognized symptom of sleep apnea and other occult sleep disorders. *Arch Intern Med*. 1996 Mar 11;156(5):545-50. X-1C
1689. Primus G. Maximal electrical stimulation in neurogenic detrusor hyperactivity: experiences in multiple sclerosis. *Eur J Med*. 1992 May;1(2):80-2. X-1F
1690. Primus G and Kramer G. Maximal external electrical stimulation for treatment of neurogenic or non-neurogenic urgency and/or urge incontinence. *Neurourol Urodyn*. 1996;15(3):187-94. X-7
1691. Primus G and Pummer K. Oxybutynin hydrochloride in the management of detrusor instability. *Int Urol Nephrol*. 1990;22(3):243-8. X-7
1692. Pron G, Bennett J, Common A, et al. The Ontario Uterine Fibroid Embolization Trial. Part 2. Uterine fibroid reduction and symptom relief after uterine artery embolization for fibroids. *Fertil Steril*. 2003 Jan;79(1):120-7. X-1I
1693. Pron G, Cohen M, Soucie J, et al. The Ontario Uterine Fibroid Embolization Trial. Part 1. Baseline patient characteristics, fibroid burden, and impact on life. *Fertil Steril*. 2003 Jan;79(1):112-9. X-1I
1694. Propert KJ, Schaeffer AJ, Brensinger CM, et al. A prospective study of interstitial cystitis: results of longitudinal followup of the interstitial cystitis data base cohort. The Interstitial Cystitis Data Base Study Group. *J Urol*. 2000 May;163(5):1434-9. X-1D
1695. Pryor JL and Jenkins AD. Use of double-pigtail stents in extracorporeal shock wave lithotripsy. *J Urol*. 1990 Mar;143(3):475-8. X-1I
1696. Pryor JL, Langley MJ and Jenkins AD. Comparison of symptom characteristics of indwelling ureteral catheters. *J Urol*. 1991 Apr;145(4):719-22. X-1A, X-7
1697. Quarto G, Autorino R, Gallo A, et al. Quality of life in women with multiple sclerosis and overactive bladder syndrome. *Int Urogynecol J Pelvic Floor Dysfunct*. 2007 Feb;18(2):189-94. X-1I
1698. Quinn M. Obstetric denervation-gynaecological reinnervation: disruption of the inferior hypogastric plexus in childbirth as a source of gynaecological symptoms. *Med Hypotheses*. 2004;63(3):390-3. X-1I, X-2
1699. Quinn P, Goka J and Richardson H. Assessment of an electronic daily diary in patients with overactive bladder. *BJU Int*. 2003 May;91(7):647-52. X-1H
1700. Qvist N, Kristensen ES, Nielsen KK, et al. Detrusor instability in children with recurrent urinary tract infection and/or enuresis. I. Clinical conditions and symptomatology. *Urol Int*. 1986;41(3):196-8. X-1I, X-4
1701. Qvist N, Nielsen KK, Kristensen ES, et al. Detrusor instability in children with recurrent urinary tract infection and/or enuresis. II. Treatment. *Urol Int*. 1986;41(3):199-201. X-1I, X-4
1702. Rabey JM, Moriel EZ, Farkas A, et al. Detrusor hyperreflexia in multiple sclerosis. Alleviation by a combination of imipramine and propantheline, a clinico-laboratory study. *Eur Neurol*. 1979;18(1):33-7. X-1F
1703. Rackley R, Weiss JP, Rovner ES, et al. Nighttime dosing with tolterodine reduces overactive bladder-related nocturnal micturitions in patients with overactive bladder and nocturia. *Urology*. 2006 Apr;67(4):731-6; discussion 736. X-4
1704. Radej M, Hitrec V, Krivec O, et al. Functional and tonometrical investigation of bladder and urethra in patients with stress incontinence. *J Urol*. 1976 May;115(5):551-4. X-1B, X-1H
1705. Radley SC, Chapple CR, Mitsogiannis IC, et al. Transurethral implantation of macropastique for the treatment of female stress urinary incontinence secondary to urethral sphincter deficiency. *Eur Urol*. 2001 Apr;39(4):383-9. X-1B
1706. Radley SC, Rosario DJ, Chapple CR, et al. Conventional and ambulatory urodynamic findings in women with symptoms suggestive of bladder overactivity. *J Urol*. 2001 Dec;166(6):2253-8. X-1H
1707. Radomski SB, Caley B, Reiz JL, et al. Preliminary evaluation of a new controlled-release oxybutynin in urinary incontinence. *Curr Med Res Opin*. 2004;20(2):249-53. X-7
1708. Radomski SB and Herschorn S. Laparoscopic Burch bladder neck suspension: early results. *J Urol*. 1996 Feb;155(2):515-8. X-1B
1709. Raes A, Hoebeke P, Segaert I, et al. Retrospective analysis of efficacy and tolerability of tolterodine in children with overactive bladder. *Eur Urol*. 2004 Feb;45(2):240-4. X-4
1710. Rahmanou P, Chaliha C, Kulinskaya E, et al. Reliability testing of urodynamics, pressure flow studies and cough leak point pressure in women with urodynamic stress incontinence with and without detrusor overactivity. *Int Urogynecol J Pelvic Floor Dysfunct*. 2008 Jul;19(7):933-8. X-1H, X-7
1711. Rajan L. The impact of obstetric procedures and analgesia/anaesthesia during labour and delivery on breast feeding. *Midwifery*. 1994 Jun;10(2):87-103. X-1A

1712. Rajkumar GN, Small DR, Mustafa AW, et al. A prospective study to evaluate the safety, tolerability, efficacy and durability of response of intravesical injection of botulinum toxin type A into detrusor muscle in patients with refractory idiopathic detrusor overactivity. *BJU Int*. 2005 Oct;96(6):848-52. X-7
1713. Ramahi AJ and Richardson DA. A practical approach to the painful bladder syndrome. *J Reprod Med*. 1990 Aug;35(8):805-9. X-1D
1714. Ramamurthy HR and Kanitkar M. Recurrent urinary tract infection and functional voiding disorders. *Indian Pediatr*. 2008 Aug;45(8):689-91. X-II
1715. Ramon J, Mekras J and Webster GD. Transvaginal needle suspension procedures for recurrent stress incontinence. *Urology*. 1991 Dec;38(6):519-22. X-1B
1716. Ramsay IN, Clancy S and Hilton P. Subtrigonal phenol injections in the treatment of idiopathic detrusor instability in the female--a long-term urodynamic follow-up. *Br J Urol*. 1992 Apr;69(4):363-5. X-7
1717. Ramsay-Olocco K, Alexandrova L, Nellore R, et al. Pre-clinical and clinical evaluation of solution and soft gelatin capsule formulations for a BCS class 3 compound with atypical physicochemical properties. *J Pharm Sci*. 2004 Sep;93(9):2214-21. X-1G, X-2
1718. Ramsden PD, Smith JC, Dunn M, et al. Distension therapy for the unstable bladder: later results including an assessment of repeat distensions. *Br J Urol*. 1976;48(7):623-9. X-7
1719. Raney B, Jr., Heyn R, Hays DM, et al. Sequelae of treatment in 109 patients followed for 5 to 15 years after diagnosis of sarcoma of the bladder and prostate. A report from the Intergroup Rhabdomyosarcoma Study Committee. *Cancer*. 1993 Apr 1;71(7):2387-94. X-1A
1720. Ransmayr GN, Holliger S, Schletterer K, et al. Lower urinary tract symptoms in dementia with Lewy bodies, Parkinson disease, and Alzheimer disease. *Neurology*. 2008 Jan 22;70(4):299-303. X-1F
1721. Rapidi CA, Petropoulou K, Galata A, et al. Neuropathic bladder dysfunction in patients with motor complete and sensory incomplete spinal cord lesion. *Spinal Cord*. 2008 Oct;46(10):673-8. X-1F
1722. Rapp DE, Lucioni A, Katz EE, et al. Use of botulinum-A toxin for the treatment of refractory overactive bladder symptoms: an initial experience. *Urology*. 2004 Jun;63(6):1071-5. X-7
1723. Rardin CR, Rosenblatt PL, Kohli N, et al. Release of tension-free vaginal tape for the treatment of refractory postoperative voiding dysfunction. *Obstet Gynecol*. 2002 Nov;100(5 Pt 1):898-902. X-1B
1724. Rasmussen KL, Krue S, Johansson LE, et al. Obesity as a predictor of postpartum urinary symptoms. *Acta Obstet Gynecol Scand*. 1997 Apr;76(4):359-62. X-1A
1725. Rasuli P, Hammond I, Al-Mutairi B, et al. Spherical versus conventional polyvinyl alcohol particles for uterine artery embolization. *J Vasc Interv Radiol*. 2008 Jan;19(1):42-6. X-II
1726. Raza-Khan F, Graziano S, Kenton K, et al. Peripartum urinary incontinence in a racially diverse obstetrical population. *Int Urogynecol J Pelvic Floor Dysfunct*. 2006 Sep;17(5):525-30. X-1A
1727. Reese PR, Pleil AM, Okano GJ, et al. Multinational study of reliability and validity of the King's Health Questionnaire in patients with overactive bladder. *Qual Life Res*. 2003 Jun;12(4):427-42. X-II
1728. Reeves P, Irwin D, Kelleher C, et al. The current and future burden and cost of overactive bladder in five European countries. *Eur Urol*. 2006 Nov;50(5):1050-7. X-6
1729. Reid GC, DeLancey JO, Hopkins MP, et al. Urinary incontinence following radical vulvectomy. *Obstet Gynecol*. 1990 May;75(5):852-8. X-1A
1730. Reiertsen O and Kulseng-Hanssen S. Simultaneous urethrocytometry in gynaecological patients with lower urinary tract dysfunction. *Ann Chir Gynaecol*. 1983;72(5):278-81. X-IH
1731. Reinberg Y, Crocker J, Wolpert J, et al. Therapeutic efficacy of extended release oxybutynin chloride, and immediate release and long acting tolterodine tartrate in children with diurnal urinary incontinence. *J Urol*. 2003 Jan;169(1):317-9. X-II, X-4
1732. Reitz A, Denys P, Fermanian C, et al. Do repeat intradetrusor botulinum toxin type a injections yield valuable results? Clinical and urodynamic results after five injections in patients with neurogenic detrusor overactivity. *Eur Urol*. 2007 Dec;52(6):1729-35. X-1F, X-7
1733. Reitz A, Stohrer M, Kramer G, et al. European experience of 200 cases treated with botulinum-A toxin injections into the detrusor muscle for urinary incontinence due to neurogenic detrusor overactivity. *Eur Urol*. 2004 Apr;45(4):510-5. X-1F
1734. Resnick NM, Yalla SV and Laurino E. The pathophysiology of urinary incontinence among institutionalized elderly persons. *N Engl J Med*. 1989 Jan 5;320(1):1-7. X-1H, X-5
1735. Rezapour M and Ulmsten U. Tension-Free vaginal tape (TVT) in women with mixed urinary incontinence--a long-term follow-up. *Int Urogynecol J Pelvic Floor Dysfunct*. 2001;12 Suppl 2:S15-18. X-1B
1736. Ricci JA, Baggish JS, Hunt TL, et al. Coping strategies and health care-seeking behavior in a US national sample of adults with symptoms suggestive of overactive bladder. *Clin Ther*. 2001 Aug;23(8):1245-59. X-II
1737. Richmond D. The incontinent woman: 2. *Br J Hosp Med*. 1993 Oct 20-Nov 2;50(8):490-2. X-II, X-2, X-6
1738. Richter HE, Diokno A, Kenton K, et al. Predictors of treatment failure 24 months after surgery for stress urinary incontinence. *J Urol*. 2008 Mar;179(3):1024-30. X-1B
1739. Richter HE, Goode PS, Brubaker L, et al. Two-year outcomes after surgery for stress urinary incontinence in older compared with younger women. *Obstet Gynecol*. 2008 Sep;112(3):621-9. X-1A
1740. Richter HE, Varner RE, Sanders E, et al. Effects of pubovaginal sling procedure on patients with urethral hypermobility and intrinsic sphincteric deficiency: would they do it again? *Am J Obstet Gynecol*. 2001 Jan;184(2):14-9. X-1B
1741. Ridley DM. Urinary urge incontinence--selection of patients for surgery. *S Afr Med J*. 1988 May 7;73(9):537-9. X-7
1742. Rigby D. Promoting continence with electrostimulation. *Prof Nurse*. 1996 Apr;11(7):431-434. X-2

1743. Riss PA and Koelbl H. The predictive values of single and combined urodynamic parameters. *Acta Obstet Gynecol Scand.* 1988;67(6):505-11. X-1H
1744. Ritchie RW, Lindfield DM, Lockyer CR, et al. Ocular side-effects of urological pharmacy. *BJU Int.* 2008 Jun;101(11):1336-8. X-1I
1745. Riva D and Casolati E. Oxybutynin chloride in the treatment of female idiopathic bladder instability. Results from double blind treatment. *Clin Exp Obstet Gynecol.* 1984;11(1-2):37-42. X-7
1746. Robb-Nicholson C. By the way, doctor. I'm 80 and suffer from an overactive bladder that hasn't responded to the usual treatments. My urologist is recommending an Interstim neuromodulator. What are the pros and cons of this therapy? *Harv Womens Health Watch.* 2008 Nov;16(3):8. X-2
1747. Robertson AS, Davies JB, Webb RJ, et al. Bladder augmentation and replacement. Urodynamic and clinical review of 25 patients. *Br J Urol.* 1991 Dec;68(6):590-7. X-1A, X-7
1748. Robertson AS, Griffiths C and Neal DE. Conventional urodynamics and ambulatory monitoring in the definition and management of bladder outflow obstruction. *J Urol.* 1996 Feb;155(2):506-11. X-1A X-1H, X-4
1749. Robertson AS, Griffiths CJ, Ramsden PD, et al. Bladder function in healthy volunteers: ambulatory monitoring and conventional urodynamic studies. *Br J Urol.* 1994 Mar;73(3):242-9. X-1H, X-7
1750. Robinson D, Anders K, Cardozo L, et al. Outcome measures in urogynaecology: the clinicians' perspective. *Int Urogynecol J Pelvic Floor Dysfunct.* 2007 Mar;18(3):273-9. X-1H
1751. Robinson D, Anders K, Cardozo L, et al. Can ultrasound replace ambulatory urodynamics when investigating women with irritative urinary symptoms? *BJOG.* 2002 Feb;109(2):145-8. X-1H
1752. Robinson JM and Brocklehurst JC. Emepronium bromide and flavoxate hydrochloride in the treatment of urinary incontinence associated with detrusor instability in elderly women. *Br J Urol.* 1983 Aug;55(4):371-6. X-7
1753. Robson WL and Leung AK. Extraordinary urinary frequency syndrome. *Urology.* 1993 Sep;42(3):321-4. X-4, X-7
1754. Rodhe N, Englund L, Molstad S, et al. Bacteriuria is associated with urge urinary incontinence in older women. *Scand J Prim Health Care.* 2008 Mar;26(1):35-9. X-1I
1755. Rodrigues P, Hering F, Meler A, et al. Pubo-fascial versus vaginal sling operation for the treatment of stress urinary incontinence: a prospective study. *Neurourol Urodyn.* 2004;23(7):627-31. X-1B
1756. Rodriguez LV, Berman J and Raz S. Polypropylene sling for treatment of stress urinary incontinence: an alternative to tension-free vaginal tape. *Tech Urol.* 2001 Jun;7(2):87-9. X-1B
1757. Rodriguez LV, Bukkapatnam R, Shah SM, et al. Transvaginal paravaginal repair of high-grade cystocele central and lateral defects with concomitant suburethral sling: report of early results, outcomes, and patient satisfaction with a new technique. *Urology.* 2005 Nov;66(5 Suppl):57-65. X-1E
1758. Rodriguez LV, de Almeida F, Dorey F, et al. Does Valsalva leak point pressure predict outcome after the distal urethral polypropylene sling? Role of urodynamics in the sling era. *J Urol.* 2004 Jul;172(1):210-4. X-1B
1759. Rohr G, Kragstrup J, Gaist D, et al. Genetic and environmental influences on urinary incontinence: a Danish population-based twin study of middle-aged and elderly women. *Acta Obstet Gynecol Scand.* 2004 Oct;83(10):978-82. X-6
1760. Romancik M, Lutter I, Goncalves F, et al. Valsalva leak point pressure predicts outcome after transobturator suburethral tape implantation--fact or fiction? *Bratisl Lek Listy.* 2006;107(11-12):426-9. X-1B, X-1I
1761. Romanzi LJ, Chaikin DC and Blaivas JG. The effect of genital prolapse on voiding. *J Urol.* 1999 Feb;161(2):581-6. X-1E, X-1H
1762. Romanzi LJ, Groutz A, Heritz DM, et al. Involuntary detrusor contractions: correlation of urodynamic data to clinical categories. *Neurourol Urodyn.* 2001;20(3):249-57. X-1F, X-1H
1763. Rombis V, Triantafyllidis A, Balaxis E, et al. Nocturnal enuresis in children. A four-year experience in outpatient clinics of pediatric urology. *Folia Med (Plovdiv).* 2005;47(2):24-8. X-1I, X-4
1764. Roongruangsilp U, Lertsithichai P, Kochakarn W, et al. Correlation between symptoms and urodynamic findings in Thai female patients with urinary incontinence. *J Med Assoc Thai.* 2005 Mar;88(3):364-70. X-1H
1765. Roovers JP, van der Bom JG, Huub van der Vaart C, et al. Does mode of hysterectomy influence micturition and defecation? *Acta Obstet Gynecol Scand.* 2001 Oct;80(10):945-51. X-1A
1766. Roovers JP, van der Vaart CH, van der Bom JG, et al. A randomised controlled trial comparing abdominal and vaginal prolapse surgery: effects on urogenital function. *BJOG.* 2004 Jan;111(1):50-6. X-1E
1767. Roovers JP, van Laar JO, Loffeld C, et al. Does urodynamic investigation improve outcome in patients undergoing prolapse surgery? *Neurourol Urodyn.* 2007;26(2):170-5. X-1A, X-7
1768. Rortveit G, Daltveit AK, Hannestad YS, et al. Vaginal delivery parameters and urinary incontinence: the Norwegian EPINCONT study. *Am J Obstet Gynecol.* 2003 Nov;189(5):1268-74. X-1I
1769. Rortveit G, Hannestad YS, Daltveit AK, et al. Age- and type-dependent effects of parity on urinary incontinence: the Norwegian EPINCONT study. *Obstet Gynecol.* 2001 Dec;98(6):1004-10. X-6E
1770. Rosario DJ, Smith DJ, Radley SC, et al. Pharmacodynamics of anticholinergic agents measured by ambulatory urodynamic monitoring: a study of methodology. *Neurourol Urodyn.* 1999;18(3):223-33; discussion 223-4. X-1H, X-7
1771. Rose AE, Payne CK and Azevedo K. Pilot study of the feasibility of in-office bladder distention using electromotive drug administration (EMDA). *Neurourol Urodyn.* 2005;24(3):254-60. X-1D
1772. Rosen LS, Brown J, Laxa B, et al. Phase I study of TLK286 (glutathione S-transferase P1-1 activated glutathione analogue) in advanced refractory solid malignancies. *Clin Cancer Res.* 2003 May;9(5):1628-38. X-1A

1773. Rosen RC, Coyne KS, Henry D, et al. Beyond the cluster: methodological and clinical implications in the Boston Area Community Health survey and EPIC studies. *BJU Int.* 2008 May;101(10):1274-8. X-II
1774. Rosenberg LJ, Griffiths DJ and Resnick NM. Factors that distinguish continent from incontinent older adults with detrusor overactivity. *J Urol.* 2005 Nov;174(5):1868-72. X-6
1775. Rosenberg MJ, Roachat RW, Akbar J, et al. Sterilization in Bangladesh: mortality, morbidity, and risk factors. *Int J Gynaecol Obstet.* 1982 Aug;20(4):283-91. X-II
1776. Rosenblum N, Scarpero HM and Nitti VW. Voiding dysfunction in young, nulliparous women: symptoms and urodynamic findings. *Int Urogynecol J Pelvic Floor Dysfunct.* 2004 Nov-Dec;15(6):373-7; discussion 377. X-II
1777. Ross JW. Techniques of laparoscopic repair of total vault eversion after hysterectomy. *J Am Assoc Gynecol Laparosc.* 1997 Feb;4(2):173-83. X-II
1778. Ross JW. Laparoscopic Burch colposuspension and overlapping sphincteroplasty for double incontinence. *JSL.S.* 2001 Jul-Sep;5(3):203-9. X-1B
1779. Rosser CJ, Slaton JW, Izawa JI, et al. Clinical presentation and outcome of high-grade urinary bladder leiomyosarcoma in adults. *Urology.* 2003 Jun;61(6):1151-5. X-II
1780. Rost A, Fiedler U and Fester C. Comparative analysis of the results of suspension-urethroplasty according to Marshall-Marchetti-Krantz and of urethrovesicopexy with adhesive. *Urol Int.* 1979;34(3):167-75. X-1B
1781. Rotman M, Aziz H, Porrazzo M, et al. Treatment of advanced transitional cell carcinoma of the bladder with irradiation and concomitant 5-fluorouracil infusion. *Int J Radiat Oncol Biol Phys.* 1990 May;18(5):1131-7. X-II
1782. Roupret M, Chartier-Kastler E, Almeras C, et al. Sacral neuromodulation for refractory detrusor overactivity in women with an artificial urinary sphincter. *J Urol.* 2004 Jul;172(1):236-9. X-1A, X-7
1783. Rovner ES, Rackley R, Nitti VW, et al. Tolterodine extended release is efficacious in continent and incontinent subjects with overactive bladder. *Urology.* 2008 Sep;72(3):488-93. X-4
1784. Ruarte AC, Podesta ML and Medel R. Detrusor after-contractions in children with normal urinary tracts. *BJU Int.* 2002 Aug;90(3):286-93. X-II, X-4
1785. Ruffion A, Capelle O, Paparel P, et al. What is the optimum dose of type A botulinum toxin for treating neurogenic bladder overactivity? *BJU Int.* 2006 May;97(5):1030-4. X-1F
1786. Rufford J and Cardozo L. Urethral diverticula: a diagnostic dilemma. *BJU Int.* 2004 Nov;94(7):1044-7. X-II
1787. Rufford J, Hextall A, Cardozo L, et al. A double-blind placebo-controlled trial on the effects of 25 mg estradiol implants on the urge syndrome in postmenopausal women. *Int Urogynecol J Pelvic Floor Dysfunct.* 2003 Jun;14(2):78-83. X-7
1788. Ruud Bosch JL and Groen J. Treatment of refractory urge urinary incontinence with sacral spinal nerve stimulation in multiple sclerosis patients. *Lancet.* 1996 Sep 14;348(9029):717-9. X-1F
1789. Sackley CM, Rodriguez NA, van den Berg M, et al. A phase II exploratory cluster randomized controlled trial of a group mobility training and staff education intervention to promote urinary continence in UK care homes. *Clin Rehabil.* 2008 Aug;22(8):714-21. X-II, X-7
1790. Safarinejad MR and Hosseini SY. Safety and efficacy of tramadol in the treatment of idiopathic detrusor overactivity: a double-blind, placebo-controlled, randomized study. *Br J Clin Pharmacol.* 2006 Apr;61(4):456-63. X-7
1791. Safir MH, Gousse AE, Rovner ES, et al. 4-Defect repair of grade 4 cystocele. *J Urol.* 1999 Feb;161(2):587-94. X-1B
1792. Sahai A, Khan MS and Dasgupta P. Efficacy of botulinum toxin-A for treating idiopathic detrusor overactivity: results from a single center, randomized, double-blind, placebo controlled trial. *J Urol.* 2007 Jun;177(6):2231-6. X-7
1793. Sahai A, Khan MS, Le Gall N, et al. Urodynamic assessment of poor responders after botulinum toxin-A treatment for overactive bladder. *Urology.* 2008 Mar;71(3):455-9. X-II, X-7
1794. Saini J, Axelrod FB, Maayan C, et al. Urinary incontinence in familial dysautonomia. *Int Urogynecol J Pelvic Floor Dysfunct.* 2003 Aug;14(3):209-13; discussion 213. X-1F
1795. Saito M, Kawatani M, Kinoshita Y, et al. Effectiveness of an anti-inflammatory drug, loxoprofen, for patients with nocturia. *Int J Urol.* 2005 Aug;12(8):779-82. X-7
1796. Saito M, Kondo A, Kato K, et al. Bladder dysfunction due to human T-lymphotropic virus type I associated myelopathy. *Br J Urol.* 1991 Oct;68(4):365-8. X-1H, X-II, X-7
1797. Saito M, Kondo A, Kato T, et al. Frequency-volume charts: comparison of frequency between elderly and adult patients. *Br J Urol.* 1993 Jul;72(1):38-41. X-1H
1798. Saito M, Tabuchi F, Otsubo K, et al. Treatment of overactive bladder with modified intravesical oxybutynin chloride. *Neurourol Urodyn.* 2000;19(6):683-8. X-7
1799. Saito M, Watanabe T, Tabuchi F, et al. Urodynamic effects and safety of modified intravesical oxybutynin chloride in patients with neurogenic detrusor overactivity: 3 years experience. *Int J Urol.* 2004 Aug;11(8):592-6. X-1F
1800. Sakakibara R, Hattori T, Boku K, et al. Micturitional disturbance in neuro-Behcet's syndrome. *Auton Neurosci.* 2000 Sep 1;83(1-2):86-9. X-1F
1801. Sakakibara R, Hattori T, Kita K, et al. Stress induced urinary incontinence in patients with spinocerebellar degeneration. *J Neurol Neurosurg Psychiatry.* 1998 Mar;64(3):389-91. X-1B
1802. Sakakibara R, Hattori T, Kuwabara S, et al. Micturitional disturbance in patients with Guillain-Barre syndrome. *J Neurol Neurosurg Psychiatry.* 1997 Nov;63(5):649-53. X-1F
1803. Sakakibara R, Hattori T, Kuwabara S, et al. Micturitional disturbance in patients with chronic inflammatory demyelinating polyneuropathy. *Neurology.* 1998 Apr;50(4):1179-82. X-1F
1804. Sakakibara R, Hattori T, Uchiyama T, et al. Urinary dysfunction in Brown-Sequard syndrome. *Neurourol Urodyn.* 2001;20(6):661-7. X-1F

1805. Sakakibara R, Hattori T, Uchiyama T, et al. Neurogenic failures of the external urethral sphincter closure and relaxation; a videourodynamic study. *Auton Neurosci*. 2001 Jan 14;86(3):208-15. X-1F
1806. Sakakibara R, Hattori T, Yasuda K, et al. Micturition disturbance in acute transverse myelitis. *Spinal Cord*. 1996 Aug;34(8):481-5. X-1F
1807. Sakakibara R, Hattori T, Yasuda K, et al. Micturitional disturbance after acute hemispheric stroke: analysis of the lesion site by CT and MRI. *J Neurol Sci*. 1996 Apr;137(1):47-56. X-1F
1808. Sakakibara R, Hattori T, Yasuda K, et al. Micturitional disturbance and the pontine tegmental lesion: urodynamic and MRI analyses of vascular cases. *J Neurol Sci*. 1996 Sep 15;141(1-2):105-10. X-1F
1809. Sakakibara R, Hattori T, Yasuda K, et al. Micturitional disturbance in acute disseminated encephalomyelitis (ADEM). *J Auton Nerv Syst*. 1996 Sep 12;60(3):200-5. X-1F
1810. Sakakibara R, Hattori T, Yasuda K, et al. Micturitional disturbance in syringomyelia. *J Neurol Sci*. 1996 Nov;143(1-2):100-6. X-1F
1811. Sakakibara R, Ito T, Uchiyama T, et al. Lower urinary tract function in dementia of Lewy body type. *J Neurol Neurosurg Psychiatry*. 2005 May;76(5):729-32. X-1F
1812. Sakakibara R, Ito T, Uchiyama T, et al. Effects of milnacipran and paroxetine on overactive bladder due to neurologic diseases: a urodynamic assessment. *Urol Int*. 2008;81(3):335-9. X-1F, X-7
1813. Sakakibara R, Kanda T, Sekido T, et al. Mechanism of bladder dysfunction in idiopathic normal pressure hydrocephalus. *Neurourol Urodyn*. 2008;27(6):507-10. X-1F, X-7
1814. Sakakibara R, Shinotoh H, Uchiyama T, et al. Questionnaire-based assessment of pelvic organ dysfunction in Parkinson's disease. *Auton Neurosci*. 2001 Sep 17;92(1-2):76-85. X-1F
1815. Sakakibara R, Uchiyama T, Arai K, et al. Lower urinary tract dysfunction in Machado-Joseph disease: a study of 11 clinical-urodynamic observations. *J Neurol Sci*. 2004 Mar 15;218(1-2):67-72. X-1F
1816. Sakakibara R, Uchiyama T, Awa Y, et al. Psychogenic urinary dysfunction: a uro-neurological assessment. *Neurourol Urodyn*. 2007;26(4):518-24. X-1H, X-7
1817. Sakakibara R, Uchiyama T, Liu Z, et al. Nocturnal polyuria with abnormal circadian rhythm of plasma arginine vasopressin in post-stroke patients. *Intern Med*. 2005 Apr;44(4):281-4. X-II, X-7
1818. Sakakibara R, Uchiyama T, Yamanishi T, et al. Urinary function in patients with corticobasal degeneration; comparison with normal subjects. *Neurourol Urodyn*. 2004;23(2):154-8. X-1F
1819. Sakakibara R, Uchiyama T, Yoshiyama M, et al. Preliminary communication: urodynamic assessment of donepezil hydrochloride in patients with Alzheimer's disease. *Neurourol Urodyn*. 2005;24(3):273-5. X-1F
1820. Sale PG and Wyman JF. Achievement of goals associated with bladder training by older incontinent women. *Appl Nurs Res*. 1994 May;7(2):93-6. X-4
1821. Salonia A, Zanni G, Nappi RE, et al. Sexual dysfunction is common in women with lower urinary tract symptoms and urinary incontinence: results of a cross-sectional study. *Eur Urol*. 2004 May;45(5):642-8; discussion 648. X-1A
1822. Salvatore S, Khullar V, Cardozo L, et al. Evaluating ambulatory urodynamics: a prospective study in asymptomatic women. *BJOG*. 2001 Jan;108(1):107-11. X-1H
1823. Sammour ZM, Gomes CM, Duarte RJ, et al. Voiding dysfunction and the Williams-Beuren syndrome: a clinical and urodynamic investigation. *J Urol*. 2006 Apr;175(4):1472-6. X-1F
1824. Samuel M, Boddy SA and Wang K. What happens to the bladder at night? Overnight urodynamic monitoring in children with neurogenic vesical dysfunction. *J Urol*. 2001 Jun;165(6 Pt 2):2335-40. X-1F, X-4
1825. Sand PK. Pelvic floor stimulation in the treatment of mixed incontinence complicated by a low-pressure urethra. *Obstet Gynecol*. 1996 Nov;88(5):757-60. X-1B, X-7
1826. Sand PK and Appell R. Disruptive effects of overactive bladder and urge urinary incontinence in younger women. *Am J Med*. 2006 Mar;119(3 Suppl 1):16-23. X-2
1827. Sand PK, Bowen LW and Ostergard DR. The prognostic significance of augmentation of urethral closure pressure and functional length. *Int J Gynaecol Obstet*. 1990 Oct;33(2):135-9. X-1B
1828. Sand PK, Brubaker LT and Novak T. Simple standing incremental cystometry as a screening method for detrusor instability. *Obstet Gynecol*. 1991 Mar;77(3):453-7. X-1H
1829. Sand PK, Hill RC and Ostergard DR. Supine urethroscopic and standing cystometry as screening methods for the detection of detrusor instability. *Obstet Gynecol*. 1987 Jul;70(1):57-60. X-1H
1830. Sand PK, Hill RC and Ostergard DR. Incontinence history as a predictor of detrusor stability. *Obstet Gynecol*. 1988 Feb;71(2):257-60. X-1H
1831. Sandvik H, Hunskaar S, Vanvik A, et al. Diagnostic classification of female urinary incontinence: an epidemiological survey corrected for validity. *J Clin Epidemiol*. 1995 Mar;48(3):339-43. X-1H
1832. Sandvik H, Kveine E and Hunskaar S. Female urinary incontinence--psychosocial impact, self care, and consultations. *Scand J Caring Sci*. 1993;7(1):53-6. X-II
1833. Sangthawan D, Wathanaarpornchai S and Phunggrassami T. Randomized double blind, placebo-controlled study of pilocarpine administered during head and neck irradiation to reduce xerostomia. *J Med Assoc Thai*. 2001 Feb;84(2):195-203. X-II
1834. Satoh A, Hanawa Y and Nakamura S. Clinical study of bladder cancer: Proteinuria as a predictor of recurrence and efficacy of intravesical bacille Calmette-Guerin therapy. *Int J Urol*. 2004 Jul;11(7):476-82. X-II
1835. Sau G, Siracusano S, Aiello I, et al. The usefulness of the somatosensory evoked potentials of the pudendal nerve in diagnosis of probable multiple sclerosis. *Spinal Cord*. 1999 Apr;37(4):258-63. X-1F, X-1H

1836. Sau GF, Aiello I, Siracusano S, et al. Pudendal nerve somatosensory evoked potentials in probable multiple sclerosis. *Ital J Neurol Sci.* 1997 Oct;18(5):289-91. X-1F, X-1H
1837. Saw AH. Management of urinary tract infection and vesico-ureteric reflux in children. *Singapore Med J.* 1990 Jun;31(3):266-8. X-1I, X-4
1838. Scarpero HM, Dmochowski RR and Nitti VW. Repeat urethrolysis after failed urethrolysis for iatrogenic obstruction. *J Urol.* 2003 Mar;169(3):1013-6. X-1A
1839. Schaer GN, Koechli OR, Schuessler B, et al. Improvement of perineal sonographic bladder neck imaging with ultrasound contrast medium. *Obstet Gynecol.* 1995 Dec;86(6):950-4. X-1B, X-7
1840. Schaer GN, Koechli OR, Schuessler B, et al. Usefulness of ultrasound contrast medium in perineal sonography for visualization of bladder neck funneling--first observations. *Urology.* 1996 Mar;47(3):452-3. X-1H, X-1B
1841. Schaffer JI, Bloom SL, Casey BM, et al. A randomized trial of the effects of coached vs uncoached maternal pushing during the second stage of labor on postpartum pelvic floor structure and function. *Am J Obstet Gynecol.* 2005 May;192(5):1692-6. X-1I
1842. Scheepens WA, de Bie RA, Weil EH, et al. Unilateral versus bilateral sacral neuromodulation in patients with chronic voiding dysfunction. *J Urol.* 2002 Nov;168(5):2046-50. X-7
1843. Scheepens WA, Jongen MM, Nieman FH, et al. Predictive factors for sacral neuromodulation in chronic lower urinary tract dysfunction. *Urology.* 2002 Oct;60(4):598-602. X-1F
1844. Scheepens WA, Van Koevinge GA, De Bie RA, et al. Long-term efficacy and safety results of the two-stage implantation technique in sacral neuromodulation. *BJU Int.* 2002 Dec;90(9):840-5. X-7
1845. Schewe J, Brands FH and Pannek J. Voiding dysfunction in children: role of urodynamic studies. *Urol Int.* 2002;69(4):297-301. X-1I, X-4
1846. Schimpf MO, O'Sullivan DM, LaSala CA, et al. Anterior vaginal wall prolapse and voiding dysfunction in urogynecology patients. *Int Urogynecol J Pelvic Floor Dysfunct.* 2007 Jul;18(7):721-5. X-1E
1847. Schober JM, Lipman R, Haltigan JD, et al. The impact of monosymptomatic nocturnal enuresis on attachment parameters. *Scand J Urol Nephrol.* 2004;38(1):47-52. X-1C, X-4, X-7
1848. Schofield EC, Clausen JA, Burcher E, et al. GAP-43 immunoreactivity of subepithelial and detrusor muscle nerve fibres in patients with refractory idiopathic detrusor overactivity. *Neurourol Urodyn.* 2005;24(4):325-33. X-1G, X-7
1849. Scholtmeijer RJ. Treatment of vesicoureteric reflux: results after 3 years in a prospective study. *Child Nephrol Urol.* 1991;11(1):29-32. X-1I, X-4
1850. Scholtmeijer RJ. Treatment of vesicoureteric reflux. Results of a prospective study. *Br J Urol.* 1993 Mar;71(3):346-9. X-1I, X-4
1851. Scholtmeijer RJ and Griffiths DJ. Treatment of vesicoureteric reflux. Preliminary report of a prospective study. *Br J Urol.* 1988 Mar;61(3):205-9. X-1I, X-4
1852. Scholtmeijer RJ and van Mastrigt R. The effect of oxyphenonium bromide and oxybutynin hydrochloride on detrusor contractility and reflux in children with vesicoureteral reflux and detrusor instability. *J Urol.* 1991 Aug;146(2 (Pt 2)):660-2. X-4
1853. Schostak M, Miller K, Muller M, et al. Transvaginal bone anchors in female stress urinary incontinence: poor results. *Gynecol Obstet Invest.* 2002;54(3):154-8. X-1B
1854. Schrepferman CG, Griebeling TL, Nygaard IE, et al. Resolution of urge symptoms following sling cystourethropexy. *J Urol.* 2000 Nov;164(5):1628-31. X-1B
1855. Schulman SL, Zderic S and Kaplan P. Increased prevalence of urinary symptoms and voiding dysfunction in Williams syndrome. *J Pediatr.* 1996 Sep;129(3):466-9. X-1I, X-7
1856. Schulte-Baukloh H, Knispel HH, Stolze T, et al. Repeated botulinum-A toxin injections in treatment of children with neurogenic detrusor overactivity. *Urology.* 2005 Oct;66(4):865-70; discussion 870. X-1F, X-4
1857. Schulte-Baukloh H, Schobert J, Stolze T, et al. Efficacy of botulinum-A toxin bladder injections for the treatment of neurogenic detrusor overactivity in multiple sclerosis patients: an objective and subjective analysis. *Neurourol Urodyn.* 2006;25(2):110-5. X-1F, X-7
1858. Schulte-Baukloh H, Weiss C, Stolze T, et al. Botulinum-A toxin detrusor and sphincter injection in treatment of overactive bladder syndrome: objective outcome and patient satisfaction. *Eur Urol.* 2005 Dec;48(6):984-90; discussion 990. X-7
1859. Schulte-Baukloh H, Weiss C, Stolze T, et al. Botulinum-A toxin for treatment of overactive bladder without detrusor overactivity: urodynamic outcome and patient satisfaction. *Urology.* 2005 Jul;66(1):82-7. X-7
1860. Schultheiss D, Hofner K, Oelke M, et al. Does bone anchor fixation improve the outcome of percutaneous bladder neck suspension in female stress urinary incontinence? *Br J Urol.* 1998 Aug;82(2):192-5. X-1B, X-7
1861. Schurch B, de Seze M, Denys P, et al. Botulinum toxin type a is a safe and effective treatment for neurogenic urinary incontinence: results of a single treatment, randomized, placebo controlled 6-month study. *J Urol.* 2005 Jul;174(1):196-200. X-1F
1862. Schurch B, Denys P, Kozma CM, et al. Reliability and validity of the Incontinence Quality of Life questionnaire in patients with neurogenic urinary incontinence. *Arch Phys Med Rehabil.* 2007 May;88(5):646-52. X-1F
1863. Schurch B, Denys P, Kozma CM, et al. Botulinum toxin A improves the quality of life of patients with neurogenic urinary incontinence. *Eur Urol.* 2007 Sep;52(3):850-8. X-1F
1864. Schurch B, Reitz A and Tenti G. Electromotive drug administration of lidocaine to anesthetize the bladder before botulinum-A toxin injections into the detrusor. *Spinal Cord.* 2004 Jun;42(6):338-41. X-1F, X-7
1865. Schurch B, Stohrer M, Kramer G, et al. Botulinum-A toxin for treating detrusor hyperreflexia in spinal cord injured patients: a new alternative to anticholinergic drugs? Preliminary results. *J Urol.* 2000 Sep;164(3 Pt 1):692-7. X-1F, X-7

1866. Scotti RJ, Angell G, Flora R, et al. Antecedent history as a predictor of surgical cure of urgency symptoms in mixed incontinence. *Obstet Gynecol.* 1998 Jan;91(1):51-4. X-1B
1867. Sears CL, Wright J, O'Brien J, et al. The racial distribution of female pelvic floor disorders in an equal access health care system. *J Urol.* 2009 Jan;181(1):187-92. X-8
1868. Seckiner I, Yesilli C, Mungan NA, et al. Correlations between the ICIQ-SF score and urodynamic findings. *Neurourol Urodyn.* 2007;26(4):492-4. X-7
1869. Segal J, Steele A, Vassallo B, et al. Various surgical approaches to treat voiding dysfunction following anti-incontinence surgery. *Int Urogynecol J Pelvic Floor Dysfunct.* 2006 Jun;17(4):372-7. X-II
1870. Segal JL, Vassallo B, Kleeman S, et al. Prevalence of persistent and de novo overactive bladder symptoms after the tension-free vaginal tape. *Obstet Gynecol.* 2004 Dec;104(6):1263-9. X-1A
1871. Segal JL, Vassallo BJ, Kleeman SD, et al. The efficacy of the tension-free vaginal tape in the treatment of five subtypes of stress urinary incontinence. *Int Urogynecol J Pelvic Floor Dysfunct.* 2006 Feb;17(2):120-4. X-1B
1872. Seim A, Sivertsen B, Eriksen BC, et al. Treatment of urinary incontinence in women in general practice: observational study. *BMJ.* 1996 Jun 8;312(7044):1459-62. X-1B
1873. Seki N, Masuda K, Kinukawa N, et al. Risk factors for febrile urinary tract infection in children with myelodysplasia treated by clean intermittent catheterization. *Int J Urol.* 2004 Nov;11(11):973-7. X-1A, X-4
1874. Sekido N, Hinotsu S, Kawai K, et al. How many uncomplicated male and female overactive bladder patients reveal detrusor overactivity during urodynamic study? *Int J Urol.* 2006 Oct;13(10):1276-9. X-7
1875. Sellors JW, Pickard L, Gafni A, et al. Effectiveness and efficiency of selective vs universal screening for chlamydial infection in sexually active young women. *Arch Intern Med.* 1992 Sep;152(9):1837-44. X-1A
1876. Sen I, Onaran M, Tan MO, et al. Evaluation of sexual function in women with overactive bladder syndrome. *Urol Int.* 2007;78(2):112-5. X-7
1877. Sensirivatana R, Watana D, Sornmani W, et al. Diagnostic study of urinary frequency in children. *Urology.* 1987 Jul;30(1):50-2. X-4, X-7
1878. Serels S and Stein M. Prospective study comparing hyoscyamine, doxazosin, and combination therapy for the treatment of urgency and frequency in women. *Neurourol Urodyn.* 1998;17(1):31-6. X-7
1879. Serels SR, Rackley RR and Appell RA. Surgical treatment for stress urinary incontinence associated with valsalva induced detrusor instability. *J Urol.* 2000 Mar;163(3):884-7. X-1B
1880. Serra DB, Afrime MB, Bedigian MP, et al. QT and QTc interval with standard and suprathreshold doses of darifenacin, a muscarinic M3 selective receptor antagonist for the treatment of overactive bladder. *J Clin Pharmacol.* 2005 Sep;45(9):1038-47. X-II
1881. Sethia KK and Smith JC. The effect of pH and lignocaine on detrusor instability. *Br J Urol.* 1987 Dec;60(6):516-8. X-7
1882. Sethia KK, Webb RJ and Neal DE. Urodynamic study of ileocystoplasty in the treatment of idiopathic detrusor instability. *Br J Urol.* 1991 Mar;67(3):286-90. X-7
1883. Setiati S, Santoso BI and Istanti R. Estimating the annual cost of overactive bladder in Indonesia. *Acta Med Indones.* 2006 Oct-Dec;38(4):189-92. X-6
1884. Sevestre S, Ciofu C, Deval B, et al. Results of the tension-free vaginal tape technique in the elderly. *Eur Urol.* 2003 Jul;44(1):128-31. X-1B
1885. Shafik A, El Sibai O, Shafik AA, et al. Vesical pacing: pacing parameters required for normalization of vesical electric activity in patients with overactive bladder. *Front Biosci.* 2004 Jan 1;9:995-9. X-7
1886. Shafik A, El-Sibai O, Shafik AA, et al. The electrovesicogram in the overactive bladder: role in determining pathogenesis and diagnostic significance. *Urol Res.* 2004 Aug;32(4):290-3. X-1H
1887. Shafik A, Shafik I, el-Sibai O, et al. Vesical pacing in patients with overactive bladder: technique and results. *Int Urol Nephrol.* 2004;36(1):29-34. X-7
1888. Shafik A and Shafik IA. Overactive bladder inhibition in response to pelvic floor muscle exercises. *World J Urol.* 2003 May;20(6):374-7. X-1H, X-7
1889. Shaker HS and Hassouna M. Sacral nerve root neuromodulation: an effective treatment for refractory urge incontinence. *J Urol.* 1998 May;159(5):1516-9. X-7
1890. Shapiro T, Dalton M, Hammock J, et al. The prevalence of urinary tract infections and sexually transmitted disease in women with symptoms of a simple urinary tract infection stratified by low colony count criteria. *Acad Emerg Med.* 2005 Jan;12(1):38-44. X-II
1891. Sharma S, Albertazzi P and Bottazzi M. The long-term effect of raloxifene on the genitourinary tract. *Climacteric.* 2007 Jun;10(3):244-8. X-1B, X-IH
1892. Shaw C, Gupta RD, Bushnell DM, et al. The extent and severity of urinary incontinence amongst women in UK GP waiting rooms. *Fam Pract.* 2006 Oct;23(5):497-506. X-6
1893. Shaya FT, Blume S, Gu A, et al. Persistence with overactive bladder pharmacotherapy in a Medicaid population. *Am J Manag Care.* 2005 Jul;11(4 Suppl):S121-9. X-II
1894. Shelbaia AM and Hashish NM. Limited experience in early management of genitourinary tract fistulas. *Urology.* 2007 Mar;69(3):572-4. X-II
1895. Sheremata WA, Harrington WJ, Jr., Bradshaw PA, et al. Association of '(tropical) ataxic neuropathy' with HTLV-II. *Virus Res.* 1993 Jul;29(1):71-7. X-1F
1896. Sherman ND, Jamison MG, Webster GD, et al. Sacral neuromodulation for the treatment of refractory urinary urge incontinence after stress incontinence surgery. *Am J Obstet Gynecol.* 2005 Dec;193(6):2083-7. X-1B
1897. Shibasaki H, Endo C, Kuroda Y, et al. Clinical picture of HTLV-I associated myelopathy. *J Neurol Sci.* 1988 Oct;87(1):15-24. X-II
1898. Shima H, Mori Y, Nojima M, et al. Lower urinary tract problems in patients with enuresis. *Eur Urol.* 1998;33 Suppl 3:37-40. X-II, X-4

1899. Shimabukuro T and Naito K. Evaluation of lower urinary tract symptoms and how bothersome it was with or without urinary incontinence in apparently healthy persons of both sexes. *Hinyokika Kiyo*. 2007 Mar;53(3):157-62. X-6
1900. Shimonovitz S, Monga AK and Stanton SL. Does the menstrual cycle influence cystometry? *Int Urogynecol J Pelvic Floor Dysfunct*. 1997;8(4):213-5; discussion 215-6. X-1H
1901. Shin JC, Kim YW, Park CI, et al. Effect of the intravesical resiniferatoxin instillation evaluated by the ice provocative urodynamic study. *Spinal Cord*. 2006 May;44(5):309-14. X-1F, X-5, X-7
1902. Shinopulos NM and Jacobson J. Relationship between health promotion lifestyle profiles and patient outcomes of biofeedback therapy for urinary incontinence. *Urol Nurs*. 1999 Dec;19(4):249-53. X-1I, X-7
1903. Shoukry MS, El Salmy S, Aly GA, et al. Urodynamic predictors of upper tract deterioration in children with myelodysplasia. *Scand J Urol Nephrol*. 1998 Apr;32(2):94-7. X-1F, X-4
1904. Shull BL and Baden WF. A six-year experience with paravaginal defect repair for stress urinary incontinence. *Am J Obstet Gynecol*. 1989 Jun;160(6):1432-9; discussion 1439-40. X-1B
1905. Siami P, Seidman LS and Lama D. A multicenter, prospective, open-label study of tolterodine extended-release 4 mg for overactive bladder: the speed of onset of therapeutic assessment trial (STAT). *Clin Ther*. 2002 Apr;24(4):616-28. X-4
1906. Siddiqui K, Raj H, Flynn RJ, et al. Minimally invasive treatment of female stress urinary incontinence: 100 cases using SPARC sling. *Ir J Med Sci*. 2008 Mar;177(1):39-42. X-1B
1907. Siddiqui MA, Perry CM and Scott LJ. Oxybutynin extended-release: a review of its use in the management of overactive bladder. *Drugs*. 2004;64(8):885-912. X-2
1908. Siegel SW, Richardson DA, Miller KL, et al. Pelvic floor electrical stimulation for the treatment of urge and mixed urinary incontinence in women. *Urology*. 1997 Dec;50(6):934-40. X-7
1909. Sievert KD, Nagele U, Pannek J, et al. Subcutaneous tunneling of the temporary testing electrode significantly improves the success rate of subchronic sacral nerve modulation (SNM). *World J Urol*. 2007 Dec;25(6):607-12. X-7
1910. Siltberg H, Larsson G and Victor A. Frequency/volume chart: the basic tool for investigating urinary symptoms. *Acta Obstet Gynecol Scand Suppl*. 1997;166:24-7. X-1H, X-2
1911. Silva C, Avelino A, Souto-Moura C, et al. A light- and electron-microscopic histopathological study of human bladder mucosa after intravesical resiniferatoxin application. *BJU Int*. 2001 Sep;88(4):355-60. X-4, X-7
1912. Silva C, Ribeiro MJ and Cruz F. The effect of intravesical resiniferatoxin in patients with idiopathic detrusor instability suggests that involuntary detrusor contractions are triggered by C-fiber input. *J Urol*. 2002 Aug;168(2):575-9. X-7
1913. Silva C, Rio ME and Cruz F. Desensitization of bladder sensory fibers by intravesical resiniferatoxin, a capsaicin analog: long-term results for the treatment of detrusor hyperreflexia. *Eur Urol*. 2000 Oct;38(4):444-52. X-7
1914. Silva C, Silva J, Ribeiro MJ, et al. Urodynamic effect of intravesical resiniferatoxin in patients with neurogenic detrusor overactivity of spinal origin: results of a double-blind randomized placebo-controlled trial. *Eur Urol*. 2005 Oct;48(4):650-5. X-1F
1915. Silva-Filho AL, Candido EB, Noronha A, et al. Comparative study of autologous pubovaginal sling and synthetic transobturator (TOT) SAFYRE sling in the treatment of stress urinary incontinence. *Arch Gynecol Obstet*. 2006 Feb;273(5):288-92. X-1B, X-7
1916. Simeonova Z and Bengtsson C. Prevalence of urinary incontinence among women at a Swedish primary health care centre. *Scand J Prim Health Care*. 1990 Dec;8(4):203-6. X-6
1917. Singh G and Thomas DG. Artificial urinary sphincter in patients with neurogenic bladder dysfunction. *Br J Urol*. 1996 Feb;77(2):252-5. X-1F
1918. Sinha D, Blackwell A and Moran PA. Outcome measures after TVT for mixed urinary incontinence. *Int Urogynecol J Pelvic Floor Dysfunct*. 2008 Jul;19(7):927-31. X-1A, X-7
1919. Sinha D, Nallaswamy V and Arunkalaivanan AS. Value of leak point pressure study in women with incontinence. *J Urol*. 2006 Jul;176(1):186-8; discussion 188. X-1H
1920. Sivaslioglu AA, Caliskan E, Dolen I, et al. A randomized comparison of transobturator tape and Burch colposuspension in the treatment of female stress urinary incontinence. *Int Urogynecol J Pelvic Floor Dysfunct*. 2007 Sep;18(9):1015-9. X-1B
1921. Skala C, Emons G, Krauss T, et al. Postoperative funneling after anti-incontinence surgery--a prognostic indicator?--Part 1: colposuspension. *Neurourol Urodyn*. 2004;23(7):636-42. X-1B
1922. Skeil D and Thorpe AC. Transcutaneous electrical nerve stimulation in the treatment of neurological patients with urinary symptoms. *BJU Int*. 2001 Dec;88(9):899-908. X-1F
1923. Slors FJ, van Zuijlen PP and van Dijk GJ. Sexual and bladder dysfunction after total mesorectal excision for benign diseases. *Scand J Gastroenterol Suppl*. 2000(232):48-51. X-1A
1924. Smet PJ, Moore KH and Jonavicius J. Distribution and colocalization of calcitonin gene-related peptide, tachykinins, and vasoactive intestinal peptide in normal and idiopathic unstable human urinary bladder. *Lab Invest*. 1997 Jul;77(1):37-49. X-1G, X-7
1925. Smith CP, Nishiguchi J, O'Leary M, et al. Single-institution experience in 110 patients with botulinum toxin A injection into bladder or urethra. *Urology*. 2005 Jan;65(1):37-41. X-4
1926. Smith DA and Ouslander JG. Pharmacologic management of urinary incontinence in older adults. *Topics in Geriatric Rehabilitation*. 2000;16(1):54-60. X-2

1927. Smith DB, Boileau MA and Buan LD. A self-directed home biofeedback system for women with symptoms of stress, urge, and mixed incontinence. *J Wound Ostomy Continence Nurs.* 2000 Jul;27(4):240-6. X-7
1928. Smith DN, Appell RA, Winters JC, et al. Collagen injection therapy for female intrinsic sphincteric deficiency. *J Urol.* 1997 Apr;157(4):1275-8. X-II
1929. Smith EM and Elder JS. Double antimicrobial prophylaxis in girls with breakthrough urinary tract infections. *Urology.* 1994 May;43(5):708-12; discussion 712-3. X-II, X-4
1930. Smith JC, Dunn M and Ardran GM. Hydrostatic bladder distension - physiology and clinical application. *Eur Urol.* 1975;1(2):75-7. X-1G
1931. Smith JJ, 3rd. Intravaginal stimulation randomized trial. *J Urol.* 1996 Jan;155(1):127-30. X-7
1932. Smith P, Heimer G, Lindskog M, et al. Oestradiol-releasing vaginal ring for treatment of postmenopausal urogenital atrophy. *Maturitas.* 1993 Mar;16(2):145-54. X-II
1933. Smulders RA, Kuipers ME and Krauwinkel WJ. Multiple doses of the antimuscarinic agent solifenacin do not affect the pharmacodynamics or pharmacokinetics of warfarin or the steady-state pharmacokinetics of digoxin in healthy subjects. *Br J Clin Pharmacol.* 2006 Aug;62(2):210-7. X-II, X-4
1934. Snape J, Castleden CM, Duffin HM, et al. Long-term follow-up of habit retraining for bladder instability in elderly patients. *Age Ageing.* 1989 May;18(3):192-4. X-7
1935. Sobhgol SS and Charandabee SM. Related factors of urge, stress, mixed urinary incontinence and overactive bladder in reproductive age women in Tabriz, Iran: a cross-sectional study. *Int Urogynecol J Pelvic Floor Dysfunct.* 2008 Mar;19(3):367-73. X-1A
1936. Soligo M, Salvatore S, Milani R, et al. Double incontinence in urogynecologic practice: a new insight. *Am J Obstet Gynecol.* 2003 Aug;189(2):438-43. X-II
1937. Song C, Park SH, Han JY, et al. Identification of the optimal time to treat urgency after a midurethral sling procedure for stress urinary incontinence. *Int Urogynecol J Pelvic Floor Dysfunct.* 2008 Apr;19(4):573-6. X-1B
1938. Soomro NA, Khadra MH, Robson W, et al. A crossover randomized trial of transcutaneous electrical nerve stimulation and oxybutynin in patients with detrusor instability. *J Urol.* 2001 Jul;166(1):146-9. X-7
1939. Soontrapa S, Ruksakul W, Nonthasood B, et al. The efficacy of Thai capsaicin in management of overactive bladder and hypersensitive bladder. *J Med Assoc Thai.* 2003 Sep;86(9):861-7. X-7
1940. Sousa-Escandon A, Lema Grille J, Rodriguez Gomez JJ, et al. Externally readjustable device to regulate sling tension in stress urinary incontinence: preliminary results. *J Endourol.* 2003 Sep;17(7):515-21. X-1A
1941. Soygur T, Arikan N, Tokatli Z, et al. The role of video-urodynamic studies in managing non-neurogenic voiding dysfunction in children. *BJU Int.* 2004 Apr;93(6):841-3. X-1H, X-4
1942. Soyupek F, Soyupek S, Akkus S, et al. The coexistence of the fibromyalgia syndrome and the overactive bladder syndrome. *Journal of Musculoskeletal Pain.* 2007;15(3):31-7. X-II
1943. Sozubir S, Celik A, Emir N, et al. Consistency of urodynamic parameters in children with detrusor instability: how many times should the bladder be filled? *Urol Int.* 2005;75(2):129-32. X-1H, X-4
1944. Spinelli M, Giardiello G, Gerber M, et al. New sacral neuromodulation lead for percutaneous implantation using local anesthesia: description and first experience. *J Urol.* 2003 Nov;170(5):1905-7. X-7
1945. Spinelli M, Malaguti S, Giardiello G, et al. A new minimally invasive procedure for pudendal nerve stimulation to treat neurogenic bladder: description of the method and preliminary data. *Neurourol Urodyn.* 2005;24(4):305-9. X-1F
1946. Sprem M, Milicic D, Oreskovic S, et al. Intravesically administered ketoprofen in treatment of detrusor instability: cross-over study. *Croat Med J.* 2000 Dec;41(4):423-7. X-7
1947. Spruijt J, Vierhout M, Verstraeten R, et al. Vaginal electrical stimulation of the pelvic floor: a randomized feasibility study in urinary incontinent elderly women. *Acta Obstet Gynecol Scand.* 2003 Nov;82(11):1043-8. X-7
1948. Stach-Lempinen B, Hakala AL, Laippala P, et al. Severe depression determines quality of life in urinary incontinent women. *Neurourol Urodyn.* 2003;22(6):563-8. X-7
1949. Stage P, Fischer-Rasmussen W and Hansen RI. The value of colpo-cysto-urethrography in female stress- and urge incontinence and following operation. *Acta Obstet Gynecol Scand.* 1986;65(5):401-4. X-1H
1950. Stanton SL. Stress incontinence: why and how operations work. *Clin Obstet Gynaecol.* 1985 Jun;12(2):369-77. X-1B
1951. Stanton SL, Cardozo L, Williams JE, et al. Clinical and urodynamic features of failed incontinence surgery in the female. *Obstet Gynecol.* 1978 May;51(5):515-20. X-1A
1952. Stanton SL, Hilton P, Norton C, et al. Clinical and urodynamic effects of anterior colporrhaphy and vaginal hysterectomy for prolapse with and without incontinence. *Br J Obstet Gynaecol.* 1982 Jun;89(6):459-63. X-1A, X-7
1953. Stanton SL, Kerr-Wilson R and Harris VG. The incidence of urological symptoms in normal pregnancy. *Br J Obstet Gynaecol.* 1980 Oct;87(10):897-900. X-II
1954. Stanton SL and Monga AK. Incontinence in elderly women: is periurethral collagen an advance? *Br J Obstet Gynaecol.* 1997 Feb;104(2):154-7. X-1B, X-7
1955. Starer P and Libow LS. The measurement of residual urine in the evaluation of incontinent nursing home residents. *Arch Gerontol Geriatr.* 1988 Mar;7(1):75-81. X-5
1956. Starkman JS, Duffy JW, 3rd, Wolter CE, et al. The evolution of obstruction induced overactive bladder symptoms following urethrolysis for female bladder outlet obstruction. *J Urol.* 2008 Mar;179(3):1018-23. X-1A, X-7
1957. Starkman JS, Duffy JW, 3rd, Wolter CE, et al. Refractory overactive bladder after urethrolysis for bladder outlet obstruction: management with sacral neuromodulation. *Int Urogynecol J Pelvic Floor Dysfunct.* 2008 Feb;19(2):277-82. X-1A, X-7
1958. Starkman JS, Wolter CE, Scarpero HM, et al. Management of refractory urinary urge incontinence following urogynecological surgery with sacral neuromodulation. *Neurourol Urodyn.* 2007;26(1):29-35; discussion 36. X-1A, X-7

1959. Staskin DR, Choe JM and Breslin DS. The Gore-tex sling procedure for female sphincteric incontinence: indications, technique, and results. *World J Urol.* 1997;15(5):295-9. X-1B
1960. Staskin DR and Te AE. Short- and long-term efficacy of solifenacin treatment in patients with symptoms of mixed urinary incontinence. *BJU Int.* 2006 Jun;97(6):1256-61. X-1B
1961. Stav K, Dwyer PL, Rosamilia A, et al. Urinary symptoms before and after female urethral diverticulectomy--can we predict de novo stress urinary incontinence? *J Urol.* 2008 Nov;180(5):2088-90. X-1B
1962. Steel SA, Cox C and Stanton SL. Long-term follow-up of detrusor instability following the colposuspension operation. *Br J Urol.* 1986 Apr;58(2):138-42. X-1A, X-7
1963. Steele AC, Walsh P, Bentley M, et al. A randomized, double-blind placebo-controlled trial of the effects of the 5-hydroxytryptamine(4) agonist cisapride on the female urinary bladder. *Am J Obstet Gynecol.* 2001 Jul;185(1):62-4. X-1I
1964. Steers W. Pathogenesis of the overactive bladder and its attendant risk factors. *BJU Int.* 2000 May;85 Suppl 3:69; discussion 70-1. X-2
1965. Steers WD, Meythaler JM, Haworth C, et al. Effects of acute bolus and chronic continuous intrathecal baclofen on genitourinary dysfunction due to spinal cord pathology. *J Urol.* 1992 Dec;148(6):1849-55. X-1F, X-7
1966. Stein M, Discippio W, Davia M, et al. Biofeedback for the treatment of stress and urge incontinence. *J Urol.* 1995 Mar;153(3 Pt 1):641-3. X-1B, X-7
1967. Stein RJ, Santos S, Nagatomi J, et al. Cool (TRPM8) and hot (TRPV1) receptors in the bladder and male genital tract. *J Urol.* 2004 Sep;172(3):1175-8. X-1G, X-4, X-5, X-6, X-7
1968. Steinauer JE, Waetjen LE, Vittinghoff E, et al. Postmenopausal hormone therapy: does it cause incontinence? *Obstet Gynecol.* 2005 Nov;106(5 Pt 1):940-5. X-II
1969. Stenberg A, Heimer G, Ulmsten U, et al. Prevalence of genitourinary and other climacteric symptoms in 61-year-old women. *Maturitas.* 1996 May;24(1-2):31-6. X-1A
1970. Stevens LA, Chapple CR and Chess-Williams R. Human idiopathic and neurogenic overactive bladders and the role of M2 muscarinic receptors in contraction. *Eur Urol.* 2007 Aug;52(2):531-8. X-1G
1971. Stewart DJ, Green R, Futter N, et al. Phase I and pharmacology study of intravesical mitoxantrone for recurrent superficial bladder tumors. *J Urol.* 1990 Apr;143(4):714-6. X-1A, X-7
1972. Stimpel H, Aagaard J and Reuther K. Repeated bladder distension by the Cystomat in the treatment of detrusor instability. *Br J Urol.* 1984 Jun;56(3):285-8. X-7
1973. Stohrer M, Murtz G, Kramer G, et al. Propiverine compared to oxybutynin in neurogenic detrusor overactivity--results of a randomized, double-blind, multicenter clinical study. *Eur Urol.* 2007 Jan;51(1):235-42. X-1F
1974. Stothers L, Goldenberg SL and Leone EF. Complications of periurethral collagen injection for stress urinary incontinence. *J Urol.* 1998 Mar;159(3):806-7. X-1B
1975. Su X, Leon LA, Wu CW, et al. Modulation of bladder function by prostaglandin EP3 receptors in the central nervous system. *Am J Physiol Renal Physiol.* 2008 Oct;295(4):F984-94. X-1G
1976. Subak LL, Quesenberry CP, Posner SF, et al. The effect of behavioral therapy on urinary incontinence: a randomized controlled trial. *Obstet Gynecol.* 2002 Jul;100(1):72-8. X-7
1977. Subak LL, Wing R, West DS, et al. Weight loss to treat urinary incontinence in overweight and obese women. *N Engl J Med.* 2009 Jan 29;360(5):481-90. X-1B
1978. Suchowersky O, Furtado S and Rohs G. Beneficial effect of intranasal desmopressin for nocturnal polyuria in Parkinson's disease. *Mov Disord.* 1995 May;10(3):337-40. X-1A, X-1C, X-7
1979. Sugaya K, Nishijima S, Oda M, et al. Change of blood viscosity and urinary frequency by high water intake. *Int J Urol.* 2007 May;14(5):470-2. X-1A
1980. Sugaya K, Nishijima S, Oda M, et al. Biochemical and body composition analysis of nocturia in the elderly. *Neurourol Urodyn.* 2008;27(3):205-11. X-1C
1981. Sugaya K, Nishizawa O, Satoh T, et al. Bladder-pumping therapy for the treatment of low-capacity or low-compliance bladders. *Neurourol Urodyn.* 2000;19(1):19-28. X-7
1982. Sugaya K, Onaga T, Nishijima S, et al. Relationship between serum cholinesterase level and urinary bladder activity in patients with or without overactive bladder and/or neurogenic bladder. *Biomedical Research.* 2007;28(6):287-294. X-1I
1983. Sugaya K, Onaga T, Nishijima S, et al. Relationship between serum cholinesterase level and urinary bladder activity in patients with or without overactive bladder and/or neurogenic bladder. *Biomed Res.* 2008 Dec;28(6):287-94. X-1G
1984. Suhel P, Kralj B and Plevnik S. Advances in nonimplantable electrical stimulators for correction of urinary incontinence. *TIT J Life Sci.* 1978;8(1-2):11-6. X-2
1985. Sui GP, Wu C, Severs N, et al. The association between T-type Ca²⁺ current and outward current in isolated human detrusor cells from stable and overactive bladders. *BJU Int.* 2007 Feb;99(2):436-41. X-1H
1986. Summitt RL, Jr., Stovall TG, Bent AE, et al. Urinary incontinence: correlation of history and brief office evaluation with multichannel urodynamic testing. *Am J Obstet Gynecol.* 1992 Jun;166(6 Pt 1):1835-40; discussion 1840-4. X-1H
1987. Susset J, Galea G, Manbeck K, et al. A predictive score index for the outcome of associated biofeedback and vaginal electrical stimulation in the treatment of female incontinence. *J Urol.* 1995 May;153(5):1461-6. X-7
1988. Sussman DO, Kraus SR, Carlsson M, et al. Onset of efficacy of tolterodine extended release in patients with overactive bladder. *Curr Med Res Opin.* 2007 Apr;23(4):777-81. X-1H
1989. Sutherst JR and Brown MC. Comparison of single and multichannel cystometry in diagnosing bladder instability. *Br Med J (Clin Res Ed).* 1984 Jun 9;288(6432):1720-2. X-1H

1990. Sutherst JR, Brown MC and Richmond D. Analysis of the pattern of urine loss in women with incontinence as measured by weighing perineal pads. *Br J Urol.* 1986 Jun;58(3):273-8. X-1H
1991. Suzuki S, Dennerstein L, Greenwood KM, et al. Sleeping patterns during pregnancy in Japanese women. *J Psychosom Obstet Gynaecol.* 1994 Mar;15(1):19-26. X-1I
1992. Svigos JM and Matthews CD. Assessment and treatment of female urinary incontinence by cystometrogram and bladder retraining programs. *Obstet Gynecol.* 1977 Jul;50(1):9-12. X-7
1993. Svihra J, Kurca E, Luptak J, et al. Neuromodulative treatment of overactive bladder--noninvasive tibial nerve stimulation. *Bratisl Lek Listy.* 2002;103(12):480-3. X-7
1994. Swann J. Simple steps for promoting urinary continence. *Nursing & Residential Care.* 2005;7(10):464-466. X-2
1995. Swanson SG, Drosman S, Helmond FA, et al. Tibolone for the treatment of moderate to severe vasomotor symptoms and genital atrophy in postmenopausal women: a multicenter, randomized, double-blind, placebo-controlled study. *Menopause.* 2006 Nov-Dec;13(6):917-25. X-1I
1996. Swart PJ, Krauwinkel WJ, Smulders RA, et al. Pharmacokinetic effect of ketoconazole on solifenacin in healthy volunteers. *Basic Clin Pharmacol Toxicol.* 2006 Jul;99(1):33-6. X-1A, X-7
1997. Sweat SD, Itano NB, Clemens JQ, et al. Polypropylene mesh tape for stress urinary incontinence: complications of urethral erosion and outlet obstruction. *J Urol.* 2002 Jul;168(1):144-6. X-1B
1998. Swierzewski SJ, 3rd and McGuire EJ. Pubovaginal sling for treatment of female stress urinary incontinence complicated by urethral diverticulum. *J Urol.* 1993 May;149(5):1012-4. X-1B
1999. Swift SE. The reliability of performing a screening cystometrogram using a fetal monitoring device for the detection of detrusor instability. *Obstet Gynecol.* 1997 May;89(5 Pt 1):708-12. X-1H
2000. Swift SE and Yoon EA. Test-retest reliability of the cough stress test in the evaluation of urinary incontinence. *Obstet Gynecol.* 1999 Jul;94(1):99-102. X-1B
2001. Swithinbank LV, James M, Shepherd A, et al. Role of ambulatory urodynamic monitoring in clinical urological practice. *Neurourol Urodyn.* 1999;18(3):215-22. X-1H
2002. Switzer D and Hendriks O. Interferential therapy for the treatment of stress and urge incontinence. *Ir Med J.* 1988 Nov;81(1):30-1. X-7
2003. Syrjala KL, Langer SL, Abrams JR, et al. Late effects of hematopoietic cell transplantation among 10-year adult survivors compared with case-matched controls. *J Clin Oncol.* 2005 Sep 20;23(27):6596-606. X-1I
2004. Sze EH, Jones WP, Ferguson JL, et al. Prevalence of urinary incontinence symptoms among black, white, and Hispanic women. *Obstet Gynecol.* 2002 Apr;99(4):572-5. X-6
2005. Sze EH, Miklos JR, Partoll L, et al. Sacrospinous ligament fixation with transvaginal needle suspension for advanced pelvic organ prolapse and stress incontinence. *Obstet Gynecol.* 1997 Jan;89(1):94-6. X-1B
2006. Tadic SD, Griffiths D, Schaefer W, et al. Abnormal connections in the supraspinal bladder control network in women with urge urinary incontinence. *Neuroimage.* 2008 Feb 15;39(4):1647-53. X-1G, X-1H, X-7
2007. Taekema-Roelvink ME, Swart PJ, Kuipers ME, et al. Pharmacokinetic interaction of solifenacin with an oral contraceptive containing ethinyl estradiol and levonorgestrel in healthy women: a double-blind, placebo-controlled study. *Clin Ther.* 2005 Sep;27(9):1403-10. X-1A
2008. Tahseen S, Reid PC and Charan P. Short-term complications of the trans-obturator foramen procedure for urinary stress incontinence. *J Obstet Gynaecol.* 2007 Jul;27(5):500-2. X-1B
2009. Tai HL, Chang YJ, Chang SC, et al. The epidemiology and factors associated with nocturnal enuresis and its severity in primary school children in Taiwan. *Acta Paediatr.* 2007 Feb;96(2):242-5. X-1I, X-4
2010. Takahashi R, Nishimura J, Hirano K, et al. Ca²⁺ sensitization in contraction of human bladder smooth muscle. *J Urol.* 2004 Aug;172(2):748-52. X-1G
2011. Takazawa K and Arisawa K. Relationship between the type of urinary incontinence and falls among frail elderly women in Japan. *J Med Invest.* 2005 Aug;52(3-4):165-71. X-5
2012. Takei M and Homma Y. Long-term safety, tolerability and efficacy of extended-release tolterodine in the treatment of overactive bladder in Japanese patients. *Int J Urol.* 2005 May;12(5):456-64. X-4
2013. Tan TL, Bergmann MA, Griffiths D, et al. Which stop test is best? Measuring detrusor contractility in older females. *J Urol.* 2003 Mar;169(3):1023-7. X-1I
2014. Tan TL, Bergmann MA, Griffiths D, et al. Stop test or pressure-flow study? Measuring detrusor contractility in older females. *Neurourol Urodyn.* 2004;23(3):184-9. X-1H
2015. Tanagho EA. Urodynamics of female urinary incontinence with emphasis on stress incontinence. *J Urol.* 1979 Aug;122(2):200-4. X-1B, X-1G
2016. Tanaka H, Matsuda M, Moriya K, et al. Ultrasonographic measurement of bladder wall thickness as a risk factor for upper urinary tract deterioration in children with myelodysplasia. *J Urol.* 2008 Jul;180(1):312-6; discussion 316. X-1F, X-1H, X-4
2017. Tanaka ST, Stone AR and Kurzrock EA. Transverse myelitis in children: long-term urological outcomes. *J Urol.* 2006 May;175(5):1865-8; discussion 1868. X-1F, X-1I, X-4
2018. Tannenbaum C, Bachand G, Dubeau CE, et al. Experience of an incontinence clinic for older women: no apparent age limit for potential physical and psychological benefits. *J Womens Health Gend Based Med.* 2001 Oct;10(8):751-6. X-7
2019. Tantanasis T, Giannoulis C, Daniilidis A, et al. Anterior vaginal wall reconstruction: anterior colporrhaphy reinforced with tension free vaginal tape underneath bladder base. *Acta Obstet Gynecol Scand.* 2008;87(4):464-8. X-1E
2020. Tapp A, Fall M, Norgaard J, et al. Terodiline: a dose titrated, multicenter study of the treatment of idiopathic detrusor instability in women. *J Urol.* 1989 Oct;142(4):1027-31. X-9

2021. Tapp AJ, Cardozo LD, Versi E, et al. The treatment of detrusor instability in post-menopausal women with oxybutynin chloride: a double blind placebo controlled study. *Br J Obstet Gynaecol.* 1990 Jun;97(6):521-6. X-7
2022. Tarhan F, Erbay E, Eryildirim B, et al. The effect of intravesical sodium nitroprusside on idiopathic detrusor overactivity. *Urol Res.* 2004 Jun;32(3):200-3. X-7
2023. Taskinen S, Fagerholm R and Rintala R. Mini-invasive collagen sling in the treatment of urinary incontinence due to sphincteric incompetence. *Int Braz J Urol.* 2007 May-Jun;33(3):395-400; discussion 400-6. X-II
2024. Taskinen S, Valanne L and Rintala R. Effect of spinal cord abnormalities on the function of the lower urinary tract in patients with anorectal abnormalities. *J Urol.* 2002 Sep;168(3):1147-9. X-1F, X-1H, X-7
2025. Taylor CM, Corkery JJ and White RH. Micturition symptoms and unstable bladder activity in girls with primary vesicoureteric reflux. *Br J Urol.* 1982 Oct;54(5):494-8. X-II, X-4
2026. Taylor PH and Sussman DO. Contemporary treatment options for overactive bladder. *JAAPA.* 2005 Nov;Suppl:3-13; quiz 14-5. X-2
2027. Taylor T, Smith AN and Fulton M. Effects of hysterectomy on bowel and bladder function. *Int J Colorectal Dis.* 1990 Dec;5(4):228-31. X-II
2028. Tebyani N, Patel H, Yamaguchi R, et al. Percutaneous needle bladder neck suspension for the treatment of stress urinary incontinence in women: long-term results. *J Urol.* 2000 May;163(5):1510-2. X-1B, X-7
2029. Tegerstedt G, Maehle-Schmidt M, Nyren O, et al. Prevalence of symptomatic pelvic organ prolapse in a Swedish population. *Int Urogynecol J Pelvic Floor Dysfunct.* 2005 Nov-Dec;16(6):497-503. X-1E
2030. Tegerstedt G, Sjoberg B and Hammarstrom M. Clinical outcome or abdominal urethropexy-colposuspension: a long-term follow-up. *Int Urogynecol J Pelvic Floor Dysfunct.* 2001;12(3):161-5. X-1B
2031. Teichman JM and Nielsen-Omeis BJ. Potassium leak test predicts outcome in interstitial cystitis. *J Urol.* 1999 Jun;161(6):1791-4; discussion 1794-6. X-1D
2032. Teleman P, Gunnarsson M, Lidfeldt J, et al. Urodynamic characterisation of women with naive urinary incontinence: a population-based study in subjectively incontinent and healthy 53-63 years old women. *Eur Urol.* 2002 Dec;42(6):583-9. X-1G
2033. Teleman P, Lidfeldt J, Nerbrand C, et al. Lower urinary tract symptoms in middle-aged women--prevalence and attitude towards mild urinary incontinence: a community-based population study. *Acta Obstet Gynecol Scand.* 2005 Nov;84(11):1108-12. X-II
2034. Teleman PM, Lidfeldt J, Nerbrand C, et al. Overactive bladder: prevalence, risk factors and relation to stress incontinence in middle-aged women. *BJOG.* 2004 Jun;111(6):600-4. X-6
2035. Temml C, Heidler S, Ponholzer A, et al. Prevalence of the overactive bladder syndrome by applying the International Continence Society definition. *Eur Urol.* 2005 Oct;48(4):622-7. X-6
2036. Tennstedt SL, Fitzgerald MP, Nager CW, et al. Quality of life in women with stress urinary incontinence. *Int Urogynecol J Pelvic Floor Dysfunct.* 2007 May;18(5):543-9. X-1B
2037. Teo JS, Briffa NK, Devine A, et al. Do sleep problems or urinary incontinence predict falls in elderly women? *Aust J Physiother.* 2006;52(1):19-24. X-II
2038. Teo RE, Yisa S, Mayne C, et al. A comparison of urodynamic diagnoses and detrusor muscle function in white and South Indian Asian women. *J Urol.* 2005 Jul;174(1):184-6. X-1A
2039. ter Meulen PH, Heesakkers JP and Janknegt RA. A study on the feasibility of vesicomatomy in patients with motor urge incontinence. *Eur Urol.* 1997;32(2):166-9. X-7
2040. Thakar R, Ayers S, Clarkson P, et al. Outcomes after total versus subtotal abdominal hysterectomy. *N Engl J Med.* 2002 Oct 24;347(17):1318-25. X-1A
2041. Thase ME, Shelton RC and Khan A. Treatment with venlafaxine extended release after SSRI nonresponse or intolerance: a randomized comparison of standard- and higher-dosing strategies. *J Clin Psychopharmacol.* 2006 Jun;26(3):250-8. X-II
2042. Theodorou C, Floratos D, Katsifotis C, et al. Transvaginal incisionless bladder neck suspension. A simplified technique for female genuine stress incontinence. *Int Urol Nephrol.* 1998;30(3):273-8. X-1B
2043. Theodorou C, Katsifotis C, Bocos J, et al. Urodynamics prior to renal transplantation--its impact on treatment decision and final results. *Scand J Urol Nephrol.* 2003;37(4):335-8. X-1G, X-1I
2044. Theofrastous JP, Wyman JF, Bump RC, et al. Effects of pelvic floor muscle training on strength and predictors of response in the treatment of urinary incontinence. *Neurourol Urodyn.* 2002;21(5):486-90. X-7
2045. Theofrastous JP, Wyman JF, Bump RC, et al. Relationship between urethral and vaginal pressures during pelvic muscle contraction. The Continence Program for Women Research Group. *Neurourol Urodyn.* 1997;16(6):553-8. X-1H
2046. Theoharides TC, Sant GR, el-Mansoury M, et al. Activation of bladder mast cells in interstitial cystitis: a light and electron microscopic study. *J Urol.* 1995 Mar;153(3 Pt 1):629-36. X-1D
2047. Thiede HA and Thiede FK. A glance at the urodynamic database. *J Reprod Med.* 1990 Oct;35(10):925-31. X-1B, X-1G
2048. Thompson JA, O'Sullivan PB, Briffa NK, et al. Comparison of transperineal and transabdominal ultrasound in the assessment of voluntary pelvic floor muscle contractions and functional manoeuvres in continent and incontinent women. *Int Urogynecol J Pelvic Floor Dysfunct.* 2007 Jul;18(7):779-86. X-1H
2049. Thyssen H, Sander P and Lose G. A vaginal device (continence guard) in the management of urge incontinence in women. *Int Urogynecol J Pelvic Floor Dysfunct.* 1999;10(4):219-22. X-7
2050. Tibaek S, Gard G and Jensen R. Pelvic floor muscle training is effective in women with urinary incontinence after stroke: a randomised, controlled and blinded study. *Neurourol Urodyn.* 2005;24(4):348-57. X-1F, X-7
2051. Tincello DG, Adams EJ, Sutherst JR, et al. Oxybutynin for detrusor instability with adjuvant salivary stimulant pastilles to improve compliance: results of a multicentre, randomized controlled trial. *BJU Int.* 2000 Mar;85(4):416-20. X-II

2052. Tincello DG and Richmond DH. The Larsson frequency/volume chart is not a substitute for cystometry in the investigation of women with urinary incontinence. *Int Urogynecol J Pelvic Floor Dysfunct.* 1998;9(6):391-6. X-1H
2053. Toaff R and Toaff ME. Diagnosis of impending late abortion. *Obstet Gynecol.* 1974 May;43(5):756-9. X-II
2054. Tomoe H, Kondo A, Takei M, et al. Quality of life assessments in women operated on by tension-free vaginal tape (TVT). *Int Urogynecol J Pelvic Floor Dysfunct.* 2005 Mar-Apr;16(2):114-8; discussion 108. X-1B
2055. Tomoe H, Sekiguchi Y, Horiguchi M, et al. Questionnaire survey on female urinary frequency and incontinence. *Int J Urol.* 2005 Jul;12(7):621-30. X-6, X-7
2056. Topping J, Petersen T, Klemar B, et al. Selective sacral rootlet neurectomy in the treatment of detrusor hyperreflexia. Technique and long-term results. *J Neurosurg.* 1988 Feb;68(2):241-5. X-7
2057. Touloupidis S, Papatsoris AG, Thanopoulos C, et al. Tension-free vaginal tape for the treatment of stress urinary incontinence in geriatric patients. *Gerontology.* 2007;53(3):125-7. X-1B
2058. Tow AM, Toh KL, Chan SP, et al. Botulinum toxin type A for refractory neurogenic detrusor overactivity in spinal cord injured patients in Singapore. *Ann Acad Med Singapore.* 2007 Jan;36(1):11-7. X-1F, X-7
2059. Triantafyllidis A, Charalambous S, Papatsoris AG, et al. Management of nocturnal enuresis in Greek children. *Pediatr Nephrol.* 2005 Sep;20(9):1343-5. X-II, X-4
2060. Trienekens TA, Stobberingh EE, Winkens RA, et al. Different lengths of treatment with co-trimoxazole for acute uncomplicated urinary tract infections in women. *BMJ.* 1989 Nov 25;299(6711):1319-22. X-1A
2061. Trifillis AL, Cui X, Jacobs S, et al. Culture of bladder epithelium from cystoscopic biopsies of patients with interstitial cystitis. *J Urol.* 1995 Jan;153(1):243-8. X-1D, X-1G, X-7
2062. Trockman BA, Leach GE, Hamilton J, et al. Modified Pereyra bladder neck suspension: 10-year mean followup using outcomes analysis in 125 patients. *J Urol.* 1995 Nov;154(5):1841-7. X-1B
2063. Trsinar B and Kraij B. Maximal electrical stimulation in children with unstable bladder and nocturnal enuresis and/or daytime incontinence: a controlled study. *Neurourol Urodyn.* 1996;15(2):133-42. X-II, X-4
2064. Truss MC, Stief CG, Uckert S, et al. Initial clinical experience with the selective phosphodiesterase-I isoenzyme inhibitor vinpocetine in the treatment of urge incontinence and low compliance bladder. *World J Urol.* 2000 Dec;18(6):439-43. X-7
2065. Tse V, Wills E, Szonyi G, et al. The application of ultrastructural studies in the diagnosis of bladder dysfunction in a clinical setting. *J Urol.* 2000 Feb;163(2):535-9. X-1H, X-7
2066. Tseng LH, Liang CC, Lo HP, et al. The prevalence of urinary incontinence and associated risk factors in Taiwanese women with lower urinary tract symptoms. *Chang Gung Med J.* 2006 Nov-Dec;29(6):596-602. X-1H, X-6
2067. Tsivian A, Kessler O, Mogutin B, et al. Tape related complications of the tension-free vaginal tape procedure. *J Urol.* 2004 Feb;171(2 Pt 1):762-4. X-1B
2068. Tsui KP, Ng SC, Yeh GP, et al. Outcomes of autologous fascial slingplasty procedure for treating female urinary incontinence. *Int Urogynecol J Pelvic Floor Dysfunct.* 2008 Jul;19(7):949-54. X-1A, X-1B
2069. Tsuzuki T and Epstein JI. Fibroepithelial polyp of the lower urinary tract in adults. *Am J Surg Pathol.* 2005 Apr;29(4):460-6. X-1A, X-7
2070. Tyagi S, Tyagi P, Van-le S, et al. Qualitative and quantitative expression profile of muscarinic receptors in human urothelium and detrusor. *J Urol.* 2006 Oct;176(4 Pt 1):1673-8. X-1G
2071. Uchiyama T, Sakakibara R, Hattori T, et al. Short-term effect of a single levodopa dose on micturition disturbance in Parkinson's disease patients with the wearing-off phenomenon. *Mov Disord.* 2003 May;18(5):573-8. X-1F
2072. Ueda T, Tamaki M, Ogawa O, et al. Improvement of interstitial cystitis symptoms and problems that developed during treatment with oral IPD-1151T. *J Urol.* 2000 Dec;164(6):1917-20. X-1D, X-7
2073. Ueda T, Tamaki M, Ogawa O, et al. Over expression of platelet-derived endothelial cell growth factor/thymidine phosphorylase in patients with interstitial cystitis and bladder carcinoma. *J Urol.* 2002 Jan;167(1):347-51. X-1A, X-1D
2074. Uekado Y, Hirano A, Shinka T, et al. The effects of intravesical chemoimmunotherapy with epirubicin and bacillus Calmette-Guerin for prophylaxis of recurrence of superficial bladder cancer: a preliminary report. *Cancer Chemother Pharmacol.* 1994;35 Suppl:S65-8. X-II, X-7
2075. Uemura S and Homma Y. Reliability and validity of King's Health Questionnaire in patients with symptoms of overactive bladder with urge incontinence in Japan. *Neurourol Urodyn.* 2004;23(2):94-100. X-II
2076. Ukkonen M, Elovaara I, Dastidar P, et al. Urodynamic findings in primary progressive multiple sclerosis are associated with increased volumes of plaques and atrophy in the central nervous system. *Acta Neurol Scand.* 2004 Feb;109(2):100-5. X-1F
2077. Ulmsten U, Andersson KE and Persson CG. Diagnostic and therapeutic aspects of urge urinary incontinence in women. *Urol Int.* 1977;32(2-3):88-96. X-7
2078. Ulmsten U, Ekman G and Andersson KE. The effect of terodiline treatment in women with motor urge incontinence. Results from a double-blind study and long-term treatment. *Am J Obstet Gynecol.* 1985 Nov 15;153(6):619-22. X-7
2079. Ulmsten U, Henriksson L and Iosif S. The unstable female urethra. *Am J Obstet Gynecol.* 1982 Sep 1;144(1):93-7. X-1G, X-1H
2080. Ulmsten U and Petros P. Intravaginal slingplasty (IVS): an ambulatory surgical procedure for treatment of female urinary incontinence. *Scand J Urol Nephrol.* 1995 Mar;29(1):75-82. X-1B
2081. Ulshofer B, Bihl AM, Bodeker RH, et al. Randomised, double-blind, placebo-controlled study on the efficacy and tolerance of trospium chloride in patients with motor urge incontinence. *Clinical Drug Investigation.* 2001;21(8):563-569. X-7

2082. Uluocak N, Oktar T, Acar O, et al. Positional changes in voiding dynamics of children with non-neurogenic bladder dysfunction. *Urology*. 2008 Sep;72(3):530-4; discussion 534-5. X-1H, X-4
2083. Unsal A, Saglam R and Cimentepe E. Extracorporeal magnetic stimulation for the treatment of stress and urge incontinence in women--results of 1-year follow-up. *Scand J Urol Nephrol*. 2003;37(5):424-8. X-7
2084. Urwitz-Lane R and Ozel B. Sexual function in women with urodynamic stress incontinence, detrusor overactivity, and mixed urinary incontinence. *Am J Obstet Gynecol*. 2006 Dec;195(6):1758-61. X-8
2085. Ushiroyama T, Ikeda A and Ueki M. Clinical efficacy of clenbuterol and propiverine in menopausal women with urinary incontinence: improvement in quality of life. *J Med*. 2000;31(5-6):311-9. X-7
2086. Vahtera T, Haaranen M, Viramo-Koskela AL, et al. Pelvic floor rehabilitation is effective in patients with multiple sclerosis. *Clin Rehabil*. 1997 Aug;11(3):211-9. X-1F
2087. Valaitis SR and Stanton SL. Sacrocolpopexy: a retrospective study of a clinician's experience. *Br J Obstet Gynaecol*. 1994 Jun;101(6):518-22. X-1E, X-7
2088. Valencic M, Spanjol J, Maricic A, et al. Cystocoele and sensory urgency--our experience. *Coll Antropol*. 2008 Oct;32 Suppl 2:207-9. X-1E
2089. Van Arendonk KJ, Austin JC, Boyt MA, et al. Frequency of wetting is predictive of response to anticholinergic treatment in children with overactive bladder. *Urology*. 2006 May;67(5):1049-53; discussion 1053-4. X-4
2090. van Balen FA, Touw-Otten FW and de Melker RA. Single-dose pefloxacin versus five-days treatment with norfloxacin in uncomplicated cystitis in women. *J Antimicrob Chemother*. 1990 Oct;26 Suppl B:153-60. X-1I
2091. van Balken MR, Vandoninck V, Gisolf KW, et al. Posterior tibial nerve stimulation as neuromodulative treatment of lower urinary tract dysfunction. *J Urol*. 2001 Sep;166(3):914-8. X-7
2092. van Balken MR, Vandoninck V, Messelink BJ, et al. Percutaneous tibial nerve stimulation as neuromodulative treatment of chronic pelvic pain. *Eur Urol*. 2003 Feb;43(2):158-63; discussion 163. X-1A, X-7
2093. van Balken MR, Vergunst H and Bemelmans BL. Prognostic factors for successful percutaneous tibial nerve stimulation. *Eur Urol*. 2006 Feb;49(2):360-5. X-4
2094. van Balken MR, Vergunst H and Bemelmans BL. Sexual functioning in patients with lower urinary tract dysfunction improves after percutaneous tibial nerve stimulation. *Int J Impot Res*. 2006 Sep-Oct;18(5):470-5; discussion 476. X-4
2095. van Brummen HJ, Bruinse HW, van de Pol G, et al. Botherome lower urinary tract symptoms 1 year after first delivery: prevalence and the effect of childbirth. *BJU Int*. 2006 Jul;98(1):89-95. X-1A
2096. Van Brummen HJ, Bruinse HW, Van de Pol G, et al. What is the effect of overactive bladder symptoms on woman's quality of life during and after first pregnancy? *BJU Int*. 2006 Feb;97(2):296-300. X-1I
2097. van Brummen HJ, Bruinse HW, van de Pol G, et al. The effect of vaginal and cesarean delivery on lower urinary tract symptoms: what makes the difference? *Int Urogynecol J Pelvic Floor Dysfunct*. 2007 Feb;18(2):133-9. X-1A
2098. van Brummen HJ, Bruinse HW, van der Bom JG, et al. How do the prevalences of urogenital symptoms change during pregnancy? *Neurourol Urodyn*. 2006;25(2):135-9. X-1I
2099. van Brummen HJ, Heintz AP and van der Vaart CH. The association between overactive bladder symptoms and objective parameters from bladder diary and filling cystometry. *Neurourol Urodyn*. 2004;23(1):38-42. X-1H
2100. van Brummen HJ, van de Pol G, Aalders CI, et al. Sacrospinous hysteropexy compared to vaginal hysterectomy as primary surgical treatment for a descensus uteri: effects on urinary symptoms. *Int Urogynecol J Pelvic Floor Dysfunct*. 2003 Nov;14(5):350-5; discussion 355. X-1E
2101. van der Pal F, van Balken MR, Heesakkers JP, et al. Percutaneous tibial nerve stimulation in the treatment of refractory overactive bladder syndrome: is maintenance treatment necessary? *BJU Int*. 2006 Mar;97(3):547-50. X-7
2102. van der Pal F, van Balken MR, Heesakkers JP, et al. Correlation between quality of life and voiding variables in patients treated with percutaneous tibial nerve stimulation. *BJU Int*. 2006 Jan;97(1):113-6. X-7
2103. van der Vaart CH, de Leeuw JR, Roovers JP, et al. Measuring health-related quality of life in women with urogenital dysfunction: the urogenital distress inventory and incontinence impact questionnaire revisited. *Neurourol Urodyn*. 2003;22(2):97-104. X-1I
2104. van der Vaart CH, van der Bom JG, de Leeuw JR, et al. The contribution of hysterectomy to the occurrence of urge and stress urinary incontinence symptoms. *BJOG*. 2002 Feb;109(2):149-54. X-1A, X-1B
2105. van Gool JD and de Jonge GA. Urge syndrome and urge incontinence. *Arch Dis Child*. 1989 Nov;64(11):1629-34. X-4
2106. Van Kampen M, Bogaert G, Akinwuntan EA, et al. Long-term efficacy and predictive factors of full spectrum therapy for nocturnal enuresis. *J Urol*. 2004 Jun;171(6 Pt 2):2599-602; discussion 2602. X-1C
2107. van Kerrebroeck P, Abrams P, Lange R, et al. Duloxetine versus placebo in the treatment of European and Canadian women with stress urinary incontinence. *BJOG*. 2004 Mar;111(3):249-57. X-1B
2108. Van Kerrebroeck PE, Amarenco G, Thuroff JW, et al. Dose-ranging study of tolterodine in patients with detrusor hyperreflexia. *Neurourol Urodyn*. 1998;17(5):499-512. X-7
2109. Van Laecke E, Wille S, Vande Walle J, et al. The daytime alarm: a useful device for the treatment of children with daytime incontinence. *J Urol*. 2006 Jul;176(1):325-7. X-4
2110. van Meel TD, de Wachter S and Wyndaele JJ. Repeated ice water tests and electrical perception threshold determination to detect a neurologic cause of detrusor overactivity. *Urology*. 2007 Oct;70(4):772-6. X-1H

2111. van Melick HH, Gisolf KW, Eckhardt MD, et al. One 24-hour frequency-volume chart in a woman with objective urinary motor urge incontinence is sufficient. *Urology*. 2001 Aug;58(2):188-92. X-1H
2112. Van Pienbroek E, Hermans J, Kaptein AA, et al. Fosfomycin trometamol in a single dose versus seven days nitrofurantoin in the treatment of acute uncomplicated urinary tract infections in women. *Pharm World Sci*. 1993 Dec 17;15(6):257-62. X-1A
2113. van Venrooij GE and Boon TA. Extensive urodynamic investigation: interaction among diuresis, detrusor instability, urethral relaxation, incontinence and complaints in women with a history of urge incontinence. *J Urol*. 1994 Nov;152(5 Pt 1):1535-8. X-1H
2114. Van Voskuilen AC, Oerlemans DJ, Weil EH, et al. Medium-term experience of sacral neuromodulation by tined lead implantation. *BJU Int*. 2007 Jan;99(1):107-10. X-7
2115. van Waalwijk van Doorn ES and Ambergen AW. Diagnostic assessment of the overactive bladder during the filling phase: the detrusor activity index. *BJU Int*. 1999 Mar;83 Suppl 2:16-21. X-1H
2116. van Waalwijk van Doorn ES, Ambergen AW and Janknegt RA. Detrusor activity index: quantification of detrusor overactivity by ambulatory monitoring. *J Urol*. 1997 Feb;157(2):596-9. X-1H
2117. van Waalwijk van Doorn ES, Remmers A and Janknegt RA. Extramural ambulatory urodynamic monitoring during natural filling and normal daily activities: evaluation of 100 patients. *J Urol*. 1991 Jul;146(1):124-31. X-1H
2118. van Waalwijk van Doorn ES, Remmers A and Janknegt RA. Conventional and extramural ambulatory urodynamic testing of the lower urinary tract in female volunteers. *J Urol*. 1992 May;147(5):1319-25; discussion 1326. X-1H
2119. Vandoninck V, Van Balken MR, Finazzi Agro E, et al. Posterior tibial nerve stimulation in the treatment of urge incontinence. *Neurourol Urodyn*. 2003;22(1):17-23. X-7
2120. Vandoninck V, van Balken MR, Finazzi Agro E, et al. Percutaneous tibial nerve stimulation in the treatment of overactive bladder: urodynamic data. *Neurourol Urodyn*. 2003;22(3):227-32. X-4
2121. Vardan S, Smulyan H, Mookherjee S, et al. Effects of tiodazosin, a new antihypertensive, hemodynamics and clinical variables. *Clin Pharmacol Ther*. 1983 Sep;34(3):290-6. X-1A, X-7
2122. Vasavada SP, Comiter CV and Raz S. Incisionless pubovaginal fascial sling using transvaginal bone anchors for the treatment of stress urinary incontinence. *ScientificWorldJournal*. 2004 Jun 7;4 Suppl 1:357-63. X-1B
2123. Vasavada SP, Rackley RR and Appell RA. In situ anterior vaginal wall sling formation with preservation of the endopelvic fascia for treatment of stress urinary incontinence. *Int Urogynecol J Pelvic Floor Dysfunct*. 1998;9(6):379-84. X-1B, X-7
2124. Velez JB. Behavior therapy for urge incontinence in older women. *J Fam Pract*. 1999 Mar;48(3):168-9. X-2
2125. Vella M, Robinson D, Cardozo L, et al. Predicting detrusor overactivity using a physician-based scoring system. *Int Urogynecol J Pelvic Floor Dysfunct*. 2008 Sep;19(9):1223-7. X-1H
2126. Venn S and Mundy T. Bladder reconstruction: urothelial augmentation, trauma, fistula. *Curr Opin Urol*. 2002 May;12(3):201-3. X-2
2127. Venn SN and Mundy AR. Long-term results of augmentation cystoplasty. *Eur Urol*. 1998;34 Suppl 1:40-2. X-4
2128. Vereecken RL. The after-contraction: a true detrusor contraction or a late dyssynergic urethral sphincter contraction? *BJU Int*. 2000 Feb;85(3):246-8. X-1H
2129. Vereecken RL and Das J. Urethral instability: related to stress and/or urge incontinence? *J Urol*. 1985 Oct;134(4):698-701. X-1G
2130. Vereecken RL and Grisar P. Perineal electromyographic patterns in urge incontinence. *Arch Gynecol*. 1986;237(4):235-42. X-1H
2131. Vereecken RL and Proesmans W. Extensive surgery on the trigone for complete ureteral duplication does not cause incontinence or voiding problems. *Urology*. 2000 Feb;55(2):267-70; discussion 270-1. X-1A, X-4
2132. Vereecken RL and Proesmans W. Urethral instability as an important element of dysfunctional voiding. *J Urol*. 2000 Feb;163(2):585-8. X-1I, X-4
2133. Versi E, Griffiths DJ and Harvey MA. A new external urethral occlusive device for female urinary incontinence. *Obstet Gynecol*. 1998 Aug;92(2):286-91. X-1B
2134. Versi E, Orrego G, Hardy E, et al. Evaluation of the home pad test in the investigation of female urinary incontinence. *Br J Obstet Gynaecol*. 1996 Feb;103(2):162-7. X-1H, X-6
2135. Vervest HA, Kiewiet de Jonge M, Vervest TM, et al. Micturition symptoms and urinary incontinence after non-radical hysterectomy. *Acta Obstet Gynecol Scand*. 1988;67(2):141-6. X-1I
2136. Vervest HA, van Venrooij GE, Barents JW, et al. Non-radical hysterectomy and the function of the lower urinary tract. I: Urodynamic quantification of changes in storage function. *Acta Obstet Gynecol Scand*. 1989;68(3):221-9. X-1I
2137. Videla FL and Wall LL. Stress incontinence diagnosed without multichannel urodynamic studies. *Obstet Gynecol*. 1998 Jun;91(6):965-8. X-1B, X-1H
2138. Viereck V, Bader W, Krauss T, et al. Intra-operative introital ultrasound in Burch colposuspension reduces post-operative complications. *BJOG*. 2005 Jun;112(6):791-6. X-1A
2139. Viereck V, Bader W, Skala C, et al. Determination of bladder neck position by intraoperative introital ultrasound in colposuspension: outcome at 6-month follow-up. *Ultrasound Obstet Gynecol*. 2004 Aug;24(2):186-91. X-1B
2140. Viereck V, Pauer HU, Bader W, et al. Introital ultrasound of the lower genital tract before and after colposuspension: a 4-year objective follow-up. *Ultrasound Obstet Gynecol*. 2004 Mar;23(3):277-83. X-1B, X-1H

2141. Viereck V, Pauer HU, Hesse O, et al. Urethral hypermobility after anti-incontinence surgery - a prognostic indicator? *Int Urogynecol J Pelvic Floor Dysfunct.* 2006 Nov;17(6):586-92. X-1B
2142. Vierhout ME and Mulder AF. De novo detrusor instability after Burch colposuspension. *Acta Obstet Gynecol Scand.* 1992 Aug;71(6):414-6. X-1A
2143. Vijverberg MA, Elzinga-Plomp A, Messer AP, et al. Bladder rehabilitation, the effect of a cognitive training programme on urge incontinence. *Eur Urol.* 1997;31(1):68-72. X-4
2144. Viktrup L. The risk of lower urinary tract symptoms five years after the first delivery. *Neurourol Urodyn.* 2002;21(1):2-29. X-6
2145. Viktrup L and Lose G. Lower urinary tract symptoms 5 years after the first delivery. *Int Urogynecol J Pelvic Floor Dysfunct.* 2000 Dec;11(6):336-40. X-6
2146. Vine MF, Calingaert B, Berchuck A, et al. Characterization of prediagnostic symptoms among primary epithelial ovarian cancer cases and controls. *Gynecol Oncol.* 2003 Jul;90(1):75-82. X-1A
2147. Vinson RK and Diokno AC. Uninhibited neurogenic bladder in adults. *Urology.* 1976 Apr;7(4):376-8. X-1F
2148. Virtanen H, Makinen J, Tenho T, et al. Effects of abdominal hysterectomy on urinary and sexual symptoms. *Br J Urol.* 1993 Dec;72(6):868-72. X-1I
2149. Visco AG, Weidner AC, Cundiff GW, et al. Observed patient compliance with a structured outpatient bladder retraining program. *Am J Obstet Gynecol.* 1999 Dec;181(6):1392-4. X-1I
2150. Viseshsindh W, Kochakarn W, Waikakul W, et al. A randomized controlled trial of pubovaginal sling versus vaginal wall sling for stress urinary incontinence. *J Med Assoc Thai.* 2003 Apr;86(4):308-15. X-1B
2151. von Gontard A, Lettgen B, Olbing H, et al. Behavioural problems in children with urge incontinence and voiding postponement: a comparison of a paediatric and child psychiatric sample. *Br J Urol.* 1998 May;81 Suppl 3:100-6. X-4
2152. von Gontard A, Mauer-Mucke K, Pluck J, et al. Clinical behavioral problems in day- and night-wetting children. *Pediatr Nephrol.* 1999 Oct;13(8):662-7. X-1I, X-4
2153. Voorham-van der Zalm PJ, Pelger RC, Stiggelbout AM, et al. Effects of magnetic stimulation in the treatment of pelvic floor dysfunction. *BJU Int.* 2006 May;97(5):1035-8. X-7
2154. Wada Y, Ando Y, Kai N, et al. Lower urinary tract dysfunction in type 1 familial amyloidotic polyneuropathy in Kumamoto, Japan. *Int J Urol.* 2006 Dec;13(12):1475-8. X-1F
2155. Wadie BS. Primary nocturnal enuresis persistent to adulthood, functional evaluation. *Neurourol Urodyn.* 2004;23(1):54-7. X-1I
2156. Wadie BS, Edwan A and Nabeeh AM. Autologous fascial sling vs polypropylene tape at short-term followup: a prospective randomized study. *J Urol.* 2005 Sep;174(3):990-3. X-1B
2157. Wagener N, Reitz A, Pfitzenmaier J, et al. Adherens junctions of the human detrusor. *BJU Int.* 2005 Oct;96(6):843-7. X-1G, X-7
2158. Wagg A, Bayliss M, Ingham NJ, et al. Urodynamic variables cannot be used to classify the severity of detrusor instability. *Br J Urol.* 1998 Oct;82(4):499-502. X-1I
2159. Wagg A, Wyndaele JJ and Sieber P. Efficacy and tolerability of solifenacin in elderly subjects with overactive bladder syndrome: a pooled analysis. *Am J Geriatr Pharmacother.* 2006 Mar;4(1):14-24. X-2
2160. Wagg AS, Lieu PK, Ding YY, et al. A urodynamic analysis of age associated changes in urethral function in women with lower urinary tract symptoms. *J Urol.* 1996 Dec;156(6):1984-8. X-1G
2161. Wagner TH and Hu TW. Economic costs of urinary incontinence in 1995. *Urology.* 1998 Mar;51(3):355-61. X-1I
2162. Walker J and Rickwood AM. Daytime urinary frequency in children. *BMJ.* 1988 Aug 13;297(6646):455. X-4
2163. Wall LL, Copas P and Galloway NT. Use of a pedicled rectus abdominis muscle flap sling in the treatment of complicated stress urinary incontinence. *Am J Obstet Gynecol.* 1996 Dec;175(6):1460-4; discussion 1464-6. X-1B
2164. Wall LL and Hewitt JK. Urodynamic characteristics of women with complete posthysterectomy vaginal vault prolapse. *Urology.* 1994 Sep;44(3):336-41; discussion 341-2. X-1E, X-1G
2165. Wall LL and Stanton SL. Transvesical phenol injection of pelvic nerve plexuses in females with refractory urge incontinence. *Br J Urol.* 1989 May;63(5):465-8. X-7
2166. Wall LL, Wiskind AK and Taylor PA. Simple bladder filling with a cough stress test compared with subtracted cystometry for the diagnosis of urinary incontinence. *Am J Obstet Gynecol.* 1994 Dec;171(6):1472-7; discussion 1477-9. X-1H
2167. Wallwiener D, Grischke EM, Rimbach S, et al. Endoscopic retropubic colposuspension: "Retziusscopy" versus laparoscopy--a reasonable enlargement of the operative spectrum in the management of recurrent stress incontinence? *Endosc Surg Allied Technol.* 1995 Apr-Jun;3(2-3):115-8. X-1B
2168. Walsh IK, Nambirajan T, Donellan SM, et al. Cadaveric fascia lata pubovaginal slings: early results on safety, efficacy and patient satisfaction. *BJU Int.* 2002 Sep;90(4):415-9. X-1B, X-7
2169. Walter S. Detrusor hyperreflexia in female urinary incontinence treated pharmacologically. *Urol Int.* 1978;33(5):316-21. X-1F
2170. Walter S, Meyhoff HH, Gerstenberg T, et al. Urinary incontinence in the female. A long-term study of the effect of anticholinergics on overactive detrusor function. *Acta Obstet Gynecol Scand.* 1984;63(2):159-61. X-9
2171. Walter S and Olesen KP. Urinary incontinence and genital prolapse in the female: clinical, urodynamic and radiological examinations. *Br J Obstet Gynaecol.* 1982 May;89(5):393-401. X-1E
2172. Walter U, Dressler D, Wolters A, et al. Overactive bladder in Parkinson's disease: alteration of brainstem raphe detected by transcranial sonography. *Eur J Neurol.* 2006 Dec;13(12):1291-7. X-1F

2173. Walters MD and Shields LE. The diagnostic value of history, physical examination, and the Q-tip cotton swab test in women with urinary incontinence. *Am J Obstet Gynecol.* 1988 Jul;159(1):145-9. X-1H
2174. Walters MD, Taylor S and Schoenfeld LS. Psychosexual study of women with detrusor instability. *Obstet Gynecol.* 1990 Jan;75(1):22-6. X-1I, X-7
2175. Wan EC, Gordon TP and Jackson MW. Autoantibodies to calcium channels in type 1 diabetes mediate autonomic dysfunction by different mechanisms in colon and bladder and are neutralized by antiidiotypic antibodies. *J Autoimmun.* 2008 Aug;31(1):66-72. X-1G
2176. Wang AC. Burch colposuspension vs. Stamey bladder neck suspension. A comparison of complications with special emphasis on detrusor instability and voiding dysfunction. *J Reprod Med.* 1996 Jul;41(7):529-33. X-1B
2177. Wang AC. Bladder-sphincter biofeedback as treatment of detrusor instability in women who failed to respond to oxybutynin. *Chang Gung Med J.* 2000 Oct;23(10):590-9. X-7
2178. Wang AC, Wu RC, Lin CT, et al. A microbiological and immunohistochemical analysis of periurethral and vaginal tissue in women with de novo urge symptoms after mid-urethral sling procedures--a prospective case-controlled study. *Int Urogynecol J Pelvic Floor Dysfunct.* 2008 Aug;19(8):1145-50. X-1A
2179. Wang HZ, Brink PR and Christ GJ. Gap junction channel activity in short-term cultured human detrusor myocyte cell pairs: gating and unitary conductances. *Am J Physiol Cell Physiol.* 2006 Dec;291(6):C1366-76. X-1G
2180. Wang KH, Neimark M and Davila GW. Voiding dysfunction following TVT procedure. *Int Urogynecol J Pelvic Floor Dysfunct.* 2002 Nov;13(6):353-7; discussion 358. X-1B
2181. Wanigasuriya JK, Wijewickrama ES, Jayasekara WM, et al. Lower urinary tract symptoms; an undetected cause for morbidity in adults. *Ceylon Med J.* 2007 Sep;52(3):86-9. X-4
2182. Warne SA, Godley ML and Wilcox DT. Surgical reconstruction of cloacal malformation can alter bladder function: a comparative study with anorectal anomalies. *J Urol.* 2004 Dec;172(6 Pt 1):2377-81; discussion 2381. X-1I
2183. Warren JW. Is interstitial cystitis an infectious disease? *Med Hypotheses.* 1994 Sep;43(3):183-6. X-1D, X-2, X-6, X-7
2184. Washington BB and Hines BJ. Implant infection after two-stage sacral nerve stimulator placement. *Int Urogynecol J Pelvic Floor Dysfunct.* 2007 Dec;18(12):1477-80. X-7
2185. Watanabe T, Yokoyama T, Sasaki K, et al. Intravesical resiniferatoxin for patients with neurogenic detrusor overactivity. *Int J Urol.* 2004 Apr;11(4):200-5. X-1F, X-7
2186. Wattanayingcharoenchai R, Manonai J, Vannatim N, et al. Impact of stress urinary incontinence and overactive bladder on quality of life in Thai women attending the urogynecology clinic. *J Med Assoc Thai.* 2007 Jan;90(1):26-31. X-1I
2187. Webb RJ, Ramsden PD and Neal DE. Ambulatory monitoring and electronic measurement of urinary leakage in the diagnosis of detrusor instability and incontinence. *Br J Urol.* 1991 Aug;68(2):148-52. X-1H
2188. Weber AM, Abrams P, Brubaker L, et al. The standardization of terminology for researchers in female pelvic floor disorders. *Int Urogynecol J Pelvic Floor Dysfunct.* 2001;12(3):178-86. X-2
2189. Weber AM and Walters MD. Burch procedure compared with sling for stress urinary incontinence: a decision analysis. *Obstet Gynecol.* 2000 Dec;96(6):867-73. X-1B
2190. Weber AM and Walters MD. Cost-effectiveness of urodynamic testing before surgery for women with pelvic organ prolapse and stress urinary incontinence. *Am J Obstet Gynecol.* 2000 Dec;183(6):1338-46; discussion 1346-7. X-1B, X-1E
2191. Webster DC. Interstitial cystitis: women at risk for psychiatric misdiagnosis. *AWHONNS Clin Issues Perinat Womens Health Nurs.* 1993;4(2):236-43. X-1D
2192. Webster DC. Recontextualizing sexuality in chronic illness: women and interstitial cystitis. *Health Care Women Int.* 1997 Nov-Dec;18(6):575-89. X-1D
2193. Webster DC and Brennan T. Use and effectiveness of physical self-care strategies for interstitial cystitis. *Nurse Pract.* 1994 Oct;19(10):55-61. X-1D
2194. Webster GD, Koefoot RB, Jr. and Sihelnik S. Urodynamic abnormalities in neurologically normal children with micturition dysfunction. *J Urol.* 1984 Jul;132(1):74-7. X-1G, X-4
2195. Weerasekera DS and Gunawardena KK. The Stamey procedure: a retrospective analysis of clinical outcome in stress incontinence. *Ceylon Med J.* 1995 Sep;40(3):101-2. X-1B
2196. Weerasinghe N and Malone PS. The value of videourodynamics in the investigation of neurologically normal children who wet. *Br J Urol.* 1993 May;71(5):539-42. X-1H, X-4
2197. Weese DL, Roskamp DA, Leach GE, et al. Intravesical oxybutynin chloride: experience with 42 patients. *Urology.* 1993 Jun;41(6):527-30. X-7
2198. Weidner AC, Myers ER, Visco AG, et al. Which women with stress incontinence require urodynamic evaluation? *Am J Obstet Gynecol.* 2001 Jan;184(2):20-7. X-1B
2199. Weil A, Miege B, Rottenberg R, et al. Clinical significance of urethral instability. *Obstet Gynecol.* 1986 Jul;68(1):106-10. X-1I
2200. Weil EH, Ruiz-Cerda JL, Eerdmans PH, et al. Sacral root neuromodulation in the treatment of refractory urinary urge incontinence: a prospective randomized clinical trial. *Eur Urol.* 2000 Feb;37(2):161-71. X-7
2201. Weil EH, Ruiz-Cerda JL, Eerdmans PH, et al. Clinical results of sacral neuromodulation for chronic voiding dysfunction using unilateral sacral foramen electrodes. *World J Urol.* 1998;16(5):313-21. X-7
2202. Wein AJ. Overactive bladder: defining the disease. *Am J Manag Care.* 2000 Jul;6(11 Suppl):S559-64; discussion S607-19. X-2
2203. Wein AJ. Urgency: the cornerstone symptom of overactive bladder. *J Urol.* 2005 Jun;173(6):2054. X-2
2204. Wein AJ, Khullar V, Wang JT, et al. Achieving continence with antimuscarinic therapy for overactive bladder: effects of baseline incontinence severity and bladder diary duration. *BJU Int.* 2007 Feb;99(2):360-3. X-2

2205. Wein AJ and Rovner ES. Definition and epidemiology of overactive bladder. *Urology*. 2002 Nov;60(5 Suppl 1):7-12; discussion 12. X-2, X-6
2206. Weinberger MW, Goodman BM and Carnes M. Long-term efficacy of nonsurgical urinary incontinence treatment in elderly women. *J Gerontol A Biol Sci Med Sci*. 1999 Mar;54(3):M117-21. X-7
2207. Weinberger MW and Ostergard DR. Long-term clinical and urodynamic evaluation of the polytetrafluoroethylene suburethral sling for treatment of genuine stress incontinence. *Obstet Gynecol*. 1995 Jul;86(1):92-6. X-1B
2208. Weiss GR, O'Donnell MA, Loughlin K, et al. Phase 1 study of the intravesical administration of recombinant human interleukin-12 in patients with recurrent superficial transitional cell carcinoma of the bladder. *J Immunother*. 2003 Jul-Aug;26(4):343-8. X-1A, X-7
2209. Weiss JP, Blaivas JG, Jones M, et al. Age related pathogenesis of nocturia in patients with overactive bladder. *J Urol*. 2007 Aug;178(2):548-51; discussion 551. X-II
2210. Weiss JP, Blaivas JG, Stember DS, et al. Nocturia in adults: etiology and classification. *Neurourol Urodyn*. 1998;17(5):467-72. X-1C
2211. Wells TJ, Brink CA and Diokno AC. Urinary incontinence in elderly women: clinical findings. *J Am Geriatr Soc*. 1987 Oct;35(10):933-9. X-6
2212. Wen JG and Tong EC. Cystometry in infants and children with no apparent voiding symptoms. *Br J Urol*. 1998 Mar;81(3):468-73. X-1H, X-4, X-7
2213. Wen JG, Yeung CK, Chu WC, et al. Video cystometry in young infants with renal dilation or a history of urinary tract infection. *Urol Res*. 2001 Aug;29(4):249-55. X-1G, X-4
2214. Wennberg AL, Edlund C, Fall M, et al. Stamey's abdominovaginal needle colposuspension for the correction of female genuine stress urinary incontinence--long-term results. *Scand J Urol Nephrol*. 2003;37(5):419-23. X-1B
2215. Werner M, Schmid DM and Schussler B. Efficacy of botulinum-A toxin in the treatment of detrusor overactivity incontinence: a prospective nonrandomized study. *Am J Obstet Gynecol*. 2005 May;192(5):1735-40. X-7
2216. Wester C, FitzGerald MP, Brubaker L, et al. Validation of the clinical bulbocavernosus reflex. *Neurourol Urodyn*. 2003;22(6):589-91; discussion 591-2. X-1H
2217. Westney OL, Amundsen CL and McGuire EJ. Bladder endometriosis: conservative management. *J Urol*. 2000 Jun;163(6):1814-7. X-1A, X-7
2218. Westney OL, Lee JT, McGuire EJ, et al. Long-term results of Ingelman-Sundberg denervation procedure for urge incontinence refractory to medical therapy. *J Urol*. 2002 Sep;168(3):1044-7. X-7
2219. Wheelahan JB. The pressure-volume plot and prediction of treatment outcome in female incontinence. *Br J Urol*. 1986 Oct;58(5):514-9. X-1H
2220. Wheelahan JB. Long-term results of colposuspension. *Br J Urol*. 1990 Apr;65(4):329-32. X-1B
2221. Whelan CM and McKenna PH. Urologic applications of biofeedback therapy. *Contemporary Urology*. 2004;16(10):23. X-2
2222. White WM, Pickens RB, Doggweiler R, et al. Short-term efficacy of botulinum toxin a for refractory overactive bladder in the elderly population. *J Urol*. 2008 Dec;180(6):2522-6. X-7
2223. Whiteside JL, Hijaz A, Imrey PB, et al. Reliability and agreement of urodynamics interpretations in a female pelvic medicine center. *Obstet Gynecol*. 2006 Aug;108(2):315-23. X-1H
2224. Whitler ET, Feltner F, Owens J, et al. Kentucky Homeplace Defeat Diabetes Screening Test: an analysis of rural Kentucky's challenge to overcome the growing diabetes epidemic. *J Ky Med Assoc*. 2005 Jul;103(7):307-14. X-II
2225. Wiener JS, Scales MT, Hampton J, et al. Long-term efficacy of simple behavioral therapy for daytime wetting in children. *J Urol*. 2000 Sep;164(3 Pt 1):786-90. X-4
2226. Wiener LB, Baum NH and Suarez GM. New method for management of detrusor instability: transdermal scopolamine. *Urology*. 1986 Sep;28(3):208-10. X-7
2227. Wijma J, Tinga DJ and Visser GH. Compensatory mechanisms which prevent urinary incontinence in aging women. *Gynecol Obstet Invest*. 1992;33(2):102-4. X-1B, X-1H, X-7
2228. Wilkinson E. Urinary incontinence in women. *Nurs Times*. 2006 Nov 28-Dec 4;102(48):23-4. X-2
2229. Wilson S, Quek ML and Ginsberg DA. Transurethral injection of bulking agents for stress urinary incontinence following orthotopic neobladder reconstruction in women. *J Urol*. 2004 Jul;172(1):244-6. X-1B
2230. Winder A. Antimuscarinic management of overactive bladder. *Br J Community Nurs*. 2005 Jul;10(7):313-6. X-2
2231. Winge K, Friberg L, Werdelin L, et al. Relationship between nigrostriatal dopaminergic degeneration, urinary symptoms, and bladder control in Parkinson's disease. *Eur J Neurol*. 2005 Nov;12(11):842-50. X-1F
2232. Winge K, Nielsen KK, Stimpel H, et al. Lower urinary tract symptoms and bladder control in advanced Parkinson's disease: effects of deep brain stimulation in the subthalamic nucleus. *Mov Disord*. 2007 Jan 15;22(2):220-5. X-1F
2233. Winge K, Skau AM, Stimpel H, et al. Prevalence of bladder dysfunction in Parkinson's disease. *Neurourol Urodyn*. 2006;25(2):116-22. X-1F
2234. Winiiecka W, Zoch-Zwiercz W, Wasilewska A, et al. Evaluation of bladder instability in children with recurrent urinary tract infections. *Med Sci Monit*. 2002 Jan;8(1):CR19-23. X-1I, X-4
2235. Winkler HA and Sand PK. Treatment of detrusor instability with oxybutynin rectal suppositories. *Int Urogynecol J Pelvic Floor Dysfunct*. 1998;9(2):100-2. X-7
2236. Winters JC and Appell RA. Collagen injection therapy in the treatment of urinary incontinence. *Tech Urol*. 1996 Summer;2(2):59-64. X-II
2237. Wise BG and Cardozo LD. Urinary urgency in women. *Br J Hosp Med*. 1993 Sep 1-14;50(5):243-50. X-2
2238. Wise BG, Cardozo LD, Cutner A, et al. Prevalence and significance of urethral instability in women with detrusor instability. *Br J Urol*. 1993 Jul;72(1):26-9. X-1G

2239. Wiseman PA, Malone-Lee J and Rai GS. Terodiline with bladder retraining for treating detrusor instability in elderly people. *BMJ*. 1991 Apr 27;302(6783):994-6. X-7
2240. Wiskind AK, Miller KF and Wall LL. One hundred unstable bladders. *Obstet Gynecol*. 1994 Jan;83(1):108-12. X-1H
2241. Wong T, Lau BY, Mak HL, et al. Changing prevalence and knowledge of urinary incontinence among Hong Kong Chinese women. *Int Urogynecol J Pelvic Floor Dysfunct*. 2006 Nov;17(6):593-7. X-6E
2242. Woodman PJ, Misko CA and Fischer JR. The use of short-form quality of life questionnaires to measure the impact of imipramine on women with urge incontinence. *Int Urogynecol J Pelvic Floor Dysfunct*. 2001;12(5):312-5; discussion 315-6. X-7
2243. Woodtli A. Stress incontinence: clinical identification and validation of defining characteristics. *Nurs Diagn*. 1995 Jul-Sep;6(3):115-22. X-1B
2244. Woodtli MA. Assessing urge incontinence in elderly women. *Geriatr Nurs*. 1993 Jan-Feb;14(1):19-22. X-1H, X-7
2245. Wraige E and Borzyskowski M. Investigation of daytime wetting: when is spinal cord imaging indicated? *Arch Dis Child*. 2002 Aug;87(2):151-5. X-1H, X-4
2246. Wu CC, Huang MY, Kapoor R, et al. Metabolism of omega-6 polyunsaturated fatty acids in women with dysmenorrhea. *Asia Pac J Clin Nutr*. 2008;17 Suppl 1:216-9. X-11
2247. Wu EQ, Birnbaum H, Kang YJ, et al. A retrospective claims database analysis to assess patterns of interstitial cystitis diagnosis. *Curr Med Res Opin*. 2006 Mar;22(3):495-500. X-1D
2248. Wu EQ, Birnbaum H, Marynchenko M, et al. Employees with overactive bladder: work loss burden. *J Occup Environ Med*. 2005 May;47(5):439-46. X-11
2249. Wu JM, Williams KS, Hundley AF, et al. Microscopic hematuria as a predictive factor for detecting bladder cancer at cystoscopy in women with irritative voiding symptoms. *Am J Obstet Gynecol*. 2006 May;194(5):1423-6. X-11
2250. Wu WJ, Chen MT, Huang CN, et al. Experience in the morbidity associated with double-J catheter indwelling and its management. *Gaoxiong Yi Xue Ke Xue Za Zhi*. 1993 Sep;9(9):532-9. X-11
2251. Wujanto R and O'Reilly PH. Stamey needle suspension for stress urinary incontinence. A prospective study of 40 patients. *Br J Urol*. 1989 Feb;63(2):162-4. X-1B
2252. Wyman JF, Choi SC, Harkins SW, et al. The urinary diary in evaluation of incontinent women: a test-retest analysis. *Obstet Gynecol*. 1988 Jun;71(6 Pt 1):812-7. X-11
2253. Wyman JF, Elswick RK, Jr., Ory MG, et al. Influence of functional, urological, and environmental characteristics on urinary incontinence in community-dwelling older women. *Nurs Res*. 1993 Sep-Oct;42(5):270-5. X-11
2254. Wyman JF and Fantl JA. Bladder training in ambulatory care management of urinary incontinence. *Urol Nurs*. 1991 Sep;11(3):11-7. X-2
2255. Wyman JF, Fantl JA, McClish DK, et al. Quality of life following bladder training in older women with urinary incontinence. *Int Urogynecol J Pelvic Floor Dysfunct*. 1997;8(4):223-9. X-7
2256. Wyman JF, Harkins SW, Choi SC, et al. Psychosocial impact of urinary incontinence in women. *Obstet Gynecol*. 1987 Sep;70(3 Pt 1):378-81. X-7
2257. Wyndaele JJ, Van Meel TD and De Wachter S. Detrusor overactivity. Does it represent a difference if patients feel the involuntary contractions? *J Urol*. 2004 Nov;172(5 Pt 1):1915-8. X-1G, X-1H
2258. Yalcin I, Versi E, Benson JT, et al. Validation of a clinical algorithm to diagnose stress urinary incontinence for large studies. *J Urol*. 2004 Jun;171(6 Pt 1):2321-5. X-1B
2259. Yalcin OT, Hassa H and Sarac I. Short-term intravaginal maximal electrical stimulation for refractive detrusor instability. *Int J Gynaecol Obstet*. 2002 Dec;79(3):241-4. X-7
2260. Yalcin OT, Yildirim A and Hassa H. The effects of severe cystocele on urogynecologic symptoms and findings. *Acta Obstet Gynecol Scand*. 2001 May;80(5):423-7. X-11
2261. Yamada T, Arai G, Masuda H, et al. The correction of type 2 stress incontinence with a polytetrafluoroethylene patch sling: 5-year mean followup. *J Urol*. 1998 Sep;160(3 Pt 1):746-9. X-1B, X-7
2262. Yamada T, Nishimura M and Mita H. Increased number of apoptotic endothelial cells in bladder of interstitial cystitis patients. *World J Urol*. 2007 Aug;25(4):407-13. X-1D, X-1G, X-7
2263. Yamaguchi C, Sakakibara R, Uchiyama T, et al. Bladder sensation in peripheral nerve lesions. *Neurourol Urodyn*. 2006;25(7):763-9. X-1H
2264. Yamamoto T, Sakakibara R, Uchiyama T, et al. When is Onuf's nucleus involved in multiple system atrophy? A sphincter electromyography study. *J Neurol Neurosurg Psychiatry*. 2005 Dec;76(12):1645-8. X-1F
2265. Yamamoto T, Sakakibara R, Uchiyama T, et al. Neurological diseases that cause detrusor hyperactivity with impaired contractile function. *Neurourol Urodyn*. 2006;25(4):356-60. X-1F
2266. Yamanishi T, Sakakibara R, Uchiyama T, et al. Comparative study of the effects of magnetic versus electrical stimulation on inhibition of detrusor overactivity. *Urology*. 2000 Nov 1;56(5):777-81. X-7
2267. Yamanishi T, Yasuda K, Hamano S, et al. Urethral obstruction in patients with nighttime wetting: urodynamic evaluation and outcome of surgical incision. *Neurourol Urodyn*. 2000;19(3):241-8. X-11, X-4
2268. Yamanishi T, Yasuda K, Murayama N, et al. Biofeedback training for detrusor overactivity in children. *J Urol*. 2000 Nov;164(5):1686-90. X-4, X-7
2269. Yamanishi T, Yasuda K, Sakakibara R, et al. Randomized, double-blind study of electrical stimulation for urinary incontinence due to detrusor overactivity. *Urology*. 2000 Mar;55(3):353-7. X-7
2270. Yamanishi T, Yasuda K, Sakakibara R, et al. Detrusor overactivity and penile erection in patients with lower lumbar spine lesions. *Eur Urol*. 1998 Oct;34(4):360-4. X-1F, X-4, X-5, X-6, X-7

2271. Yamanishi T, Yasuda K, Suda S, et al. Effect of functional continuous magnetic stimulation for urinary incontinence. *J Urol*. 2000 Feb;163(2):456-9. X-7
2272. Yamao Y, Kawauchi A, Tanaka Y, et al. Systematic treatment for nocturnal urinary frequency following a sleep-micturition chart. *Psychiatry Clin Neurosci*. 1999 Apr;53(2):277-8. X-1C
2273. Yamashita H and Kumazawa J. Voiding dysfunction: patients with human T-lymphotropic-virus-type-1-associated myelopathy. *Urol Int*. 1991;47 Suppl 1:69-71. X-1F
2274. Yang JM and Huang WC. Discrimination of bladder disorders in female lower urinary tract symptoms on ultrasonographic cystourethrography. *J Ultrasound Med*. 2002 Nov;21(11):1249-55. X-1H
2275. Yang JM and Huang WC. Implications of abdominal straining in women with lower urinary tract symptoms. *Urology*. 2002 Sep;60(3):428-33. X-1A, X-1G, X-1H
2276. Yang JM and Huang WC. Bladder wall thickness on ultrasonographic cystourethrography: affecting factors and their implications. *J Ultrasound Med*. 2003 Aug;22(8):777-82. X-1G
2277. Yang JM, Yang SH and Huang WC. A surgical technique to adjust bladder neck suspension in laparoscopic Burch colposuspension. *J Minim Invasive Gynecol*. 2006 Jul-Aug;13(4):289-95. X-1B
2278. Yantiss RK, Clement PB and Young RH. Endometriosis of the intestinal tract: a study of 44 cases of a disease that may cause diverse challenges in clinical and pathologic evaluation. *Am J Surg Pathol*. 2001 Apr;25(4):445-54. X-1I
2279. Yeung CK, Godley ML, Duffy PG, et al. Natural filling cystometry in infants and children. *Br J Urol*. 1995 Apr;75(4):531-7. X-1G, X-H, X-4
2280. Yeung CK, Godley ML, Ho CK, et al. Some new insights into bladder function in infancy. *Br J Urol*. 1995 Aug;76(2):235-40. X-1I, X-4
2281. Yeung CK, Sihoe JD, Sit FK, et al. Urodynamic findings in adults with primary nocturnal enuresis. *J Urol*. 2004 Jun;171(6 Pt 2):2595-8. X-1I
2282. Yeung CK, Sit FK, To LK, et al. Reduction in nocturnal functional bladder capacity is a common factor in the pathogenesis of refractory nocturnal enuresis. *BJU Int*. 2002 Aug;90(3):302-7. X-1I, X-4
2283. Yeung CK, Sreedhar B, Leung YF, et al. Correlation between ultrasonographic bladder measurements and urodynamic findings in children with recurrent urinary tract infection. *BJU Int*. 2007 Mar;99(3):651-5. X-1G, X-4
2284. Yiangou Y, Facer P, Ford A, et al. Capsaicin receptor VR1 and ATP-gated ion channel P2X3 in human urinary bladder. *BJU Int*. 2001 Jun;87(9):774-9. X-1G, X-7
2285. Yip SK, Chan A, Pang S, et al. The impact of urodynamic stress incontinence and detrusor overactivity on marital relationship and sexual function. *Am J Obstet Gynecol*. 2003 May;188(5):1244-8. X-7
2286. Yip SK and Pang MW. Tension-free vaginal tape sling procedure for the treatment of stress urinary incontinence in Hong Kong women with and without pelvic organ prolapse: 1-year outcome study. *Hong Kong Med J*. 2006 Feb;12(1):15-20. X-1B
2287. Yip SK, Sahota D, Chang AM, et al. Four-year follow-up of women who were diagnosed to have postpartum urinary retention. *Am J Obstet Gynecol*. 2002 Sep;187(3):648-52. X-1A
2288. Yokoyama T, Fujita O, Nishiguchi J, et al. Extracorporeal magnetic innervation treatment for urinary incontinence. *Int J Urol*. 2004 Aug;11(8):602-6. X-7
2289. Yokoyama T, Kumon H and Nagai A. Correlation of urinary nerve growth factor level with pathogenesis of overactive bladder. *NeuroUrol Urodyn*. 2008;27(5):417-20. X-1G, X-7
2290. Yokoyama T, Nozaki K, Fujita O, et al. Role of C afferent fibers and monitoring of intravesical resiniferatoxin therapy for patients with idiopathic detrusor overactivity. *J Urol*. 2004 Aug;172(2):596-600. X-7
2291. Yono M, Yohida M, Wada Y, et al. Pharmacological effects of tolterodine on human isolated urinary bladder. *Eur J Pharmacol*. 1999 Mar 5;368(2-3):223-30. X-1G
2292. Yoon HS, Song HH and Ro YJ. A comparison of effectiveness of bladder training and pelvic muscle exercise on female urinary incontinence. *Int J Nurs Stud*. 2003 Jan;40(1):45-50. X-1B
2293. Yoon SM, Jung JK, Lee SB, et al. Treatment of female urethral syndrome refractory to antibiotics. *Yonsei Med J*. 2002 Oct;43(5):644-51. X-1I
2294. Yoshimura K and Terai A. Classification and distribution of symptomatic nocturia with special attention to duration of time in bed: a patient-based study. *BJU Int*. 2005 Jun;95(9):1259-62. X-1C
2295. Yossepowitch O, Gillon G, Baniel J, et al. The effect of cholinergic enhancement during filling cystometry: can edrophonium chloride be used as a provocative test for overactive bladder? *J Urol*. 2001 May;165(5):1441-5. X-1H, X-7
2296. Young AE, Fine PM, McCrery R, et al. Spanish language translation of pelvic floor disorders instruments. *Int Urogynecol J Pelvic Floor Dysfunct*. 2007 Oct;18(10):1171-8. X-1I
2297. Young SB, Howard AE and Baker SP. Mersilene mesh sling: short- and long-term clinical and urodynamic outcomes. *Am J Obstet Gynecol*. 2001 Jul;185(1):32-40. X-1B
2298. Yu HJ, Lee WC, Lee KL, et al. Voiding dysfunction in women with systemic lupus erythematosus. *Arthritis Rheum*. 2004 Jan;50(1):166-72. X-1I
2299. Yu LC, Rohner TJ, Kaltreider DL, et al. Profile of urinary incontinent elderly in long-term care institutions. *J Am Geriatr Soc*. 1990 Apr;38(4):433-9. X-5
2300. Yu YF, Nichol MB, Yu AP, et al. Persistence and adherence of medications for chronic overactive bladder/urinary incontinence in the California Medicaid program. *Value Health*. 2005 Jul-Aug;8(4):495-505. X-1I
2301. Yucel S, Akkaya E, Guntekin E, et al. Can alpha-blocker therapy be an alternative to biofeedback for dysfunctional voiding and urinary retention? A prospective study. *J Urol*. 2005 Oct;174(4 Pt 2):1612-5; discussion 1615. X-4, X-7
2302. Yucel S, Akkaya E, Guntekin E, et al. Should we switch over to tolterodine in every child with non-neurogenic daytime urinary incontinence in whom oxybutynin failed? *Urology*. 2005 Feb;65(2):369-73. X-4

2303. Yucel S, Kutlu O, Kukul E, et al. Impact of urodynamics in treatment of primary nocturnal enuresis persisting into adulthood. *Urology*. 2004 Nov;64(5):1020-5; discussion 1025. X-1C, X-7
2304. Zachoval R, Pitha J, Medova E, et al. Augmentation cystoplasty in patients with multiple sclerosis. *Urol Int*. 2003;70(1):21-6; discussion 26. X-1F
2305. Zahariou A, Karamouti M, Karagiannis G, et al. Maximal bladder capacity is a positive predictor of response to desmopressin treatment in patients with MS and nocturia. *Int Urol Nephrol*. 2008;40(1):65-9. X-1F
2306. Zajackowska M, Moulhee NM, Piechuta L, et al. Dysfunctional voiding and urodynamic disorders in children with recurrent urinary tract infection. *Ann Univ Mariae Curie Sklodowska [Med]*. 2004;59(2):385-91. X-1I, X-4
2307. Zanoschi C and Carauleanu A. The use of the suprapubic fibrous complex in isthmus hysteropexies. *Rev Med Chir Soc Med Nat Iasi*. 2003 Jul-Sep;107(3):627-32. X-1E
2308. Zarina B, Juwita S and Nor GRM. Prevalence and factors associated with urinary incontinence in adult women attending family medicine clinic. *International Medical Journal*. 2005 Dec;12(4):303-10. X-6
2309. Zeidman EJ, Helfrick B, Pollard C, et al. Bacillus Calmette-Guerin immunotherapy for refractory interstitial cystitis. *Urology*. 1994 Jan;43(1):121-4. X-1D
2310. Zeng Y, Wu XX, Homma Y, et al. Uroplakin III-delta4 messenger RNA as a promising marker to identify nonulcerative interstitial cystitis. *J Urol*. 2007 Oct;178(4 Pt 1):1322-7; discussion 1327. X-1D
2311. Zermann DH, Löffler U, Reichelt O, et al. Bladder dysfunction and end stage renal disease. *Int Urol Nephrol*. 2003;35(1):93-7. X-1I
2312. Zhang CO, Li ZL and Kong CZ. APF, HB-EGF, and EGF biomarkers in patients with ulcerative vs. non-ulcerative interstitial cystitis. *BMC Urol*. 2005;5:7. X-1D
2313. Zhang J, Haselden P and Tepe R. A case series of reduced urinary incontinence in elderly patients following chiropractic manipulation. *Journal of Chiropractic Medicine*. 2006 Fall;5(3):88-91. X-7
2314. Zhao J, Bai J, Zhou Y, et al. Posterior tibial nerve stimulation twice a week in patients with interstitial cystitis. *Urology*. 2008 Jun;71(6):1080-4. X-1D
2315. Zink S, Freitag CM and von Gontard A. Behavioral comorbidity differs in subtypes of enuresis and urinary incontinence. *J Urol*. 2008 Jan;179(1):295-8; discussion 298. X-4
2316. Zinner N, Harnett M, Sabounjian L, et al. The overactive bladder-symptom composite score: a composite symptom score of toilet voids, urgency severity and urge urinary incontinence in patients with overactive bladder. *J Urol*. 2005 May;173(5):1639-43. X-1I
2317. Zobrist RH, Quan D, Thomas HM, et al. Pharmacokinetics and metabolism of transdermal oxybutynin: in vitro and in vivo performance of a novel delivery system. *Pharm Res*. 2003 Jan;20(1):103-9. X-1I
2318. Zollner-Nielsen M and Samuelsson SM. Maximal electrical stimulation of patients with frequency, urgency and urge incontinence. Report of 38 cases. *Acta Obstet Gynecol Scand*. 1992 Dec;71(8):629-31. X-7
2319. Zorn BH, Montgomery H, Pieper K, et al. Urinary incontinence and depression. *J Urol*. 1999 Jul;162(1):82-4. X-1B
2320. Zoubek J, Bloom DA and Sedman AB. Extraordinary urinary frequency. *Pediatrics*. 1990 Jun;85(6):1112-4. X-4
2321. Zoubek J, McGuire EJ, Noll F, et al. The late occurrence of urinary tract damage in patients successfully treated by radiotherapy for cervical carcinoma. *J Urol*. 1989 Jun;141(6):1347-9. X-1I
2322. Zozikov B, Kunchev SI and Varlev C. Application of clenbuterol in the treatment of urinary incontinence. *Int Urol Nephrol*. 2001;33(3):413-6. X-9
2323. Zullo MA, Plotti F, Calcagno M, et al. Vaginal estrogen therapy and overactive bladder symptoms in postmenopausal patients after a tension-free vaginal tape procedure: a randomized clinical trial. *Menopause*. 2005 Jul-Aug;12(4):421-7. X-1B
2324. Zullo MA, Ruggiero A, Plotti F, et al. Anterior colporrhaphy plus inside-out tension-free vaginal tape for associated stress urinary incontinence and cystocele. *J Minim Invasive Gynecol*. 2008 Jul-Aug;15(4):446-51. X-1A
2325. Zupi E, Marconi D, Sbracia M, et al. Directed laparoscopic cryomyolysis for symptomatic leiomyomata: one-year follow up. *J Minim Invasive Gynecol*. 2005 Jul-Aug;12(4):343-6. X-1A, X-7
2326. Zupi E, Piredda A, Marconi D, et al. Directed laparoscopic cryomyolysis: a possible alternative to myomectomy and/or hysterectomy for symptomatic leiomyomas. *Am J Obstet Gynecol*. 2004 Mar;190(3):639-43. X-1A, X-7
2327. Zuraw BL, Sugimoto S, Parsons CL, et al. Activation of urinary kallikrein in patients with interstitial cystitis. *J Urol*. 1994 Sep;152(3):874-8. X-1D

Appendix E. List of Peer Reviewers

We extend our appreciation to the members of the Technical Expert Panel (TEP) who provided advice and input during our research process. We identified technical experts on the topic of OAB in the fields of urology, urogynecology, gynecology, primary care, nursing, and patient advocacy to provide assistance during the project. The TEP was both an additional resource and a sounding board during the project. The TEP included twelve members serving as technical or clinical experts, including an AUA representative. To ensure robust, scientifically relevant work, we called on the TEP to provide reactions to work in progress and advice on substantive issues or possibly overlooked areas of research. TEP members are listed below (* also a peer reviewer):

Roger Dmochowski, M.D.*

Professor
Vanderbilt University, Urologic surgery

Catherine DuBeau, M.D.*

Associate Professor, University of Chicago
Director, Geriatric Continence Center

Heddy Hubbard, Ph.D., M.P.H., R.N., F.A.A.N.*

Director of Guidelines
American Urological Association

Melissa Lavender, M.B.A.

Founder, Executive Director, Women's Health Foundation

Susan Meikle, M.D., M.S.P.H.

NICHD, Project Scientist, Pelvic Floor Disorders Network

Diane K. Newman, R.N.C., M.S.N., C.R.N.P., F.A.A.N.*

Co-Director, Penn Center for Continence and Pelvic Health
University of Pennsylvania

Richard S. Pelman, M.D.*

American Urological Association
Bellevue Urology Associates

Maureen G. Phipps, M.D., M.P.H.

Associate Professor of Obstetrics/Gynecology
Women & Infants Hospital of Rhode Island

Neil Resnick, M.D.

Professor and Chief, University of Pittsburgh Medical Center, Division of Geriatric Medicine
Director, Institute on Aging

Holly Richter, M.D., Ph.D.*

Chief, Women's Pelvic Medicine & Reconstructive Surgery
University of Alabama-Birmingham

Eric S. Rovner, M.D.*

Associate Professor, Medical University of South Carolina Urology Services

J. Christian Winters, M.D.

Chairman and Professor, Department of Urology
LSU Health Sciences Center

Jennifer Wu, M.D., M.P.H.*

Assistant Professor, Division of Urogynecology
Duke University Medical Center ,